

MicroRNA-9 regulates the expression of peroxisome proliferator-activated receptor δ in human monocytes during the inflammatory response

PETRA THULIN^{1*}, TIANLING WEI^{1*}, OLIVERA WERNGREN¹, LOUISA CHEUNG¹, RACHEL M. FISHER¹, DAN GRANDÉR², MARTIN CORCORAN² and EWA EHRENBORG¹

¹Atherosclerosis Research Unit, Department of Medicine, Center for Molecular Medicine, Karolinska Institutet, Karolinska University Hospital L8:03, SE-171 76 Stockholm; ²Department of Oncology-Pathology, Cancer Center Karolinska (CCK), Karolinska Institutet, Karolinska University Hospital R8:03, SE-171 76 Stockholm, Sweden

Received December 12, 2012; Accepted January 30, 2013

DOI: 10.3892/ijmm.2013.1311

Abstract. PPAR δ is involved in the inflammatory response and its expression is induced by cytokines, however, limited knowledge has been produced regarding its regulation. Since recent findings have shown that microRNAs, which are small non-coding RNAs that regulate gene expression, are involved in the immune response, we set out to investigate whether PPAR δ can be regulated by microRNAs expressed in monocytes. Bioinformatic analysis identified a putative miR-9 target site within the 3'-UTR of PPAR δ that was subsequently verified to be functional using reporter constructs. Primary human monocytes stimulated with LPS showed a downregulation of PPAR δ and its target genes after 4 h while the expression of miR-9 was induced. Analysis of pro-inflammatory (M1) and anti-inflammatory (M2) macrophages showed that human PPAR δ mRNA as well as miR-9 expression was higher in M1 compared to M2 macrophages. Furthermore, treatment with the PPAR δ agonist, GW501516, induced the expression of PPAR δ target genes in the pro-inflammatory M1 macrophages while no change was observed in the anti-inflammatory M2 macrophages. Taken together, these data suggest that PPAR δ is regulated by miR-9 in monocytes and that activation of PPAR δ may be of importance in M1 pro-inflammatory but not in M2 anti-inflammatory macrophages in humans.

Introduction

Peroxisome proliferator-activated receptor δ (PPAR δ) is a ubiquitously expressed member of the ligand-activated nuclear receptor superfamily. PPAR δ has not only been shown to regulate genes involved in lipid and glucose homeostasis but also seems to play an important role in inflammation and innate immunity (1,2). Upon ligand binding, PPAR δ is activated and releases the anti-inflammatory transcriptional suppressor protein B cell lymphoma-6 (BCL-6), which leads to suppression of inflammatory gene expression (3,4). Furthermore, activation of PPAR δ has been shown to inhibit tumor necrosis factor α (TNF α)-induced nuclear factor of κ light polypeptide gene enhancer in B-cells 1 (NF- κ B) activation (5). PPAR δ has lately attracted attention for its involvement in different macrophage phenotypes in mice where it has been reported to induce a switch from pro-inflammatory M1 macrophages to the anti-inflammatory M2 phenotype in metabolically important organs such as liver and adipose tissue (6,7). The pro-inflammatory M1 macrophage phenotype is driven by Th1 cytokines e.g., interferon γ (IFN γ) and TNF α while the anti-inflammatory M2 macrophage phenotype is driven by Th2 cytokines e.g., interleukin 4 (IL-4) and IL-13. During atherosclerosis monocytes infiltrate the vessel wall, where these two subpopulations of macrophages can be detected (8).

Bone marrow-specific deletion of *Ppar δ* in mice renders adipose tissue and liver macrophages incapable of transition to the M2 phenotype, which in turn causes inflammation and metabolic derangement in adipocytes as well as hepatic dysfunction and systemic insulin resistance, respectively (6,7). These data demonstrate the importance of PPAR δ for the inflammatory response and the dependence of PPAR δ for maintaining the alternative macrophage phenotype in mice. In contrast, a recent human study investigating PPAR δ activation during monocyte differentiation did not detect any increased expression of PPAR δ in IL-4 induced M2 macrophages compared to untreated macrophages (9). While research on the function of PPAR δ has mainly focussed on its activation, considerably less is known about the regulation of expression of human PPAR δ . In the last few years

Correspondence to: Dr Ewa Ehrenborg, Atherosclerosis Research Unit, Department of Medicine, Center for Molecular Medicine, Karolinska Institutet, Karolinska University Hospital L8:03, SE-171 76 Stockholm, Sweden
E-mail: ewa.ehrenborg@ki.se

*Contributed equally

Key words: peroxisome proliferator-activated receptor δ , microRNA-9, gene regulation, monocytes, macrophages, inflammatory response

microRNAs (miRNAs) have emerged as an important class of fine-tuning regulators of gene and protein expression. These transcripts are endogenous non-coding single-stranded RNAs ~22 nucleotides in length that bind to the 3'-untranslated region (3'-UTR) of their target mRNA and suppress expression either by inducing mRNA degradation or inhibiting protein translation (10). It is known that ~50% of all mammal protein-coding genes are directly regulated by miRNAs (11) and there are over two thousand known human miRNAs described today. MiRNAs are implicated in processes such as development and differentiation but many of them have also been reported to be aberrantly expressed in different forms of cancers. Computational studies have shown that recurrent networks exist consisting of specific transcription factors and specific miRNAs that both appear to regulate one another, thus coupling transcriptional and post-transcriptional regulation in order to create flexible expression (12). Identified miRNAs that have been shown to play key roles in monocytes and/or macrophages during inflammation include microRNA-9 (miR-9) and microRNA-155 (miR-155). MiR-9 is involved in the immune response by fine tuning the expression of a key member of the NF- κ B family in monocytes and polymorphonuclear neutrophils (4) while the expression of miR-155 is increased during inflammation and has been implicated in macrophage polarization, where miR-155 modulates the switch between pro-inflammatory M1 and anti-inflammatory M2 phenotypes. MiR-155 directly inhibits the expression of IL-13 receptor α 1 and thereby downregulates the anti-inflammatory IL-13 pathway in macrophages (13).

Of the human PPARs, both PPAR α and PPAR γ have been shown to be regulated by miRNAs (14–19), but until now no data are available regarding the regulation of PPAR δ by miRNAs. Considering the important role that PPAR δ plays in immunity, we explored the regulation of human PPAR δ expression by miRNAs expressed in primary human monocytes during the inflammatory response and during macrophage polarization to M1 and M2 phenotypes, respectively.

Materials and methods

Bioinformatic sequence analysis. Prediction of miRNA binding to the PPAR δ 3'-UTR was performed by computer-aided algorithms obtained from TargetScan (<http://www.targetscan.org>), PicTar (<http://pictar.mdc-berlin.de>), miRanda (<http://www.microrna.org>) and miRWalk (<http://www.umm.uniheidelberg.de/apps/zmf/mirwalk>). Approximately 3 kb of the miR-9 promoter was analyzed for putative PPRES using the MatInspector software (<http://www.genomatix.de>).

Reporter gene constructs. Firefly luciferase reporter plasmids containing the 3'-UTR of the PPAR δ gene and empty luciferase vector were obtained from Promega (Madison, WI, USA). The 36 nucleotides of PPAR δ 3'-UTR containing the miR-9 binding site were cloned into psiCHECK-2 (Promega) using the restriction sites *Xho*I/*Not*I. The following oligonucleotides were utilized (the mutated sequences are shown in bold): miR-9WT-forw: 5'-tcgaTGTCTTCAGAGCAAAGACTTGAGCCATCCAAAGAA-3'; miR-9WT-rev: 5'-ggccTTCTTTGGATGGCTCAAGTCTTTTGCTCTGAAGACA-3'; PPAR δ -mir9Mut-forw: 5'-tcgaTGTCTTCAGAGCAAAGACTTGA

GCCATTAGCGTAA-3'; PPAR δ -mir9Mut-rev: 5'-ggccTTACGCTAATGGCTCAAGTCTTTTGCTCTGAAGACA-3'. The plasmids were sequenced and purified using the Endofree Plasmid Maxi kit (Qiagen, Düsseldorf, Germany).

Cell culture and transfection. HEK293 cells were cultured in DMEM Glutamax (1 g/ml glucose, Invitrogen) containing 10% FBS, penicillin (100 U/ml) and streptomycin (100 μ g/ml) at 37°C in 5% CO₂. For luciferase assays, HEK293 cells were plated in 24-well tissue-culture dishes 24 h prior to transfection at a density of 80,000 cells per well. Cells were transfected with the luciferase reporters, 50 ng per well (Promega), together with pre-miR-9, 10 nM per well, or miRNA mimics negative control no. 1 (pre-miR-CON, Ambion, Foster City, CA, USA), 10 nM per well; 50 nM LNA-based anti-miR-9 (Exiqon, Vedbaek, Denmark) or 50 nM universal LNA-based negative control (anti-miR-CON) (Exiqon). All transfections were carried out in triplicates with Lipofectamine 2000 (Invitrogen, Carlsbad, CA, USA). Cells were lysed 24 h post-transfection using passive lysis buffer (Promega). Luciferase activity was determined using Dual Luciferase[®] Reporter assay system (Promega) following the manufacturer's instructions. Relative luciferase activity was determined by the ratio of renilla luciferase signal intensity to that of firefly luciferase for normalization.

Human monocytes were isolated from buffy coats as previously described (20). In brief, human peripheral blood mononuclear cells (PBMCs) were isolated from buffy coats by endotoxin-free Ficoll density gradient centrifugation. Monocytes were then separated from lymphocytes by high-density hyper-osmotic Percoll density gradient centrifugation and separated from platelets and dead cells on a low-density iso-osmotic Percoll density gradient. Monocytes were cultured in RPMI-1640 medium (Invitrogen) supplemented with penicillin-streptomycin, L-glutamine (2.05 mM) and 10% human AB-serum (Invitrogen). Monocytes were transfected with miRNA oligonucleotides as described above.

Monocytes were transfected with 50 nM anti-miR-9 or 50 nM anti-miR-CON (Exiqon) using Lipofectamine RNAiMAX (Invitrogen), following the manufacturer's instructions. At 24 h after transfection, monocytes were stimulated with LPS (100 ng/ml) for 4 h.

Pro-inflammatory M1 and anti-inflammatory M2 macrophages were obtained by stimulating freshly isolated monocytes with recombinant human IFN- γ (20 ng/ml) and IL-4 (15 ng/ml) for 7 days, respectively. The human embryonic kidney cell line HEK293 was purchased from ATCC (Rockville, MD, USA). The cells were cultured in Dulbecco's modified Eagle's medium (1 g/ml glucose, Invitrogen) supplemented with 10% newborn calf serum, 100 U/ml penicillin and 100 μ g/ml streptomycin (Invitrogen). All the cells were cultured at 37°C in 5% CO₂. All cytokines were purchased from Peprotech (Rocky Hill, NJ, USA). GW501516 was synthesized by Synthelec AB, Sweden as described (21).

RNA extraction, reverse transcription and quantitative real-time PCR. Total RNA was prepared using miRNeasy kit (Qiagen) according to the manufacturer's instructions. The RNA concentration was determined by spectrophotometry. Total RNA (0.5 μ g) was reverse transcribed (RT) into cDNA

in a 20- μ l reaction by a poly-dT primer using Superscript III™ (Invitrogen).

Quantification of miRNAs by TaqMan® real-time PCR was carried out as described by the manufacturer (Applied Biosystems, Foster City, CA, USA). Briefly, 10 ng of template RNA was reverse transcribed using the TaqMan MicroRNA Reverse Transcription kit and miRNA-specific stem-loop primers (Applied Biosystems). RT product (1.5 μ l) was introduced into 20- μ l PCR reactions which were incubated in 96-well plates on the ABI PRISM® 7900HT Sequence Detection System (Applied Biosystems) at 95°C for 10 min, followed by 40 cycles of 95°C for 15 sec and 60°C for 1 min. Target gene expression was normalized between different samples based on the values of U48 RNA expression.

The cDNA was amplified by real-time PCR as described (22). For the quantification, 15 ng of cDNA were amplified per reaction in the presence of TaqMan universal master mix (Applied Biosystems) and TaqMan Gene Expression Assays for PPAR δ (Hs04187066_g1), PLIN2 (Hs00605340_m1), CPT1A (Hs00912676_m1), ANGPTL4 (Hs01101127_m1), TNF α (Hs01113624_g1), STAT1 (Hs01013996_m1), MRC1 (Hs00267207_m1) and STAT6 (Hs00598625_m1), all purchased from Applied Biosystems. Gene-specific PCR products were measured by means of the ABI PRISM® 7900HT Sequence Detection System (Applied Biosystems). Target gene expression was normalized based on the values of the expression of cyclophilin A, PPIA (Hs04194521_s1) and 18s (Hs99999901_s1) obtained from Applied Biosystems.

Statistical analysis. Wilcoxon rank-sum and signed-rank tests were employed to determine statistical differences between means of quantitative real-time PCR and luciferase assay data. $P < 0.05$ was considered statistically significant.

Results

PPAR δ is the direct target of miR-9. To investigate whether the expression of PPAR δ in monocytes could be regulated by miRNAs during the inflammatory response and human macrophage polarization, bioinformatic analyses were performed. We screened the human PPAR δ 3'-UTR for putative target sites of candidate miRNAs expressed in monocytes and/or macrophages (4,13,23). Four different computational prediction programs; TargetScan, PicTar, miRanda and miRWalk, were used. One putative miRNA target site for miR-9 was identified by all four programs. The identified target site for miR-9 in PPAR δ 3'-UTR is highly conserved in many mammals including human, mouse, rat and dog (Fig. 1A).

In order to verify whether PPAR δ is a direct target of miR-9, we performed 3'-UTR luciferase activity assays. Accordingly, 36 nucleotides encompassing the putative miR-9 binding site in the 3'-UTR of the PPAR δ gene were cloned into a reporter plasmid containing the renilla luciferase gene. This reporter construct was transiently co-transfected with miR-9 mimic (pre-miR-9) or the specific inhibitor of miR-9 (anti-miR-9) as well as their respective control oligonucleotides into HEK293 cells and relative luciferase activities were determined 24 h after transfection (Fig. 1B). Overexpression of miR-9 reduced the luciferase activity down to 64% ($P < 0.001$), whereas inhibition of endogenous miR-9 increased the luciferase activity




Figure 1. PPAR δ is the direct target of miR-9. (A) DNA sequence alignment of the evolutionary conserved miR-9 binding site in the PPAR δ 3'-UTR of mammals. The miR-9 seed match region is highlighted in red capital letters. (B) The relative expression obtained from reporter constructs containing the wild-type (WT) PPAR δ 3'-UTR that were transfected into HEK293 cells together with pre-miR-9 or miR-9 inhibitor oligonucleotide (anti-miR-9). The luciferase activity was measured 24 h after transfection. (C) The nucleotides changed in the PPAR δ 3'-UTR mutant construct are indicated in red lower case. (D) The relative expression obtained when HEK293 cells were transfected with reporter constructs containing wild-type (WT) or mutant (MUT) PPAR δ 3'-UTR or empty renilla expression vector (vector) together with miR-9 precursor RNA (pre-miR-9) or miRNA precursor control (pre-miR-ctrl). The luciferase activity was measured 24 h after transfection. The experiments were repeated 3 times. A representative result is shown. Each bar represents the mean $\pm$ SD for 3 wells. *** $P < 0.001$.

up to 130% ($P < 0.001$) compared with scrambled pre-miR or anti-miR control nucleotides, respectively (Fig. 1B), indicating that miR-9 directly targets PPAR δ .

To examine whether the effects on transcription are mediated by the predicted miR-9 target site in the 3'-UTR of PPAR δ , we changed 6 nucleotides within the miR-9 seed-matching sequence of the 3'-UTR of PPAR δ to generate a construct named MUT (Fig. 1C). Mutation of the miR-9 seed-matching sequence led to a complete restoration of luciferase activity and reversed the inhibitory effect of miR-9 in the 3'-UTR of




Figure 2. PPAR δ mRNA expression is suppressed by miR-9 in monocytes after LPS treatment. Human primary monocytes were seeded in 6-well plates (2×10^6 cells/well) and treated with LPS (100 ng/ml) for 2, 4, 8 and 24 h. The expression of (A) miR-9, (B) PPAR δ , (C) PLIN2 and (D) CPT1A was then measured by real-time PCR. The experiment was repeated 3 times. A representative result is shown. miR-9 expression was normalized to U48 expression while PPIA was used for normalization of PPAR δ , PLIN2 and CPT1A expression. Each bar represents the mean $\pm$ SD for 3 wells. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

PPAR δ (Fig. 1D), which shows that the effects are mediated through the identified miR-9 target site. Taken together, these results demonstrate that miR-9 directly regulates PPAR δ expression by binding to the target site in the 3'-UTR of PPAR δ mRNA.

PPAR δ mRNA expression is regulated by miR-9 in monocytes after LPS treatment. Since miR-9 has been shown to play an important role in the inflammatory response in monocytes, where it serves as a feedback controller of inflammation by suppressing NF κ B1 signaling (4), the relevance of miR-9 regulation in relation to PPAR δ expression was investigated in monocytes. To this end, primary monocytes were isolated from PBMC obtained from healthy donors followed by stimulation with LPS for different time periods. Expression of miR-9 and PPAR δ was evaluated by qRT-PCR. To estimate the corresponding effects due to changes in human PPAR δ protein expression, we analysed PPAR δ target genes, such as perilipin-2 (PLIN2), carnitine palmitoyltransferase 1 α (CPT1A) and angiopoietin-related protein 4 (ANGPTL4), none of which had putative miR-9 target sites as evaluated by bioinformatic analyses (24).

In line with the findings by Bazzoni *et al* (4), miR-9 levels increased rapidly after 2 h and remained increased until 24 h after treatment with LPS (Fig. 2A). The PPAR δ mRNA level, on the other hand, was unchanged until 4 h after LPS treatment when it was suppressed by 60% compared to control cells treated with vehicle only. The suppression was abolished after 8 h and increased at 24 h of LPS stimulation (Fig. 2B). Analysis of the PPAR δ target genes PLIN2 and CPT1A showed 80 and 76%, respectively, reduced expression levels compared to unstimulated control cells after 4 h of LPS treatment and this reduction remained until 8 h, followed by an increase at 24 h (Fig. 2C and D). Analysis of another PPAR δ target gene, ANGPTL4, showed the same trend as PLIN2 and CPT1A (data not shown). Thus, PLIN2 was chosen as the PPAR δ target gene in the following analyses due to its relatively high abundance. Taken together, the results suggest a model in which upregulation of miR-9 upon LPS stimulation in monocytes results in a decrease in PPAR δ expression thereby suppressing its corresponding target genes at an early time-point (4–8 h), indicating that PPAR δ is regulated by miR-9 in monocytes after LPS treatment.

Suppression of miR-9 upregulates the mRNA expression of PPAR δ and its target gene PLIN2 in human primary monocytes. The influence of miR-9 on PPAR δ expression in monocytes during the inflammatory response is unknown. Thus, we further evaluated regulation by miR-9 of PPAR δ expression in human primary monocytes stimulated with the pro-inflammatory agent LPS. Human primary monocytes were transfected with the specific anti-miR-9 or its control oligonucleotides for 24 h, followed by LPS stimulation for 4 h. In order to measure the effect of miR-9 the mRNA expression of PPAR δ and its target gene PLIN2 was quantified by qRT-PCR. As shown in Fig. 3, specific inhibition of miR-9 significantly increased PPAR δ and PLIN2 mRNA expression 40 and 70%, respectively, compared to scrambled control oligonucleotides (anti-miR-ctrl). This result suggests that miR-9 downregulates the expression of PPAR δ and its target gene PLIN2 and further confirms the regulatory link between PPAR δ and miR-9.

Both PPAR δ mRNA and miR-9 expression are higher in M1 than in M2 macrophages. PPAR δ has been suggested to play a role in the switch from pro-inflammatory M1 phenotype to




Figure 3. Suppression of miR-9 upregulates the mRNA expression of PPAR δ and its target gene PLIN2. Human primary monocytes were transfected with anti-miR-9 or miRNA inhibitor control (anti-miR-ctrl) for 24 h, followed by LPS treatment for 4 h. The expression of PPAR δ and its target gene PLIN2 was then examined by real-time PCR. The experiment was repeated 3 times. A representative result is shown. Each bar represents the mean $\pm$ SD for 3 wells. *** P <0.001.

anti-inflammatory M2 phenotype in macrophages. To study whether miR-9 expression was involved in the polarization of M1 and M2 macrophage phenotypes through modulation of PPAR δ expression, human primary monocytes were differentiated into M1 and M2 macrophages, respectively. Hence, human primary monocytes were cultured in the presence of either IFN γ to promote a shift to pro-inflammatory M1 macrophages, or IL-4 to polarize the macrophages into the anti-inflammatory M2 phenotype. As expected the M2 marker, mannose receptor C type 1 (MRC1), showed higher expression in M2 compared with M1 or untreated macrophages using FACS analysis (data not shown). TaqMan analysis revealed that the expression of TNF α and signal transducer and activator of transcription 1 (STAT1), which are markers of M1, were higher in M1 compared to M2 macrophages while the mRNA expression of MRC1 and STAT6 were higher in M2 compared to M1 macrophages (Fig. 4). In addition, PPAR γ mRNA was also analysed since its expression has been shown to correlate with markers for M2 macrophages. As shown in Fig. 4B, PPAR γ mRNA levels were significantly increased in M2 compared with M1 macrophages. These results demonstrate that two distinct macrophage phenotypes, M1 and M2, were obtained.

Analyses of the expression of PPAR δ mRNA and miR-9 showed that their levels are significantly increased in M1 compared with M2 macrophages (Fig. 5), which suggest that PPAR δ and miR-9 might be of importance in modulating the pro-inflammatory M1 human macrophage phenotype.

Expression of miR-9 in relation to PPAR δ agonist treatment. The polarization towards the M1 phenotype is important for macrophages exposed to pro-inflammatory stress. To further investigate the interplay between PPAR δ and miR-9 in monocytes and macrophages during the inflammatory response, we examined whether miR-9 can be regulated by PPAR δ . Since pri-mir-9-1 is the only primary miR-9 transcript induced by LPS (4), we set out to analyse whether a PPAR response element (PPRE) exists in the pri-mir-9-1 promoter region. *In silico* analysis of the promoter region of pri-mir-9-1 revealed five putative PPREs within 2.5-kb upstream of the transcriptional start site (data not shown).




Figure 4. Expression of markers for M1 and M2 macrophages. Human primary monocytes were seeded in 6-well plates (2×10^6 cells/well). M1 and alternative M2 macrophages were obtained by stimulating the monocytes with recombinant human IFN- γ (20 ng/ml) and IL-4 (15 ng/ml) for 7 days, respectively. The cells were harvested and proceeded to quantitative real-time PCR for measurement of (A) M1 (STAT1 and TNF α) and (B) M2 (STAT6, PPAR γ and MRC1) markers. The experiment was repeated 4 times. A representative result is shown. STAT1, TNF α , STAT6, PPAR γ and MRC1 expression were normalized to 18s expression. Each bar represents the mean $\pm$ SD for 3 wells. * P <0.05, ** P <0.01, *** P <0.001.




Figure 5. Both the expression of PPAR δ mRNA and miR-9 are higher in M1 than in M2 macrophages. The expression of miR-9 and PPAR δ mRNA was measured in macrophages, after 7 days of treatment with IFN- γ or IL-4 to obtain the M1 and M2 phenotypes, respectively. The experiment was repeated 5 times using monocytes from 6 different donors. Mir-9 expression was normalized to U48 expression and PPAR δ expression was normalized to 18s expression. Means $\pm$ SD of 5-6 independent experiments are shown. * P <0.05.

Next, we explored whether PPAR δ activation could induce miR-9 expression in human monocytes and/or macrophages using the specific PPAR δ agonist, GW501516. Human




Figure 6. The miR-9 expression is not regulated by PPAR δ in monocytes. (A) Human primary monocytes were seeded in 6-well plates (2×10^6 cells/well) and treated with 100 nM PPAR δ agonist (GW501516) for 2 h followed by isolation of RNA from the cells. To verify the potency of GW501516, mRNA expression of the PPAR δ target gene, PLIN2, was analysed by real-time PCR. (B) The expression of miR-9 is shown from the same cells. (C) The expression of miR-9 in M1 and M2 macrophages, respectively, is shown after 4 h of treatment with GW501516. The experiment was repeated in monocytes from two individuals and in macrophages from two individuals. Representative results from these experiments are shown. White bars show results from no GW501516 treated cells and the results were set to one. PLIN2 expression was normalized to PPIA expression and miR-9 expression was normalized to U48 expression. Each bar represents the mean $\pm$ SD for 3 wells. * $P < 0.05$.

monocytes were isolated from PBMC and stimulated with GW501516 for 2, 4, 8 and 24 h. Since PLIN2 gene expression is regulated by PPAR δ , PLIN2 mRNA levels were measured at the different time-points to serve as a positive control and to ensure that GW501516 had activated PPAR δ . As expected PLIN2 mRNA expression was upregulated at the different time-points analysed after adding GW501516 (Fig. 6A). However, the miR-9 expression was not induced at any of the different time-points of treatment with GW501516 (Fig. 6B), nor was there any difference of miR-9 expression in M1 or M2 macrophages after GW501516 treatment (Fig. 6C).




Figure 7. GW501516 induces PLIN2 mRNA expression in M1 but not in M2 macrophages. Pro-inflammatory M1 or anti-inflammatory M2 macrophages were obtained after 7 days of stimulation with IFN- γ or IL-4, respectively, followed by treatment with the PPAR δ agonist, GW501516, for 4 h. The relative mRNA expression of the PPAR δ target genes, (A) PLIN2 and (B) CPT1A, was analyzed in the two phenotypes of macrophages. A representative result is shown. PLIN2 and CPT1 expression were normalized to 18s expression. Each bar represents the mean $\pm$ SD for 3 wells. * $P < 0.05$.

PPAR δ agonist treatment results in upregulation of PPAR δ target genes in M1 but not in M2 macrophages. In order to evaluate the influence of PPAR δ on the expression of its target genes, M1 and M2 macrophages were subjected to GW501516 treatment for 4 h. Human primary monocytes that were differentiated into macrophages of the pro-inflammatory M1 phenotype responded to the treatment with GW501516 by upregulation of PLIN2 and CPT1A mRNA while macrophages differentiated into the anti-inflammatory M2 phenotype did not (Fig. 7). Another PPAR δ target gene that was analysed, ANGPTL4, also showed induction of mRNA expression upon GW501516 treatment only in M1 but not in M2 macrophages (data not shown).

Discussion

It has been suggested that the primary regulation of the human PPAR δ gene is not at the transcriptional level. Instead, PPAR δ is known to be regulated at the post-translational level by the presence of ligands along with specific cofactors that are of major importance for PPAR δ activation and function. In this study, we show that PPAR δ is also regulated at the post-transcriptional level by miR-9. Upregulation of miR-9 results in the direct repression of PPAR δ , the mRNA levels of which are found to be higher in pro-inflammatory M1 than in anti-inflammatory M2 macrophages.

MiR-9 has been shown to be of importance during the immune response (4), post-traumatic stress (25), neuronal differentiation (26), different forms of cancers (27,28) and

exocytosis of insulin from pancreatic islets (29). Here, we show that miR-9 is upregulated in human monocytes after LPS treatment while PPAR δ and its target genes PLIN2, CPT1A and ANGPTL4 are downregulated 4-8 h after LPS-treatment compared with untreated cells. The suppression, intriguingly, was abolished after 8-24 h of LPS stimulation despite the presence of continued high levels of miR-9. One possible explanation might be that the effect and activity of miR-9 are influenced due to interactions with RNA-binding proteins (30). Another reason might be that LPS triggers other factors that in turn could affect the expression of PPAR δ . Indeed, Tan *et al* (31) have shown that pro-inflammatory cytokines, such as TNF α , can both increase PPAR δ expression via the stress kinase signaling pathway and trigger the production of ligands for this receptor, which is in agreement with our findings that PPAR δ is upregulated after 24 h of LPS treatment. The inhibition of PPAR δ by miR-9 at the early time-points might be a mechanism to delay the effect of PPAR δ action early in inflammation to prevent PPAR δ from suppression of NF- κ B since NF- κ B and PPAR δ have been shown to be able to cross-talk and inhibit the function of each other (32). The inhibitory effect by miR-9 on PPAR δ expression was confirmed by transfection of anti-miR-9 that sequesters mature miR-9 thus inhibiting its biologic function into monocytes, which resulted in induction of PPAR δ and PLIN2 mRNA levels.

Hence, the miR-9 mediated inhibition of PPAR δ expression in monocytes may constitute a negative feedback loop, modulating the levels of the receptor during inflammation. This finding regarding miR-9 expression in monocytes is in agreement with a previous study, which showed that the expression of miR-9 is dramatically increased after treatment with LPS (4). Bazzoni *et al* showed that the miR-9 targeting of NF- κ B at the mRNA level constitutes a feedback loop of the inflammatory response. Other studies have confirmed the regulatory function of miR-9 on the mRNA of NF- κ B in both ovarian and gastric cancers where miR-9 has been shown to act as a tumor suppressor (33,34). Of note, in the monocytic cell line THP1, miR-9 is abundantly expressed and there is no induction of miR-9 upon LPS stimulation (unpublished data), therefore all the current experiments were carried out in human primary monocytes or macrophages.

Here, the regulation of PPAR δ by miR-9 was examined in relation to the pro-inflammatory M1 and anti-inflammatory M2 macrophage phenotypes. The role of PPAR δ in the pro-inflammatory M1 macrophages has not been investigated before while PPAR δ has been suggested to be of importance in macrophages with the anti-inflammatory M2 phenotype, both in adipose tissue and liver of mice (6,7) although this seems not to be the case in humans (9).

Monocytes were differentiated to macrophages of the M1 and M2 phenotypes by stimulation with IFN γ and IL-4, respectively, for 7 days. To confirm this, several markers that are commonly used to distinguish between the two distinct M1 and M2 macrophage phenotypes were analyzed. In agreement with other reports the expression of STAT1 and TNF α were higher in M1 than M2 macrophages while STAT6, mannose receptor type C and PPAR γ were higher in M2 than in M1 macrophages (35).

In this study, both the expression of PPAR δ and miR-9 were higher in pro-inflammatory M1 than in anti-inflammatory M2

macrophages, indicating the potential involvement of PPAR δ and miR-9 in modulating the M1 macrophage phenotype. As expected TNF α expression was higher in M1 than in M2 macrophages and the PPAR δ ligand, GW501516, did not influence the expression of miR-9 in M1 macrophages. These results indicate that the pro-inflammatory cytokines are responsible for the induction of both miR-9 and PPAR δ expression, which is in agreement with previous studies (4,36,37). In addition, the expression of PPAR δ target genes was induced by GW501516 in macrophages of the M1, but not the M2 phenotype, which probably is due to the fact that PPAR δ is mainly expressed in M1 macrophages while PPAR γ is predominantly expressed in M2 macrophages in humans (6-9).

Our data show the regulation of PPAR δ by miR-9 in monocytes. Whether this regulation might play a role in foam cell formation and initiation of atherosclerosis remains to be elucidated. However, activation of PPAR δ does not seem to promote the polarization into M2 macrophages (unpublished data). In contrast to the studies by Kang *et al* and Odegaard *et al* (6,7), our study of human macrophages showed a higher PPAR δ mRNA expression in M1 macrophages compared to M2 macrophages. This result supports the data from Bouhrel *et al* (9), who showed that PPAR δ expression was lower in M2 macrophages compared to resting (untreated) macrophages. These data highlight differences between human and mouse macrophages and suggest that they may be regulated by different factors.

The importance of miR-9 in monocytes and macrophages as a response to proatherogenic factors is further demonstrated by a recent report showing that miR-9 is significantly upregulated in monocytes and macrophages after exposure to oxLDL (38). One purpose of the inhibition of PPAR δ by miR-9 in monocytes could be to induce a block in monocyte expansion. Whether PPAR δ inhibits or stimulates cell proliferation is still debated (39), but during the inflammatory response cells must focus on repair and disposal of causal pathogens rather than undergoing cell divisions. In this context, PPAR δ has been shown to function as a sensor of apoptotic cells in macrophages by promoting the clearance of apoptotic cells and suppressing autoreactive immune responses (40).

Importantly, a miRNA does not only target one gene, but frequently rather modestly regulates a number of genes belonging to the same network. The long list of putative miR-9 targets identified by computational prediction programs contains genes working in a close network with PPAR δ , for example PPAR α , RXR α , PGC1 α , FOXO and BCL-6. If all these genes are miR-9 regulated, the outcome of miR-9 regulation would be due to effects on the whole network rather than only to PPAR δ . Putative target genes of PPAR δ can also be found on the miR-9 target list, such as ABCA1, ABCD1, GOT1 and PDK4. The effects of targeting these genes would cause similar results as targeting of PPAR δ itself, but resulting in a larger net effect. To draw firm conclusions, however, a functional approach for each gene would be required in order to validate whether these genes are true miR-9 targets, which although relevant, was outside the scope of this study.

In conclusion, we have identified miR-9 as a regulator of PPAR δ expression in monocytes through direct targeting of a specific sequence with the 3'-UTR of PPAR δ . These data show that miR-9 and PPAR δ are involved in central signalling

pathways during the inflammatory response in monocytes. Interestingly, activation of PPAR δ by GW501516 showed that the PPAR δ target genes expressed in the pro-inflammatory M1 macrophages were induced while the expression of the same genes were unaffected in the anti-inflammatory M2 macrophages, which may suggest that PPAR δ is of importance in M1 pro-inflammatory macrophages.

Acknowledgements

This study was supported by the Swedish Research Council, the Swedish Heart-Lung Foundation, the Swedish Diabetes Foundation, the Novo Nordisk Foundation, the Swedish Cancer Foundation, Radiumhemmets Forskningsfonder and the Karolinska Institutet Funds. P.T. was supported by a doctoral research fellowship from the Swedish Heart-Lung Foundation and E.E. was supported by a senior researcher fellowship from the Swedish Heart-Lung Foundation.

References

- Ehrenborg E and Krook A: Regulation of skeletal muscle physiology and metabolism by peroxisome proliferator-activated receptor δ . *Pharmacol Rev* 61: 373-393, 2009.
- Yessoufou A and Wahli W: Multifaceted roles of peroxisome proliferator-activated receptors (PPARs) at the cellular and whole organism levels. *Swiss Med Wkly* 140: w13071, 2010.
- Barish GD, Narkar VA and Evans RM: PPAR δ : a dagger in the heart of the metabolic syndrome. *J Clin Invest* 116: 590-597, 2006.
- Bazzoni F, Rossato M, Fabbri M, *et al*: Induction and regulatory function of miR-9 in human monocytes and neutrophils exposed to proinflammatory signals. *Proc Natl Acad Sci USA* 106: 5282-5287, 2009.
- Barroso E, Eyre E, Palomer X and Vázquez-Carrera M: The peroxisome proliferator-activated receptor β/δ (PPAR β/δ) agonist GW501516 prevents TNF- α -induced NF- κ B activation in human HaCaT cells by reducing p65 acetylation through AMPK and SIRT1. *Biochem Pharmacol* 81: 534-543, 2011.
- Kang K, Reilly SM, Karabacak V, *et al*: Adipocyte-derived Th2 cytokines and myeloid PPAR δ regulate macrophage polarization and insulin sensitivity. *Cell Metab* 7: 485-495, 2008.
- Odegaard JI, Ricardo-Gonzalez RR, Red Eagle A, *et al*: Alternative M2 activation of kupffer cells by PPAR δ ameliorates obesity-induced insulin resistance. *Cell Metab* 7: 496-507, 2008.
- Chinetti-Gbaguidi G, Baron M, Bouhrel MA, *et al*: Human atherosclerotic plaque alternative macrophages display low cholesterol handling but high phagocytosis because of distinct activities of the PPARgamma and LXRalpha pathways. *Circ Res* 108: 985-995, 2011.
- Bouhrel MA, Brozek J, Derudas B, *et al*: Unlike PPARgamma, PPARalpha or PPARdelta activation does not promote human monocyte differentiation toward alternative macrophages. *Biochem Biophys Res Commun* 386: 459-462, 2009.
- Bartel DP: MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116: 281-297, 2004.
- Chekulaeva M and Filipowicz W: Mechanisms of miRNA-mediated post-transcriptional regulation in animal cells. *Curr Opin Cell Biol* 21: 452-460, 2009.
- Dahan O, Gingold H and Pilpel Y: Regulatory mechanisms and networks couple the different phases of gene expression. *Trends Genet* 27: 316-322, 2011.
- Martinez-Nunez RT, Louafi F and Sanchez-Elsner T: The interleukin 13 (IL-13) pathway in human macrophages is modulated by microRNA-155 via direct targeting of interleukin 13 receptor alpha1 (IL13Ralpha1). *J Biol Chem* 286: 1786-1794, 2011.
- Martinelli R, Nardelli C, Pilone V, *et al*: miR-519d overexpression is associated with human obesity. *Obesity* 18: 2170-2176, 2010.
- Zhou J, Wang K-C, Wu W, *et al*: MicroRNA-21 targets peroxisome proliferator-activated receptor-alpha in an autoregulatory loop to modulate flow-induced endothelial inflammation. *Proc Natl Acad Sci USA* 108: 10355-10360, 2011.
- Tong JL, Zhang CP, Nie F, *et al*: MicroRNA 506 regulates expression of PPAR alpha in hydroxycamptothecin-resistant human colon cancer cells. *FEBS Lett* 585: 3560-3568, 2011.
- Karbiener M, Fischer C, Nowitsch S, *et al*: microRNA miR-27b impairs human adipocyte differentiation and targets PPARgamma. *Biochem Biophys Res Commun* 390: 247-251, 2009.
- Kim SY, Kim AY, Lee HW, *et al*: miR-27a is a negative regulator of adipocyte differentiation via suppressing PPARgamma expression. *Biochem Biophys Res Commun* 392: 323-328, 2010.
- Lee EK, Lee MJ, Abdelmohsen K, *et al*: miR-130 suppresses adipogenesis by inhibiting peroxisome proliferator-activated receptor gamma expression. *Mol Cell Biol* 31: 626-638, 2011.
- Repnik U, Knezevic M and Jeras M: Simple and cost-effective isolation of monocytes from buffy coats. *J Immunol Methods* 278: 283-292, 2003.
- Glinghammar B, Skogsberg J, Hamsten A and Ehrenborg E: PPAR δ activation induces COX-2 gene expression and cell proliferation in human hepatocellular carcinoma cells. *Biochem Biophys Res Commun* 308: 361-368, 2003.
- Lundell K, Thulin P, Hamsten A and Ehrenborg E: Alternative splicing of human peroxisome proliferator-activated receptor delta (PPARdelta): effects on translation efficiency and trans-activation ability. *BMC Mol Biol* 8: 70, 2007.
- Contreras J and Rao DS: MicroRNAs in inflammation and immune responses. *Leukemia* 26: 404-413, 2012.
- Wagner K-D and Wagner N: Peroxisome proliferator-activated receptor beta/delta (PPAR δ) acts as regulator of metabolism linked to multiple cellular functions. *Pharmacol Ther* 125: 423-435, 2010.
- Zhang Y, Liao Y, Wang D, *et al*: Altered expression levels of miRNAs in serum as sensitive biomarkers for early diagnosis of traumatic injury. *J Cell Biochem* 112: 2435-2442, 2011.
- Krichevsky AM, Sonntag K-C, Isacson O and Kosik KS: Specific microRNAs modulate embryonic stem cell-derived neurogenesis. *Stem Cells* 24: 857-864, 2006.
- Nie K, Gomez M, Landgraf P, *et al*: MicroRNA-mediated down-regulation of PRDM1/Blimp-1 in Hodgkin/Reed-Sternberg cells: a potential pathogenetic lesion in Hodgkin lymphomas. *Am J Pathol* 173: 242-252, 2008.
- Ma L, Young J, Prabhala H, *et al*: miR-9, a MYC/MYCIN-activated microRNA, regulates E-cadherin and cancer metastasis. *Nat Cell Biol* 12: 247-256, 2010.
- Plaisance V, Abderrahmani A, Perret-Menoud V, Jacquemin P, Lemaigre F and Regazzi R: MicroRNA-9 controls the expression of granuphilin/Slp4 and the secretory response of insulin-producing cells. *J Biol Chem* 281: 26932-26942, 2006.
- Agami R: microRNAs, RNA binding proteins and cancer. *Eur J Clin Invest* 40: 370-374, 2010.
- Tan NS, Michalik L, Noy N, *et al*: Critical roles of PPAR beta/delta in keratinocyte response to inflammation. *Genes Dev* 15: 3263-3277, 2001.
- Delerive P, De Bosscher K, Besnard S, *et al*: Peroxisome proliferator-activated receptor alpha negatively regulates the vascular inflammatory gene response by negative cross-talk with transcription factors NF-kappaB and AP-1. *J Biol Chem* 274: 32048-32054, 1999.
- Guo L-M, Pu Y, Han Z, *et al*: MicroRNA-9 inhibits ovarian cancer cell growth through regulation of NF- κ B1. *FEBS J* 276: 5537-5546, 2009.
- Wan H-Y, Guo L-M, Liu T, Liu M, Li X and Tang H: Regulation of the transcription factor NF-kappaB1 by microRNA-9 in human gastric adenocarcinoma. *Mol Cancer* 9: 16, 2010.
- Wolfs IM, Donners MM and de Winther MP: Differentiation factors and cytokines in the atherosclerotic plaque micro-environment as a trigger for macrophage polarisation. *Thromb Haemost* 106: 763-771, 2011.
- Lee CH, Chawla A, Urbizondo N, *et al*: Transcriptional repression of atherogenic inflammation: modulation by PPARdelta. *Science* 302: 453-457, 2003.
- Kharroubi I, Lee CH, Hekerman P, *et al*: BCL-6: a possible missing link for anti-inflammatory PPAR- δ signalling in pancreatic beta cells. *Diabetologia* 49: 2350-2358, 2006.
- Chen T, Huang Z, Wang L, *et al*: MicroRNA-125a-5p partly regulates the inflammatory response, lipid uptake, and ORP9 expression in oxLDL-stimulated monocyte/macrophages. *Cardiovasc Res* 83: 131-139, 2009.
- Peters JM and Gonzalez FJ: Sorting out the functional role(s) of peroxisome proliferator-activated receptor- β/δ (PPAR β/δ) in cell proliferation and cancer. *Biochim Biophys Acta* 1796: 230-241, 2009.
- Mukundan L, Odegaard JI, Morel CR, *et al*: PPAR-[delta] senses and orchestrates clearance of apoptotic cells to promote tolerance. *Nat Med* 15: 1266-1272, 2009.