

Mitochondrial DNA mutations and essential hypertension (Review)

YU DING^{1,2}, BOHOU XIA³, JINFANG YU⁴, JIANHANG LENG^{1,2} and JINYU HUANG^{2,5}

¹Central Laboratory, Hangzhou First People's Hospital; ²Affiliated Hangzhou Hospital, Nanjing Medical University, Hangzhou, Zhejiang; ³Department of Pharmacy, Hunan University of Chinese Medicine, Changsha, Hunan; ⁴Department of Cardiology, Xiaoshan First People's Hospital; ⁵Department of Cardiology, Hangzhou First People's Hospital, Nanjing Medical University, Hangzhou, Zhejiang, P.R. China

Received May 1, 2013; Accepted July 22, 2013

DOI: 10.3892/ijmm.2013.1459

Abstract. Essential hypertension (EH) is a frequent, chronic, age-related disorder, which remains a major modifiable risk factor for cardiovascular disease despite important advances in our understanding of its pathophysiology. Previous studies have noted a consistent maternal effect on blood pressure (BP). Consequently, mutations in mitochondrial DNA (mtDNA) have become an additional target of investigations on the missing BP heritability. Among these mutations, mt-transfer RNA (tRNA) is a hot mutational spot for pathogenic mutations associated with EH. Mutant mtDNA aggravates mitochondrial dysfunction, pivotally contributing to the clinical phenotype. Moreover, the damaged mitochondria, due to their inability to provide the high-energy requirements for cells, generate reactive oxygen species (ROS) and induce mitochondrial-mediated cell death pathways. Therefore, mitochondrial dysfunction plays a critical role in the pathogenesis of EH. This review summarizes the basic knowledge of mitochondrial genetics and EH-associated mtDNA mutations and further discusses the molecular mechanisms behind these mtDNA mutations in clinical manifestations of EH.

Contents

1. Introduction
2. Mitochondrial genome and mitochondrial genetics
3. Mitochondrial function and dysfunction
4. Oxidative stress and hypertension
5. Role of mtDNA mutations in EH
6. Molecular mechanisms behind mtDNA mutations associated with EH
7. Conclusion

Correspondence to: Dr Jinyu Huang, Department of Cardiology, Hangzhou First People's Hospital, Nanjing Medical University, 261 Huan Sha Road, Hangzhou, Zhejiang 310006, P.R. China
E-mail: hjyu41@sohu.com

Key words: mitochondria, mutation, essential hypertension, dysfunction

1. Introduction

Hypertension is an established risk factor for coronary heart disease, stroke, congestive heart failure and renal dysfunction. Despite significant advances in our understanding of the pathophysiology of hypertension, it remains to be one of the world's greatest public health issues (1). It is estimated that one third of the world's adult population will be hypertensive by 2025 (2). In particular, in industrialized countries, the risk of becoming hypertensive [blood pressure (BP) >140/90 mmHg] during a lifetime exceeds 90% (3). Essential hypertension (EH), or hypertension due to undetermined causes, accounts for >90% of cases of hypertension. It is a heterogeneous disorder, with different patients having different causal factors that lead to high BP.

To date, the etiology of EH is not well understood due to multifactorial causes. It is generally believed that EH is a multifactorial trait involving interactions among genetic, environmental and demographic factors (4). Of these, hereditary factors account for 30 to 50% of BP variability (5). Variations in a variety of genes have shown an association with hypertension in certain studies; however, these associations are often not reproducible in studies on other populations. Improved techniques of genetic analysis, particularly genome-wide linkage analysis, have enabled the search for genes that contribute to the development of primary hypertension in the population (6,7). However, the majority of the reported genetic variants were identified in studies of the nuclear genome (8-10); only limited insights have been gained from the investigation of the mitochondrial genome.

This review provides a detailed introduction of the mitochondrial genome, summarizes the results of studies on the role of mitochondrial DNA (mtDNA) mutations associated with EH published thus far, and highlights some of the general conclusions that have become apparent.

2. Mitochondrial genome and mitochondrial genetics

Mitochondria are bacterium-sized organelles found in all nucleated cells (11). Uniquely, they contain their own genome (mtDNA) and it is widely accepted that mitochondria originate from aerobic bacteria engulfed by an anaerobic eukaryotic cell.




Figure 1. Genetic map of the human mitochondrial genome. The outer circle represents the heavy (H) strand and the inner circle the light (L) strand.

Mammalian mtDNA encodes 13 proteins that are subunits of the oxidative phosphorylation (OXPHOS) system and 22 transfer RNAs (tRNAs) and 2 ribosomal RNAs (rRNAs) (Fig. 1). The mitochondrial OXPHOS complexes are assembled from genes distributed between mtDNA and nuclear DNA (nDNA) (12). Unlike nDNA, mtDNAs are maternally inherited and are present in multiple copies/cell. The numbers vary according to the bioenergetic needs of each unique tissue and can range over three orders of magnitude, depending on the cell type (13).

Each mammalian cell contains hundreds of mitochondria and thousands of mtDNAs. Since mtDNA is in the proximity of reactive oxygen species (ROS) generation sites and mitochondria have less sophisticated DNA protection and repair systems, mtDNA is hence vulnerable to a high mutation rate (14). The most prominent is polyplasmcy. Polyplasmcy is the basis of heteroplasmcy (alternations of mtDNA may be present in some of the mtDNA molecules) and homoplasmcy (mutations in all of the molecules). Neutral polymorphisms are usually homoplasmcy, whereas pathogenic mutations are usually heteroplasmcy in nature, and EH-associated mtDNA mutations are commonly homoplasmcy or almost homoplasmcy.

3. Mitochondrial function and dysfunction

Mitochondria exert both vital and lethal functions in physiological and pathological conditions. On the one hand, they provide the majority of cellular energy in the form of adenosine-5'-triphosphate (ATP) through OXPHOS (15). Additionally, they are involved in a variety of processes, including regulation of the cell cycle, cell signaling, apoptosis and calcium (Ca^{2+}) buffering (12,16,17).

Oxygen radicals, such as ROS, are also generated during OXPHOS as a toxic byproduct in the mitochondria, which may damage the mitochondrial and cellular DNA, protein, lipids and other molecules, leading to oxidative stress and

mitochondrial dysfunction (18,19). An impairment of normal mitochondrial function leads to an excessive production of ROS and a general decrease in ATP levels. Moreover, there is a concomitant loss of mitochondrial transmembrane potential (20). Excessive ROS and Ca^{2+} production lead to mitochondrial outer membrane permeabilization and to the release into the cytosol of cytotoxic proteins normally confined within the mitochondrial intermembrane space. As a result, this process induces apoptosis or necrosis (21).

4. Oxidative stress and hypertension

In animal models, oxidative stress has been observed in spontaneously hypertensive rats (22), renovascular hypertension (23) and salt-sensitive hypertension (24). Although its pathogenesis is complex and multifactorial, human hypertension is considered as a state of increased oxidative stress (25). An excessive endothelial production of ROS and nitric oxide synthase (NOS) may lead to impaired endothelium-dependent vasorelaxation in human internal mammary arteries and the saphenous vein (26), contributing to vascular pathophysiology by promoting increased vascular tone, cell growth, as well as the activation of matrix metalloproteinases and the deposition of extracellular matrix proteins, which are processes associated with the vascular phenotype of hypertension (27).

5. Role of mtDNA mutations in EH

12S rRNA A1555G mutation. The A1555G mutation in the 12S rRNA gene has been associated with aminoglycoside-induced and non-syndromic hearing loss in various ethnic populations (28-31). Chen *et al* (32) described two Han Chinese families with hearing loss and hypertension carrying the homoplasmic A1555G mutation. An A to G transition at this position in the 12S rRNA gene has been predicted to encode an aminoglycoside binding based on sequence similarity to *Escherichia coli* (33) and alters mitochondrial ribosomal function and translation (34-36), which in turn causes OXPHOS defects that are thought to contribute to the clinical pathology of hypertension. Insufficient metabolism caused by mitochondrial dysfunction may lead to the elevation of systolic BP and may be involved in the development of hypertension (37).

ND1 T3308C mutation. The homoplasmic ND1 T3308C mutation is a known disease-associated mutation. This mutation has been suggested to contribute to the higher penetrance of hearing loss in a large African family than Japanese and French pedigrees carrying the tRNA^{Ser(UCN)} T7511C mutation (38,39). Liu *et al* (40) reported a Han Chinese family carrying the ND1 T3308C mutation with EH. The T to C transition at position 3308 causes translation-initiating methionine with a threonine in ND1. Thus, the ND1 mRNA is shortened by two amino acids (41). Moreover, the T3308C mutation is also located in two nucleotides adjacent to the 3' end of the tRNA^{Leu(UUR)} gene; the T3308C mutation also affects the processing of the H-strand polycistronic RNA precursors (42).

ND5 T12338C mutation. The well-known T12338C mutation in the ND5 gene, combined with tRNA^{Leu(CUN)} A12330G mutation, has been found in a three generation Han Chinese

family with high penetrance of EH (43). Moreover, this mutation was also shown to be present in a Chinese family with hypertrophic cardiomyopathy (44). In fact, the *ND5* T12338C mutation, which is similar to the *ND1* T3308C mutation, causes a replacement of the first amino acid, translation-initiating methionine with a threonine in the *ND5* polypeptide, and decreases *ND5* mRNA levels and alters the processing of RNA precursors. In addition, this mutation is located in two nucleotides adjacent to the 3' end of the tRNA^{Leu(CUN)}, and it is anticipated that the T12338C mutation will lead to a reduction in tRNA^{Leu(CUN)} levels, whereas the A12330G mutation disrupts the highly conserved base pairing (6T-67A) in the acceptor arm of tRNA^{Leu(CUN)} (45). Therefore, the combination of T12338C and A12330G mutations may have contributed to the high penetrance of EH in this Chinese family (43).

ND6 T14484C mutation. The T14484C mutation in the *ND6* gene is one of the primary mutations associated with Leber's hereditary optic neuropathy (LHON) (46). In a recent study, a large Han Chinese family with maternally transmitted EH but not presenting any LHON phenotype, was found to be associated with the T14484C mutation (47). Analysis of the complete mtDNA sequence of the proband showed the absence of additional pathogenic mutations, apart from the T14484C mutation. Moreover, analysis of the mitochondrial function of lymphoblastoid cell lines established from the family members showed that mitochondrial respiration rate and membrane potential were significantly reduced when compared with the control cell lines. In addition, there was an increase in the levels of ROS and mitochondrial mass in the mutant cell lines. These data suggest that *ND6* T14484C also plays an important role in the pathogenesis of EH and is also a pathogenic mutation associated with EH (47).

CyB G15059A mutation. The heteroplasmic G15059A mutation in *CyB* gene was first described in a patient with mitochondrial myopathy (48). This substitution results in the replacement of a glycine at amino acid position 190 of *CyB* with a stop codon leading to a truncated protein that misses 244 amino acids at the C-terminus of *CyB* (49). Nikitin *et al* (50) examined the role of the *CyB* G15059A mutation in patients with type 2 diabetes (T2D) with EH. In addition, the G15059A heteroplasmy level exceeding 39% was associated with an increased risk of EH, indicating a direct pathogenic role for this mutation in EH (50).

50-bp deletion. The 50-bp deletion (m.298_347del150) in the mtDNA control region removes the conserved sequence block II (CSBII) and the replication primer location (51-53). This deletion, co-occurring with two novel mutations (*ND1* C3519T and *ND5* G13204A), accounted for the complex clinical traits, including hypertension, T2D and coronary artery disease (CAD) in an Indian family (54). Of note, two short homologous direct repeats of CCAAACCCC flanked the 50-bp deletion. The CSBII directs transcription termination and primer formation in mtDNA replication (55). Thus, it can be predicted that the 50-bp deletion may reduce the mtDNA copy number and decrease the levels of cellular energy. However, conflicting reports have shown that this deletion does not affect the mtDNA copy number (53); its functional role requires further elucidation in future studies.

tRNA^{Ile} A4295G mutation. The homoplasmic A4295G mutation in the tRNA^{Ile} gene was identified in a three generation Han Chinese family with maternally inherited EH (56). An A to G transition at nucleotide position 4295 was shown to be highly evolutionarily conserved, and was not present in the healthy controls (57). The A4295G mutation is localized at the 3' end adjacent to the anticodon (position 37) of tRNA^{Ile} (45); nucleotide at position 37 is responsible for the stabilization of functional tRNA (58). Thus, it has been suggested that this mutation reduces the efficiency with which tRNA^{Ile} can be processed by 3'-tRNase, reducing the level of functional tRNA^{Ile} (59). The functional characterization of the A4295G mutation shows that this mutation induces a significant decrease in complex III protein levels, leading to a decrease in the activity of this complex, which is reflected by the decline in mitochondrial respiration (60).

tRNA^{Ile} A4263G mutation. The A4263G mutation changes the stop codon TAA of the *ND1* mRNA to an equivalent TAG stop codon. This mutation causes an A to G transition at the 5' end of the tRNA^{Ile} gene (61,62). Cybrid cells derived from the proband carrying A4263G mutation show a reduction in tRNA^{Ile} steady-state levels, as well as in the rate of mitochondrial protein translation. Increased ROS and the decreased efficiency of 5' end processing of tRNA^{Ile} precursor indicated that this mutation caused mitochondrial dysfunction that was responsible for EH in this Han Chinese family (61).

tRNA^{Ile} T4291C mutation. The homoplasmic T4291C mutation in the tRNA^{Ile} gene has been associated with a cluster of metabolic defects, including hypertension, hypercholesterolemia and hypomagnesaemia in a large family (63). The T4291C mutation occurs immediately 5' to the tRNA^{Ile} anticodon (position 33), and is conserved in every sequenced tRNA^{Ile} from bacteria to human mitochondria (64). Biochemical studies with anticodon stem-loop analogs of tRNAs have been performed and have indicated that the substitution of cytidine for uridine at this position markedly impairs ribosomal binding (65), providing evidence of the functional importance of this mutation.

tRNA^{Met} A4435G mutation. The presence of the A4435G mutation with chronic progressive external ophthalmoplegia (CPEO) was initially reported in the study by Jaksch *et al* (66). Later on, this mutation was found in patients with LHON (67), as well as in a Japanese subject with diabetes (68). In addition, the East Asian haplogroup G2a1-specific A4435G mutation (69) has also been associated with EH in two Chinese families (70,71). The homoplasmic A4435G mutation, which is located at immediately 3' end to the anticodon of tRNA^{Met} (nucleotide position 37), is extremely conserved from bacteria to human mitochondria (45). In fact, as shown in previous studies, the A4435G mutation causes ~40-50% reduction in the steady-state level of tRNA^{Met} and consequently results in the failure of mt-tRNA metabolism (67,70). Impaired mt-tRNA metabolism subsequently worsens the mitochondrial protein synthesis, decreases ATP production and increases ROS levels. Thus, mitochondrial dysfunction may contribute to the development of EH in these families carrying the A4435G point mutation (37,63,72).

Table I. Summary of cardiovascular diseases associated mt-tRNA mutations.

Position	tRNA species	Allele	Homoplasmy/ heteroplasmy	Clinical features	Refs.
29	tRNA ^{Val}	C1628T	Homoplasmy	Cardiomyopathy	(80)
14	tRNA ^{Leu(UUR)}	A3243G	Heteroplasmy	Cardiomyopathy	(81)
29	tRNA ^{Leu(UUR)}	A3260G	Heteroplasmy	Myopathy, cardiomyopathy, MELAS	(82-84)
72	tRNA ^{Leu(UUR)}	C3303T	Homoplasmy	Cardiomyopathy, myopathy, HCM	(85-87)
1	tRNA ^{Ile}	A4263G	Homoplasmy	Hypertension	(61)
7	tRNA ^{Ile}	A4269G	Homoplasmy	Cardiomyopathy	(88)
15	tRNA ^{Ile}	T4277C	Homoplasmy	Cardiomyopathy	(89)
33	tRNA ^{Ile}	T4291C	Homoplasmy	Hypertension, hypercholesterolemia, hypomagnesaemia	(63)
37	tRNA ^{Ile}	A4295G	Homoplasmy	HCM, hypertension	(56,57)
42	tRNA ^{Ile}	A4300G	Homoplasmy	HCM	(90)
37	tRNA ^{Met}	A4435G	Homoplasmy	LHON, CPEO, hypertension	(66-68,70,71)
1	tRNA ^{Gln} and tRNA ^{Met}	A4401G	Homoplasmy	LVH, hypertension	(73,74)
49	tRNA ^{Gln}	T4353C	Homoplasmy	Hypertension	(76)
58	tRNA ^{Ala}	A5600T	Heteroplasmy	DCM	(91)
54	tRNA ^{Lys}	A8348G	Heteroplasmy	Cardiomyopathy	(92)
7	tRNA ^{Gly}	T9997C	Heteroplasmy	HCM	(93)
59	tRNA ^{His}	G12192A	Homoplasmy	Cardiomyopathy	(94)
67	tRNA ^{Leu(CUN)}	A12330G	Homoplasmy	Hypertension	(43)
37	tRNA ^{Glu}	T14709C	Homoplasmy	Cardiomyopathy	(95)
2	tRNA ^{Thr}	T15889C	Heteroplasmy	DCM	(91)
16	tRNA ^{Thr}	A15902G	Heteroplasmy	DCM	(91)
38	tRNA ^{Thr}	T15924C	Homoplasmy	DCM	(96)
51	tRNA ^{Thr}	A15935G	Heteroplasmy	DCM	(91)

MELAS, mitochondrial encephalomyopathy, lactic acidosis and stroke-like episodes; HCM, hypertrophic cardiomyopathy; LVH, left ventricular hypertrophy; LHON, Leber's hereditary optic neuropathy; CPEO, chronic progressive external ophthalmoplegia; DCM, dilated cardiomyopathy.




Figure 2. Cloverleaf structure of mt-tRNA with standard nucleotide numbering. The nucleotides in the variable stem have the prefix 'e' and are located between positions 45 and 46, obeying the base-pairing rules. The nucleotides in the 54-strand and 34-strand are numbered by e11, e12, e13; the second digit identifies the base-pair.

A4401G mutation in tRNA^{Met} and tRNA^{Gln}. The homoplasmic A4401G mutation was originally described in a three generation Chinese family with left ventricular hypertrophy (LVH) (73). This mutation was also present in a five generation Chinese pedigree with EH (74). The A4401G mutation is localized at the junction of tRNA^{Met} at the H-strand and tRNA^{Gln} at the L-strand (75). Thus, it is anticipated that this mutation may lead to defective tRNA^{Met} 5' end processing in the H-strand transcripts and reduce the efficiency of tRNA^{Gln} precursor 5' end cleavage in the L-strand transcripts. Functional characterization of cell lines derived from the proband carrying the A4401G mutation shows virtually ~30% reduction in the levels of tRNA^{Ile} and tRNA^{Gln} (74). In addition, the mutant cell lines present a significant decrease in the oxygen consumption rate (73). These findings indicate that the A4401G mutation is involved in the pathogenesis of EH in Han Chinese families.

T4353C mutation in tRNA^{Gln}. The T4353C mutation, in conjunction with the tRNA^{Trp} C593T mutation, has been shown to account for the high penetrance of EH in a Han Chinese family (76). Clinical evaluation of this family showed a typical

maternally transmitted pattern. Analysis of the complete mtDNA sequence identified the homoplasmic T4353C mutation. This mutation alters a conservative base-pairing (49A-65U) on the T arm of tRNA^{Gln}, thereby affecting tRNA metabolism (45). Moreover, the T4353C mutation reduces the steady-state level of tRNA^{Gln}, mutant tRNA^{Gln} and tRNA^{Phe} may be metabolically less stable and more subject to degradation. The cybrid cell lines carrying the T4353C mutation present mitochondrial protein synthesis defects. As a result, this mutation causes the mitochondrial dysfunction responsible for EH.

6. Molecular mechanisms behind mtDNA mutations associated with EH

In previous studies, we noticed that several hypertension-associated mitochondrial pathogenic mutations were located in tRNA genes. Mt-tRNA mutations have structural and functional effects, including destabilization of the tRNA tertiary structure, altered processing of RNA precursors, loss of nucleotide modification and deficient aminoacylation (Fig. 2 and Table I). In particular, these pathogenic mutations may lead to deficiencies in tRNA 3' end metabolism including 3' end cleavage, CCA addition, aminoacylation or impairment of critical subunits of respiratory chain functions. Failures in mt-tRNA metabolism subsequently lead to the impairment of mitochondrial protein (77-79). However, mutations in protein coding genes may have the potential to affect OXPHOS and in turn cellular death, resulting in a failure in mitochondrial protein synthesis. The increased bioavailability of ROS results in oxidative stress, which leads to cardiovascular and renal damage, thus, contributing to EH.

7. Conclusion

EH is a multifactorial syndrome that is characterized by abnormal energy metabolism and high BP. Recent studies have identified the mitochondria as the target and origin of major pathogenic pathways which lead to the progression of hypertension (97-99). Although existing therapies have been beneficial, there is a clear need for new approaches to treatment. Pharmacological targeting of the cellular stresses underlying mitochondrial dysfunction is showing promise. In addition, screening for mtDNA mutations for the clinical expression of hypertension may provide further insight into the understanding of the pathophysiology for maternally inherited hypertension.

Acknowledgements

This study was supported by grants from the Ministry of Public Health of Zhejiang Province (2013KYA158), Nanjing Medical University (no. 2010NJMU011), Hangzhou Bureau of Science and Technology (no. 20120533Q03) and the Specialized Research Fund for the Doctoral Program of Higher Education (no. 20124323120004).

References

1. El Shamieh S, Herbeth B, Azimi-Nezhad M, *et al*: Human formyl peptide receptor 1 C32T SNP interacts with age and is associated with blood pressure levels. *Clin Chim Acta* 413: 34-38, 2012.
2. Kearney PM, Whelton M, Reynolds K, *et al*: Global burden of hypertension: analysis of worldwide data. *Lancet* 365: 217-223, 2005.
3. Messerli FH, Williams B and Ritz E: Essential hypertension. *Lancet* 370: 591-560, 2007.
4. Kunes J and Zicha J: Developmental windows and environment as important factors in the expression of genetic information: a cardiovascular physiologist's view. *Clin Sci* 111: 295-305, 2006.
5. Timberlake DS, O'Connor DT and Parmer RJ: Molecular genetics of essential hypertension: recent results and emerging strategies. *Curr Opin Nephrol Hypertens* 10: 71-79, 2001.
6. Fan JB, Chen X, Halushka MK, *et al*: Parallel genotyping of human SNPs using generic high-density oligonucleotide tag arrays. *Genome Res* 10: 853-860, 2000.
7. Oliphant A, Barker DL, Stuelpnagel JR and Chee MS: BeadArray technology: enabling an accurate, cost-effective approach to high-throughput genotyping. *Biotechniques* 56 (Suppl 56-58): 60-61, 2002.
8. Levy D, Ehret GB, Rice K, *et al*: Genome-wide association study of blood pressure and hypertension. *Nat Genet* 41: 677-687, 2009.
9. Tobin MD, Tomaszewski M, Braund PS, *et al*: Common variants in genes underlying monogenic hypertension and hypotension and blood pressure in the general population. *Hypertension* 51: 1658-1664, 2008.
10. Padmanabhan S, Melander O, Johnson T, *et al*: Genome-wide association study of blood pressure extremes identifies variant near UMOD associated with hypertension. *PLoS Genet* 6: e1001177, 2010.
11. Schon EA: Mitochondria. In: *Encyclopedia of Human Biology*. Dulbecco R (ed). Vol 5. 2nd edition. Academic Press, London, pp713-724, 1997.
12. Wallace DC: Mitochondrial DNA mutations in disease and aging. *Environ Mol Mutagen* 51: 440-450, 2010.
13. Garcia-Rodriguez LJ: Appendix 1. Basic properties of mitochondria. *Methods Cell Biol* 80: 809-812, 2007.
14. DiMauro S and Schon EA: Mitochondrial DNA mutations in human disease. *Am J Med Genet* 106: 18-26, 2001.
15. Cadenas E and Davies KJ: Mitochondrial free radical generation, oxidative stress, and aging. *Free Radic Biol Med* 29: 222-230, 2000.
16. Schon EA, DiMauro S and Hirano M: Human mitochondrial DNA: roles of inherited and somatic mutations. *Nat Rev Genet* 13: 878-890, 2012.
17. McBride HM, Neuspiel M and Wasiak S: Mitochondria: more than just a powerhouse. *Curr Biol* 16: R551-R560, 2006.
18. Choksi KB, Boylston WH, Rabek JP, *et al*: Oxidatively damaged proteins of heart mitochondrial electron transport complexes. *Biochim Biophys Acta* 1688: 95-101, 2004.
19. Murphy MP: Induction of mitochondrial ROS production by electrophilic lipids: a new pathway of redox signaling? *Am J Physiol Heart Circ Physiol* 290: H1754-H1755, 2006.
20. Abou-Sleiman PM, Muqit MM and Wood NW: Expanding insights of mitochondrial dysfunction in Parkinson's disease. *Nat Rev Neurosci* 7: 207-219, 2006.
21. Li P, Nijhawan D and Wang X: Mitochondrial activation of apoptosis. *Cell* 116 (Suppl 2): S57-S59, 2004.
22. Wu L and Juurlink BH: Increased methylglyoxal and oxidative stress in hypertensive rat vascular smooth muscle cells. *Hypertension* 39: 809-814, 2002.
23. Lerman LO, Nath KA, Rodriguez-Porcel M, *et al*: Increased oxidative stress in experimental renovascular hypertension. *Hypertension* 37: 541-546, 2001.
24. Trolliet MR, Rudd MA and Loscalzo J: Oxidative stress and renal dysfunction in salt-sensitive hypertension. *Kidney Blood Press Res* 24: 116-123, 2001.
25. Romero JC and Reckelhoff JF: State-of-the-Art lecture. Role of angiotensin and oxidative stress in essential hypertension. *Hypertension* 34: 943-949, 1999.
26. Hamilton CA, Berg G, McIntyre M, *et al*: Effects of nitric oxide and superoxide on relaxation in human artery and vein. *Atherosclerosis* 133: 77-86, 1997.
27. Paravicini TM and Touyz RM: Redox signaling in hypertension. *Cardiovasc Res* 71: 247-258, 2006.
28. Prezant TR, Agopian JV, Bohlman MC, *et al*: Mitochondrial ribosomal RNA mutation associated with both antibiotic-induced and non-syndromic deafness. *Nat Genet* 4: 289-294, 1993.
29. Estivill X, Govea N, Barceló E, *et al*: Familial progressive sensorineural deafness is mainly due to the mtDNA A1555G mutation and is enhanced by treatment of aminoglycosides. *Am J Hum Genet* 62: 27-35, 1998.

30. Guan MX: Molecular pathogenetic mechanism of maternally inherited deafness. *Ann NY Acad Sci* 1011: 259-271, 2004.
31. Van Camp G and Smith RJ: Maternally inherited hearing impairment. *Clin Genet* 57: 409-414, 2000.
32. Chen H, Zheng J, Xue L, *et al*: The 12S rRNA A1555G mutation in the mitochondrial haplogroup D5a is responsible for maternally inherited hypertension and hearing loss in two Chinese pedigrees. *Eur J Hum Genet* 20: 607-612, 2012.
33. Hamasaki K and Rando RR: Specific binding of aminoglycosides to a human rRNA construct based on a DNA polymorphism which causes aminoglycoside-induced deafness. *Biochemistry* 36: 12323-12328, 1997.
34. Cotney J, McKay SE and Shadel GS: Elucidation of separate, but collaborative functions of the rRNA methyltransferase-related human mitochondrial transcription factors B1 and B2 in mitochondrial biogenesis reveals new insight into maternally inherited deafness. *Hum Mol Genet* 18: 2670-2682, 2009.
35. Hobbie SN, Bruell CM, Akshay S, *et al*: Mitochondrial deafness alleles confer misreading of the genetic code. *Proc Natl Acad Sci USA* 105: 3244-3249, 2008.
36. Guan MX, Fischel-Ghodsian N and Attardi G: Nuclear background determines biochemical phenotype in the deafness-associated mitochondrial 12S rRNA mutation. *Hum Mol Genet* 10: 573-580, 2001.
37. Bernal-Mizrachi C, Gates AC, Weng S, *et al*: Vascular respiratory uncoupling increases blood pressure and atherosclerosis. *Nature* 435: 502-506, 2005.
38. Li X, Fischel-Ghodsian N, Schwartz F, *et al*: Biochemical characterization of the mitochondrial tRNA^{Ser(UCN)} T7511C mutation associated with nonsyndromic deafness. *Nucleic Acids Res* 32: 867-877, 2004.
39. Li R, Ishikawa K, Deng JH, *et al*: Maternally inherited nonsyndromic hearing loss is associated with the T7511C mutation in the mitochondrial tRNA^{Ser(UCN)} gene in a Japanese family. *Biochem Biophys Res Commun* 328: 32-37, 2005.
40. Liu Y, Li Z, Yang L, *et al*: The mitochondrial ND1 T3308C mutation in a Chinese family with the secondary hypertension. *Biochem Biophys Res Commun* 368: 18-22, 2008.
41. Anderson S, Bankier AT, Barrell BG, *et al*: Sequence and organization of the human mitochondrial genome. *Nature* 290: 457-465, 1981.
42. Guan MX, Enriquez JA, Fischel-Ghodsian N, *et al*: The deafness-associated mitochondrial DNA mutation at position 7445, which affects tRNA^{Ser(UCN)} precursor processing, has long-range effects on NADH dehydrogenase subunit ND6 gene expression. *Mol Cell Biol* 18: 5868-5879, 1998.
43. Teng L, Zheng J, Leng J and Ding Y: Clinical and molecular characterization of a Han Chinese family with high penetrance of essential hypertension. *Mitochondrial DNA* 23: 461-465, 2012.
44. Liu Z, Song Y, Gu S, *et al*: Mitochondrial ND5 12338T>C variant is associated with maternally inherited hypertrophic cardiomyopathy in a Chinese pedigree. *Gene* 506: 339-343, 2012.
45. Florentz C, Sohm B, Tryoen-Tóth P, *et al*: Human mitochondrial tRNAs in health and disease. *Cell Mol Life Sci* 60: 1356-1375, 2003.
46. Yu-Wai-Man P, Griffiths PG, Hudson G and Chinnery PF: Inherited mitochondrial optic neuropathies. *J Med Genet* 46: 145-158, 2009.
47. Guo H, Zhuang XY, Zhang AM, *et al*: Presence of mutation m.14484T>C in a Chinese family with maternally inherited essential hypertension but no expression of LHON. *Biochim Biophys Acta* 1822: 1535-1543, 2012.
48. Andreu AL, Hanna MG, Reichmann H, *et al*: Exercise intolerance due to mutations in the cytochrome b gene of mitochondrial DNA. *N Engl J Med* 341: 1037-1044, 1999.
49. Andreu AL, Bruno C, Dunne TC, *et al*: A nonsense mutation (G15059A) in the cytochrome b gene in a patient with exercise intolerance and myoglobinuria. *Ann Neurol* 45: 127-130, 1999.
50. Nikitin AG, Lavrikova EY and Chistiakov DA: The heteroplasmic 15059G>A mutation in the mitochondrial cytochrome b gene and essential hypertension in type 2 diabetes. *Diabetes Metab Syndr* 6: 150-156, 2012.
51. Chang DD and Clayton DA: Priming of human mitochondrial DNA replication occurs at the light-strand promoter. *Proc Natl Acad Sci USA* 82: 351-355, 1985.
52. Kang D, Miyako K, Kai Y, *et al*: In vivo determination of replication origins of human mitochondrial DNA by ligation-mediated polymerase chain reaction. *J Biol Chem* 272: 15275-15279, 1997.
53. Bi R, Zhang AM, Zhang W, *et al*: The acquisition of an inheritable 50-bp deletion in the human mtDNA control region does not affect the mtDNA copy number in peripheral blood cells. *Hum Mutat* 31: 538-543, 2010.
54. Elango S, Govindaraj P, Vishwanadha VP, *et al*: Analysis of mitochondrial genome revealed a rare 50 bp deletion and substitutions in a family with hypertension. *Mitochondrion* 11: 878-885, 2011.
55. Pham XH, Farge G, Shi Y, *et al*: Conserved sequence box II directs transcription termination and primer formation in mitochondria. *J Biol Chem* 281: 24647-24652, 2006.
56. Li Z, Liu Y, Yang L, *et al*: Maternally inherited hypertension is associated with the mitochondrial tRNA(Ile) A4295G mutation in a Chinese family. *Biochem Biophys Res Commun* 367: 906-911, 2008.
57. Merante F, Myint T, Tein I, *et al*: An additional mitochondrial tRNA(Ile) point mutation (A-to-G at nucleotide 4295) causing hypertrophic cardiomyopathy. *Hum Mutat* 8: 216-222, 1996.
58. Suzuki T, Nagao A and Suzuki T: Human mitochondrial tRNAs: biogenesis, function, structural aspects, and diseases. *Annu Rev Genet* 45: 299-329, 2011.
59. Levinger L, Giegé R and Florentz C: Pathology-related substitutions in human mitochondrial tRNA(Ile) reduce precursor 3' end processing efficiency in vitro. *Nucleic Acids Res* 31: 1904-1912, 2003.
60. Gutiérrez Cortés N, Pertuiset C, Dumon E, *et al*: Novel mitochondrial DNA mutations responsible for maternally inherited nonsyndromic hearing loss. *Hum Mutat* 33: 681-689, 2012.
61. Wang S, Li R, Fettermann A, *et al*: Maternally inherited essential hypertension is associated with the novel 4263A>G mutation in the mitochondrial tRNA^{Ile} gene in a large Han Chinese family. *Circ Res* 108: 862-870, 2011.
62. Zhu HY, Wang SW, Liu L, *et al*: Genetic variants in mitochondrial tRNA genes are associated with essential hypertension in a Chinese Han population. *Clin Chim Acta* 410: 64-69, 2009.
63. Wilson FH, Hariri A, Farhi A, *et al*: A cluster of metabolic defects caused by mutation in a mitochondrial tRNA. *Science* 306: 1190-1194, 2004.
64. Sprinzl M, Steegborn C, Hübel F and Steinberg S: Compilation of tRNA sequences and sequences of tRNA genes. *Nucleic Acids Res* 24: 68-72, 1996.
65. Ashraf SS, Sochacka E, Cain R, *et al*: Single atom modification (O>S) of tRNA confers ribosome binding. *RNA* 5: 188-194, 1999.
66. Jaksch M, Kleinle S, Scharfe C, *et al*: Frequency of mitochondrial transfer RNA mutations and deletions in 225 patients presenting with respiratory chain deficiencies. *J Med Genet* 38: 665-673, 2001.
67. Qu J, Li R, Zhou X, *et al*: The novel A4435G mutation in the mitochondrial tRNA^{Met} may modulate the phenotypic expression of the LHON-associated ND4 G11778A mutation. *Invest Ophthalmol Vis Sci* 47: 475-483, 2006.
68. Guo LJ, Oshida Y, Fuku N, *et al*: Mitochondrial genome polymorphisms associated with type-2 diabetes or obesity. *Mitochondrion* 5: 15-33, 2005.
69. Kong QP, Bandelt HJ, Sun C, *et al*: Updating the East Asian mtDNA phylogeny: a prerequisite for the identification of pathogenic mutations. *Hum Mol Genet* 15: 2076-2086, 2006.
70. Liu Y, Li R, Li Z, *et al*: Mitochondrial transfer RNAMet 4435A>G mutation is associated with maternally inherited hypertension in a Chinese pedigree. *Hypertension* 53: 1083-1090, 2009.
71. Lu Z, Chen H, Meng Y, *et al*: The tRNA^{Met} 4435A>G mutation in the mitochondrial haplogroup G2a1 is responsible for maternally inherited hypertension in a Chinese pedigree. *Eur J Hum Genet* 19: 1181-1186, 2011.
72. Postnov YV, Orlov SN, Budnikov YY, *et al*: Mitochondrial energy conversion disturbance with decrease in ATP production as a source of systemic arterial hypertension. *Pathophysiology* 14: 195-204, 2007.
73. Zhu HY, Wang SW, Liu L, *et al*: A mitochondrial mutation A4401G is involved in the pathogenesis of left ventricular hypertrophy in Chinese hypertensives. *Eur J Hum Genet* 17: 172-178, 2009.
74. Li R, Liu Y, Li Z, *et al*: Failures in mitochondrial tRNA^{Met} and tRNA^{Gln} metabolism caused by the novel 4401A>G mutation are involved in essential hypertension in a Han Chinese Family. *Hypertension* 54: 329-337, 2009.
75. Ojala D, Montoya J and Attardi G: tRNA punctuation model of RNA processing in human mitochondria. *Nature* 290: 470-474, 1981.
76. Qiu Q, Li R, Jiang P, *et al*: Mitochondrial tRNA mutations are associated with maternally inherited hypertension in two Han Chinese pedigrees. *Hum Mutat* 33: 1285-1293, 2012.

77. Levinger L, Mörl M and Florentz C: Mitochondrial tRNA 3' end metabolism and human disease. *Nucleic Acids Res* 32: 5430-5441, 2004.
78. Kelley SO, Steinberg SV and Schimmel P: Functional defects of pathogenic human mitochondrial tRNAs related to structural fragility. *Nat Struct Biol* 7: 862-865, 2000.
79. Kelley SO, Steinberg SV and Schimmel P: Fragile T-stem in disease-associated human mitochondrial tRNA sensitizes structure to local and distant mutations. *J Biol Chem* 276: 10607-10611, 2001.
80. Arredondo JJ, Gallardo ME, García-Pavía P, *et al*: Mitochondrial tRNA valine as a recurrent target for mutations involved in mitochondrial cardiomyopathies. *Mitochondrion* 12: 357-362, 2012.
81. Vilarinho L, Santorelli FM, Rosas MJ, *et al*: The mitochondrial A3243G mutation presenting as severe cardiomyopathy. *J Med Genet* 34: 607-609, 1997.
82. Zeviani M, Gellera C, Antozzi C, *et al*: Maternally inherited myopathy and cardiomyopathy: association with mutation in mitochondrial DNA tRNA(Leu)(UUR). *Lancet* 338: 143-147, 1991.
83. Sweeney MG, Brockington M, Weston MJ, *et al*: Mitochondrial DNA transfer RNA mutation Leu(UUR)A→G 3260: a second family with myopathy and cardiomyopathy. *Q J Med* 86: 435-438, 1993.
84. Nishino I, Komatsu M, Kodama S, *et al*: The 3260 mutation in mitochondrial DNA can cause mitochondrial myopathy, encephalopathy, lactic acidosis, and stroke-like episodes (MELAS). *Muscle Nerve* 19: 1603-1604, 1996.
85. Goldstein JD, Shanske S, Bruno C and Perszyk AA: Maternally inherited mitochondrial cardiomyopathy associated with a C-to-T transition at nucleotide 3303 of mitochondrial DNA in the tRNA(Leu)(UUR) gene. *Pediatr Dev Pathol* 2: 78-85, 1999.
86. Silvestri G, Santorelli FM, Shanske S, *et al*: A new mtDNA mutation in the tRNA(Leu)(UUR) gene associated with maternally inherited cardiomyopathy. *Hum Mutat* 3: 37-43, 1994.
87. Palecek T, Tesarova M, Kuchynka P, *et al*: Hypertrophic cardiomyopathy due to the mitochondrial DNA mutation m.3303C>T diagnosed in an adult male. *Int Heart J* 53: 383-387, 2012.
88. Taniike M, Fukushima H, Yanagihara I, *et al*: Mitochondrial tRNA(Ile) mutation in fatal cardiomyopathy. *Biochem Biophys Res Commun* 186: 47-53, 1992.
89. Giordano C, Perli E, Orlandi M, *et al*: Cardiomyopathies due to homoplasmic mitochondrial tRNA mutations: morphologic and molecular features. *Hum Pathol* 44: 1262-1270, 2013.
90. Taylor RW, Giordano C, Davidson MM, *et al*: A homoplasmic mitochondrial transfer ribonucleic acid mutation as a cause of maternally inherited hypertrophic cardiomyopathy. *J Am Coll Cardiol* 41: 1786-1796, 2003.
91. Arbustini E, Diegoli M, Fasani R, *et al*: Mitochondrial DNA mutations and mitochondrial abnormalities in dilated cardiomyopathy. *Am J Pathol* 153: 1501-1510, 1998.
92. Terasaki F, Tanaka M, Kawamura K, *et al*: A case of cardiomyopathy showing progression from the hypertrophic to the dilated form: association of Mt8348A→G mutation in the mitochondrial tRNA(Lys) gene with severe ultrastructural alterations of mitochondria in cardiomyocytes. *Jpn Circ J* 65: 691-694, 2001.
93. Merante F, Tein I, Benson L and Robinson BH: Maternally inherited hypertrophic cardiomyopathy due to a novel T-to-C transition at nucleotide 9997 in the mitochondrial tRNA(glycine) gene. *Am J Hum Genet* 55: 437-446, 1994.
94. Shin WS, Tanaka M, Suzuki J, *et al*: A novel homoplasmic mutation in mtDNA with a single evolutionary origin as a risk factor for cardiomyopathy. *Am J Hum Genet* 67: 1617-1620, 2000.
95. Van Hove JL, Freehauf C, Miyamoto S, *et al*: Infantile cardiomyopathy caused by the T14709C mutation in the mitochondrial tRNA glutamic acid gene. *Eur J Pediatr* 167: 771-776, 2008.
96. Ruppert V, Nolte D, Aschenbrenner T, *et al*: Novel point mutations in the mitochondrial DNA detected in patients with dilated cardiomyopathy by screening the whole mitochondrial genome. *Biochem Biophys Res Commun* 318: 535-543, 2004.
97. Dikalov S: Cross talk between mitochondria and NADPH oxidases. *Free Radic Biol Med* 51: 1289-1301, 2011.
98. Leone TC and Kelly DP: Transcriptional control of cardiac fuel metabolism and mitochondrial function. *Cold Spring Harb Symp Quant Biol* 76: 175-182, 2011.
99. Cottrill KA and Chan SY: Metabolic dysfunction in pulmonary hypertension: the expanding relevance of the Warburg effect. *Eur J Clin Invest* 43: 855-865, 2013.