

Unusual clotting dynamics of plasma supplemented with iron(III)

JERZY JANKUN¹⁻³, PHILIP LANDETA¹, ETHERESIA PRETORIUS⁴,
EWA SKRZYPCZAK-JANKUN¹ and BOGUSŁAW LIPINSKI⁵

¹Urology Research Center, Department of Urology, The University of Toledo - Health Science Campus, Toledo, OH, USA;

²Protein Research Chair, Department of Biochemistry, College of Sciences, King Saud University,

Riyadh, Kingdom of Saudi Arabia; ³Department of Clinical Nutrition, Medical University of Gdańsk, Gdańsk, Poland;

⁴Department of Physiology, Faculty of Health Sciences, University of Pretoria, Arcadia, Pretoria,

Republic of South Africa; ⁵Joslin Diabetes Center, Harvard Medical School, Boston, MA, USA

Received November 8, 2013; Accepted December 5, 2013

DOI: 10.3892/ijmm.2013.1585

Abstract. Iron salts are used in the treatment of iron deficiency anemia. Diabetic patients are frequently anemic and treatment includes administration of iron. Anemic patients on hemodialysis are at an increased risk of thromboembolic coronary events associated with the formation of dense fibrin clots resistant to fibrinolysis. Moreover, in chronic kidney disease patients, high labile plasma iron levels associated with iron supplementation are involved in complications found in dialyzed patients such as myocardial infarction. The aim of the present study was to investigate whether iron treatment is involved in the formation of the fibrin clots. Clotting of citrated plasma supplemented with Fe³⁺ was investigated by thromboelastometry and electron microscopy. The results revealed that iron modifies coagulation in a complex manner. FeCl₃ stock solution underwent gradual chemical modification during storage and altered the coagulation profile over 29 days, suggesting that Fe³⁺ interacts with both proteins of the coagulation cascade as well as the hydrolytic Fe³⁺ species. Iron extends clotting of plasma by interacting with proteins of the coagulation cascade. Fe³⁺ and/or its hydrolytic species interact with fibrinogen and/or fibrin changing their morphology and properties. In general FeCl₃ weakens the fibrin clot while at the same time precipitating plasma proteins immediately after application. Fe³⁺ or its derivatives induced the formation of insoluble coagulums in non-enzymatic reactions including albumin and transferrin. Iron plays a role in coagulation and can precipitate plasma proteins. The formation of coagulums resistant to lysis in non-enzymatic reactions can increase the risk of thrombosis, and extending clotting of plasma can prolong bleeding.

Introduction

Iron salts are used in the treatment of iron deficiency anemia, as a supplemental intake of iron during pregnancy and in multivitamin preparations. In the majority of cases it is safe to use but toxic effects begin to appear at doses >10-20 mg/kg of elemental iron, and ingestions of >50 mg/kg are associated with severe toxicity or lethality (1,2). Traditionally ferric chloride was used in an arterial thrombosis model in rats to induce vessel damage resulting in blood clotting (3). In the 19th century, ferrous salts and ferrous chloride in particular were considered the most effective agents in stanching the flow of blood from wounds (4). Moreover, it was demonstrated that ferric chloride treatment of mouse aorta *ex vivo* caused endothelial denudation, collagen exposure and when injected intravenously formed occlusive thrombi (5,6). Diabetic patients are frequently anemic and treatment may include oral or intravenous iron administration (7). Undas *et al* observed that anemic patients on hemodialysis due to chronic kidney disease (CKD) are at an increased risk of thromboembolic coronary events associated with the formation of dense fibrin clots resistant to fibrinolysis (8). Moreover, in CKD patients a high labile plasma iron level (LPI) associated with iron supplementation is involved in complications in dialyzed patients such as myocardial infarction and bacterial infection (9).

The role of iron treatment in the formation of fibrin clots should therefore be investigated. Of note is that in humans divalent iron, Fe²⁺, is rapidly oxidized to trivalent iron, Fe³⁺, by ferroxidase (10). Additionally, ferric chloride and in general Fe³⁺ ions show a markedly complex chemistry producing a multiplicity of compounds over 29 days as shown in the examples (11): FeCl₃ + 3H₂O ⇌ Fe(OH)Cl₂ + HCl + 2H₂O ⇌ Fe(OH)₂Cl + 2HCl + H₂O ⇌ Fe(OH)₃ + 3HCl. Feng and Nansheng provide additional species distribution of three simple low-molecular-weight Fe³⁺ hydroxy complexes (12): Fe³⁺ + H₂O → Fe(OH)²⁺ + H⁺; Fe³⁺ + 2H₂O → Fe(OH)₂⁺ + 2H⁺; Fe³⁺ + 2H₂O → Fe₂(OH)₂⁴⁺ + 2H⁺. Moreover, reactive free radicals are produced in the presence of ferric ions alone by the Fenton reaction (10): Fe³⁺ + HO⁻ → Fe²⁺ + HO. These changes can be analyzed by UV spectroscopy since different iron

Correspondence to: Professor Jerzy Jankun, Urology Research Center, Department of Urology, The University of Toledo - Health Science Campus, 3000 Arlington, Toledo, OH 43614, USA
E-mail: jerzy.jankun@utoledo.edu

Key words: coagulation, fibrin, iron, plasma

chemicals have distinct λ_{\max} , for example: $\text{Fe}(\text{H}_2\text{O})_6^{3+}$ absorbs λ_{\max} at 240 nm, $\text{Fe}(\text{OH})_2^{2+}$ shows λ_{\max} at 205 and 297 nm, and $\text{Fe}_2(\text{OH})_2^{4+}$ comes into view at λ_{\max} at 335 nm (12).

In the present study, we investigated clotting of citrated plasma supplemented with Fe^{3+} (and calcium Ca^{2+} to initiate clotting) by thromboelastometry and electron microscopy. The results showed that iron changes plasma clotting characteristics, kinetics and the dynamics of clot formation in plasma. More changes were observed as the time of storing stock solution of FeCl_3 increased, possibly due to different derivatives of Fe^{3+} being formatted over 29 days. Additionally, the morphology of clotted fibrin in the Ca^{2+} - and Fe^{3+} -treated plasma was different than the untreated, normal, control-clotted fibrin.

Materials and methods

Chemicals, plasticware and proteins. Kaolin, CaCl_2 solution, pins and cups were purchased from Haemoscope Co. (Neils, IL, USA). Fully active human tissue plasminogen activator (tPA), product number HTPA-TC was purchased from Molecular Innovations, Inc. (Novi, MI, USA). Ferric chloride, fibrin and thrombin were purchased from Sigma-Aldrich Co. LLC (St. Louis, MO, USA).

Preparation of plasma. Lyophilized specialty assayed reference plasma, cat. no. 5185 (S.A.R.P., 10x1 ml) purchased from Helena Laboratories (Beaumont, TX, USA) was prepared from a frozen pool of citrated plasma obtained from healthy donors. S.A.R.P. has normal PT and aPTT clotting times and may be used as reference data based on the following parameters: fibrinogen**, factor II*, factorV**, factor VII*, factor VIII*, factor IX*, factor X*, factor XI**, ristocetin cofactor*, vWF:Ag*, factor XII, protein C*, protein S - total, free) where (*) denotes samples standardized according to World Health Organization (WHO) regulations, and (**) denotes samples calibrated against ISTH reference material. Plasma was stored at 4°C and reconstituted by adding 1 ml of deionized water, followed by a 3-min rest. Plasma for electron microscopy experiments was obtained from healthy subjects aged between 20 and 25 years, both males and females. Ethical approval was obtained from the University of Pretoria Human Ethics Committee, and this study conforms to the principles of the Declaration of Helsinki.

Analysis of plasma clot formation with thromboelastography. Thromboelastography allows measurement of a total coagulation profile and yields data on the kinetics and dynamics of clot formation in plasma (13). The essential part of the TEG® 5000 Thrombelastograph® Hemostasis Analyzer System (Haemonetics Corporation, Braintree, MA, USA) is a pin hanging on a torsion wire and inserted in a cup holding a sample (360 μl) (13,14). This pin oscillates at 6 rpm at a 4°45' angle at 37°C. When plasma viscosity changes during clot formation, the pin motion is progressively restrained by the clot and the cup. Sodium-citrated, reconstituted plasma was used for TEG assays by mixing 1 ml of plasma with 20 μl of kaolin and in some samples a constant amount of tPA was added [10 μl of tPA (2.1 mg/ml in 0.4 M HEPES, 0.1 M NaCl, pH 7.4)] as a fibrinolytic agent (15) to measure proteolysis under controlled conditions (16,17). Subsequently, 320 μl of the mixture was transferred to each cup and 20 μl of CaCl_2




Figure 1. UV/VIS spectra of FeCl_3 stock solution diluted at 1:1,000 at day 0 until day 29.

(0.2 M) and/or FeCl_3 (0.2 M) was added. In a separate experiment 1% of DMSO was added to the stock solution and plasma clotting was analyzed as described above. The critical parameters of clotting measured by TEG were: R was the time from initiation of the reaction until a measurable clot was detected, K was the time from the R point until a certain clot firmness was achieved, (α) was the maximum angle representing kinetics of clotting and LY30 (percentage) represented clot lysis 30 min after MA (maximum amplitude) (13,18,19).

Electron microscopy. Purified fibrinogen (cat. no. F3879-250MG; Sigma-Aldrich), human albumin (cat. no. A9511, Sigma-Aldrich) samples were treated with 5 μl 0.2 M CaCl_2 , followed by the addition of 5 μl of freshly prepared 0.2 M FeCl_3 . After mixing, thrombin was added, to create an extensive fibrin network. Human platelet rich plasma (PRP) samples were treated (addition of CaCl_2 and FeCl_3) in the same manner, but without thrombin. The samples were fixed immediately in 2.5% glutaraldehyde/formaldehyde in PBS solution, pH 7.4, for 30 min. The samples were then left for 16 min and 3 h, followed by fixing in order to obtain a time-dependent analysis of the effect of FeCl_3 and CaCl_2 on PRP. Smears were then fixed followed by rinsing three times with PBS for 5 min prior to being fixed for 30 min with 1% osmium tetroxide (OsO_4). The samples were again rinsed three times with PBS for 5 min and were dehydrated serially with 30, 50, 70 and 90% ethanol, and three times with 100% ethanol. The material was mounted and coated with carbon. A Zeiss ULTRA plus FEG-SEM with InLens capabilities (Microscopy and Microanalysis Unit of the University of Pretoria, Pretoria, South Africa) was used to study the surface morphology of fibrin and micrographs were taken at 1 kV.

UV/VIS spectrometry. FeCl_3 water solution was diluted at 1:1,000 from 0.2 M stock solution with or without DMSO and analyzed on a UV/VIS spectrometer at a range of 230–800 nm. Samples were analyzed at day 0 and periodically up to day 29 after FeCl_3 preparation.

Results

UV/VIS spectrometry. UV/VIS spectra of FeCl_3 were altered over the 29 days (Fig. 1). In general an increase in




Figure 2. Typical thromboelastogram of clotted plasma at day 0: (a) plasma treated with Ca^{2+} , control, solid blue line; plasma treated with Ca^{2+} and Fe^{3+} , dashed orange line; plasma treated with Ca^{2+} , Fe^{3+} and tPA at $0.5 \mu\text{g/ml}$, dashed dotted green line. Thromboelastogram of plasma at day 29: (b) plasma treated with Ca^{2+} , control, solid blue line; plasma treated with Ca^{2+} and Fe^{3+} , dashed orange line.

the absorption of $\sim 260 \text{ nm}$, and a slight increase of $\sim 290 \text{ nm}$ was observed. On day 14 and 29 the opposite changes were detected. An increase of absorption when approaching 330 nm was observed over the 29 days.

Analysis of plasma clot formation with thromboelastography. There are no normal ranges of TEG parameters for control plasma. However, the results yielded in this study were very consistent: R (sec): 408, K (sec): 84, An ($^\circ$): 70.6, MA (mm): 27.2, LY30 (%): 0 (all parameters $\pm 10\%$) (20). Observed parameters for all the controls were within these values. The addition of freshly prepared FeCl_3 was manifested by an immediate increase of viscosity/precipitation of plasma proteins [R (sec) ~ 10 , K (sec) N/A, MA (mm) ~ 15], followed by lysing as per classical thromboelastography (Fig. 2). Classic enzymatic coagulation appeared to be normal, with the exception of extended R time ($\sim 1,050 \text{ sec}$). The remaining parameters were normal (K, An, MA, LY30) although they were not recorded by the TEG instrument as it is not designed to register parameters for the second peak. The addition of tPA to plasma treated with FeCl_3 resulted in coagulation of proteins immediately after measurement but no enzymatic coagulation (Fig. 2). FeCl_3 was stored and it was observed that over time initial MA increased, while a secondary peak following initiation of enzymatic coagulation was observed. By contrast, the strength of the clot as measured by MA decreased. These changes were gradually more evident over time (Fig. 2).

While FeCl_3 stock solution with DMSO was used all described effects of Fe^{3+} were less evident confirming our previous suggestion that the free radicals were playing role in coagulation (10).

Electron microscopy. Electron microscopy images revealed that iron-treated plasma forms structures different from those of the control fibrin where fibrin strands formed a solid and thick mesh (Fig. 3). Proteins precipitated immediately after the addition of Fe^{3+} did not contain any typical fibrin fibers (Fig. 3a) but rather large aggregates with circular surface depressions, as is evident in the morphology of clotted plasma at $\sim 960 \text{ sec}$ (time when the secondary peak on the thromboelastogram was detected). Fig. 3b shows some scattered fiber strands typical for fibrin in addition to flat, irregular protein




Figure 3. Morphology of plasma treated with Ca^{2+} and Fe^{3+} at (a) ~ 0 , (b) ~ 960 and (c) $\sim 4000 \text{ sec}$. (d) Control plasma treated with Ca^{2+} . (e) Pure fibrinogen treated with thrombin and Ca^{2+} and Fe^{3+} . (f) Human serum albumin treated with Ca^{2+} and Fe^{3+} .

bodies. Images captured at the end of clotting show numerous granules covering the fiber strands (Fig. 3c).

Discussion

General. The aim of this study was to investigate the effects of Fe^{3+} on coagulation. However, during initial experiments we observed that the stock solution of FeCl_3 changed color and some changes were evident in the thromboelastogram. Therefore, we prepared FeCl_3 stock solution and analyzed the samples obtained by spectroscopy from day 0 to 29. Spectrometric data strongly indicated that FeCl_3 undergoes gradual chemical modification. Our results suggest that, for example, the concentration of $\text{Fe}_2(\text{OH})_2^{4+}$ (λ_{max} at 335 nm) increases while that of $\text{Fe}(\text{OH})^{2+}$ (λ_{max} at 297 nm) decreases,




Figure 4. TEG pin and cup of plasma treated with Ca^{2+} and Fe^{3+} . Arrows point to coagulums formed subsequent to Fe^{3+} addition.

which is in agreement with observations from previous studies (12). Clearly, the UV data have shown that iron hydrates or other chemicals are formed; however, they remain to be elucidated.

Fibrinogen coagulation. Simultaneously, we detected gradual changes in the plasma clotting parameters. Divalent and trivalent metals can change the clotting characteristics of plasma and blood (21). In the coagulation pathway, ions of calcium activate prothrombin to thrombin which converts fibrinogen to fibrin. Calcium is required for two distinct processes in prothrombin activation: binding factor X and prothrombin to the phospholipid surface. The first step is activated by numerous cations such as Mg^{2+} , Ca^{2+} , Sr^{2+} , Ba^{2+} , Mn^{2+} , Be^{2+} , Fe^{2+} , Fe^{3+} , Zn^{2+} (22). Replacement of calcium in this step can slow coagulation depending on the metal (22-26). However, the calcium binding sites involved in the protein-phospholipid structure, show exceptional selectivity for cations required for the protein transition, with the exception of strontium and barium, which can replace calcium in this role. The other metals form a protein-phospholipid complex with a different structure resulting in inhibition of the coagulation reactions (22,23). It is plausible that Fe^{3+} interferes in the mechanism of factor X-initiated prothrombin transformation resulting in slowing clot formation. Findings of previous studies have shown that iron prevents/slows the coagulation of normal plasma or blood while Mg^{2+} increases the clotting time of human plasma (21,27,28). However, we have found that iron modifies coagulation in a more complex manner than the simple extension of clot formation.

In the present study, we have established that coagulation parameters change as FeCl_3 storing time increases. Thus, Fe^{3+} as well as the hydrolytic Fe^{3+} species interact with proteins of the coagulation cascade. We also observed clot lysis following its initial formation. This happened a few days after the preparation of FeCl_3 stock solution and was more evident (Fig. 2b) over time. When fibrin is formed it is relatively unstable. The fibrin clot is stabilized catalytically by factor XIII and is, not only mechanically stronger than the non-cross-linked, but also less vulnerable to premature fibrinolysis degradation (10). Therefore it is possible that hydrolytic Fe^{3+} species inactivates factor XIII making possible a premature partial lysing of fibrin. Literature on iron and factor XIII is rather sparse, but it was reported that a severe iron intoxication in a 15-year-old girl resulted in numerous proteins of the fibrinolytic cascade, especially factor VIII and XIII, being affected (29).

Additionally, Fe^{3+} or its hydrolytic species interacts with fibrin or fibrinogen-changing morphology (Fig. 3c). Tightly bound fibrin fibers and spherical structures are clearly visible and this image differs significantly from that of the normal clot (Fig. 3d). Similar dense matted deposits and some spherical structures were observed even with lower concentrations of Fe^{3+} (30). Furthermore, it has been found that human fibrinogen directly recognizes iron ions and changes in the morphology of fibrin may be a result of this modification (31). The experiments conducted on the animals revealed that iron induces coagulopathy in a dose-dependent manner. It prolonged the prothrombin, thrombin, and partial thromboplastin time in animals as well as and in the human plasma. It was found that thrombin was markedly inhibited by iron in its clotting effect on fibrinogen. The inhibitory effect was reversible subsequent to iron removal by EDTA chelation and gel filtration. Additionally, amidolytic activity of thrombin, factor Xa, kallikrein, and trypsin were reversibly inhibited by Fe^{3+} . The coagulopathy was likely induced by Fe^{3+} as serine proteases are capable of binding Fe^{3+} ion(s) (28).

Free radicals are known to affect coagulation and fibrinolysis, and free radical scavengers normalize these processes (32) as was evident from results of our experiments with DMSO. It was reported that hydroxyl radical-induced modification of fibrin(ogen) molecules makes them resistant to fibrinolytic degradation (33). Subsequently, we treated plasma with Ca^{2+} , Fe^{3+} and tPA. Non-fibrinogen coagulation was identified when Fe^{3+} was added as expected. However, a fibrin clot was not formed, which may be attributed to delayed fibrin formation in the presence of Fe^{3+} , and degradation of fibrinogen and fibrin by plasmin activated by tPA (34-36). However, in that experiment we identified some residual but not lysed clots, which may be explained by the presence of fibrinogen molecules resistant to fibrinolytic degradation, as described by Lipinski *et al* (33).

Non-enzymatic coagulation/precipitation. In the Fe^{3+} -treated samples instantaneous formation of insoluble coagulums was observed. This effect was more prominent over time and was the effect of Fe^{3+} and its hydrolytic species (Fig. 2a and b). The thromboelastograms show that after protein(s) precipitation these coagulums were lysed, which may be an artifact. It seems that initially formed large aggregates with circular surface depressions (Fig. 3a) were self-aggregated to form some scattered fiber strands typical for fibrin in addition to spherical and flat, irregular protein bodies. Additionally, after removal of the pin from the TEG cup a reddish-colored clump was present in all Fe^{3+} -treated samples (Fig. 4), which may be due to initially formed, loosely connected, precipitated viscous proteins being clumped by oscillation of the pin inside of cup. This clump of proteins were rotated inside the cup with less resistance resulting in instrument interpretation of this as proteolysis.

We also attempted to identify the proteins that were precipitated following iron addition. Albumin is the most abundant protein in the circulation and represents 52-60% of the total plasma protein. It plays an important role in the transportation and storage of hormones, fatty acids and drugs, and in the transportation of essential metal ions. Both Fe^{2+} and Fe^{3+} ions bind to heme serum albumin through the heme iron

complex but only Fe^{3+} binds to heme-free albumin. Fe^{3+} ions are transported in plasma mainly by a non-heme iron-binding glycoprotein transferrin, which composes ~7-10% of plasma protein (37). Iron can denature proteins in general and albumin in particular (38,39). The two proteins constitute up to 70% of total plasma proteins. In a separate experiment we therefore show that Fe^{3+} precipitates albumin (Fig. 3f). The reddish color observed suggests that transferrin possibly co-precipitates among the other proteins incorporated into these particles (40,41).

FeCl_3 is used in animal models to study early arterial thrombus formation as a result of rapid endothelial injury, and the associated thrombotic formation. FeCl_3 application is a valuable model for investigation into thrombosis and atherosclerosis. However, caution should be applied since iron interacts with various proteins from the coagulation cascade and its effects depend on storage of the stock solution (42).

In conclusion, trivalent iron is involved in coagulation in a complex manner. It extends the clotting of plasma by interacting with proteins of the coagulation cascade. Fe^{3+} and/or its hydrolytic species interact with fibrinogen and/or fibrin, changing their morphology and properties. Moreover, when stored, FeCl_3 produces derivatives that potentiate changes in plasma clotting, some of which can be attributed to free radicals formed during FeCl_3 storage. In general FeCl_3 is able to weaken the fibrin clot while precipitating plasma proteins immediately after application. This property can be exploited therapeutically in stanching the flow of blood from wounds when optimum concentrations of FeCl_3 are found.

Acknowledgements

This study was supported in part by grants from the Frank Stranahan Endowed Chair and Children Miracle Network.

References

- Chang TP and Rangan C: Iron poisoning: a literature-based review of epidemiology, diagnosis, and management. *Pediatr Emerg Care* 27: 978-985, 2011.
- Mills KC and Curry SC: Acute iron poisoning. *Emerg Med Clin North Am* 12: 397-413, 1994.
- Elg M, Gustafsson D and Carlsson S: Antithrombotic effects and bleeding time of thrombin inhibitors and warfarin in the rat. *Thromb Res* 94: 187-197, 1999.
- Augustus F, Barnard P and Guyot A: *New Universal Cyclopedia: A Scientific and Popular Treasury of Useful Knowledge*. Vol. 2, 1st edition, A. J. Johnson & Co., New York, p82, 1881.
- Eckly A, Hechler B, Freund M, *et al*: Mechanisms underlying FeCl_3 -induced arterial thrombosis. *J Thromb Haemost*: 779-789, 2011.
- Woollard KJ, Sturgeon S, Chin-Dusting JP, Salem HH and Jackson SP: Erythrocyte hemolysis and hemoglobin oxidation promote ferric chloride-induced vascular injury. *J Biol Chem* 284: 13110-13118, 2009.
- O'Mara NB: Anemia in patients with chronic kidney disease. *Diab Spectr* 21: 12-19, 2008.
- Undas A, Kolarz M, Kopec G and Tracz W: Altered fibrin clot properties in patients on long-term haemodialysis: relation to cardiovascular mortality. *Nephrol Dial Transplant* 23: 2010-2015, 2008.
- Esposito BP, Breuer W, Slotki I and Cabantchik ZI: Labile iron in parenteral iron formulations and its potential for generating plasma nontransferrin-bound iron in dialysis patients. *Eur J Clin Invest* 32 (Suppl 1): 42-49, 2002.
- Lipinski B and Pretorius E: Novel pathway of iron-induced blood coagulation: implications for diabetes mellitus and its complications. *Pol Arch Med Wewn* 122: 115-122, 2012.
- Chlorek zelaza(III). Wikipedia, 2013. http://pl.wikipedia.org/wiki/Chlorek_%C5%BCelaza%28III%29. Wikipedia, 2013 (In Polish).
- Feng W and Nansheng D: Photochemistry of hydrolytic iron (III) species and photoinduced degradation of organic compounds. A minireview. *Chemosphere* 48: 1137-1147, 2000.
- Evans PA, Hawkins K, Lawrence M, Barrow MS, Williams PR and Williams RL: Studies of whole blood coagulation by oscillatory shear, thromboelastography and free oscillation rheometry. *Clin Hemorheol Microcirc* 38: 267-277, 2008.
- Gallimore MJ, Harris SL, Tappenden KA, Winter M and Jones DW: Urokinase induced fibrinolysis in thromboelastography: a model for studying fibrinolysis and coagulation in whole blood. *J Thromb Haemost* 3: 2506-2513, 2005.
- Carr ME Jr, Krishnamurti C and Alving BM: Effect of plasminogen activator inhibitor-1 on tissue-type plasminogen activator-induced fibrinolysis. *Thromb Haemost* 67: 106-110, 1992.
- Jankun J, Aleem AM, Selman SH, *et al*: Highly stable plasminogen activator inhibitor type one (VLHL PAI-1) protects fibrin clots from tissue plasminogen activator-mediated fibrinolysis. *Int J Mol Med* 20: 683-687, 2007.
- Jankun J, Aleem AM, Selman SH, Basrur V and Skrzypczak-Jankun E: VLHL plasminogen activator inhibitor spontaneously reactivates from the latent to active form. *Int J Mol Med* 23: 57-63, 2009.
- Jankun J, Keck R, Selman SH and Skrzypczak-Jankun E: Systemic or topical application of plasminogen activator inhibitor with extended half-life (VLHL PAI-1) reduces bleeding time and total blood loss. *Int J Mol Med* 26: 501-504, 2010.
- Jankun J, Selman SH, Keck RW, Lysiak-Szydłowska W and Skrzypczak-Jankun E: Very long half-life plasminogen activator inhibitor type 1 reduces bleeding in a mouse model. *BJU Int* 105: 1469-1476, 2010.
- Jankun J, Skotnicka M, Lysiak-Szydłowska W, Al-Senaidy A and Skrzypczak-Jankun E: Diverse inhibition of plasminogen activator inhibitor type 1 by theaflavins of black tea. *Int J Mol Med* 27: 525-529, 2011.
- Jankun J, Skrzypczak-Jankun E and Lipinski B: Complex function of magnesium in blood clot formation and lysis. *Cent Eur J Immunol* 38: 149-153, 2013.
- Nelsestuen GL, Broderius M and Martin G: Role of gamma-carboxyglutamic acid. Cation specificity of prothrombin and factor X-phospholipid binding. *J Biol Chem* 251: 6886-6893, 1976.
- Soriano-Garcia M, Padmanabhan K, de Vos AM and Tulinsky A: The Ca^{2+} ion and membrane binding structure of the Gla domain of Ca-prothrombin fragment 1. *Biochemistry* 31: 2554-2566, 1992.
- Urano T, Ihara H, Suzuki Y, Takada Y and Takada A: Coagulation-associated enhancement of fibrinolytic activity via a neutralization of PAI-1 activity. *Semin Thromb Hemost* 26: 39-42, 2000.
- Urano T, Ihara H, Takada Y, Nagai N and Takada A: The inhibition of human factor Xa by plasminogen activator inhibitor type 1 in the presence of calcium ion, and its enhancement by heparin and vitronectin. *Biochim Biophys Acta* 1298: 199-208, 1996.
- Urano T, Nagai N, Matsuura M, Ihara H, Takada Y and Takada A: Human thrombin and calcium bound factor Xa significantly shorten tPA-induced fibrin clot lysis time via neutralization of plasminogen activator inhibitor type 1 activity. *Thromb Haemost* 80: 161-166, 1998.
- Abou-Shady EA, Farrag HE, el-Damarawy NA, Mohamed FA, Kamel AM and Massoud AA: In vitro effects of trace elements on blood clotting and platelet function. A--Iron, copper, and gold. *J Egypt Public Health Assoc* 66: 21-48, 1991.
- Rosenmund A, Haerberli A and Straub PW: Blood coagulation and acute iron toxicity. Reversible iron-induced inactivation of serine proteases in vitro. *J Lab Clin Med* 103: 524-533, 1984.
- Henriksson P, Nilsson L, Nilsson IM and Stenberg P: Fatal iron intoxication with multiple coagulation defects and degradation of factor VIII and factor XIII. *Scand J Haematol* 22: 235-240, 1979.
- Pretorius E and Lipinski B: Differences in morphology of fibrin clots induced with thrombin and ferric ions and its pathophysiological consequences. *Heart Lung Circ* 22: 447-449, 2013.
- Orino K: Functional binding analysis of human fibrinogen as an iron- and heme-binding protein. *Biomaterials* 26: 789-794, 2013.
- Mishina M, Komaba Y, Kobayashi S, *et al*: Administration of free radical scavenger edaravone associated with higher frequency of hemorrhagic transformation in patients with cardiogenic embolism. *Neurol Med Chir (Tokyo)* 48: 292-297, 2008.

33. Lipinski B, Pretorius E, Oberholzer HM and Van Der Spuy WJ: Iron enhances generation of fibrin fibers in human blood: implications for pathogenesis of stroke. *Microsc Res Tech* 75: 1185-1190, 2012.
34. Dempfle CE, Argiriou S, Alesci S, *et al*: Fibrin formation and proteolysis during anocrod treatment. Evidence for des-A-profibrin formation and thrombin independent factor XIII activity. *Ann NY Acad Sci* 936: 210-214, 2001.
35. Drinane MC, Sherman JA, Hall AE, Simons M and Mulligan-Kehoe MJ: Plasminogen and plasmin activity in patients with coronary artery disease. *J Thromb Haemost* 4: 1288-1295, 2006.
36. Talbot K, Meixner SC and Pryzdial EL: Enhanced fibrinolysis by proteolysed coagulation factor Xa. *Biochim Biophys Acta* 1804: 723-730, 2010.
37. Xu X, Zhang L, Shen D, Wu H and Liu Q: Oxygen-dependent oxidation of Fe(II) to Fe(III) and interaction of Fe(III) with bovine serum albumin, leading to a hysteretic effect on the fluorescence of bovine serum albumin. *J Fluoresc* 18: 193-201, 2008.
38. Davies MJ, Gilbert BC and Haywood RM: Radical-induced damage to bovine serum albumin: role of the cysteine residue. *Free Radic Res Commun* 18: 353-367, 1993.
39. Zardeneta G, Milam SB and Schmitz JP: Iron-dependent generation of free radicals: plausible mechanisms in the progressive deterioration of the temporomandibular joint. *J Oral Maxillofac Surg* 58: 302-308, 2000.
40. Crichton RR: Proteins of iron storage and transport. *Adv Protein Chem* 40: 281-363, 1990.
41. Crichton RR and Charlotiaux-Wauters M: Iron transport and storage. *Eur J Biochem* 164: 485-506, 1987.
42. Tseng MT, Dozier A, Haribabu B and Graham UM: Transendothelial migration of ferric ion in FeCl₃ injured murine common carotid artery. *Thromb Res* 118: 275-280, 2006.