

The copy number and integration site analysis of IGF-1 transgenic goat

JIAN LIN, QIANG ZHANG, LI Q. ZHU, QING H. YU and QIAN YANG

Key Laboratory of Animal Physiology and Biochemistry, Ministry of Agriculture,
Nanjing Agricultural University, Nanjing, Jiangsu 210095, P.R. China

Received February 2, 2014; Accepted June 12, 2014

DOI: 10.3892/ijmm.2014.1841

Abstract. Transgenic animals have been used previously to study gene function, produce important proteins, and generate models for the study of human diseases. As the number of transgenic species increases, reliable detection and molecular characterization of integration sites and copy number are crucial for confirming transgene expression and genetic stability, as well as for safety evaluation and to meet commercial demands. In this study, we generated four transgenic goats by somatic cell nuclear transfer (SCNT). After birth, the cloned goat contained transferred insulin-like growth factor I (*IGF-1*) gene was initially confirmed using a polymerase chain reaction (PCR)-based method. The four cloned goats were identified as IGF-1 transgenic goats by southern blotting. The number of copies of the *IGF-1* gene in each of the transgenic goats was determined. Additionally, four integration sites of the transgene in the transgenic goats with a modified thermal asymmetric interlaced (TAIL)-PCR method were identified. The four different integration sites were located on chromosomes 2, 11, 16 and 18. The present study identified the copy number and integration sites using quantitative PCR (qPCR) and TAIL-PCR, enabling the bio-safety evaluation of the transgenic goats.

Introduction

Transgenic animals are a powerful tool used in the study of the function and regulation of genes *in vivo*, the production of important pharmaceutical proteins, and the creation of pathologic models for human disease therapy (1-3). When new transgenic animals are generated, one essential step is to identify the transgenic livestock. Novel approaches to improve the molecular characterization of transgenic livestock would have considerable

economic and commercial benefits. Commonly used transgenic techniques such as somatic cell nuclear transfer always result in the random integration of multiple copies of target genes in the host genome (4,5). Thus detecting the existence and expression of target genes in transgenic animals, as well as determining the copy number and insertion site in transgenic animals is crucial, as these factors can greatly affect the expression level and genetic stability of targeted genes (6-8). The random insertion of multiple copies may have marked effects, such as the inactivation of an endogenous gene following transgene insertion. Different levels of transgene expression and even silencing of the transgene may result when inserted into a heterochromatic region due to chromosome position effects (9,10).

Based on the difference between the transgenic and non-transgenic animal at the DNA level (target gene-, promoter-, marker gene- and construct-specific), a number of DNA-based methods have been utilized for transgene detection, such as polymerase chain reaction (PCR), quantitative PCR (qPCR) (11,12) and thermal asymmetric interlaced (TAIL)-PCR (13-15). The PCR-based detection strategies have become the core method due to their high specificity, efficiency and sensitivity (13). The number of transgene copies has traditionally been estimated by southern blotting techniques, which are tedious and time-consuming methods and require a large amount of DNA sample for each assay. Moreover, quantification using those methods is not accurate (when multiple copies of transgenes insert into one or more loci) and yield ambiguous results (16). The emergence of the sensitivity and accuracy of qPCR technology allows for the copy numbers in transgene animals can easily be determined (7,11). qPCR technology has been applied to analyze the copy number of transgenic mice (17-19), swine (7) and livestock (5).

Chromosome walking and flanking sequence cloning procedures were employed to detect the integration site of the target gene. The potential of inserting mutation renders the identification of the transgene location critical (20). Currently, a number of PCR-based methods are available for these purposes, including inverse PCR (I-PCR) (21), ligation-mediated PCR (22) and event-specific PCR (23). However, each of these methods has drawbacks when considered for wide use in transgenic animal analysis (24). Recently, improved methodologies have been developed to address these issues. The TAIL-PCR is the most successful method currently used to precisely identify transgene flanking sequences. Additionally,

Correspondence to: Professor Qian Yang, Key Laboratory of Animal Physiology and Biochemistry, Ministry of Agriculture, Nanjing Agricultural University, 1 Wei gang, Nanjing, Jiangsu 210095, P.R. China
E-mail: zxyq@njau.edu.cn

Key words: transgenic goat, copy number, quantitative polymerase chain reaction, integration site, thermal asymmetric interlaced-PCR

TAIL-PCR has been widely used in identifying the insert sites of transgenic animals (20). To obtain precise flanking fragments rapidly, we have developed a novel efficient method that combined the restriction method from southern blotting with TAIL-PCR procedures to analyze the integration sites.

In this study, we generated the transgenic-cloned goat that specifically expresses insulin-like growth factor I (IGF-1) in milk (25). To stimulate the molecular analysis of cloned goats, we initially used PCR and southern blotting to identify the cloned goats as IGF-1 transgenic goats. The exact integration site and copy numbers of the four individual IGF-1 transgenic goats were detected by qPCR and TAIL-PCR methods.

Materials and methods

Production of IGF-1 transgenic goats. Transgenic goats were generated by somatic cell nuclear transfer (SCNT) with *SalI/PvuI*-linearized plasmid pIN, which contained 6.1 kb of the goat β -casein proximal promoter region, the coding region of goat *IGF-1* gene and 2.3 kb of the 3' β -casein region (26). Briefly, *SalI/PvuI*-linearized plasmid was transferred into fibroblasts by Lipofectame 2000 and screened with neomycin. Healthy transferred pIN donor cells (fibroblasts) were then cultured in starved medium and injected into the perivitelline space of enucleated oocytes with a beveled micropipette. In addition, the activated reconstructed couplets were cultured in the oviducts of synchronized temporary intermediate recipients or transferred into the uterine tubes of each recipient. After 5 days of culture *in vivo* of the reconstructed embryo, the developed morulae and blastocytes were surgically transferred into the uteri of synchronized final recipients. At day 35, the surrogates were scanned with B-ultrasound scanner to detect pregnant goats. Four cloned goats were eventually born.

DNA extraction and genomic PCR identification

DNA extraction. Whole goat blood genomic DNA was extracted from the four IGF-1 transgenic goats and three non-transgenic goats with the TIANamp Blood DNA Midi kit [Tiangen Biotech (Beijing) Co., Ltd, Beijing, China] according to the manufacturer's instructions. The extracted DNA was kept at -20°C until use. This animal study was approved by the Institutional Animal Care and Use Committee of Nanjing Agricultural University. The extracted genomic DNA was quantified by electrophoresis. Gels were prepared with 1% agarose in TAE buffer with ethidium bromide (EtBr).

Primer design and PCR identification. Three pairs of specific primers were designed to evaluate the presence of the transgene. Primers were designed according to Fig. 1A and are listed in Table I. PCR amplification was performed with the genomic DNA template, recombinant Taq polymerase, 10X buffer, and primers. The conditions for PCR were: 94°C for 5 min, 94°C for 30 sec, 60°C for 30 sec, and 72°C for 90 sec for 30 cycles, with a final extension at 72°C for 10 min. The PCR product was isolated and linked to plasmid pMD19-T for sequencing.

Southern blot analysis. Southern blotting hybridization was carried out with the DIG system (F. Hoffmann-La Roche AG, Basel, Switzerland) according to the manufacturer's instructions. Briefly, 30 μ g genomic DNA from four cloned goats

Table I. qPCR primers designed for detecting copy number.

Name	Sequence (5'-3')	Size (bp)
1F	ACATCCTCCTCGCATCTCTTC	804
1R	CCTTCTTAGGTTTGTTATTCTTAGCC	804
2F	CATTGTTTGATCATATGCACCTC'	1441
2R	CCTTCTTAGGTTTGTTATTCTTAGCC	1441
3F	CTCTGGTTCCTCTGCCTTTTTC	1184
3R	ATCTCCTGTTCATCTCACCTTGC	1184

PCR, polymerase chain reaction.

and three control goats were digested with *HindIII/EcoRI* and separated on a 1.0% agarose gel. Plasmid pIN was also digested as a positive control. The gels were blotted onto nylon-N⁺ membrane (Whatman plc., Maidstone, England) overnight in 20X SSC buffer. As a probe, 804 bp amplified fragments were labeled with DIG-11-dUTP using the PCR DIG Probe Synthesis kit and hybridized to the membrane at 45°C overnight in DIG Easy Hyb solution (both from F. Hoffmann-La Roche AG). Detection was performed with the DIG Wash and Block Buffer Set (F. Hoffmann-La Roche AG). The DIG-labeled probe was detected with anti-digoxigenin AP Fab fragment and visualized with nitroblue tetrazolium chloride and 5-bromo-4-chloro-3-indolyl-phosphate (NBT/BCIP) solution (both from F. Hoffmann-La Roche AG).

Detection of qPCR. To examine the copy numbers of the *IGF-1* gene in transgenic goat, it was essential to construct a standard curve (7). First, a series of standard samples containing 1, 4, 16, 64, 256 and 1,024 copies of the *IGF-1* gene, respectively, were prepared by mixing the wild-type genome of a Saanen dairy goat with plasmid pIN. A standard curve was drawn by plotting Δ Ct (Δ Ct = Ct_{IGF-1} - Ct_{GAPDH}) against the log of *IGF-1* gene copies of corresponding standard samples. qPCR was performed with SYBR Premix Ex Taq (Takara Bio Inc., Otsu, Japan) on a 7500 real-time PCR System (Applied Biosystems, Foster City, CA, USA) as follows: 95°C for 10 sec, followed by 40 two-step cycles at 95°C for 5 sec and at 60°C for 34 sec. Primers for the *IGF-1* and *GAPDH* genes (*GAPDH* was amplified concurrently as an endogenous control) are shown in Table II. A total volume of PCR (20 μ l), containing 2.0 μ l 10X PCR buffer, 0.5X SYBR-Green I deoxyribonucleoside triphosphates, 0.4 mmol/l dNTP, 1 unit Taq DNA polymerase, 2.0 μ l primers and 6 μ l 20X diluted cDNA as the template was also utilized. The Ct value was calculated by the Sequence Detection System software (Applied Biosystem). The amount of target normalized to reference was calculated by $2^{-\Delta\Delta$ Ct in qPCR. For each DNA sample (four IGF-1 transgenic goats and two non-transgenic goats), both the target and reference genes were amplified independently on the same plate and in the same experimental run in triplicate. The values are presented as mean \pm SEM. The Ct value was calculated by the Sequence Detection System software (Applied Biosystems).

TAIL-PCR analysis of the integration site. To analyze the integration sites, TAIL-PCR was performed employing three

Table II. qPCR primers designed for detecting copy numbers.

Gene	Sense	Antisense
<i>IGF-1</i>	ATGCCAGTCACATCCTCCTC	CTCCAGCCTCCTCAGATCAC
<i>GAPDH</i>	GGGTGTTGTTATACTTCTCGTGGTT	GTGATGCTGGTGGCTGAGTATGTG

PCR, polymerase chain reaction; IGF-1, insulin-like growth factor I.




Figure 1. The production of IGF-1 transgenic goats and its identification. (A) The schematic diagram of vector pIN and primer design map (β -casein 5, β -casein promoter genes of goat; IGF-1, insulin-like growth factor I gene of goat; β -casein 3, β -casein genomic fragment; neo, neomycin resistance gene). (B) IGF-1 transgenic goats (four healthy kids were born and survived to maturity). (C) The identification of IGF-1 from IGF-1 to 3' β -casein; (C1, C2 and C3, control goats; T1, T2, T3 and T4, IGF-1 transgenic goats; +, mammary gland-specific expression vector pIN; -, negative; M, Marker DNA/Trans2K Plus). (D) The identification of IGF-1 from 3' β -casein to 5' β -casein; (E) the identification of neo from 3' β -casein to neo.

transgene-specific primers and five arbitrary degenerate primers (AD1, AD2, AD3, AD4 and AW). Fig. 2A shows the schematic relationship between the specific primers and arbitrary degenerate primers with the target genomic sequence or flanking sequence of IGF-1 transgenic insertion. All the primers used in this study were purchased from Invitrogen Co., Ltd. (Shanghai, China) and are listed in Table III. The TAIL-PCR process included three sequential PCR reactions (Table IV). TAIL-PCR protocol was performed according to the method described by Liu *et al* (27). Briefly, the primary PCR reaction contained recovered *HindIII/EcoRI*-digested template (two types of template were used; one was a fragment <4 kb, while the other was a fragment >4 kb), 2.5 mM each dNTPs, 0.6 μ M of SP1 primer, 2 μ m of the AD primer, and 1 unit of Taq polymerase in 20 μ l reaction buffer. In the secondary or tertiary round, 1 μ l of 10-fold diluted first or secondary products were used as templates and supplemented in the reaction. The thermal cycling conditions are shown in Table V. The

products of tertiary TAIL-PCR reaction were separated on a 1.0% agarose gel. Specific bands for each set were purified using a gel purification kit (Omega Bio-Tek, Inc., Norcross, GA, USA) and sequenced. Sequencing results were analyzed using the BLAST and NCBI databases for bovine genomic DNA to identify the specific chromosomal integration sites.

Results

Generation of transgenic goats. In total, 50 donor goats were used in this study, with 388 *in vivo* matured oocytes being recovered. In addition, 375 oocytes were enucleated and produced karyoplast-cytoplasmic couplets using prepared donor cells. Subsequent to fusion, a total of 221 reconstructed embryos were obtained. Reconstructed pronuclear-stage embryos were transferred into 46 synchronized recipients, which produced 23 (50%) pregnancies at day 35. Four healthy kids were born and survived to maturity (Fig. 1B).

Table III. Primers designed for TAIL-PCR.

Primer	Sequence
AD1	TGCACCACTGGACTGAG CGGCCG CVNVNNGGAA
AD2	TGCACCACTGGACTGAG CGGCCG CBNBNNNGGTT
AD3	TGCACCACTGGACTGAG CGGCCG CVNVNNNCCAA
AD4	TGCACCACTGGACTGAG CGGCCG CVBNBNNCCGGT
AW	TGCACCACTGGACTGA
Special primers	
<i>Sal</i> II3	GAGAAGCGTTCAGAGGAAAGCGATC
<i>Sal</i> II2	CTCAAAGAGCAGCGAGAAGCGTTC
<i>Sal</i> II1	CAGGCCGTTCTATGATTCTGTCATTC
<i>Pvu</i> I3	CGCCGCATACACTATTCTCAGAAATG
<i>Pvu</i> I2	CAACTCGGTTCGCCGCATACACTATT
<i>Pvu</i> I1	CTTTTAAAGTTCTGCTATGTGGCGCG

TAIL-PCR, thermal asymmetric interlaced polymerase chain reaction. Bold letters indicate the core sequences in the AD primers.

Table IV. The reaction system of TAIL-PCR.

Reaction composition	First round	Second round	Third round
10X ExTaq buffer	2	2	5
2.5 Mmol/l dNTP	1	1	2
10 μ mol/l AD	2	1	2
10 μ mol/l SP	0.6	1	2
Ex Taq/(5 U/ μ l)	0.2	0.2	0.5
ddH ₂ O	Add to 20 μ l	Add to 25 μ l	Add to 50 μ l
Template	Genomic DNA digested with <i>Hind</i> III/ <i>Eco</i> RI	1 μ l of 10-fold diluted the first products	1 μ l of 10-fold diluted the secondary products

TAIL-PCR, thermal asymmetric interlaced polymerase chain reaction.

PCR detecting of transgenic goats. When four cloned goats were previously generated by SCNT, the first step was to identify whether these cloned goats were IGF-1 transgenic goat. PCR was carried out with three pairs of primers to analyze the cloned goats. Results showed that the four goats that survived to maturity were IGF-1 transgenic. Fig. 1C and D show that the *IGF-1* gene was integrated into the genome of four cloned goats, while there was no target band in the non-transgenic control goats. Sequencing results showed that the inserted IGF-1 was consistent with the plasmid pIN, without any mutations. Fig. 1E shows that four clone goats contained the *IGF-1* gene in their genome, and included the resistance gene (neo).

Southern blotting identification of transgenic goats. After PCR detection, we verified that plasmid pIN was integrated into the genomic DNA of transgenic goats using southern blotting. Southern blotting, a highly accurate and sensitive technology, has been widely used in the identification of genetically modified products, especially when the transferred genes were highly homologous with endogenous genomic DNA. For detection of southern blotting, a probe was designed and amplified (Fig. 2A).

Fig. 3A shows a clear band on the NC member only in the four IGF-1 transgenic goat lanes and positive plasmid pIN lane. This result, consistent with the PCR results, proved that the four clone goats were IGF-1 transgenic goats.

Copy number analysis of the IGF-1 gene in transgenic goats. Since the copy number of transgenes may greatly affect the expression of the target genes, we determined the copy numbers by qPCR. The absolute quantitative standard curve was drawn by plotting Δ Ct (Δ Ct = Ct_{IGF-1} - Ct_{GAPDH}) against the log of *IGF-1* gene copies of corresponding standard samples (Table VI). The standard curve was calculated as: \log_2 N (copy number) = -1.0244 Δ Ct + 5.3576 ($R^2=0.9963$) (Fig. 3C). Following construction of the standard curve, we sequentially detected the copy numbers of transgenic goats with qPCR. The reported Ct values were averaged for the triplicates (Table VII). Using the equation, the number of IGF-1 transgene copies of the four transgenic goats were estimated (Table VI). Results determined that four transgenic goats contained approximately the same number of copies of the *IGF-1* gene (7.89-9 copies), while no copies were identified for the non-transgenic goats.

Table V. The thermal cycling conditions of TAIL-PCR.

Reaction steps	First reaction		Second reaction		Third reaction	
	Temperature	time	Temperature	time	Temperature	time
1	94	1 min	94	1 min	94	1 min
2	98	1 min	94	30 sec	94	30 sec
3	94	30 sec	64	1 min	64	1 min
4	68	1 min	72	2 min	72	2 min
5	72	2 min	94	30 sec	94	30 sec
6	Go to step 3	5 times	64	1 min	64	1 min
7	94	30 sec	72	2min	72	2 min
8	25	3 min	94	30 sec	94	30 sec
9	72	2 min	44	1 min	44	1 min
10	94	30 sec	72	2 min	72	2 min
11	64	1 min	Go to step 2	15 times	Go to step 2	15 times
12	72	2 min	72	10 min	72	10 min
13	94	30 sec				
14	64	1 min				
15	72	2 min				
16	94	30 sec				
17	44	1 min				
18	72	2 min				
19	Go to step 10	15 times				
20	72	10 min				

TAIL-PCR, thermal asymmetric interlaced polymerase chain reaction.

Table VI. Standard curve conversion.

Copy no.	Ct (IGF-1)	Ct (GAPDH)	$\Delta C(t)$
1	25.51	20.07	5.45
4	23.38	20.09	3.28
16	21.34	20.26	1.08
64	19.4	20.3	-0.9
256	17.81	20.33	-2.53
1,024	16.19	20.48	-4.29

IGF-1, insulin-like growth factor I.

Integration sites analysis of IGF-1 gene in transgenic goats.

Since the transgene integration site and the resulting perturbation may greatly affect the expression of the inserted gene and its neighbors, we performed TAIL-PCR with three transgene specific primers and five arbitrary degenerate primers to clone the flanking sequence. For T1 transgenic goat, two specific fragments of ~1,500 and 2,000 bp in size were amplified when using the combination of arbitrary primer AD2 and special primer *SalI3* at the third TAIL-PCR, while TAIL-PCR amplified three specific fragments of ~500, 1,200 and 2,000 bp with primer AD2 and special primer *PvuI3* in T1 transgenic

goat (Fig. 2B). No such specific bands were amplified in the parallel negative control experiments using non-transgenic goat genomic DNA as a template. For the T2 transgenic goat, a unique fragment of ~500 bp was amplified with the arbitrary primer AD2 and special primer *SalI3* and three special fragments of ~500, 1,500 and 2,000 bp were amplified with primer AD2 and special primer *PvuI3* at the third TAIL-PCR (Fig. 2B). These amplified DNA fragments were cloned into plasmid pMD19-T and sequenced. The sequenced DNA fragments were analyzed by Bio-edit. Results show that the cloned bands from the *SalI* side contained a 69 bp sequence of exogenous plasmid pIN *SalI* border region running into a goat genomic DNA sequence. The amplified bands from the *PvuI* side were consistent with the 160-bp sequence of exogenous plasmid pIN *PvuI* border region in contiguity with the goat genomic DNA sequence. These results suggested that the junction sequences spanning the transgenic integration sites were correctly revealed.

The BLAST results showed that two distinct sites of integration were identified on the *SalI* border region, one each on chromosomes 18 (GenBank ID: NM-3104464.1; range, 823,389-823,645, 92%) and 16 (GenBank ID: NM-003104439.1; range, 9,355,733-9,355,900, 88%) (Fig. 4). For the *PvuI* border region, we identified another two distinct integration sites mapped to chromosome 11 (GenBank ID: NM-001492908.3; range, 4,167,112-4,167,267, 85%) and 2 (GenBank ID: NM-003103851.1; range, 11,847,579-11,847,774, 91%) (Fig. 5).

Table VII. Copy numbers of IGF-1 in transgenic goats.

Goat names	Ct (IGF-1)	Ct (GAPDH)	$\Delta C(t)$	log ₂ N (copy no.)	Copy no.
Wild-type 1	33.62	21.2	12.42	-7.36	
Wild-type 2	32.99	21.65	11.34	-6.25	
IGF-1-1	23.78	21.64	2.14	3.17	9.00
IGF-1-2	23.64	21.44	2.2	3.1	8.57
IGF-1-3	24.46	22.17	2.29	3.01	8.06
IGF-1-4	24.12	21.8	2.32	2.98	7.89

IGF-1, insulin-like growth factor I.




Figure 2. The design of thermal asymmetric-interlaced polymerase chain reaction (TAIL-PCR) primers and the electrophoresis of TAIL-PCR products. (A) The design map of TAIL-PCR primers [The schematic relationship between the specific primers and arbitrary degenerate primers with the target genomic sequence or flanking sequence of insulin-like growth factor I (IGF-1) transgenic insertion]. (B) The electrophoresis results of TAIL-PCR in *SalI* (C1, control goats; T1 and T2, IGF-1 transgenic goats; M, DL2000 marker); (C) The electrophoresis results of TAIL-PCR in *PvuI*; (C1, control goats; T1 and T2, IGF-1 transgenic goats; M, DL2000 marker).





Figure 3. Southern blotting identification result of the cloned goat (genomic DNA was digested with *HindIII/EcoRI*, plasmid pIN was digested as positive control. A total of 804 bp fragments from IGF-1 to 3' β -casein was amplified as a probe). (A) Southern blotting identification result of the cloned goat (lanes C1, C2 and C3, control goats; lanes T1, T2, T3 and T4, IGF-1 transgenic goats; lane +, positive control (mammary gland-specific expression vector pIN); M, DL1000 bp marker). (B) The PCR products of absolute quantitative primers (lanes C1 and C2, control goats; lanes T1 and T2, IGF-1 transgenic goats; lane +, positive control (mammary gland-specific expression vector pIN); M: DL2000 bp marker). (C) Establishment of the absolute quantitative standard curve. The standard samples containing 1, 4, 16, 64, 256 and 1,024 copies of the IGF-1 gene were prepared. The absolute quantitative standard curve was drawn by plotting ΔC_t ($\Delta C_t = C_{tIGF-1} - C_{tGAPDH}$) against the log of IGF-1 gene copies of corresponding standard samples.

Discussion

The first transgenic livestock was generated almost 30 years ago (28). Currently, a common method for producing

transgenic livestock involves using genetically modified cells in SCNT. Compared with conventional transgenic technology, SCNT technology allows the evaluation of the transferred gene *in vitro*, which may greatly improve the safety of genetically



B Alignments

Bos taurus breed Hereford chromosome 18 genomic scaffold, Bos_taurus_UMD_3.1, whole genome shotgun sequence
Sequence ID: ref|NW_003104464.1| Length: 31414493 Number of Matches: 1
Range 1: 823389 to 823645

Score	Expect	Identities	Gaps	Strand	Frame
372 bits(412)	2e-100()	237/257(92%)	3/257(1%)	Plus/Minus	

Features:
159588 bp at 5' side: uncharacterized protein LOC100847370106858 bp at 3' side: forkhead box protein F1

Query	66	CTGTCATTGAGCCCAATTTTCATCTCAACGTCACATGGTCTCACTGTCACATCCCAG	125
Sbjct	823645	CTGTCATTGAGCCCAATTTTCATCTCAACGTCACATGGTCTCACTGTCACATCCCAG	823586
Query	126	CCAGCCTGCTTGACCATAGCTCTCTGCCCTTGGGGGAGGCCCTGGGCAGAGTCACTT	185
Sbjct	823585	CCAGCCTGCTTGACCATAGCTCTCTGCCCTTGGGGGAGGCCCTGGGCAGAGTCACTT	823526
Query	186	AGCAACTGTGGAAATCCACCTGAGCGAATCAGTAGCTTCCITACCCAGGCAAGCCTTATG	245
Sbjct	823525	AGCAACTGTGGAAATCCACCTGAGCGAATCAGTAGCTTCCITACCCAGGCAAGCCTTATG	823466
Query	246	CTTCTCTGTGCTCCGCTCCCGCTGCTGGGGGAGCCGGTAGACG--GGGATGCTG	302
Sbjct	823465	CTTCTCTGTGCTCCGCTCCCGCTGCTGGGGGAGCCAGTAGATGGACGGGATGCTG	823406
Query	303	AGCCCGGTGCCTGCCCC	319
Sbjct	823405	AGCCCGGTGCCTGCCCC	823389

C Alignments


Bos taurus breed Hereford chromosome 16 genomic scaffold, Bos_taurus_UMD_3.1, whole genome shotgun sequence
Sequence ID: ref|NW_003104439.1| Length: 16624581 Number of Matches: 5
Range 1: 9355733 to 9355900

Score	Expect	Identities	Gaps	Strand	Frame
214 bits(236)	1e-52()	148/168(88%)	0/168(0%)	Plus/Plus	

Features:
2381 bp at 5' side: PRAME family member 9/1522945 bp at 3' side: LOW QUALITY PROTEIN: PRAME family member 5-like

Query	413	GCAACCGCGCCTAAGTGTGGGTGGCTGCGACCAGCACACTAAGCAGGGTGGAGGAG	472
Sbjct	9355733	GCAACCGCGCCTAAGTGTGGGTGGCTGCGACCAGCACACTAAGCAGGGTGGAGGAG	9355792
Query	473	CTaaccacgtctcagggtcaggggcagaagctgggaggacctcATGCTGAAGGGTGGCG	532
Sbjct	9355793	CGACCCACGTCAGGTCAGGGGCGAAGGCCAGAGGACCTCATGCCAAGGGCGGCA	9355852
Query	533	gtcaagaggagttaccocatgtccagggtcaggggcagtgagctgggag	580
Sbjct	9355853	GCCAAGAGGAGTTACCCACGTCAGGTCAGGGGAGTGGCCAGAG	9355900

Figure 4. The BLAST results of thermal asymmetric-interlaced polymerase chain reaction (TAIL-PCR) in *SalI* side. (A) Sequence alignment results among primer *SalI*3, *SalI* border region and TAIL-PCR product with Bio-edit software. (B) The alignment results of flank sequence from *SalI* side. (C) The alignment results of the flank sequence from *SalI* side.



B Alignments


Bos taurus breed Hereford unplaced genomic scaffold, alternate assembly Btau_4.6.1 ChrUn.004-5.11787.scaffold1
 Sequence ID: ref|NW_001501683.1| Length: 1014 Number of Matches: 1
 Range 1: 656 to 962

Score	Expect	Identities	Gaps	Strand	Frame
219 bits(242)	3e-54()	237/309(77%)	5/309(1%)	Plus/Plus	
Features:					
Query 295	CCGCTCACAAATCCACACAACATACGAGCCGGAAGCATAAAGTGTAAAGCCTGGGGTGCC				354
Sbjct 656	CCGCCACAAAACACACAACAGACGAGCCGGAAGCACAAAAGCGCAAAGCCGGGGGGCC				715
Query 355	TAAAGTGTAGCTAACTCACATTAATTGCGTTGCGCTCACTGCCCGCTTCCAGTCGGG				414
Sbjct 716	CAAAGAGCGAGCNAACACACAAATGCGCAGCGCCAC-GGCCCGCCACCAGGGCGG				774
Query 415	AAACCTGTCGTGCCAGCTGCATTAATGAATCGGCCAACCGGC-GGGAGAGGC-GTTTTCG				472
Sbjct 775	AAACAGGCGCGCCAGCAGCAAAAAGAAACGCCAACCGCGGGGAGAGGCGCCCGCG				834
Query 473	TATTGGGCGTCTTCCGCTTCTCGCTCACTGACTCGCTGCGCTCGGTTCGGTTCG				532
Sbjct 835	AAAAGGGCGC-ACNACCGCACCCGCGCACACAGACACGACGCGCAGGACGNACGGCCGC				893
Query 533	GGCGAGCGGCACCAGCTCACTC-AAGGTGGTAATACCGTTATCCACAGAATCAGGGGATA				591
Sbjct 894	GGCGAGCGGGAACAGCCACCCAAAGGCGGGAANACGGCCANCCACAGAACCAGGGGAAA				953
Query 592	ACGCAGGAA 600				
Sbjct 954	ACGCAGGAA 962				

Bos taurus breed Hereford chromosome 11 genomic scaffold, alternate assembly Btau_4.6.1 Chr11.scaffold8
 Sequence ID: ref|NW_001492908.3| Length: 5730378 Number of Matches: 2
 Range 1: 4167112 to 4167267

Score	Expect	Identities	Gaps	Strand	Frame
176 bits(194)	3e-41()	134/157(85%)	1/157(0%)	Plus/Plus	
Features: 147431 bp at 5' side: 3-oxo-5-alpha-steroid 4-dehydrogenase 28024 bp at 3' side: protein MEMO1					
Query 272	TGTTTCTGTGTGAAATTGTTATCCGCTCACAAATCCACACAACATACGAGCCGGAAGCA				331
Sbjct 4167112	TGTTTCTGTGTGACATTGTGATCCACTCACAAATCCACACAACATACTATCCCGATTCA				4167171
Query 332	TAAAGTGTAAAGCCTGGGGTGCCTAATGAGTGTAGCTAACTCACATTAATTGCGTTGCGCT				391
Sbjct 4167172	TACAGTGTATATGCTGACCTGCCTAATGAGTGAATAACTCACATTAATTGAGATGCGCT				4167231
Query 392	CACTGCCCGCTTTCAGTCGGGAACCTGTCGTGCC 428				
Sbjct 4167232	CACTG-CCCGCTTCCCGTGGGAATCCTGTCGTGCC 4167267				

Figure 5. The blast results of thermal asymmetric-interlaced polymerase chain reaction (TAIL-PCR) in the *PvuI* side. (A) Sequence alignment results among primer *PvuI* 3, *PvuI* border region and TAIL-PCR product with Bio-ed software. (B) The alignment results of the flank sequence from the *PvuI* side.



D Alignments

Bos taurus breed Hereford unplaced genomic scaffold, alternate assembly Btau_4.6.1 ChrUn.004-5.11787.scaffold1
 Sequence ID: ref|NW_001501683.1| Length: 1014 Number of Matches: 1
 Range 1: 656 to 1014

Score	Expect	Identities	Gaps	Strand	Frame
345 bits(382)	4e-92()	291/359(81%)	0/359(0%)	Plus/Plus	
Features:					
Query 68	CCGCTCACAAITCCACACAACATACGAGCCGGAAGCATAAAGTGTAAAGCCTGGGGTGCC	127			
Sbjct 656	CCGCCACAAAACCCACACAACAGACGAGCCGGAAGCACAAGCGCAAAGCCGGGGGGCC	715			
Query 128	TAATGAGTGAGCTAACTCACATTAATTGCGTTGCGCTCACTGCCCGCTTCCAGTCGGGA	187			
Sbjct 716	CAAAGAGCGAGCNAACACAACAATAATGCGCAGCGCCACGCGCCGCCACAGGCGGGA	775			
Query 188	AACCTGTCGTGCCAGCTGCATTAATGAATCGGCCAACCGCGGGGAGAGCGGTTTGCCT	247			
Sbjct 776	AACCAGGCGCGCCAGCAGCAAAAAGAAACGGCCAACCGCGGGGAGAGCGGCCCGCG	835			
Query 248	ATTGGCGCTCTCCGCTTCTCGCTCACTGACTCGCTCGCTCGCTCGCTCGCTCGCTGCG	307			
Sbjct 836	AAAGGGCGCAGNACCGCACCCCGCGCACACAGACCGCAGCGCACGACGACGACGACG	895			
Query 308	CGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAAC	367			
Sbjct 896	CGAGCGGGAACAGCCACCCAAAGGCGGGAAACGGCCANCCACAGAACCGGGGAAAC	955			
Query 368	GCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCG	426			
Sbjct 956	GCAGGAAAGAACAGAGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGGAAAAAGGCCG	1014			

Bos taurus breed Hereford chromosome 2 genomic scaffold, Bos_taurus_UMD_3.1, whole genome shotgun sequence
 Sequence ID: ref|NW_003103851.1| Length: 12151094 Number of Matches: 2
 Range 1: 11847579 to 11847774

Score	Expect	Identities	Gaps	Strand	Frame
260 bits(288)	1e-66()	180/197(91%)	5/197(2%)	Plus/Plus	
Features: 54450 bp at 5' side: ephrin type-A receptor 2 precursor3028 bp at 3' side: protein FAM131C					
Query 455	CCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGTGG-CGAAACCCGACAGG	513			
Sbjct 11847579	CCCCCTCCGAGC-TCACAACAATCGACGCTCAAGTCAGAGTGGCGAAACCCGACCGG	11847637			
Query 514	ACTATAAAGATACCAAGCGTTTCCCCCTGGAAGCTCCCTCGTGCCTCTCCTGTTCCGAC	573			
Sbjct 11847638	ACTATAAAGATACCAAGCGTTTCCCCCTGGAAGCTCCCTCTTGCCTCTCCTGTATCGAC	11847697			
Query 574	CCTGCCGCTTA-CGGGATACCTGICCGCCTTCTCCC-ITCGGG-ARGCGTGGCGCTTTC	630			
Sbjct 11847698	CCTGCCGCTTACCGGATACCTCGTCTCCTTCTCCCTTTCGGGAAAGCGTGGCGCTTTC	11847757			
Query 631	TCATAGCTCACGCTGTA 647				
Sbjct 11847758	TCATAGCTCACGCTGTA 11847774				

Figure 5. Continued. (C) Sequence alignment results among primer *PvuI3*, *PvuI* border region and TAIL-PCR product with Bio-edit software. (D) The alignment results of the flank sequence from the *PvuI* side.

modified animal by assessing positive donor cells, and markedly reduce the probability of the transgene being silenced in offspring (29-31). In this study, we obtained four IGF-1 transgenic goats, which were developed to adolescence and expected to be bred in September 2013. The four goats have suitably developed in body and mammary gland, without any disease.

To evaluate the transgenically cloned goat for commercial use, the transgene copies and integration sites should be identified. The transgene in the present study was a 465 bp IGF-1 fragment in the pBC1 vector. PCR and southern blotting results demonstrated that all four goats were IGF-1 transgenic goats. It is known that the transgene may randomly integrate into any site of the genome, the integration site and copies may be influenced by the insert time, plasmid form and insert position (32). Both the number of copies are presently known to integrate and the genomic context of the transgene has been proven to influence the phenotype of the transgenic animal (7,33,34). The high copy number tandem integration was thought to lead to transgene silencing (35,36) and the high copy number may decrease with aging in transgenic animals. Ballester *et al* and Vaisman detected the copies of transgenic animal using qPCR methods (11,37). The study by Kong *et al* suggested that the transgene expression level is associated with the copy number in transgenic pigs (7). In this study, we aimed to identify the copy numbers in IGF-1 transgenic goats with qPCR. qPCR is considered a simple, rapid and accurate method to estimate the transgene copy number in transgenic animals. By using this method, we revealed that four transgenic goats contained almost the same copy numbers of the transferred *IGF-1* gene in goat genome (from 7.89 to 9 copies). For a similar copy number of the four transgenic goats, the same donor cell may be an appropriate explanation.

Following evaluation of the copy numbers of IGF-1 transgenic goats, we detected the integration sites of the transgenic goats, which primarily affect the transgene expression. The integration site has been proven to affect the level and time of transgene expression (38,39). In transgenic mouse, the same transferred gene would be transcribed in different time periods for their different insert location (40). However, results of recent studies have suggested that the integration site may not always be random (42). Previously it was found that in certain sites the transgene was inclined to integrate by random transfection, such as LINE elements (41). However, random transfection may not result in random integration but in some hot integration sites and these hot integration sites may have a common character allowing the transgene to integrate. Yan *et al* demonstrated that the foreign *fad2* behaved similar to an X-linked gene and that foreign DNA molecules were inserted into the eukaryotic genome through a homologous illegitimate random integration (42).

Considering the importance of identifying the integration site, several PCR-based methods have been applied for the precise determination of the integration site of foreign DNA into native chromosomes. These methods include I-PCR, interspersed repetitive sequence PCR (IRS-PCR), linear amplification-mediated PCR (LAM-PCR) and TAIL-PCR. Of these, TAIL-PCR has obvious advantages as it may be used to identify homozygous animals (14,15). TAIL-PCR has been widely used for genome walking and flanking sequence

cloning in transgenic animals. In TAIL-PCR, enrichment of the target products depends on the difference in amplification velocity between the target and non-target products (13,24). In this study, we improved the TAIL-PCR methods. The performance of TAIL-PCR was strongly dependent on the PCR efficiency and specificity. The improved methods entailed one more digestion reaction. When we performed the TAIL-PCR, we first digested the template goat genome DNA by *HindIII* and *EcoRI*, which was used in southern blotting identification. The digested genome was run on 0.8% agarose gel and then recovered; the fragments were >4 or <4 kb. TAIL-PCR with subsequently performed using the two types of templates as described in a previous section. The digested step was used to increase the specificity of target genes. Compared with conventional TAIL-PCR, our methods divided possible integration site into two sections, increasing the efficiency of TAIL-PCR, particularly for the template <4 kb. The results suggest that the improved TAIL-PCR was more efficient than conventional TAIL-PCR for cloning flanking sequences. Using these modified methods, we identified the integration sites NM-001492908.3, NM-003103851.1, NM-003104439.1, NM-3104464.1.

In conclusion, PCR-based techniques have been widely used for the precise transgene flanking sequence and copy number identification in molecular biology research. The present study has demonstrated the successful use of PCR and southern blotting to characterize four cloned goats as IGF-1 transgenic goats. The qPCR method clarified the copy numbers of the *IGF-1* gene in transgenic goats. Furthermore, using the TAIL-PCR approach, we identified four integration sites with high specificity and provided information on the chromosomal location. The future application of TAIL-PCR to characterize transgenic animals is likely to be extremely significant.

Acknowledgements

This study was supported by the National Animal Transgenic Breeding Grand Project of China (no. 2013ZX08008-004) and a project funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions.

References

1. Houdebine LM: The methods to generate transgenic animals and to control transgene expression. *J Biotechnol* 98: 145-160, 2002.
2. Maksimenko OG, Deykin AV, Khodarovich YM and Georgiev PG: Use of transgenic animals in biotechnology: prospects and problems. *Acta Naturae* 5: 33-46, 2013.
3. Ménoret S, Tesson L, Rémy S, *et al*: Technical advances in the generation of transgenic animals and in their applications. *Transgenic Res* 22: 1065-1072, 2013.
4. Niemann H, Kues W and Carnwath JW: Transgenic farm animals: present and future. *Rev Sci Tech* 24: 285-298, 2005.
5. Zhang R, Yin Y, Zhang Y, *et al*: Molecular characterization of transgene integration by next-generation sequencing in transgenic cattle. *PLoS One* 7: e50348, 2012.
6. Chandler KJ, Chandler RL, Broeckelmann EM, Hou Y, Southard-Smith EM and Mortlock DP: Relevance of BAC transgene copy number in mice: transgene copy number variation across multiple transgenic lines and correlations with transgene integrity and expression. *Mamm Genome* 18: 693-708, 2007.
7. Kong Q, Wu M, Huan Y, *et al*: Transgene expression is associated with copy number and cytomegalovirus promoter methylation in transgenic pigs. *PLoS One* 4: e6679, 2009.

8. Pena RN, Webster J, Kwan S, Korbel J and Whitelaw BA: Transgene methylation in mice reflects copy number but not expression level. *Mol Biotechnol* 26: 215-220, 2004.
9. Garrick D, Fiering S, Martin DI and Whitelaw E: Repeat-induced gene silencing in mammals. *Nat Genet* 18: 56-59, 1998.
10. Hogan B: Molecular biology. Enhancers, chromosome position effects, and transgenic mice. *Nature* 306: 313-314, 1983.
11. Ballester M, Castelló A, Ibáñez E, Sánchez A and Folch JM: Real-time quantitative PCR-based system for determining transgene copy number in transgenic animals. *Biotechniques* 37: 610-613, 2004.
12. Mitrečić D, Huzak M, Curlin M and Gajović S: An improved method for determination of gene copy numbers in transgenic mice by serial dilution curves obtained by real-time quantitative PCR assay. *J Biochem Biophys Methods* 64: 83-98, 2005.
13. Huang H, Wang G, Zhao Y, Shi P, Luo H and Yao B: Direct and efficient cloning of full-length genes from environmental DNA by RT-qPCR and modified TAIL-PCR. *Appl Microbiol Biotechnol* 87: 1141-1149, 2010.
14. Singer T and Burke E: High-throughput TAIL-PCR as a tool to identify DNA flanking insertions. *Methods Mol Biol* 236: 241-272, 2003.
15. Thanh T, Chi VT, Abdullah MP, Omar H and Napis S: Efficiency of ligation-mediated PCR and TAIL-PCR methods for isolation of RbcS promoter sequences from green microalgae *Ankistrodesmus convolutus*. *Mol Biol (Mosk)* 46: 64-70, 2012.
16. Mason G, Provero P, Vaira AM and Accotto GP: Estimating the number of integrations in transformed plants by quantitative real-time PCR. *BMC Biotechnol* 2: 20, 2002.
17. Alexander GM, Erwin KL, Byers N, *et al*: Effect of transgene copy number on survival in the G93A SOD1 transgenic mouse model of ALS. *Brain Res Mol Brain Res* 130: 7-15, 2004.
18. Ganss R, Montoliu L, Monaghan AP and Schutz G: A cell-specific enhancer far upstream of the mouse tyrosinase gene confers high level and copy number-related expression in transgenic mice. *EMBO J* 13: 3083-3093, 1994.
19. Zimonjic DB, Ullmannova-Benson V, Factor VM, Thorgeirsson SS and Popescu NC: Recurrent and nonrandom DNA copy number and chromosome alterations in Myc transgenic mouse model for hepatocellular carcinogenesis: implications for human disease. *Cancer Genet Cytogenet* 191: 17-26, 2009.
20. Pillai MM, Venkataraman GM, Kosak S and Torok-Storb B: Integration site analysis in transgenic mice by thermal asymmetric interlaced (TAIL)-PCR: segregating multiple-integrant founder lines and determining zygosity. *Transgenic Res* 17: 749-754, 2008.
21. Triglia T, Peterson MG and Kemp DJ: A procedure for in vitro amplification of DNA segments that lie outside the boundaries of known sequences. *Nucleic Acids Res* 16: 8186, 1988.
22. Yuanxin Y, Chengcai A, Li L, Jiayu G, Guihong T and Zhangliang C: T-linker-specific ligation PCR (T-linker PCR): an advanced PCR technique for chromosome walking or for isolation of tagged DNA ends. *Nucleic Acids Res* 31: e68, 2003.
23. Liu YG and Whittier RF: Thermal asymmetric interlaced PCR: automatable amplification and sequencing of insert end fragments from P1 and YAC clones for chromosome walking. *Genomics* 25: 674-681, 1995.
24. Tonooka Y and Fujishima M: Comparison and critical evaluation of PCR-mediated methods to walk along the sequence of genomic DNA. *Appl Microbiol Biotechnol* 85: 37-43, 2009.
25. Lin J, Yu Q, Zhang Q and Yang Q: Construction of mammary gland specific expression plasmid pIN and its expression in vitro and in vivo. *Afr J Biotechnol* 11: 6946-6955, 2012.
26. Zou X, Wang Y, Cheng Y, *et al*: Generation of cloned goats (*Capra hircus*) from transfected foetal fibroblast cells, the effect of donor cell cycle. *Mol Reprod Dev* 61: 164-172, 2002.
27. Liu YG, Mitsukawa N, Oosumi T and Whittier RF: Efficient isolation and mapping of *Arabidopsis thaliana* T-DNA insert junctions by thermal asymmetric interlaced PCR. *Plant J* 8: 457-463, 1995.
28. Hammer RE, Pursel VG, Rexroad CE Jr., *et al*: Production of transgenic rabbits, sheep and pigs by microinjection. *Nature* 315: 680-683, 1985.
29. Goissis MD, Suhr ST and Cibelli JB: Effects of donor fibroblasts expressing OCT4 on bovine embryos generated by somatic cell nuclear transfer. *Cell Reprogram* 15: 24-34, 2013.
30. Niemann H and Lucas-Hahn A: Somatic cell nuclear transfer cloning: practical applications and current legislation. *Reprod Domest Anim* 47 (Suppl 5): 2-10, 2012.
31. Zhao MT, Lin H, Liu FJ, *et al*: Efficiency of human lactoferrin transgenic donor cell preparation for SCNT. *Theriogenology* 71: 376-384, 2009.
32. Eszterhas SK, Bouhassira EE, Martin DI and Fiering S: Transcriptional interference by independently regulated genes occurs in any relative arrangement of the genes and is influenced by chromosomal integration position. *Mol Cell Biol* 22: 469-479, 2002.
33. Bender B, Bodrogi L, Mayer B, *et al*: Position independent and copy-number-related expression of the bovine neonatal Fc receptor alpha-chain in transgenic mice carrying a 102 kb BAC genomic fragment. *Transgenic Res* 16: 613-627, 2007.
34. Rahman MA, Hwang GL, Razak SA, Sohm F and Maclean N: Copy number related transgene expression and mosaic somatic expression in hemizygous and homozygous transgenic tilapia (*Oreochromis niloticus*). *Transgenic Res* 9: 417-427, 2000.
35. Smith CJ, Nedwell DB, Dong LF and Osborn AM: Evaluation of quantitative polymerase chain reaction-based approaches for determining gene copy and gene transcript numbers in environmental samples. *Environ Microbiol* 8: 804-815, 2006.
36. Tang W, Newton RJ and Weidner DA: Genetic transformation and gene silencing mediated by multiple copies of a transgene in eastern white pine. *J Exp Bot* 58: 545-554, 2007.
37. Vaisman BL: Genotyping of transgenic animals by real-time quantitative PCR with TaqMan probes. *Methods Mol Biol* 1027: 233-251, 2013.
38. Bressan FF, Dos Santos Miranda M, Perecin F, *et al*: Improved production of genetically modified fetuses with homogeneous transgene expression after transgene integration site analysis and recloning in cattle. *Cell Reprogram* 13: 29-36, 2011.
39. Chawla R, Ariza-Nieto M, Wilson AJ, Moore SK and Srivastava V: Transgene expression produced by biolistic-mediated, site-specific gene integration is consistently inherited by the subsequent generations. *Plant Biotechnol J* 4: 209-218, 2006.
40. Woychik RP and Alagramam K: Insertional mutagenesis in transgenic mice generated by the pronuclear microinjection procedure. *Int J Dev Biol* 42: 1009-1017, 1998.
41. Migliaccio AR, Bengra C, Ling J, *et al*: Stable and unstable transgene integration sites in the human genome: extinction of the green fluorescent protein transgene in K562 cells. *Gene* 256: 197-214, 2000.
42. Yan B, Li D and Gou K: Homologous illegitimate random integration of foreign DNA into the X chromosome of a transgenic mouse line. *BMC Mol Biol* 11: 58, 2010.