

# miR-542-3p inhibits the growth and invasion of colorectal cancer cells through targeted regulation of cortactin

HAO-CHENG LONG<sup>1\*</sup>, XIA GAO<sup>2\*</sup>, CHANG-JIANG LEI<sup>1\*</sup>, BIN ZHU<sup>1</sup>, LEI LI<sup>1</sup>,  
CHENG ZENG<sup>1</sup>, JIAN-BIN HUANG<sup>1</sup> and JIA-RUI FENG<sup>1</sup>

Departments of <sup>1</sup>General Surgery and <sup>2</sup>Oncology, The Fifth Hospital of Wuhan, Wuhan, Hubei 430050, P.R. China

Received May 16, 2015; Accepted February 8, 2016

DOI: 10.3892/ijmm.2016.2505

**Abstract.** Colorectal cancer is one of the most common malignancies. Previous studies have reported that cortactin (*CTTN*) is often overexpressed in tumors and is associated with metastasis and poor prognosis of patients. The abnormal expression of microRNAs (miRNAs or miRs) is closely related to the development and progression of various types of cancer, including colorectal cancer. However, little is known about the miRNAs targeting cortactin. In the present study, prediction using biological software revealed that cortactin has binding sites for miR-542-3p. Transfection with miR-542-3p mimic demonstrated that miR-542-3p reduced the expression of cortactin in colorectal cancer cells. Dual luciferase reporter assays further demonstrated that miR-542-3p regulated cortactin in a targeted manner and that miR-542-3p expression was significantly downregulated in colorectal cancer cells. A cell proliferation assay and Transwell migration assay were undertaken: we noted that miR-542-3p inhibited the proliferation and invasion of colorectal cancer cells while promoting their apoptosis. By contrast, cortactin acted antagonistically. When co-transfected with miR-542-3p mimic and *CTTN* overexpression vector, the inhibitory effect of miR-542-3p was blocked. This indicates that miR-542-3p regulates *CTTN* in a targeted manner to modulate the growth and invasion of colorectal cancer cells. The present study thus provides new targets for the prevention and treatment of colorectal cancer.

## Introduction

Colorectal cancer is one of the most common malignancies, and its incidence has increased worldwide over the past two decades; it is the third most common cancer in the world (1). In China, it is the fourth most common cause of mortality (2).

There is, therefore, an urgent need to study and understand the biological characteristics of colorectal cancer.

Cortactin is an actin-binding protein that is directly involved in the assembly of actin cytoskeleton. It was first identified as a substrate of oncogenic tyrosine kinase v-Src (3). Cortactin is encoded by the cortactin (*CTTN*) gene (formerly known as EMS1), which is located on chromosome 11q13; this region is amplified in various types of cancer, including head and neck, ovarian, breast, liver, and lung cancer (4). Current data has revealed that cortactin is overexpressed in many human tumors, including head and neck cancer, esophageal squamous cell carcinoma, colon cancer, gastric cancer, hepatocellular carcinoma, breast cancer and ovarian cancer (5-8). Previous studies have reported that gene amplification and increased mRNA expression of cortactin are associated with the metastasis and poor prognosis of tumors (8). A previous study compared the expression profiles of 39 metastasis-related genes, including *CTTN*, in highly and poorly metastatic liver cancer cells (9). The results showed that cortactin overexpression in poorly metastatic liver cancer cells increased the metastatic ability of cancer cells and promoted the development of metastases (9). Several studies have also demonstrated that cortactin overexpression stimulates cell migration and metastasis in both aberrant cell lines and tumor cells (8,10,11). Cortactin plays an important role in cytoskeleton remodeling. It modulates cell motility, invasiveness, synaptogenesis, phagocytosis, tumorigenesis and metastasis (12). Moreover, it also functions as a key component of the signaling pathway involved in cytoskeleton remodeling.

MicroRNAs (miRNAs or miRs) are a class of small, non-coding RNA molecules consisting of 18-25 nucleotides. They regulate the expression of target genes at the post-transcriptional level and are involved in various biological processes, such as cell proliferation, differentiation, apoptosis, metabolism and tumorigenesis (13). Although the exact mechanism and functions of miRNAs have not yet been fully characterized, experimental (14,15) and clinical studies (16,17) have reported that abnormal miRNA expression is involved in the development and progression of various cancers, including colorectal cancer. Currently, the specific expression of more than 100 miRNAs has been studied in colon cancer tissues, cell lines and normal tissues (18). miRNAs play an important role not only in the development of colon cancer, but also in its progression and metastasis; according to a retrospective study by the University of Tokyo, high miR-21 expression is associated with vascular

---

Correspondence to: Dr Hao-Cheng Long, Department of General Surgery, The Fifth Hospital of Wuhan, 122 Xianzheng Street, Hanyang, Wuhan, Hubei 430050, P.R. China  
E-mail: changjiang0118@163.com

\*Contributed equally

**Key words:** miR-542-3p, cortactin, colorectal cancer, microRNA, *CTTN*

invasion, liver metastasis and tumor staging. In addition, miR-155 expression was found to positively correlate with lymph node metastasis in colorectal cancer, and overexpression of these two miRNAs in colorectal cancer patients was linked with reduced survival (19). However, the role of cortactin and related miRNAs in colorectal cancer had not previously received much scholarly attention. With this in mind, the present study was undertaken to examine their important role in colorectal cancer.

## Materials and methods

**Cell culture.** The human colorectal cancer cell lines, LoVo, HCT116 and 293T cells were obtained from the American Type Culture Collection (ATCC, Manassas, VA, USA; www.ATCC.org) and cultured in high-glucose Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum (FBS) (both from Gibco-BRL, Gaithersburg, MD, USA). The cells were maintained at 37°C in an atmosphere of humidified air with 5% CO<sub>2</sub> in a cell culture incubator.

**Overexpression vector construction and transfection.** RNA was extracted from LoVo cells using TRIzol (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions, followed by reverse transcription-quantitative polymerase chain reaction (RT-qPCR) to amplify the coding region of *CTTN*. The products were then digested with *Bam*HI and *Eco*RI (Takara Biotechnology Co., Ltd., Dalian, China), cloned into pcDNA3.1 vectors, sequenced and verified. The primers used for *CTTN* amplification are shown in Table I. We used the biological algorithms, miRBase, miRanda, PITA and RegRNA, to predict candidate miRNAs in the 3'UTR region of the *CTTN* gene. Hsa-miR-542-3p mimic and mimic negative controls were synthesized by Shanghai GenePharma Co., Ltd. (Shanghai, China), and cortactin siRNA was purchased from Santa Cruz Biotechnology, Inc. (Dallas, TX, USA). The cells were seeded onto a 6-well plate at 1x10<sup>5</sup> cells/ml and incubated for 24 h. Transfection was performed using Lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions, when cell confluence reached approximately 70%. The concentrations of plasmid, miRNA and siRNA for transfection were 4 µg/ml, 100 nM/well and 100 nM/well, respectively.

**RT-qPCR.** Total RNA was extracted from the cells using TRIzol (Invitrogen). One microgram of total RNA was used for the first-strand cDNA synthesis with a RevertAid™ First Strand cDNA Synthesis kit (Fermentas GmbH, St. Leon-Rot, Germany). RT-qPCR was performed using 10 µl 2X SYBR-Green PCR Master Mix (Toyobo, Tokyo, Japan), with 5 µl cDNA, 0.5 µl forward primer, 0.5 µl reverse primer, and 4 µl RNase-free ddH<sub>2</sub>O contained in 20 µl of reaction mixture. The primers used for *CTTN* amplification are shown in Table I. The reaction was performed as follows: one cycle of 95°C for 5 min and 40 cycles of 95°C for 30 sec, 58°C for 30 sec and 72°C for 30 sec. Three independent experiments were conducted for each sample. Data were compared using the 2<sup>-ΔΔC<sub>t</sub></sup> method.

**Western blot analysis.** Total cellular proteins were extracted by incubating the cells in lysis buffer [1X PBS, 1% NP-40, 0.1% sodium dodecyl sulfate (SDS), 5 mM ethylenediaminetetraacetic acid (EDTA), 0.5% sodium deoxycholate, and 1 mM

sodium with protease inhibitors]. The protein concentrations in the cell lysates were determined by bicinchoninic acid assay (Pierce, Rockford, IL, USA). SDS-polyacrylamide gel electrophoresis (PAGE) was performed in 8% glycine gels (Bio-Rad, Berkeley, CA, USA) loading equal amounts of proteins per lane. Following electrophoresis, the separated proteins were transferred to nitrocellulose membranes (Pierce) and blocked with 5% non-fat milk in TBST buffer for 1 h. The membranes were then incubated with cortactin (polyclonal, rabbit, #3502) and glyceraldehyde 3-phosphate dehydrogenase (GAPDH; monoclonal, rabbit, #3683) antibodies (Cell Signaling Technology, Inc., Danvers, MA, USA) at 1:1,000 dilutions in 5% non-fat milk overnight at 4°C, and then anti-rabbit IgG monoclonal antibody conjugated with horseradish peroxidase (#7074; Cell Signaling Technology, Inc.) at 1:2,000 dilution for 1 h at room temperature. Protein bands were detected using the West Femto system (Pierce).

**Dual luciferase assay.** The *CTTN* 3'UTR was amplified from cDNA (primers are shown in Table I) and cloned into a psiCHECK-2 vector (Promega Corp., Beijing, China). A QuikChange II Site-Directed Mutagenesis kit (Stratagene, La Jolla, CA, USA) was used in order to generate mutant-type *CTTN* 3'UTR in which the seven mutated nucleotides were underlined within the seed region of the miR-542-3p binding site. A reporter vector psiCHECK-2 carrying the 3'UTR sequences of *CTTN* was assayed for luciferase expression using the Dual-Glo Luciferase assay system (Promega Corp.) according to the manufacturer's instructions. Three independent experiments were performed.

**Cell proliferation.** Cell proliferation was estimated in 96-well plates using a colorimetric immunoassay, based on the measurement of 5-bromo-2'-deoxyuridine (BrdU) incorporation during DNA synthesis (BrdU ELISA kit; Roche Diagnostics GmbH, Mannheim, Germany). Following miRNA and/or plasmid transfection for 48 h, the medium was removed and cells were labeled with BrdU (10 mM) for 3 h at 37°C. The cells were fixed and incubated with peroxidase-conjugated anti-BrdU antibody (sheep polyclonal, 11647229001) for 90 min at room temperature. The peroxidase substrate, 3,3',5,5'-tetramethylbenzidine, was then added, and BrdU incorporation was quantified by differences in absorbance at a wavelength of 370 minus 492 nm. Cell proliferation was expressed as the mean percentage of the control values (set at 100%).

**Cell cycle analysis.** The cells were pelleted (400 x g for 5 min at 4°C), washed twice with cold PBS, resuspended in 500 µl cold PBS, and then fixed for 1 h at 4°C by adding 500 µl fixation solution (2% w/v paraformaldehyde in PBS, pH 7.2). The fixed cells were pelleted, washed with cold PBS, resuspended in 1 ml of 70% ethanol added dropwise to the pellet while vortexing and then incubated overnight at 4°C. On the following day the cells were pelleted and resuspended in 1 ml propidium iodide solution (40 µg/ml with 100 µg/ml RNase A) for 30 min at 37°C in the dark and analyzed using a FACScan flow cytometer (BD Biosciences, Franklin Lakes, NJ, USA).

**Transwell Matrigel invasion assay.** Invasion was evaluated using a Transwell Matrigel invasion assay (BD Biosciences,

Table I. Sequences of primers.

Name	Sense (5'→3')	Antisense (5'→3')	Product sizes (bp)
Primer set for CTTN CDS amplification	CGGGATCCGCCACCATGT GGAAAGCTTCAGCAGG	CGGAATTCCTACTGCC GCAGCTCCACAT	
Primer set for RT-qPCR			
GAPDH	GGTATCGTGGAAGGACTC	GTAGAGGCAGGGATGATG	128
CTTN	GCCGACCGAGTAGACAAG	GTATTTGCCGCCGAAACC	102
Primer set for CTTN 3'UTR amplification	CCGCTCGAGAGCTGCGCCCT GGATCCTCA	ATAAGAATGCGGCCGCTCATA CCTATGAGGTGTGCTACAGG	

CDS, coding sequence; CTTN, cortactin. The underlining represents the restriction enzyme cutting site.




Figure 1. *CTTN* expression in LoVo and HCT116 cells. (A) LoVo and (B) HCT116 cells were transfected with miR-542-3p mimic and/or *CTTN* overexpression vector and then measured using RT-qPCR. The experiments were carried out at least in triplicate and the results are expressed as the means  $\pm$  SD (\* $P$ <0.01 vs. control). Control, untreated cells; NC, negative control.

Shanghai, China). Briefly, 200  $\mu$ l of the LoVo cells following transfection ( $1 \times 10^6$  cells/ml) and 600  $\mu$ l complete medium were added to the upper and lower compartments of the chamber, respectively. Following incubation for 48 h, the cells migrating to the lower side of the filter were fixed with 4% paraformaldehyde for 15 min at room temperature, washed with PBS, stained with crystal violet, and then observed under an inverted microscope (CKX41; Olympus Corp, Tokyo, Japan).

**Statistical analysis.** The experiments were performed at least in triplicate, and the results are expressed as the means  $\pm$  SD. Statistical analysis was performed using the SPSS statistical program version 17 (SPSS Inc., Chicago, IL, USA). A  $P$ -value <0.01 was considered to indicate a statistically significant difference.

## Results

**Prediction and validation of *CTTN* candidate miRNAs.** In the present study, we used the biological algorithms miRBase, miRanda, PITA and RegRNA to predict candidate miRNAs in the 3'UTR region of the *CTTN* gene. In so doing, we identified hsa-miR-542-3p. Although previous studies have reported that miR-542-3p was downregulated in various types of tumor tissues and cells (20,21), few studies have evaluated its functions. The

mimic of the above miRNA was synthesized and transfected into the LoVo and HCT116 cells. RT-qPCR and western blot analysis proved that hsa-miR-542-3p inhibits the expression of *CTTN* (Fig. 1 and 2). This indicates that miR-542-3p regulates *CTTN* expression by targeting this gene. However, this finding warrants further validation by dual-luciferase reporter assays.

*CTTN is a target gene of miR-542-3p.* The 3'UTR region of the *CTTN* gene was cloned into psiCHECK-2, a dual luciferase reporter vector, and the target sites of miR-542-3p were mutated (Fig. 3A, the underlined bases). The two plasmids were subsequently transfected, together with miR-542-3p mimic, into 293T cells, and the changes in luciferase expression were analyzed. Luciferase expression in the cells transfected with plasmids containing the 3'UTR of *CTTN* significantly declined by approximately 44% compared with the cells transfected with an empty reporter vector (Fig. 3B). By contrast, luciferase activity did not change significantly in the cells transfected with the plasmids containing the mutant 3'UTR of the *CTTN* gene. This indicates that miR-542-3p is strongly bound to the 3'UTR of the *CTTN* gene, suggesting that *CTTN* is a target gene of miR-542-3p.

*miR-542-3p affects colorectal cancer cell proliferation and apoptosis by targeting CTTN.* BrdU assays were performed




Figure 2. Cortactin (CTTN) expression in colorectal cancer cells. Cortactin expression in LoVo and HCT116 cells following transfection with miRNA mimic and/or CTTN overexpression vector, as measured using western blot analysis. Each bar represents the means  $\pm$  SD from three samples (\* $P < 0.01$ ). Control, untreated cells; NC, negative control.




Figure 3. CTTN is a target gene of miR-542-3p. (A) miR-542-3p target sites in the sequence of cortactin (CTTN) 3'UTR. (B) The luciferase expression level of 293T cells transfected with cloning CTTN 3'UTR vector or mutant CTTN 3'UTR vector and miR-542-3p mimic. The values are expressed as the means  $\pm$  SD from three samples (\* $P < 0.01$  vs. control). Control, cells transfected with empty reporter vector.

to evaluate the effects of increased miR-542-3p expression on human colorectal cancer cell proliferation. The results showed that miR-542-3p significantly inhibited the proliferation of colorectal cancer cells, whereas CTTN overexpression increased the proliferation of colorectal cancer cells (Fig. 4). Co-transfection with miR-542-3p mimic and the vector causing the overexpression of CTTN abolished the anti-proliferative role of miR-542-3p. Flow cytometric analysis of the cell cycle and apoptosis demonstrated that the percentage of cells in the sub-G1 peak increased significantly when LoVo cells were

transfected with miR-542-3p mimic (Fig. 5). Co-transfection with the miR-542-3p mimic and the CTTN overexpression vector overcame the pro-apoptotic effect of miR-542-3p. Tumor cells are resistant to apoptosis; however, the increased expression of miR-542-3p promoted apoptosis of tumor cells, indicating that miR-542-3p inhibits tumor growth.

*miR-542-3p inhibits the invasion of colorectal cancer cells by targeting CTTN.* Tumor cells are known to be generally invasive and metastatic (22). To determine whether




Figure 4. Effects of miR-542-3p and cortactin (CTTN) on colorectal cancer cell proliferation. The cells were transfected with miR-542-3p mimic and/or CTTN overexpression vector, and cell proliferation was examined using a BrdU assay. Each bar represents the means  $\pm$  SD from three samples (\* $P$ <0.01). Control, untreated cells; NC, negative control.




Figure 5. Effects of miR-542-3p and cortactin (CTTN) on the cell cycle distribution of LoVo cells as examined by flow cytometric analysis. The images are representative of three independent experiments with similar results. Control, untreated cells; NC, negative control.

miR-542-3p inhibits the invasive ability of colorectal cancer cells, we performed Transwell migration assays. We found that an increase in miR-542-3p expression significantly inhibited the invasive ability of LoVo cells, while CTTN overexpression increased the invasion rate (Fig. 6). Co-transfection with miR-542-3p mimic and CTTN overexpression vector overcame the anti-invasive effect of miR-542-3p. This shows that miR-542-3p inhibits the invasion of colorectal cancer cells by targeting the CTTN gene.

## Discussion

Colorectal cancer is known to be one of the most common malignancies; it is the third most common cancer and the second most deadly cancer. Its incidence and mortality are on the rise in China (1,2). Invasion and metastasis are major causes of resultant mortality in patients with colorectal cancer (18). No effective solution is currently available for controlling

colorectal metastases, and the prevention and treatment of colorectal cancer metastases has long been a challenging issue facing the medical community.

Previous research has reported that the invasion and the metastasis of malignant tumors, such as colorectal cancer, is a complex, multistep process. This complex process is possible as cancer cells are motile (23). Motility is an essential, common feature of various stages in the invasion and metastasis of cancer cells (24). Previous research has reported that during the movement of cancer cells, actin polymerization (assembly) is the power source of cancer cell motility (25). It has previously been shown that actin polymerization depends on the catalysis of the Arp 2/3 complex, and the activation of the Arp 2/3 complex requires the CTTN-encoded product, cortactin (26). Cortactin is expressed in colon cancer cells and plays an important role in the motility of cancer cells (27). Lee *et al* (5) have reported that the high expression of CTTN positively correlates with the depth of invasion and distant metastases of colorectal cancer.




Figure 6. Effects of miR-542-3p and cortactin (CTTN) on the invasive ability of LoVo cells. (A) The cells were transfected with miR-542-3p mimic and/or CTTN-overexpressing vector, and the invasion of cells was evaluated using a Transwell Matrigel invasion assay. (B) Quantitative analysis of the anti-invasive effects of miR-542-3p and cortactin. Each bar represents the means  $\pm$  SD from 3 samples (\* $P$ <0.01, vs. control). Control, untreated cells; NC, negative control.

Several studies have reported that the abnormal expression of miRNAs is closely related to the development, progression, and prognosis of colorectal cancer (28,29). Notable progress has been made in the understanding of the biogenesis and mechanisms of action of miRNAs and their role in the diagnosis, treatment, and prognosis of colorectal cancer. However, few studies have been performed on miRNAs targeting CTTN. Only Zhang *et al* (30) found that miRNA-182 inhibits the proliferation and invasion of human lung adenocarcinoma cells through its effect on human cortical actin-associated protein. Hong *et al* (31) showed that VEGF-C increased CTTN expression by downregulating Dicer-mediated maturation of miR326, thereby relieving the suppressive effect of miR326 on CTTN expression. Given the important role of CTTN and miRNAs in colorectal cancer, we performed this study to gain greater insight into this issue. We found that miR-542-3p has target sites on CTTN. Moreover, the upregulation of miR-542-3p inhibited the expression of CTTN. Dual-luciferase reporter assays further confirmed that CTTN is a target gene of miR-542-3p.

A study by Wang *et al* (32) demonstrated that the circulating miRNAs, miR-646, miR-141 and miR-542-3p, potentially serve as non-invasive biomarkers for cervical squamous cell carcinoma. Althoff *et al* (20) concluded that miR-542-3p exerts its tumor suppressive function in neuroblastoma, at least in part, by targeting survivin and the expression of miR-542-3p, and thus targeting these is a promising therapeutic strategy for treating aggressive neuroblastoma. Thus, miR-542-3p has been demonstrated to function as a tumor suppressor gene, and to be involved in the development and progression of cancer.

However, its specific mechanism of action remains unclear. In the present study, we proved that miR-542-3p inhibits the proliferation and invasion of colorectal cancer cells, and promotes their apoptosis by targeting the CTTN gene. The results of a study by Yoon *et al* (21) suggested that survivin is a direct target of miR-542-3p and growth inhibition by miR-542-3p has potential application as an anti-cancer therapy. Wang *et al* (33) have found that overexpression of miR-542-3p in cancer cells elevated p53 expression, stimulated the expression of p53 targets, and inhibited cell proliferation. It has been demonstrated by He *et al* (34) that miRNA-542-3p inhibits tumor angiogenesis by targeting angiopoietin-2. Moreover, Shen *et al* (35) have previously suggested that miRNA-542-3p suppresses the cell growth of gastric cancer cells by targeting oncogene astrocyte-elevated gene-1. Thus, miR-542-3p functions as a tumor suppressor gene by inhibiting the proliferation and invasion of tumor cells.

In conclusion, miR-542-3p inhibits CTTN expression by targeting this gene in colorectal cancer cells and hence suppresses the proliferation and invasion, as well as promoting the apoptosis, of colorectal cancer cells.

#### Acknowledgements

This study was supported by the Wuhan City Health Bureau of Medical Research project (no. WX12C27), the Wuhan Municipal Science and Technology Bureau of Applied Basic research project (no. 2015061701011630), the Education Department of Hubei Province Science and Technology

research project (no. B2015230), and the fourth batch of 'Hanyang Concert Talent Plan'.

## References

1. Siegel R, Ma J, Zou Z and Jemal A: Cancer statistics, 2014. *CA Cancer J Clin* 64: 9-29, 2014.
2. Li M and Gu J: Changing patterns of colorectal cancer in China over a period of 20 years. *World J Gastroenterol* 11: 4685-4688, 2005.
3. Rothschild BL, Shim AH, Ammer AG, Kelley LC, Irby KB, Head JA, Chen L, Varella-Garcia M, Sacks PG, Frederick B, *et al*: Cortactin overexpression regulates actin-related protein 2/3 complex activity, motility, and invasion in carcinomas with chromosome 11q13 amplification. *Cancer Res* 66: 8017-8025, 2006.
4. Schuurin E: The involvement of the chromosome 11q13 region in human malignancies: cyclin D1 and EMS1 are two new candidate oncogenes - a review. *Gene* 159: 83-96, 1995.
5. Lee YY, Yu CP, Lin CK, Nieh S, Hsu KF, Chiang H and Jin JS: Expression of survivin and cortactin in colorectal adenocarcinoma: association with clinicopathological parameters. *Dis Markers* 26: 9-18, 2009.
6. Luo ML, Shen XM, Zhang Y, Wei F, Xu X, Cai Y, Zhang X, Sun YT, Zhan QM, Wu M and Wang MR: Amplification and overexpression of CTTN (EMS1) contribute to the metastasis of esophageal squamous cell carcinoma by promoting cell migration and anoikis resistance. *Cancer Res* 66: 11690-11699, 2006.
7. Timpson P, Wilson AS, Lehrbach GM, Sutherland RL, Musgrove EA and Daly RJ: Aberrant expression of cortactin in head and neck squamous cell carcinoma cells is associated with enhanced cell proliferation and resistance to the epidermal growth factor receptor inhibitor gefitinib. *Cancer Res* 67: 9304-9314, 2007.
8. Buday L and Downward J: Roles of cortactin in tumor pathogenesis. *Biochim Biophys Acta* 1775: 263-273, 2007.
9. Chuma M, Sakamoto M, Yasuda J, Fujii G, Nakanishi K, Tsuchiya A, Ohta T, Asaka M and Hirohashi S: Overexpression of cortactin is involved in motility and metastasis of hepatocellular carcinoma. *J Hepatol* 41: 629-636, 2004.
10. Weaver AM: Cortactin in tumor invasiveness. *Cancer Lett* 265: 157-166, 2008.
11. van Rossum AG, Moolenaar WH and Schuurin E: Cortactin affects cell migration by regulating intercellular adhesion and cell spreading. *Exp Cell Res* 312: 1658-1670, 2006.
12. Ayala I, Baldassarre M, Giacchetti G, Caldieri G, Tetè S, Luini A and Buccione R: Multiple regulatory inputs converge on cortactin to control invadopodia biogenesis and extracellular matrix degradation. *J Cell Sci* 121: 369-378, 2008.
13. Esquela-Kerscher A and Slack FJ: Oncomirs - microRNAs with a role in cancer. *Nat Rev Cancer* 6: 259-269, 2006.
14. Hashimoto Y, Akiyama Y and Yuasa Y: Multiple-to-multiple relationships between microRNAs and target genes in gastric cancer. *PLoS One* 8: e62589, 2013.
15. Iwaya T, Yokobori T, Nishida N, Kogo R, Sudo T, Tanaka F, Shibata K, Sawada G, Takahashi Y, Ishibashi M, *et al*: Down-regulation of miR-144 is associated with colorectal cancer progression via activation of mTOR signaling pathway. *Carcinogenesis* 33: 2391-2397, 2012.
16. Zhang GJ, Xiao HX, Tian HP, Liu ZL, Xia SS and Zhou T: Upregulation of microRNA-155 promotes the migration and invasion of colorectal cancer cells through the regulation of claudin-1 expression. *Int J Mol Med* 31: 1375-1380, 2013.
17. Chiang Y, Song Y, Wang Z, Liu Z, Gao P, Liang J, Zhu J, Xing C and Xu H: microRNA-192, -194 and -215 are frequently down-regulated in colorectal cancer. *Exp Ther Med* 3: 560-566, 2012.
18. Goel A and Boland CR: Recent insights into the pathogenesis of colorectal cancer. *Curr Opin Gastroenterol* 26: 47-52, 2010.
19. Shibuya H, Inuma H, Shimada R, Horiuchi A and Watanabe T: Clinicopathological and prognostic value of microRNA-21 and microRNA-155 in colorectal cancer. *Oncology* 79: 313-320, 2010.
20. Althoff K, Lindner S, Odersky A, Mestdagh P, Beckers A, Karczewski S, Molenaar JJ, Bohrer A, Knauer S, Speleman F, *et al*: miR-542-3p exerts tumor suppressive functions in neuroblastoma by downregulating Survivin. *Int J Cancer* 136: 1308-1320, 2015.
21. Yoon S, Choi YC, Lee S, Jeong Y, Yoon J and Baek K: Induction of growth arrest by miR-542-3p that targets survivin. *FEBS Lett* 584: 4048-4052, 2010.
22. Condeelis J and Segall JE: Intravital imaging of cell movement in tumours. *Nat Rev Cancer* 3: 921-930, 2003.
23. Kedrin D, Wyckoff J, Sahai E, Condeelis J and Segall JE: Imaging tumor cell movement in vivo. *Curr Protoc Cell Biol* Chapter 19: Unit 19.7, 2007. doi: 10.1002/0471143030.cb1907s35.
24. Kedrin D, van Rheenen J, Hernandez L, Condeelis J and Segall JE: Cell motility and cytoskeletal regulation in invasion and metastasis. *J Mammary Gland Biol Neoplasia* 12: 143-152, 2007.
25. Pollard TD, Blanchoin L and Mullins RD: Molecular mechanisms controlling actin filament dynamics in nonmuscle cells. *Annu Rev Biophys Biomol Struct* 29: 545-576, 2000.
26. Urano T, Liu J, Zhang P, Fan YX, Egile C, Li R, Mueller SC and Zhan X: Activation of Arp2/3 complex-mediated actin polymerization by cortactin. *Nat Cell Biol* 3: 259-266, 2001.
27. Hirakawa H, Shibata K and Nakayama T: Localization of cortactin is associated with colorectal cancer development. *Int J Oncol* 35: 1271-1276, 2009.
28. Schepeler T, Reinert JT, Ostensfeld MS, Christensen LL, Silahatoglu, Dyrskjot L, Wiuf C, Sørensen FJ, Kruhoffer M, Laurberg S, *et al*: Diagnostic and prognostic microRNAs in stage II colon cancer. *Cancer Res* 68: 6416-6424, 2008.
29. Slaby O, Svoboda M, Fabian P, Smerdova T, Knoflickova D, Bednarikova M, Nenutil R and Vyzula R: Altered expression of miR-21, miR-31, miR-143 and miR-145 is related to clinicopathologic features of colorectal cancer. *Oncology* 72: 397-402, 2007.
30. Zhang L, Liu T, Huang Y and Liu J: microRNA-182 inhibits the proliferation and invasion of human lung adenocarcinoma cells through its effect on human cortical actin-associated protein. *Int J Mol Med* 28: 381-388, 2011.
31. Hong CC, Chen PS, Chiou J, Chiu CF, Yang CY, Hsiao M, Chang YW, Yu YH, Hung MC, Hsu NW, *et al*: miR326 maturation is crucial for VEGF-C-driven cortactin expression and esophageal cancer progression. *Cancer Res* 74: 6280-6290, 2014.
32. Wang WT, Zhao YN, Yan JX, Weng MY, Wang Y, Chen YQ and Hong SJ: Differentially expressed microRNAs in the serum of cervical squamous cell carcinoma patients before and after surgery. *J Hematol Oncol* 7: 6, 2014.
33. Wang Y, Huang JW, Castella M, Huntsman DG and Taniguchi T: p53 is positively regulated by miR-542-3p. *Cancer Res* 74: 3218-3227, 2014.
34. He T, Qi F, Jia L, Wang S, Song N, Guo L, Fu Y and Luo Y: MicroRNA-542-3p inhibits tumour angiogenesis by targeting angiopoietin-2. *J Pathol* 232: 499-508, 2014.
35. Shen X, Si Y, Yang Z, Wang Q, Yuan J and Zhang X: MicroRNA-542-3p suppresses cell growth of gastric cancer cells via targeting oncogene astrocyte-elevated gene-1. *Med Oncol* 32: 361, 2015.