p-CREB-1 promotes hepatic fibrosis through the transactivation of transforming growth factor- β 1 expression in rats

PEI WANG¹, LIANG DENG², CHUNBO ZHUANG³, CHUNWEI CHENG¹ and KESHU XU¹

¹Division of Gastroenterology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei 430022; ²Department of Gastroenterology, The First Affiliated Hospital of Chongqing Medical University, Chongqing 400016; ³Department of Clinical Laboratory, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei 430022, P.R. China

Received October 28, 2015; Accepted May 31, 2016

DOI: 10.3892/ijmm.2016.2630

Abstract. Phosphorylated cAMP-responsive element binding protein-1 (p-CREB-1) is an important transcription factor which has been reported to be implicated in fibrogenesis. However, the association between p-CREB-1 and transforming growth factor- β 1 (TGF- β 1)-mediated liver fibrogenesis remains poorly understood. In the present study, exogenous TGF- β 1 recombinant protein was used to activate hepatic stellate cells (HSCs), and we established a rat model of tetrachloromethane (CCl₄)-induced liver fibrosis. Loss- and gain-of-function studies were performed to examine the role of p-CREB-1 in liver fibrogenesis, and the detailed mechanism responsible for these effects was further explored using chromatin immunoprecipitation and luciferase reporter gene assays. We found that p-CREB-1 expression was significantly upregulated in a rat model of hepatic fibrosis. We also demonstrated that p-CREB-1 increased TGF- β 1 expression and auto-induction in HSCs, through directly binding to the CRE site within the TGF- β 1 promoter in order to enhance its transcriptional activity. Moreover, lentivirus-mediated CREB-1 overexpression promoted hepatic fibrogenesis in rats. These findings suggest that p-CREB-1 may function as a potent profibrogenic factor through the transactivation of TGF- β 1 expression in liver fibrosis.

Introduction

Hepatic fibrosis, characterized by the excessive production and deposition of the extracellular matrix (ECM), is a common pathologic change observed in the progression of various chronic liver diseases to cirrhosis (1-3). Transforming growth factor- β 1 (TGF- β 1), the most potent profibrogenic cytokine currently known, stimulates the activation and transformation of hepatic stellate cells (HSCs) into myofibroblasts producing abnormal and abundant ECM (4,5). It has been reported that TGF- β 1 binds to TGF- β type II receptors on the cell surface, and then recruits type I receptors; as a consequence, the type II receptor kinase phosphorylates the type I receptors and intracellular signal molecules such as Smads. Phosphorylated (p-)Smad2 and/or p-Smad3 form heterooligomers with Smad4, which translocate into the nucleus to control the ECM gene transcription (6-7). On the contrary, blocking TGF- β 1 signaling may inhibit collagen production and promote the degradation of collagen (8,9). The roles of TGF- β 1 in HSCs activation and liver fibrosis have been intensively studied; however, the transcriptional regulation of TGF- β 1 remains poorly understood.

Cyclic adenosine 3',5'-monophosphate (cAMP)-responsive element (CRE) binding protein-1 (CREB-1), a eukaryotic transcription factor, binds to CRE sites in the promoters of various cytokines to widely regulate gene transcription (10,11). The role of CREB-1 in fibrogenesis remains controversial. Several studies have reported that CREB-1 exerts anti-fibrotic effects, such as in myocardial fibrosis and pulmonary fibrosis (12,13). On the contrary, CREB-1 facilitated high glucose induced renal tubulointerstitial fibrosis in diabetic nephropathy (14). Currently, there limited research regarding the role of CREB-1 in liver fibrosis. Our previous study showed that TGF- β 3 induced CREB-1 phosphorylation in HSCs, and p-CREB-1 bound to the TGF- β 3 promoter and enhanced its transcriptional activity, suggesting that p-CREB-1 was involved in TGF- β 3 auto-regulation (15). However, the effects of p-CREB-1 in TGF- β 1-induced liver fibrosis remain unclear.

Herein, we reported that p-CREB-1 expression was significantly upregulated in the fibrotic liver tissue of rats as well as in HSCs treated with exogenous TGF- β 1. We also found that the overexpression of p-CREB-1 increased TGF- β 1 expression and auto-induction, contributing to hepatic fibrogenesis by

Correspondence to: Professor Keshu Xu, Division of Gastroenterology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, 1277 JieFang Avenue, Wuhan, Hubei 430022, P.R. China
E-mail: xuzou2015@163.com

Abbreviations: p-CREB-1, phosphorylated cyclic adenosine 3',5'-monophosphate-responsive element binding protein-1; TGF- β 1, transforming growth factor- β 1; HSCs, hepatic stellate cells; CCl₄, tetrachloromethane; RT-qPCR, reverse transcription quantitative-polymerase chain reaction; ChIP, chromatin immunoprecipitation

Key words: hepatic fibrosis, hepatic stellate cells, transforming growth factor- β 1, phosphorylated cAMP-responsive element binding protein-1, collagen I

directly binding to the CRE site within the TGF- β 1 promoter to enhance the transcriptional activity of TGF- β 1.

Materials and methods

Cell culture and treatment. The rat HSC line HSC-T6 was obtained from the Cell Bank of the Chinese Academy of Sciences (Shanghai, China). The cells were cultured in Dulbecco's modified Eagle's medium (DMEM) and supplemented with 10% fetal bovine serum (both from Gibco, New York, NY, USA) with 5% CO₂ at 37°C. Trypsin (2.5%; Gibco) was used for continuous cell culture, and cells in logarithmic growth phase were selected for subsequent experiments. The cells were seeded in 6-well plates for 24 h to ensure 80–90% confluence, and then treated with or without exogenous TGF- β 1 recombinant human protein (10 ng/ml; PeproTech, Rocky Hill, NJ, USA) for 8 h.

Construction and transfection of plasmids. The coding sequence of CREB-1 was amplified from rat liver cDNA by PCR and cloned into *Xho*I/*Eco*RI digested pIRES2-EGFP vector (Fulgen, Co., Ltd, Guangzhou, China) to generate a CREB-1 overexpression plasmid (pIRES2-CREB-1). The primers were as follows: 5'-AGACTCGAGATGACCATGGACTCTG-3' (forward) and 5'-ACGGAATTCTCCCAAATTAATCTGAC-3' (reverse). The underlined nucleotide sequences represent the restriction enzyme cutting sites. For sh-CREB-1 plasmid construction, an siRNA sequence targeting the 877–895 sequence of rat CREB-1 mRNA (5'-GAAGAAGCAGCACGAAAGA-3') was synthesized and cloned into a pGenesil-1.1 plasmid (Genesil Biotechnology Co., Ltd., Wuhan, China) at the *Bam*HI/*Hind*III sites.

Plasmid transfection was performed using Lipofectamine 2000 reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. Briefly, the HSCs were seeded in 6-well plates and allowed to reach 70–80% confluence. For each well, transfection was performed using a mixture containing 250 μ l Opti-MEM, with 4 μ g plasmid and 10 μ l Lipofectamine 2000. Following transfection for 6 h, the culture medium was replaced with fresh DMEM.

Animals. All experimental protocols and animal maintenance procedures in this study were approved by the Ethics Review Committee of the Experimental Animal Center of Tongji Medical College at the Huazhong University of Science and Technology (Wuhan, China). Male Sprague-Dawley rats (n=40) weighing 180–200 g were purchased from the Animal Experiment Center of Wuhan University and maintained under a 12-h light/dark cycle at a constant temperature of 20–22°C. The rats were randomly allocated to one of four treatment groups (n=40 for each group): i) normal group, ii) model group, iii) Lv-NC group and iv) Lv-CREB-1 group. At the beginning of the 2nd week, the rats in the Lv-NC and Lv-CREB-1 groups were tail-vein injected with 5 $\times$ 10⁸ TU Lv-NC or Lv-CREB-1 (Biowit Technologies, Shenzhen, China), and the rats in the normal and model groups were injected with an equivalent dose of sterilized saline water. From the 3rd to the 10th week, a subcutaneous injection of 40% tetrachloromethane (CCl₄; Alpha Biotechnology Co., Ltd., Wuhan, China) olive oil solution twice a week was administered to establish a rat model of hepatic fibrosis (initial dose 5 ml/kg body weight, thereafter 3 ml/kg),

and equivalent olive oil was injected into the rats of normal group as control. After 8 weeks of CCl₄ injections, the rats were intraperitoneally anesthetized with pentobarbital sodium (30 mg/kg body weight). Blood samples were collected from the heart, and serum was then separated by centrifugation. The rats were then sacrificed by exsanguination, and the liver tissues were removed and fixed with 4% phosphate-buffered paraformaldehyde for Masson's staining (Goodbio Technology Co., Ltd., Wuhan, China), or frozen for the extraction of RNA and protein. Serum TGF- β 1 levels were detected using enzyme-linked immunosorbent assay (ELISA) kits (Dakewe, Beijing, China) according to the manufacturer's instructions.

Reverse transcription-quantitative polymerase chain reaction (RT-qPCR). Total RNA was extracted using TRIzol reagent (Invitrogen) according to the manufacturer's instructions. Total cDNA was prepared by reverse transcription using PrimeScriptTM RT Master mix (Takara, Otsu, Japan). RT-qPCR was then performed using SYBR-Green PCR Master mix (Takara) in a 20 μ l volume containing 10 μ l SYBR-Green I, 2 μ l cDNA, 0.8 μ l forward primer, 0.8 μ l reverse primer and 6.4 μ l ddH₂O. All PCRs were performed on an ABI StepOneTM (Applied Biosystems, Foster City, CA, USA) under the following conditions: 95°C for 10 min, followed by 40 cycles of 95°C for 5 sec and 60°C for 1 min. The reactions were performed in triplicate with GAPDH as the internal control and the relative expression levels were calculated using the 2^{- $\Delta\Delta$ C_t} method. The specific primers were as follows: TGF- β 1 forward, 5'-CACCGCGTACCAAATGAAGA-3' and TGF- β 1 reverse, 5'-TGGTGCCCTCTGAAATGAAAG-3'; collagen I forward, 5'-GGCGAGTGTCTGTCCTTTCTG-3' and collagen I reverse, 5'-GGGTCCCTCGACTCCTATGAC-3'; CREB-1 forward, 5'-CCAAACTAGCAGTGGGCAGTATATT-3' and CREB-1 reverse, 5'-GGTACCATTGTTAGCCAGCTGTATT-3'; and GAPDH forward, 5'-GTATGACTCTACCCACGGCAAGT-3' and GAPDH reverse, 5'-TTCCCGTT GATGACCAGCTT-3'.

Western blot analysis. Proteins were extracted using RIPA lysis buffer (Beyotime Institute of Biotechnology, Jiangsu, China), and then quantified using a bicinchoninic acid kit (Applygen Technologies, Inc., Beijing, China). After separation by 10% sodium dodecyl sulfate (also known as sodium lauryl sulfate)-polyacrylamide gel electrophoresis, the proteins were transferred onto a PVDF membrane (Millipore, Billerica, MA, USA). The membrane was blocked with 8% skim milk powder in TBST for 1 h at room temperature, and then incubated with rabbit anti-rat primary antibodies CREB-1 (1:1,000, cat. no. 9197) and phospho-CREB-1 Ser133 (1:1,000, cat. no. 9198) (both from Cell Signaling Technology, Danvers, MA, USA); TGF- β 1 (1:1,000, ab92486) and collagen I (1:5,000, ab34710) (both from Abcam, Cambridge, UK); and GAPDH (1:5,000, cat. no. 5174; Cell Signaling Technology) at 4°C overnight. After washing thrice with 1% TBST for 10 min, the horseradish peroxidase-conjugated goat anti-rabbit secondary antibody (1:5,000, cat. no. 7074; Cell Signaling Technology) was added and the membrane was incubated for 1 h at room temperature. Finally, the proteins were detected using an enhanced chemiluminescence detection system (Thermo Fisher Scientific, Waltham, MA, USA). The expression of the target proteins was quantified using AlphaEaseFC software,

and the ratios of target proteins against GAPDH were calculated.

Chromatin immunoprecipitation (ChIP) assay. The HSCs were seeded in 10 cm cell culture dishes and allowed to reach 60-70% confluence, and were transfected with either a pIRES2-CREB-1 or pIRES2-EGFP vector. Forty-eight hours after transfection, a ChIP assay was performed using an EpiQuik™ ChIP kit (Epigentek, Farmingdale, NY, USA) according to the manufacturer's instructions. The antibodies used for ChIP were anti-p-CREB-1 (Cell Signaling Technology) and normal rabbit IgG. Briefly, the cells were trypsinized and cross-linked in 1% formaldehyde at room temperature for 10 min. After washing with phosphate-buffered saline, the cells were lysed and the chromatin was fragmented to approximately 200-1,000 bp by sonication. The chromatin was then incubated in the antibody-coating microwells. After washing, the immunoprecipitated DNA was retrieved from the microwell and the purified DNA was analyzed by qPCR. The amplified region was 183 bp near the transcription start site of the TGF- β 1 gene, and a distant region (164 bp) was used as a negative control. The primers for ChIP were as follows: CRE site, 5'-CGGACCTGCTGGCAATAG-3' (forward) and 5'-CCCAAGGAAAGGTAGGTGATAG-3' (reverse); distant region, 5'-GGGAGGACACAAATACGACC-3' (forward) and 5'-ACGGGACTAAACAGGGAGT-3' (reverse).

Reporter plasmid construction. To construct a TGF- β 1 promoter reporter plasmid, a 2209 bp fragment of the TGF- β 1 gene promoter (-2168/+41 relative to the transcription start site) was amplified by PCR and cloned into the pGL3-basic plasmid (Promega, Madison, WI, USA) between the *Kpn*I and *Hind*III sites, generating pGL3-2209. The primers were as follows: 5'-GAAGGTACCACTGGGGAAGAAAGGAAAC-3' (forward) and 5'-GCAAGCTTCCAAGGAAAGGTAGGTGAT-3' (reverse). The pGL3-2209 plasmid was used as template for generating several deleted TGF- β 1 promoter reporter plasmids using the same reverse primer. The forward primers for the deleted reporter plasmids were as follows: pGL3-1710, 5'-ACTGGTACCTAGCAGCCCAGGCACTC-3'; pGL3-1078, 5'-CTAGGTACCGACCCTGTTTTCTCACGA-3'; pGL3-448, 5'-AGTGGTACCTGCAAGTCAGAGACGGG-3'; and pGL3-66, 5'-AGAGGTACCCCGCGACTCCTGCTG-3'.

To generate a pGL3-mut plasmid, the predicted CREB-1 binding site (CGTCA) located in the TGF- β 1 promoter was converted to 'AATTC' by PCR using pGL3-2209 as a template. The primers were as follows: 5'-ACTCTGGTGTCTAGAGAAT TCCCGCGACTCCTGCTGC-3' (forward) and 5'-CAGCAG GAGTCGCGGAATTCTCTGACACCAGAGTG-3' (reverse).

Luciferase reporter assay. The 293T cells (purchased from the Cell Bank of the Chinese Academy of Sciences) were seeded in 96-well plates and allowed to reach 70-80% confluence. Then, 150 ng of either pIRES2-CREB-1 plasmid or pIRES2-EGFP plasmid, 50 ng of constructed reporter plasmid and 5 ng of pRL-TK *Renilla* plasmid were cotransfected into the 293T cells using Lipofectamine 2000 reagent (Invitrogen) according to the manufacturer's instructions. Forty-eight hours after transfection, the luciferase activity was measured using the Dual Luciferase Reporter assay kit (Promega).

Statistics analysis. Data were analyzed by the independent-samples t-test or one-way analysis of variance using SPSS 17.0 software (IBM Corp., Armonk, NY, USA). Correlation analysis was performed using Pearson's correlation test. A p-value <0.05 indicated a statistically significant difference.

Results

p-CREB-1 is significantly upregulated in rat fibrotic liver tissues. To examine the potential role of p-CREB-1 in liver fibrogenesis, we detected the expression of p-CREB-1 in rat liver tissues with CCl₄-induced fibrosis, and the results showed that rats with liver fibrosis had higher p-CREB-1 expression than those without liver fibrosis (1.8-fold, $p < 0.05$) (Fig. 1A). Additionally, there was a positive correlation between p-CREB-1 expression and collagen contents in a rat model of liver fibrosis (Fig. 1B). These results suggested that p-CREB-1 may play a role in liver fibrosis.

Expression of p-CREB-1 is induced by exogenous TGF- β 1 in HSCs. We then examined the regulatory mechanism responsible for the effects of p-CREB-1 in liver fibrosis. It is well known that TGF- β 1 may be induced by CCl₄ and plays a crucial role in rat liver fibrogenesis (16), thus it is of interest to explore the association between p-CREB-1 and TGF- β 1. Herein, we used recombinant human TGF- β 1 protein (10 ng/ml) to activate the HSCs, and found that the protein expression levels of p-CREB-1 were significantly increased in the TGF- β 1-treated HSCs compared with the control ($p < 0.01$) (Fig. 2D), and this was accompanied by increases in endogenous TGF- β 1 mRNA and protein (9.4-fold, $p < 0.001$; and 4.0-fold, $p < 0.01$; respectively) (Fig. 2A and D). Additionally, the TGF- β 1-treated HSCs also showed notable increases in the mRNA and protein expression of collagen I (43.3-fold, $p < 0.001$; and 3.5-fold, $p < 0.001$; respectively), compared with the HSCs treated with TGF- β 1 diluents (Fig. 2B and D). However, the mRNA and protein expression of CREB-1 was unaffected by TGF- β 1 (Fig. 2C and D).

p-CREB-1 increases TGF- β 1 expression and is required for TGF- β 1 auto-induction in HSCs. The above result indicates that the TGF- β 1 induction of p-CREB-1 is accompanied by an induction of endogenous TGF- β 1. It is of interest to determine whether p-CREB-1 promotes TGF- β 1 expression and is required for TGF- β 1 auto-induction. Herein, we performed loss- and gain-of-function studies. The HSCs were transfected with the pIRES2-CREB-1 vector in order to upregulate CREB-1, with or without exogenous TGF- β 1 treatment. As shown in Fig. 3, there were significant increases in the mRNA and protein expression of TGF- β 1 and collagen I in the pIRES2-CREB-1-transfected HSCs (TGF- β 1 mRNA, 3.1-fold, $p < 0.001$; TGF- β 1 protein, 2.2-fold, $p < 0.001$; collagen I mRNA, 11.5-fold, $p < 0.001$; collagen I protein, 3.0-fold, $p < 0.01$), as well as in the expression of p-CREB-1 which was also upregulated compared with the control; and these increases were more significant when the HSCs were additionally treated with exogenous TGF- β 1 (TGF- β 1 mRNA, 8.5-fold, $p < 0.001$; TGF- β 1 protein, 3.5-fold, $p < 0.001$; collagen I mRNA, 18.2-fold, $p < 0.001$; collagen I protein: 5.6-fold, $p < 0.001$), suggesting that the increased TGF- β 1 expression induced by exogenous TGF- β 1 was further enhanced through the upregulation of p-CREB-1 in HSCs.

Figure 1. Phosphorylated cAMP-responsive element binding protein-1 (p-CREB-1) is significantly upregulated in rat liver fibrosis tissues and in hepatic stellate cells (HSCs) stimulated by exogenous transforming growth factor- β 1 (TGF- β 1). (A) Protein expression of p-CREB-1 and collagen I were evaluated by western blot analysis of the liver tissues. The representative blot (left) and statistical analysis (right) are shown. (B) The expression of p-CREB-1 positively correlated with the expression of collagen I protein in rats of model group (n=8). * $p<0.05$, ** $p<0.01$.

Figure 2. Expression of phosphorylated cAMP-responsive element binding protein-1 (p-CREB-1), transforming growth factor- β 1 (TGF- β 1) and collagen I are upregulated by exogenous TGF- β 1. Hepatic stellate cells (HSCs) were treated with or without exogenous TGF- β 1 (10 ng/ml) for 8 h and the mRNA and protein expression of the indicated genes was detected by RT-qPCR and western blot analysis. mRNA expression of (A) TGF- β 1, (B) collagen I and (C) CREB-1. (D) Western blot analysis of CREB-1, p-CREB-1, TGF- β 1 and collagen I. The representative blot (left) and statistical analysis are shown (right). ** $p<0.01$, *** $p<0.001$ vs. control.

Moreover, we used a sh-CREB-1 plasmid to downregulate CREB-1 expression in the HSCs, and examined whether a reduction of p-CREB-1 suppressed TGF- β 1 expression and auto-induction. As shown in Fig. 3, sh-CREB-1 significantly repressed the expression of TGF- β 1 and collagen I (TGF- β 1 mRNA, 0.5-fold, $p<0.01$; TGF- β 1 protein, 0.4-fold, $p<0.001$; collagen I mRNA, 0.4-fold, $p<0.01$; collagen I protein, 0.3-fold, $p<0.01$), compared with pGenesil-1.1. Furthermore,

a reduction of p-CREB-1 decreased the exogenous TGF- β 1 induction of endogenous TGF- β 1 and collagen I expression. The above findings indicated that p-CREB-1 plays a central role in TGF- β 1 expression and auto-regulation.

p-CREB-1 transactivates the TGF- β 1 promoter through directly binding to the CRE site within the promoter. To confirm whether the transcription factor p-CREB-1 binds to the TGF- β 1

Figure 3. Phosphorylated cAMP-responsive element binding protein-1 (p-CREB-1) is required for transforming growth factor- β 1 (TGF- β 1) auto-induction. Hepatic stellate cells (HSCs) were transiently transfected with 4 μ g pIRES2-EGFP, pIRES2-CREB-1, pGenesil-1.1 or sh-CREB-1. Forty hours after transfection, the cells were treated with or without exogenous TGF- β 1 (10 ng/ml) for 8 h. The cells were then harvested and the mRNA and protein expression of the indicated genes was evaluated by RT-qPCR and western blot analysis. The mRNA expression of (A) CREB-1, (B) TGF- β 1 and (C) collagen I. (D and E) Western blot analysis of CREB-1, p-CREB-1, TGF- β 1 and collagen I. (D) The representative blot and (E) statistical analysis are shown. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

promoter, we performed ChIP assays. As shown in Fig. 4B, we found a recruitment of p-CREB-1 to the CRE site in the TGF- β 1 promoter. By contrast, when the HSCs were transfected with pIRES2-CREB-1 vector, the recruitment was significantly enhanced (2.3-fold) compared with the control, suggesting that p-CREB-1 directly bind to the TGF- β 1 promoter in HSCs.

To determine whether the upregulation of p-CREB-1 was responsible for the increased transcriptional activity of the TGF- β 1 promoter, a series of 5'-deletion mutants were constructed and transfected respectively into 293T cells, which were cotransfected with the pIRES2-CREB-1 plasmid or pIRES2-EGFP plasmid, and pRL-TK *Renilla* plasmid. Reporter gene assays were performed and showed that the luciferase activities of pGL3-2209, pGL3-1710, pGL3-1078 and pGL3-448 reporter plasmids were significantly augmented in cells transfected with the pIRES2-CREB-1 plasmid, whereas that of the pGL3-66 plasmid was not markedly increased compared with the control. These results suggested that TGF- β 1

transcription was activated by p-CREB-1, and the region between -407 and -25 was most important for the transactivation by p-CREB-1 (Fig. 4D). Furthermore, 5'-CTGCA-3' sequences in this region were identified as the putative CREB-1 binding site, and converted to 5'-AATTC-3' in the pGL3-mut plasmid. Reporter gene assays showed that the luciferase activity of the pGL3-mut plasmid was not markedly increased by CREB-1 overexpression, indicating that mutation of the site completely abrogated the transcriptional activation of the TGF- β 1 reporter by p-CREB-1 (Fig. 4D). Moreover, as shown in Fig. 4C, exogenous TGF- β 1 increased the luciferase activity of pGL3-2209 (2.3-fold higher than control), whereas pGL3-mut was unaffected. These results suggest that the CRE site is essential for the transactivation effects of p-CREB-1 on the TGF- β 1 promoter.

p-CREB-1 upregulates TGF- β 1 expression and promotes liver fibrogenesis in rats. All rats survived in the normal control group whereas 8, 9 and 7 rats survived in the model group,

Figure 4. Phosphorylated cAMP-responsive element binding protein-1 (p-CREB-1) transactivates transforming growth factor- β 1 (TGF- β 1) expression in hepatic stellate cells (HSCs). (A) Illustration of cAMP-response element (CRE) within the TGF- β 1 promoter. TGF- β 1 promoter region near the putative CREB-1 binding site is highly conserved. (B) Chromatin immunoprecipitation (ChIP) assay shows the enrichment of p-CREB-1 in the TGF- β 1 promoter. The DNA was immunoprecipitated with anti-p-CREB-1 or normal IgG. qPCR was used to quantify the enriched DNA and data was expressed as the fold of enrichment relative to input DNA. (C) Exogenous TGF- β 1 increases TGF- β 1 promoter activity. HSCs were cotransfected with either 50 ng pGL3-2209 or 50 ng pGL3-mut plasmid, and 5 ng pRL-TK plasmid. Forty-eight hours after transfection, the cells were treated with or without TGF- β 1 (10 ng/ml) for 8 h. Then the cells were harvested and luciferase activity was measured. (D) The CRE site is requisite for TGF- β 1 gene transcription. The TGF- β 1 promoter reporter plasmid, deleted or replaced mutant reporter plasmid was cotransfected with pIRES2-CREB-1 in 293T cells. Values were shown relative to the luciferase activity of pGL3-basic plasmid. ** $p < 0.01$, *** $p < 0.001$ vs. control.

Lv-NC group and Lv-CREB-1 group, respectively, until the rats were sacrificed. Masson's staining revealed that the content of collagen in liver samples from the model group is higher than that in the normal control group and lower than that in Lv-CREB-1 group (Fig. 5A), suggesting that overexpression of CREB-1 promoted CCl₄-induced hepatic fibrogenesis in rats.

Furthermore, we determined the hepatic expression of CREB-1, p-CREB-1, TGF- β 1 and collagen I by RT-qPCR and/or western blot analysis, and the results indicated that CREB-1 expression was markedly elevated in the Lv-CREB-1 group (Fig. 5C, F and G). p-CREB-1 expression was 3.7-fold and 1.8-fold upregulated in the Lv-CREB-1 group and the model group, compared with the normal control group ($p < 0.01$ and $p < 0.05$, respectively) (Fig. 5F and H). Moreover, the mRNA and protein expression of TGF- β 1 in the model group were also significantly increased compared with the normal control, and the Lv-CREB-1 groups exhibited a more significant increase (Fig. 5D, F and I). Serum TGF- β 1 levels detected by ELISA displayed similar changes among these

groups (Fig. 5B). Consistent with these findings, the mRNA and protein expression of collagen I were also markedly increased in the Lv-CREB-1 group (Fig. 5E, F and J). These results suggest that p-CREB-1 also increases the expression of TGF- β 1 and collagen I *in vivo*.

Discussion

CREB-1 is an important transcription factor in gene regulation which responds to multiple extracellular signals (17), and its transcriptional activity is positively regulated by the phosphorylation of a Ser residue, Ser133 (10). Non-phosphorylated CREB-1 is not capable of activating gene transcription. Once phosphorylated, CREB-1 translocates into the nucleus and binds to target DNA sequences, and then p-CREB-1 interacts with CREB-binding protein (CBP) to initiate the transcription of CREB-1-responsive genes (18). In the present study, we examined the expression level of activated CREB-1 and showed that p-CREB-1 expression was markedly upregulated

Figure 5. Phosphorylated cAMP-responsive element binding protein-1 (p-CREB-1) promotes transforming growth factor- β 1 (TGF- β 1)-mediated collagen I production and liver fibrogenesis in rats. Rats were tail-vein injected with 5×10^8 TU of Lv-NC or Lv-CREB-1. One week after tail-vein injection, a subcutaneous injection of 40% tetrachloromethane (CCl_4) twice/week for 8 weeks was used to induce the rat hepatic fibrosis model (initial dose 5 ml/kg body weight, thereafter 3 ml/kg). (A) Masson's staining was performed in the paraffin sections of liver tissues to visualize collagen accumulation (magnification, $\times 100$). (B) Serum TGF- β 1 levels were detected by enzyme-linked immunosorbent assay (ELISA) kits. (C-J) Expression of TGF- β 1, collagen I, CREB-1 and p-CREB-1 were evaluated by RT-qPCR and/or western blot analysis in the liver tissue. * $p < 0.05$, ** $p < 0.01$ vs. control.

in a rat model of CCl_4 -induced liver fibrosis, indicating that p-CREB-1 may play an important role in liver fibrogenesis. Our study is consistent with a recent study, which reported a similar finding; namely that CREB phosphorylation was significantly increased in alcoholic liver fibrosis (19). Furthermore, we examined the underlying regulatory mechanism as well as the role of upregulated p-CREB-1 in liver fibrosis. It is well known that TGF- β 1 is a potent cytokine which promotes the activation of HSCs and liver fibrogenesis, and is highly expressed in activated HSCs and CCl_4 -induced hepatic fibrosis (16,20,21). Herein, we explored the association between p-CREB-1 and TGF- β 1 in liver fibrosis. Firstly, we determined whether p-CREB-1 was induced by TGF- β 1. We used TGF- β 1 recombinant protein to activate the HSCs, and demonstrated that TGF- β 1 induced the protein expression of p-CREB-1 in HSCs. Similarly, another study by Chin *et al* reported that TGF- β 1 promoted CREB-1 phosphorylation through the activation of mitogen-activated protein kinase (MAPK) in neurons, resulting in the enhanced excitability of neurons (22). Jang *et al* demonstrated that TGF- β 1 induced CREB-1 phosphorylation in macrophages (23). It appears that p-CREB-1 induced by TGF- β 1 represents the

common mechanism responsible for CREB-1 activation. Moreover, we found that the TGF- β 1 induction of p-CREB-1 was accompanied by an auto-induction of TGF- β 1. This led us to explore the potential functional role of p-CREB-1 in liver fibrosis; whether p-CREB-1 is capable of promoting TGF- β 1 expression.

It has been previously established that TGF- β 1 functions as a potent pro-fibrotic factor. However, the regulatory mechanism responsible for TGF- β 1 expression in liver fibrosis remains poorly understood. Herein, the loss- and gain-of-function studies showed that p-CREB-1 upregulated TGF- β 1 expression and was required for TGF- β 1 auto-induction in HSCs. Previous studies demonstrated that TGF- β 1 expression may be regulated by several transcription factors such as AP-1, SP-1, nuclear factor (NF)- κ B, Kruppel-like factor 4 (gut) (KLF4) and signal transducer and activator of transcription 3 (STAT3) in various experimental model systems including liver fibrosis (24-27). In the present study, we reported that another transcription factor, p-CREB-1, was capable of transactivating TGF- β 1 expression. Moreover, we identified the region between -407 and -25 was crucial for the transactivation of TGF- β 1, which was consistent with the aforementioned studies. Additionally, we also

found that the overexpression of p-CREB-1 promoted hepatic fibrogenesis through the transactivation of TGF- β 1 expression in rats. Our study is consistent with the findings of a recent study, which reported that the expression of p-CREB-1 was downregulated by caffeine through its blockade of adenosine A2A receptor and the cAMP/PKA/CREB signaling pathway, thereby inhibiting the activation of HSCs and liver fibrogenesis (19). Taken together, our findings provide evidence for the pro-fibrotic role of p-CREB-1 in liver fibrogenesis.

In the present study, we found that p-CREB-1 expression was significantly upregulated in fibrotic liver tissues in rats, as well as in HSCs treated with exogenous TGF- β 1. We also found that p-CREB-1 increased TGF- β 1 expression and auto-induction, contributing to hepatic fibrogenesis by directly binding to the CRE site within the TGF- β 1 promoter to enhance the transcriptional activity of TGF- β 1. Our study characterizes p-CREB-1 as a profibrogenic factor in hepatic fibrosis, and provides a novel insight into the regulatory mechanism responsible for TGF- β 1 expression.

Acknowledgements

The authors gratefully acknowledge the financial support of the National Natural Science Foundation of China (grant nos. 30871153 and 81370525).

References

- Friedman SL: Mechanisms of hepatic fibrogenesis. *Gastroenterology* 134: 1655-1669, 2008.
- Lee UE and Friedman SL: Mechanisms of hepatic fibrogenesis. *Best Pract Res Clin Gastroenterol* 25: 195-206, 2011.
- Trautwein C, Friedman SL, Schuppan D and Pinzani M: Hepatic fibrosis: concept to treatment. *J Hepatol* 62 (Suppl 1): S15-S24, 2015.
- Gressner OA, Rizk MS, Kovalenko E, Weiskirchen R and Gressner AM: Changing the pathogenetic roadmap of liver fibrosis? Where did it start; where will it go? *J Gastroenterol Hepatol* 23: 1024-1035, 2008.
- Bissell DM, Roulot D and George J: Transforming growth factor beta and the liver. *Hepatology* 34: 859-867, 2001.
- Inagaki Y and Okazaki I: Emerging insights into transforming growth factor beta Smad signal in hepatic fibrogenesis. *Gut* 56: 284-292, 2007.
- Heldin CH, Miyazono K and ten Dijke P: TGF-beta signalling from cell membrane to nucleus through SMAD proteins. *Nature* 390: 465-471, 1997.
- Friedman SL: Mechanisms of disease: mechanisms of hepatic fibrosis and therapeutic implications. *Nat Clin Pract Gastroenterol Hepatol* 1: 98-105, 2004.
- Dooley S, Hamzavi J, Breitkopf K, Wiercinska E, Said HM, Lorenzen J, Ten Dijke P and Gressner AM: Smad7 prevents activation of hepatic stellate cells and liver fibrosis in rats. *Gastroenterology* 125: 178-191, 2003.
- Zhang X, Odom DT, Koo SH, Conkright MD, Canettieri G, Best J, Chen H, Jenner R, Herbolsheimer E, Jacobsen E, *et al*: Genome-wide analysis of cAMP-response element binding protein occupancy, phosphorylation, and target gene activation in human tissues. *Proc Natl Acad Sci USA* 102: 4459-4464, 2005.
- Altarejos JY and Montminy M: CREB and the CRTC co-activators: sensors for hormonal and metabolic signals. *Nat Rev Mol Cell Biol* 12: 141-151, 2011.
- Chan EC, Dusting GJ, Guo N, Peshavariya HM, Taylor CJ, Dilley R, Narumiya S and Jiang F: Prostacyclin receptor suppresses cardiac fibrosis: role of CREB phosphorylation. *J Mol Cell Cardiol* 49: 176-185, 2010.
- Baarsma HA, Engelbertink LH, van Hees LJ, Menzen MH, Meurs H, Timens W, Postma DS, Kerstjens HA and Gosens R: Glycogen synthase kinase-3 (GSK-3) regulates TGF- β 1-induced differentiation of pulmonary fibroblasts. *Br J Pharmacol* 169: 590-603, 2013.
- Visavadiya NP, Li Y and Wang S: High glucose upregulates upstream stimulatory factor 2 in human renal proximal tubular cells through angiotensin II-dependent activation of CREB. *Nephron Exp Nephrol* 117: e62-e70, 2011.
- Deng L, Li Y, Huang JM, Zhou G, Qian W and Xu K: Effects of p-CREB-1 on transforming growth factor- β 3 auto-regulation in hepatic stellate cells. *J Cell Biochem* 112: 1046-1054, 2011.
- Doh KO, Jung HK, Moon IJ, Kang HG, Park JH and Park JG: Prevention of CCl4-induced liver cirrhosis by ribon antisense to transforming growth factor- β 1. *Int J Mol Med* 21: 33-39, 2008.
- Wen AY, Sakamoto KM and Miller LS: The role of the transcription factor CREB in immune function. *J Immunol* 185: 6413-6419, 2010.
- Sakamoto KM and Frank DA: CREB in the pathophysiology of cancer: implications for targeting transcription factors for cancer therapy. *Clin Cancer Res* 15: 2583-2587, 2009.
- Wang Q, Dai X, Yang W, Wang H, Zhao H, Yang F, Yang Y, Li J and Lv X: Caffeine protects against alcohol-induced liver fibrosis by dampening the cAMP/PKA/CREB pathway in rat hepatic stellate cells. *Int Immunopharmacol* 25: 340-352, 2015.
- Zhang S, Sun WY, Wu JJ and Wei W: TGF- β signaling pathway as a pharmacological target in liver diseases. *Pharmacol Res* 85: 15-22, 2014.
- Purps O, Lahme B, Gressner AM, Meindl-Beinker NM and Dooley S: Loss of TGF-beta dependent growth control during HSC transdifferentiation. *Biochem Biophys Res Commun* 353: 841-847, 2007.
- Chin J, Liu RY, Cleary LJ, Eskin A and Byrne JH: TGF-beta1-induced long-term changes in neuronal excitability in aplysia sensory neurons depend on MAPK. *J Neurophysiol* 95: 3286-3290, 2006.
- Jang YS, Kim JH, Seo GY and Kim PH: TGF- β 1 stimulates mouse macrophages to express APRIL through Smad and p38MAPK/CREB pathways. *Mol Cells* 32: 251-255, 2011.
- Zhang Y, Wang Y, Liu Y, Wang N, Qi Y and Du J: Krüppel-like factor 4 transcriptionally regulates TGF- β 1 and contributes to cardiac myofibroblast differentiation. *PLoS One* 8: e63424, 2013.
- Lin W, Tsai WL, Shao RX, Wu G, Peng LF, Barlow LL, Chung WJ, Zhang L, Zhao H, Jang JY and Chung RT: Hepatitis C virus regulates transforming growth factor beta1 production through the generation of reactive oxygen species in a nuclear factor kappaB-dependent manner. *Gastroenterology* 138: 2509-2518, 2010.
- Presser LD, McRae S and Waris G: Activation of TGF- β 1 promoter by hepatitis C virus-induced AP-1 and Sp1: role of TGF- β 1 in hepatic stellate cell activation and invasion. *PLoS One* 8: e56367, 2013.
- Hosui A, Kimura A, Yamaji D, Zhu BM, Na R and Hennighausen L: Loss of STAT5 causes liver fibrosis and cancer development through increased TGF-beta and STAT3 activation. *J Exp Med* 206: 819-831, 2009.