

Prediction of key genes and miRNAs responsible for loss of muscle force in patients during an acute exacerbation of chronic obstructive pulmonary disease

YANHONG DUAN¹, MIN ZHOU², JIAN XIAO¹, CHAOMIN WU¹, LEI ZHOU¹,
FENG ZHOU¹, CHUNLING DU¹ and YUANLIN SONG³

¹Department of Respiratory Medicine, Qingpu Branch of Zhongshan Hospital, Fudan University, Shanghai 201700;

²Department of Respiratory Medicine, Jinshan Branch of The Sixth People's Hospital of Shanghai, Shanghai 201599;

³Department of Respiratory Medicine, Zhongshan Hospital, Fudan University, Shanghai 201700, P.R. China

Received December 3, 2015; Accepted August 30, 2016

DOI: 10.3892/ijmm.2016.2761

Abstract. The present study aimed to identify genes and microRNAs (miRNAs or miRs) that were abnormally expressed in the vastus lateralis muscle of patients with acute exacerbations of chronic obstructive pulmonary disease (AECOPD). The gene expression profile of GSE10828 was downloaded from the Gene Expression Omnibus database, and this dataset was comprised of 4 samples from patients with AECOPD and 5 samples from patients with stable COPD. Differentially expressed genes (DEGs) were screened using the Limma package in R. A protein-protein interaction (PPI) network of DEGs was built based on the STRING database. Module analysis of the PPI network was performed using the ClusterONE plugin and functional analysis of DEGs was conducted using DAVID. Additionally, key miRNAs were enriched using gene set enrichment analysis (GSEA) software and a miR-gene regulatory network was constructed using Cytoscape software. In total, 166 up- and 129 downregulated DEGs associated with muscle weakness in AECOPD were screened. Among them, *NCL*, *GOT1*, *TMOD1*, *TSPO*, *SOD2*, *NCL* and *PA2G4* were observed in the modules consisting of upregulated or downregulated genes. The upregulated DEGs in modules (including *KLF6* and *XRCC5*) were enriched in GO terms associated

with immune system development, whereas the downregulated DEGs were enriched in GO terms associated with cell death and muscle contraction. Additionally, 39 key AECOPD-related miRNAs were also predicted, including miR-1, miR-9 and miR-23a, miR-16 and miR-15a. In conclusion, DEGs (*NCL*, *GOT1*, *SOD2*, *KLF6*, *XRCC5*, *TSPO* and *TMOD1*) and miRNAs (such as miR-1, miR-9 and miR-23a) may be associated with the loss of muscle force in patients during an acute exacerbation of COPD which also may act as therapeutic targets in the treatment of AECOPD.

Introduction

Chronic obstructive pulmonary disease (COPD), characterized by a fixed obstruction of the airway caused by emphysema, chronic bronchitis, or both, is a common, growing public health problem that is responsible for a huge economic health burden worldwide (1). Stable COPD may lead to declines in lung functions, such as airflow obstruction, airway function decline, and respiratory muscle fatigue, which impair patient quality of life; although acute exacerbations of COPD (AECOPD) have various definitions, they are commonly characterized by worsened dyspnea and increased volumes of phlegm and phlegm purulence, usually accompanied by hypoxemia and worsened hypercapnia (2,3). Small airway lesions (due to chronic bronchiolitis) and destruction of the alveolar walls (emphysema) are the two major features of AECOPD (4). Based on available data in 2010, COPD is one of the 6 leading causes of death, and AECOPD is associated with high morbidity (5). To date, there are no methods for preventing AECOPD, and current medical therapies for AECOPD mainly involve bronchial relaxation and the use of glucocorticoids and antibiotics, which, however, are always associated with side effects leading to unsatisfactory prognosis (6).

Smoking, malnutrition, depression and drug addiction are the risk factors affecting the quality of life for patients with AECOPD (7). Furthermore, skeletal muscle dysfunction is the most severe complication of COPD, and progression is associated with oxidative stress, skeletal muscle fiber types, systemic inflammation and mitochondrial dysfunction (1),

Correspondence to: Dr Yuanlin Song, Department of Respiratory Medicine, Zhongshan Hospital, Fudan University, 180 Fenglin Road, Shanghai 201700, P.R. China
E-mail: yuanlinsongyls@163.com

Dr Chunling Du, Department of Respiratory Medicine, Qingpu Branch of Zhongshan Hospital, Fudan University, Shanghai 201700, P.R. China
Email: chunlingducl@163.com

Key words: loss of muscle force, acute exacerbation of chronic obstructive pulmonary disease, differentially expressed genes, miRNAs, miR-gene regulatory network

bringing great detrimental effects to patients. In fact, skeletal muscle depletion has been adopted as a predictor of mortality in patients with COPD (8). In COPD patients, a lack of antioxidant capacity and glutathione (GSH) indicated that oxidative stress is associated with skeletal muscle dysfunction (9). Furthermore, bacterial infection is another major cause of AECOPD (10). The increasing levels of systemic inflammatory factors, such as tumor necrosis factor α (TNF- α), interleukin (IL)-6 and IL-8, may inhibit muscle shrinkage and the protein degradation of skeletal muscle, thus leading to muscle atrophy (11). Mitochondrial dysfunction, in terms of sharply increased transmembrane potential and reduced mitochondrial density, is one of the factors associated with abnormal skeletal muscle in COPD (9). Thus, further studies are warranted in order to explore the mechanism underlying the pathogenesis of AECOPD.

Previous studies have identified a number of microRNAs (miRNAs or miRs) that may have significant regulatory functions in the progression of COPD, such as miR-223, miR-1274a and miR-15b in lung tissue (12), let-7c and miR-125b (13), as well as serum miR-20a, miR-28-3p, miR-34c-5p, miR-100 and miR-7 (14). Furthermore, the expression of miR-1 in quadriceps may be responsible for muscle dysfunction in COPD (15). To the best of our knowledge, however, there have been no reports on miRNAs contributing to skeletal muscle weakness in patients with AECOPD.

In order to identify the key genes and miRNAs that may be responsible for the loss of muscle force during an acute exacerbation in COPD patients, we downloaded a gene expression profile dataset GSE10828 (16) from the Gene Expression Omnibus (GEO) database of the National Center for Biotechnology Information (NCBI; <http://www.ncbi.nlm.nih.gov/geo>) (17), which were collected from muscle samples of patients with AECOPD or stable COPD. The authors submitting this dataset identified the differentially expressed genes (DEGs) that may be associated with loss of muscle force during AECOPD, and performed gene ontology (GO) enrichment analysis (16). In the present study, which is based on the identified DEGs, we constructed a protein-protein interaction (PPI) network and conducted subsequent module analysis in order to identify genes that play critical roles in the progression of AECOPD; we also built a miR-gene regulatory network. This study aimed to deepen our understanding of the molecular mechanisms underlying the loss of muscle force in AECOPD.

Materials and methods

Affymetrix microarray data. Only one microarray dataset (GSE10828) (16) was found to be associated with AECOPD in the GEO database of NCBI (17). The annotation platform of this dataset is the GPL2891 platform (GE Healthcare/Amersham Biosciences CodeLink™ UniSet Human 20K I Bioarray, Chalfont, UK). This dataset were collected from vastus lateralis samples from 4 male patients with acute COPD and 5 male patients with stable COPD. No significant differences were found in the basic characteristics between the patients with AECOPD and stable COPD with regard to age, body mass index (BMI), forced expiratory volume in the first second (FEV₁), FEV₁/forced vital capacity (FVC), arterial oxygen and carbon dioxide tension (PaO₂ and PaCO₂), and

maximal inspiratory mouth pressure (PI_{max}), apart from C-reactive protein (CRP) levels at admission and lower quadriceps force (16).

Data preprocessing and identification of DEGs. Firstly, the microarray data in .CEL format were converted into expression measures using the GEOquery package, a package for retrieving gene expression data sets in R/Bioconductor (Bioconductor version: Release 3.1, <http://www.bioconductor.org/packages/release/bioc/html/GEOquery.html>) (18). The Robust Multi-array Analysis (RMA) method was used to preprocess the downloaded raw data by background adjustment, quintile normalization and summarization (19). Subsequently, the differential expression values of genes between the acute COPD samples and the stable COPD samples were calculated by t-test using the Linear Models for Microarray Data (LIMMA) package (R/Bioconductor version: Release 3.1, <http://www.bioconductor.org/packages/release/bioc/html/limma.html>) (20). Multiple testing correction was conducted by the Bayesian method (21). Only genes with a false discovery rate (FDR) <0.01 and $|\log_2 \text{Fold-Change (FC)}| >1.5$ were identified as DEGs. Finally, the pheatmap package of R (<http://cran.r-project.org/web/packages/pheatmap/index.html>) was used for hierarchical clustering (22) of DEGs based on Euclidean distance (23), and the result was visualized using heat maps.

GO analysis. GO enrichment analysis was performed using the Database for Annotation, Visualization and Integrated Discovery (DAVID) online tool (version 6.7, <http://david.abcc.ncifcrf.gov/>) (24) with a p-value <0.05 as threshold.

Construction of PPI network and functional analysis of significant modules. The Search Tool for the Retrieval of Interacting Genes (STRING) (version 10, <http://string-db.org/>) database was used to predict the interactions between the encoded proteins of these up- and downregulated DEGs, respectively (25). Only PPI pairs with a combination score >0.4 were included. Cytoscape software (version 3.2.1, <http://cytoscape.org/>), an open source software platform for integrating bimolecular interaction networks with high-throughput expression data and other molecular states into a unified conceptual framework (26), was used to visualize the resulting PPIs. The top five genes with the highest connection degrees were considered to be most closely associated with AECOPD. In addition, Cluster ONE plugin (27) was used to select the significant modules of the up- and downregulated genes in the PPI network, respectively. The top three modules according to p-values were selected for further analysis.

In addition, the biological processes of DEGs in the resulting networks were functionally analyzed using DAVID (version 6.7, <http://david.abcc.ncifcrf.gov/>) (24) with a p-value <0.05.

Enrichment analysis of key miRNAs and regulatory network construction of key miRNAs. We used the gene set enrichment analysis (GSEA) software (version 2.0.14, <http://www.broadinstitute.org/gsea/index.jsp>), based on the whole genome expression profile (28), to predict the miRNAs associated with AECOPD based on the microarray data. A p-value <0.01 was selected as threshold.



A Data before normalization**B** Data after normalization

Figure 1. Cascade figure of the normalization expression value. (A) Data before normalization. (B) Data after normalization. Green represents samples from patients with acute exacerbations of chronic obstructive pulmonary disease (AECOPD); red represents samples from patients with stable COPD.

In addition, we used Cytoscape software to construct a miRNA-gene regulatory network with the miRNAs and their target DEGs. A miR-DEG regulatory network is a complicated biological system in which an miRNA functions in a variety of intracellular biological processes, such as gene regulation, cell signaling transduction, protein-protein interactions and metabolism, by regulating molecules at a transcriptional level. This provides a deeper understanding of the biological mechanisms involved (29).

Results

Screening of DEGs and hierarchical clustering analysis. The expression profiling data were preprocessed using the R GEOquery package and were normalized by the RMA method in R (Fig. 1).

With the cut-offs of $FDR < 0.01$ and $|\log_2 FCI| > 1.5$, a total of 295 DEGs were identified between the human AECOPD and stable COPD controls, including 166 upregulated genes and 129 downregulated genes (Fig. 2).

Functional annotation analysis of DEGs. According to the GO functional annotation, the top 10 enriched biological process terms included cellular respiration, generation of precursor metabolites and energy, and respiratory electron transport




Figure 2. Hierarchical clustering heat maps of differentially expressed genes (DEGs). The gradient color from green to red represents the expression level [acute chronic obstructive pulmonary disease (COPD) sample and stable COPD samples changes from upregulation to downregulation].

chain; the top 10 molecular function terms included cytoskeletal protein binding, serine-type endopeptidase inhibitor activity, protein complex binding and NADH dehydrogenase activity. Translocator protein (*TSPO*), superoxide dismutase 2 (*SOD2*) and proliferation associated protein (*PA2G4*) were the genes enriched in the significant functions (Table I).

Table I. The top 10 enriched GO terms in BP and MF categories.

Name	Term	Count	p-value	Genes
BP	GO:0015980 - energy derivation by oxidation of organic compounds	11	7.94E-05	<i>NDUFS7, NDUFS4, GOT1, UQCRC1, PPP1R1A, NDUF7, NDUFAB1, NDUFS1, PDHB, FH, SOD2</i>
BP	GO:0045333 - cellular respiration	9	1.26E-04	<i>NDUFS7, NDUFS4, UQCRC1, NDUF7, NDUFAB1, NDUFS1, PDHB, FH, SOD2</i>
BP	GO:0006091 - generation of precursor metabolites and energy	15	3.67E-04	<i>UQCRC1, TXN2, NDUF7, NDUFAB1, ATP5G3, PDHB, SOD2, NDUFS7, GOT1, NDUFS4, PKM2, PPP1R1A, ATP5I, NDUFS1, FH</i>
BP	GO:0022904 - respiratory electron transport chain	7	4.50E-04	<i>NDUFS7, NDUFS4, UQCRC1, NDUF7, NDUFAB1, NDUFS1, SOD2</i>
BP	GO:0006119 - oxidative phosphorylation	8	8.01E-04	<i>NDUFS7, NDUFS4, UQCRC1, NDUF7, NDUFAB1, ATP5I, ATP5G3, NDUFS1</i>
BP	GO:0007005 - mitochondrion organization	9	0.001355	<i>NDUFS7, FIS1, SPG7, TSPO, YWHAZ, NDUFS4, BCS1L, RRM2B, SOD2</i>
BP	GO:0042773 - ATP synthesis coupled electron transport	6	0.001691	<i>NDUFS7, NDUFS4, UQCRC1, NDUF7, NDUFAB1, NDUFS1</i>
BP	GO:0042775 - mitochondrial ATP synthesis coupled electron transport	6	0.001691	<i>NDUFS7, NDUFS4, UQCRC1, NDUF7, NDUFAB1, NDUFS1</i>
BP	GO:0022900 - electron transport chain	8	0.001937	<i>NDUFS7, NDUFS4, UQCRC1, TXN2, NDUF7, NDUFAB1, NDUFS1, SOD2</i>
BP	GO:0006120 - mitochondrial electron transport, NADH to ubiquinone	5	0.003914	<i>NDUFS7, NDUFS4, NDUF7, NDUFAB1, NDUFS1</i>
MF	GO:0004857 - enzyme inhibitor activity	16	1.89E-05	<i>CAST, PPME1, SERPING1, AZIN1, FURIN, ANXA4, ANXA2, OAZ2, SH3BP5, YWHAG, COL7A1, PPP1R1A, KAL1, SERPINE1, SERPINA3, SLPI</i>
MF	GO:0008092 - cytoskeletal protein binding	19	7.97E-04	<i>ACTN4, CNN3, TNNC1, MYL3, DIAPH3, EVL, PALLD, ANXA2, LOC729143, SPAG9, ANK1, SMTN, KATNA1, NCK1, CLIP1, ARL8B, CAP1, MYH7B, TMOD1</i>
MF	GO:0004867 - serine-type endopeptidase inhibitor activity	7	0.002949	<i>COL7A1, KAL1, SERPINE1, SERPINA3, SLPI, SERPING1, FURIN</i>
MF	GO:0019899 - enzyme binding	18	0.003008	<i>PPME1, DIAPH3, NFKBIA, FURIN, ANXA2, SPAG9, YWHAG, PA2G4, ANK1, NEDD4, IGF2R, SERPINE1, SORT1, SYTL3, AKAP1, KPNA2, PA2G4P4, DHCR24, ADAM9</i>
MF	GO:0032403 - protein complex binding	10	0.003358	<i>SYP, INSL3, YWHAZ, UQCRC1, ACTN4, TXN2, CTGF, NFKBIA, ITGB1, ADAM9</i>
MF	GO:0050136 - NADH dehydrogenase (quinone) activity	5	0.004217	<i>NDUFS7, NDUFS4, NDUF7, NDUFAB1, NDUFS1</i>
MF	GO:0008137 - NADH dehydrogenase (ubiquinone) activity	5	0.004217	<i>NDUFS7, NDUFS4, NDUF7, NDUFAB1, NDUFS1</i>
MF	GO:0003954 - NADH dehydrogenase activity	5	0.004217	<i>NDUFS7, NDUFS4, NDUF7, NDUFAB1, NDUFS1</i>
MF	GO:0016655 - oxidoreductase activity, acting on NADH or NADPH, quinone or similar compound as acceptor	5	0.006741	<i>NDUFS7, NDUFS4, NDUF7, NDUFAB1, NDUFS1</i>
MF	GO:0004866 - endopeptidase inhibitor activity	8	0.007169	<i>CAST, COL7A1, KAL1, SERPINE1, SERPINA3, SLPI, SERPING1, FURIN</i>

GO, gene ontology; BP, biological process; MF, molecular function.




Figure 3. Protein-protein interaction (PPI) network of differentially expressed genes (DEGs).

PPI network. The protein-protein network of the identified DEGs was mapped using the software STRING in order to predict the protein interactions. By integrating these correlations, interaction networks between the target genes and their interactive genes were constructed (Fig. 3). In the core of the PPI network, genes belonged to more than one module. *TSPO*, nucleolin (*NCL*), NADH dehydrogenase (ubiquinone) 1, α/β subcomplex (*NDUFAB1*), *PA2G4* and *SOD2* were the top 5 protein nodes with the highest connection degrees (Table II and Fig. 3).

Module analysis and functional analysis of DEGs in PPI network. For the upregulated DEGs, three significant modules were obtained: module 1 consisting of 11 nodes (including *NCL*) and 33 edges (p-value=3.017E-5), module 2 of 14 nodes and 16 edges (p-value=3.115E-5), and module 3 of 13 nodes and 17 edges (p-value=3.194E-5) (Fig. 4A). Moreover, 3 significant modules were also obtained for the downregulated DEGs: module 1 consisting of 16 nodes and 35 edges (p-value=1.418E-6), module 2 of 18 nodes and 21 edges (p-value=4.592E-6), and module 3 of 10 nodes and 15 edges (p-value=3.642E-4). Pyruvate dehydrogenase

Table II. Genes with the top 5 node degrees in PPI network.

Gene	Degree	Log2FC	p-value
<i>TSPO</i>	25	-2.73239	0.007907
<i>NCL</i>	14	2.021786	0.001702
<i>NDUFAB1</i>	13	-27.2233	0.006066
<i>PA2G4</i>	12	3.607543	0.006317
<i>SOD2</i>	11	21.28352	0.005172

Degree, degree of node in PPI network; Log2FC, logarithms of amplitude changes for differentially expressed gene expression values; PPI, protein-protein interaction

(lipoamide) beta (*PDHB*) was commonly detected in module 1 and 2; glutamic pyruvate transaminase (alanine aminotransferase) 2 (*GPT2*), glutamic-oxaloacetic transaminase 1 (*GOT1*) and aldehyde dehydrogenase 4 family member A1 (*ALDH4A1*) were common in module 2 and 3 (Fig. 4B).




Figure 4. Significant modules of differentially expressed genes (DEGs) in the protein-protein interaction (PPI) network. (A) Three modules for the upregulated DEGs, (B) three modules for the downregulated DEGs. Red nodes stand for the upregulated DEGs while blue nodes represent downregulated DEGs. Edges stand for the protein interaction and dot circle stand for the modules.

In addition, GO terms of the DEGs in the significant modules are presented in Table III. Functions of upregulated DEGs in module 1 were in the ribosome biogenesis. No GO

term of the upregulated DEGs in module 2 was detected. Upregulated genes in module 3, such as X-ray repair complementing defective repair in Chinese hamster cells 5 (*XRCC5*)

Table III. GO terms of the DEGs in significant modules.

Module	Term	Count	p-value	Genes
A, GO terms of the upregulated DEGs in the significant modules				
Module 1	Cluster 1 enrichment score: 4.308352643161865			
	GO:0042254 - ribosome biogenesis	5	7.66E-07	<i>DCAF13, PA2G4, UTP6, RRS1, FTSJ3</i>
	GO:0022613 - ribonucleoprotein complex biogenesis	5	3.63E-06	<i>DCAF13, PA2G4, UTP6, RRS1, FTSJ3</i>
	GO:0006364 - rRNA processing	4	2.48E-05	<i>DCAF13, PA2G4, UTP6, FTSJ3</i>
	GO:0016072 - rRNA metabolic process	4	2.82E-05	<i>DCAF13, PA2G4, UTP6, FTSJ3</i>
	GO:0034470 - ncRNA processing	4	2.05E-04	<i>DCAF13, PA2G4, UTP6, FTSJ3</i>
	GO:0034660 - ncRNA metabolic process	4	3.78E-04	<i>DCAF13, PA2G4, UTP6, FTSJ3</i>
	GO:0006396 - RNA processing	4	4.60E-03	<i>DCAF13, PA2G4, UTP6, FTSJ3</i>
Module 2	None			
Module 3	Cluster 1 enrichment score: 2.0090957996722376			
	GO:0043066 - negative regulation of apoptosis	4	2.51E-03	<i>XRCC5, HSP90B1, NFKBIA, SOD2</i>
	GO:0043069 - negative regulation of programmed cell death	4	2.61E-03	<i>XRCC5, HSP90B1, NFKBIA, SOD2</i>
	GO:0060548 - negative regulation of cell death	4	2.63E-03	<i>XRCC5, HSP90B1, NFKBIA, SOD2</i>
	GO:0042981 - regulation of apoptosis	5	2.92E-03	<i>XRCC5, HSP90B1, ACTN4, NFKBIA, SOD2</i>
	GO:0043067 - regulation of programmed cell death	5	3.03E-03	<i>XRCC5, HSP90B1, ACTN4, NFKBIA, SOD2</i>
	GO:0010941 - regulation of cell death	5	3.07E-03	<i>XRCC5, HSP90B1, ACTN4, NFKBIA, SOD2</i>
	GO:0001666 - response to hypoxia	3	5.05E-03	<i>HSP90B1, ACTN4, SOD2</i>
	GO:0070482 - response to oxygen levels	3	5.58E-03	<i>HSP90B1, ACTN4, SOD2</i>
	GO:0006916 - anti-apoptosis	3	1.16E-02	<i>HSP90B1, NFKBIA, SOD2</i>
	GO:0042592 - homeostatic process	4	2.01E-02	<i>XRCC5, HSP90B1, SERPINE1, SOD2</i>
	Cluster 2 enrichment score: 1.7537673375612617			
	GO:0030097 - hemopoiesis	3	1.50E-02	<i>XRCC5, KLF6, SOD2</i>
	GO:0048534 - hemopoietic or lymphoid organ development	3	1.81E-02	<i>XRCC5, KLF6, SOD2</i>
	GO:0002520 - immune system development	3	2.02E-02	<i>XRCC5, KLF6, SOD2</i>
	Cluster 3 enrichment score: 1.1482146347041753			
	GO:0051252 - regulation of RNA metabolic process	6	9.85E-03	<i>ZFP36, KLF6, YY1, NFKBIA, ATF1, SOD2</i>
	GO:0006357 - regulation of transcription from RNA polymerase II promoter	4	1.84E-02	<i>KLF6, YY1, NFKBIA, SOD2</i>
	GO:0006355 - regulation of transcription, DNA-dependent	5	4.53E-02	<i>KLF6, YY1, NFKBIA, ATF1, SOD2</i>
B, GO terms of the downregulated DEGs in the significant modules				
Module 1	Cluster 1 enrichment score: 8.207305207832107			
	GO:0006091 - generation of precursor metabolites and energy	10	1.12E-12	<i>NDUFS7, NDUFS4, UQCRC1, TXN2, NDUFA7, NDUFAB1, ATP5I, ATP5G3, NDUFS1, PDHB</i>
	GO:0006119 - oxidative phosphorylation	8	1.39E-12	<i>NDUFS7, NDUFS4, UQCRC1, NDUFA7, NDUFAB1, ATP5I, ATP5G3, NDUFS1</i>
	GO:0045333 - cellular respiration	7	1.92E-10	<i>NDUFS7, NDUFS4, UQCRC1, NDUFA7, NDUFAB1, NDUFS1, PDHB</i>

Table III. Continued.

Module	Term	Count	p-value	Genes
B, GO terms of the downregulated DEGs in the significant modules				
	GO:0022900 - electron transport chain	7	5.13E-10	<i>NDUFS7, NDUFS4, UQCRC1, TXN2, NDUF7, NDUFAB1, NDUFS1</i>
	GO:0042773 - ATP synthesis coupled electron transport	6	1.27E-09	<i>NDUFS7, NDUFS4, UQCRC1, NDUF7, NDUFAB1, NDUFS1</i>
	GO:0042775 - mitochondrial ATP synthesis coupled electron transport	6	1.27E-09	<i>NDUFS7, NDUFS4, UQCRC1, NDUF7, NDUFAB1, NDUFS1</i>
	GO:0015980 - energy derivation by oxidation of organic compounds	7	2.11E-09	<i>NDUFS7, NDUFS4, UQCRC1, NDUF7, NDUFAB1, NDUFS1, PDHB</i>
	GO:0022904 - respiratory electron transport chain	6	2.53E-09	<i>NDUFS7, NDUFS4, UQCRC1, NDUF7, NDUFAB1, NDUFS1</i>
	GO:0006120 - mitochondrial electron transport, NADH to ubiquinone	5	5.62E-08	<i>NDUFS7, NDUFS4, NDUF7, NDUFAB1, NDUFS1</i>
	GO:0055114 - oxidation reduction	8	6.79E-07	<i>NDUFS7, NDUFS4, UQCRC1, TXN2, NDUF7, NDUFAB1, NDUFS1, PDHB</i>
	GO:0016310 - phosphorylation	8	3.09E-06	<i>NDUFS7, NDUFS4, UQCRC1, NDUF7, NDUFAB1, ATP5I, ATP5G3, NDUFS1</i>
	GO:0006793 - phosphorus metabolic process	8	1.14E-05	<i>NDUFS7, NDUFS4, UQCRC1, NDUF7, NDUFAB1, ATP5I, ATP5G3, NDUFS1</i>
	GO:0006796 - phosphate metabolic process	8	1.14E-05	<i>NDUFS7, NDUFS4, UQCRC1, NDUF7, NDUFAB1, ATP5I, ATP5G3, NDUFS1</i>
	Cluster 2 enrichment score: 2.1705698016603754			
	GO:0010257 - NADH dehydrogenase complex assembly	3	3.82E-05	<i>NDUFS7, NDUFS4, BCS1L</i>
	GO:0032981 - mitochondrial respiratory chain complex I assembly	3	3.82E-05	<i>NDUFS7, NDUFS4, BCS1L</i>
	GO:0033108 - mitochondrial respiratory chain complex assembly	3	5.60E-05	<i>NDUFS7, NDUFS4, BCS1L</i>
	GO:0007005 - mitochondrion organization	3	7.49E-03	<i>NDUFS7, NDUFS4, BCS1L</i>
	GO:0043623 - cellular protein complex assembly	3	1.02E-02	<i>NDUFS7, NDUFS4, BCS1L</i>
	GO:0034622 - cellular macromolecular complex assembly	3	3.62E-02	<i>NDUFS7, NDUFS4, BCS1L</i>
	GO:0034621 - cellular macromolecular complex subunit organization	3	4.47E-02	<i>NDUFS7, NDUFS4, BCS1L</i>
	Cluster 3 enrichment score: 2.16913552816325			
	GO:0046034 - ATP metabolic process	3	4.40E-03	<i>ATP5I, ATP5G3, NDUFS1</i>
	GO:0009205 - purine ribonucleoside triphosphate metabolic process	3	5.44E-03	<i>ATP5I, ATP5G3, NDUFS1</i>
	GO:0009199 - ribonucleoside triphosphate metabolic process	3	5.53E-03	<i>ATP5I, ATP5G3, NDUFS1</i>
	GO:0009144 - purine nucleoside triphosphate metabolic process	3	5.90E-03	<i>ATP5I, ATP5G3, NDUFS1</i>
	GO:0009141 - nucleoside triphosphate metabolic process	3	6.77E-03	<i>ATP5I, ATP5G3, NDUFS1</i>
	GO:0009150 - purine ribonucleotide metabolic process	3	7.49E-03	<i>ATP5I, ATP5G3, NDUFS1</i>
	GO:0009259 - ribonucleotide metabolic process	3	8.46E-03	<i>ATP5I, ATP5G3, NDUFS1</i>
	GO:0006163 - purine nucleotide metabolic process	3	1.33E-02	<i>ATP5I, ATP5G3, NDUFS1</i>

Table III. Continued.

Module	Term	Count	p-value	Genes
B, GO terms of the downregulated DEGs in the significant modules				
Module 2	Cluster 1 enrichment score: 2.0431078400255807			
	GO:0008219 - cell death	6	1.12E-03	<i>MUC2, SPG7, TSPO, PKM2, DIABLO, TBP</i>
	GO:0016265 - death	6	1.15E-03	<i>MUC2, SPG7, TSPO, PKM2, DIABLO, TBP</i>
	GO:0012501 - programmed cell death	4	3.31E-02	<i>MUC2, TSPO, PKM2, DIABLO</i>
	GO:0006915 - apoptosis	3	1.58E-01	<i>MUC2, TSPO, DIABLO</i>
Module 3	Cluster 1 enrichment score: 4.256164391089315			
	GO:0006936 - muscle contraction	4	2.77E-05	<i>MYL3, TNNC1, CKMT2, MYOM1</i>
	GO:0003012 - muscle system process	4	3.66E-05	<i>MYL3, TNNC1, CKMT2, MYOM1</i>
	GO:0006941 - striated muscle contraction	3	1.68E-04	<i>MYL3, TNNC1, MYOM1</i>

GO, gene ontology; DEGs, differentially expressed genes.

and Kruppel-like factor 6 (*KLF6*), were enriched in GO terms including hemopoietic or lymphoid organ development and immune system development (Table III-A). GO functions of downregulated DEGs in module 1 were in generation of precursor metabolites and energy, whereas that in module 2 was in cell death, and that in module 3 were in muscle contraction and muscle system process (Table III-B).

Enrichment analysis of key miRNAs. In total, we identified 39 AECOPD-associated miRNAs using GSEA software at a p-value <0.01 (Table IV).

miRNA regulatory network. Thirty-nine predicted miRNAs and their target DEGs were constructed in an miR-DEG regulatory network (Fig. 5). miR-9, miR-524, miR-23a, miR-15a and miR-16 were the top 5 miRNAs with the most target DEGs (Table V), and their target DEGs included tropomodulin 1 (*TMOD1*), *GOT1*, *NR3C2* and *CPEB4* among others.

Discussion

Since AECOPD are one of the leading causes of death, there is an urgent need to investigate the mechanism underlying AECOPD and to develop an effective preventative strategy. Microarray-based studies have been performed to analyze the pathogenesis of AECOPD and to identify the AECOPD-associated genes in peripheral blood mononuclear cells (30) and skeletal muscle (16). However, no research investigating AECOPD-associated miRNAs has been reported, to the best of our knowledge. Hence, the present study was performed in order to predict AECOPD-associated mRNAs and miRNAs that may be responsible for the loss of muscle force, and to discuss the molecular mechanisms underlying the loss of muscle force during AECOPD.

NCL was observed in the Module 1 of the upregulated genes. It encodes a eukaryotic nucleolar phosphoprotein that is involved in the synthesis and maturation of ribosomes, which

is mainly located in dense, fibrillar regions of the nucleolus. Nucleolin is one of the three components consisting of a D4Z4 repeat (31), in which the number variation is frequently detected in facioscapulohumeral muscular dystrophy (32). Thus, it can be inferred that *NCL* may also play a role in the loss of muscle force in AECOPD since it is associated with muscular function. Furthermore, *NCL* was also predicted to be regulated by miR-1 in the present study. miR-1 and miR-206, another miRNA also observed to regulate the differential gene expression herein, promote myotube formation (33). Another study reported the reduced expression of miR-1 in the quadriceps of patients with COPD, suggesting that miR-1 downregulation may contribute to COPD-associated skeletal muscle dysfunction (34), and they further observed an inverse correlation between miR-1 and Akt phosphorylation levels or HDAC4 protein levels in patients. Thus, it is likely that *NCL* may be downregulated during the AECOPD due to the downregulation of miR-1. *SOD2* was another upregulated gene observed in module 3. An imbalance of the oxidation-antioxidant system in the body represents the principal cause of AECOPD (35). *SOD2* (Mn-SOD) is a key enzyme that prevents cells from damage by eliminating the endogenous free radicals in the body (36), and increased expression was found in patients with AECOPD in the present study. Considering that samples were taken from patients with an exacerbation on day 4 of hospitalization, the antioxidant system may be activated by upregulating *SOD2* in patients with AECOPD. However, this hypothesis requires further careful consideration. Additionally, Togliatto *et al* have reported that unacylated ghrelin (UnAG) induced skeletal muscle regeneration following hindlimb ischemia and was mediated by *SOD2* (37). *SOD2* may also play similar roles in muscle dysfunction during AECOPD, which suggests that *SOD2* may be used as a therapeutic target in AECOPD.

In addition, *XRCC5* and *KLF6* were also found in module 3 of the upregulated DEGs. Both genes are involved in immune system development according to GO analysis, suggesting that the two genes may be important in AECOPD. *KLF6* is

Table IV. Enrichment analysis of key miRNAs.

Name	Basic groups	Size	ES	NES	NOM	FDR
miR-23a, miR-23b	AATGTGA	343	0.278676	1.438706	0	0.089588
miR-103, miR-107	ATGCTGC	177	0.335453	1.486206	0	0.088895
miR-221, miR-222	ATGTAGC	108	0.336992	1.396569	0	0.094994
miR-320	CAGCTTT	213	0.343332	1.552349	0	0.078454
miR-520f	AAGCACT	194	0.344786	1.56579	0	0.091075
miR-183	GTGCCAT	152	0.355742	1.541346	0	0.073964
miR-524	CTTTGTA	365	0.35657	1.562035	0	0.084759
miR-493	ATGTACA	266	0.384502	1.549554	0	0.074808
miR-494	ATGTTTC	128	0.386803	1.631935	0	0.135062
miR-498	GCTTGAA	92	0.397854	1.605433	0	0.115428
miR-1, miR-206	ACATTCC	253	0.398441	1.603156	0	0.108254
miR-323	TAATGTG	131	0.400318	1.56248	0	0.087979
miR-373	TTTTGAG	194	0.406584	1.664687	0	0.125706
miR-485-3p	TGTATGA	128	0.41496	1.611675	0	0.123895
miR-9	TAGCTTT	190	0.419432	1.627284	0	0.120835
miR-17-3p	ACTGCAG	87	0.421458	1.635642	0	0.150641
miR-409-3p	AACATTC	120	0.421607	1.671052	0	0.156583
miR-422b, miR-422a	AAGTCCA	56	0.428607	1.58016	0	0.109384
miR-200a	GTAAGAT	44	0.440217	1.63578	0	0.179569
miR-518a-2	TTTGCAG	169	0.441104	1.684444	0	0.204788
miR-202	ATAGGAA	84	0.445265	1.574246	0	0.100272
miR-410	GTTATAT	76	0.466728	1.711476	0	0.295102
miR-217	ATGCAGT	95	0.477771	1.610814	0.009328	0.113216
miR-15a, miR-16, miR-15a, miR-195						
miR-424, miR-497	TGCTGCT	499	0.293906	1.427648	0.009452	0.09335
miR-126	TAATAAT	179	0.379129	1.572559	0.009524	0.09167
miR-186	ATTCTTT	234	0.390894	1.593438	0.009785	0.099291
miR-182	TTGCCAA	274	0.355565	1.520188	0.00994	0.085925
miR-519e	GGCACTT	105	0.382867	1.579575	0.00994	0.104196
miR-527	CTTTGCA	192	0.316954	1.487434	0.00996	0.089759

Basic groups: the targets of miRNAs; size, the number of target genes regulated by miRNA; ES, the integral enrichment of miRNAs; NES, standardization of the enrichment integration; NOM, standardization of p-value; FDR, false discovery rate.

Table V. miRNAs with the top 5 degrees in regulatory network.

miRNA	Degree	miRNA	Degree
miR-9	20	miR-15A	11
miR-524	13	miR-15B	11
miR-23a	10	miR-16	11
miR-23b	10	miR-182	11
miR-320	10	miR-186	11
miR-373	10	miR-195	11
miR-1	9	miR-424	11
miR-206	9	miR-497	11
miR-518a-2	9		

a member of the Kruppel-like family of transcription factors that functions as a tumor suppressor (38). Mgbemena *et al* have proven that KLF6 regulated the apoptosis of lung cells through iNOS expression during respiratory syncytial virus infection (39). In addition, KLF6 may also be involved in cell atrophy during an acute exacerbation. On the other hand, XRCC5 is an ATP-dependent DNA helicase II or DNA repair protein (40). The role of XRCC5 in COPD has not been fully elucidated. However, previous findings have revealed that DNA damage or lack of DNA repair regulated the immune response to the tissue destruction in COPD (41). Therefore, XRCC5 may be a novel target for protecting against AECOPD.

TSPO displayed downregulated expression in patients with AECOPD, and this was observed in module 2. This gene encodes a protein transformation-related 18-kDa protein that




Figure 5. The miRNA regulatory network. Blue represents miRNA; red represents upregulated differentially expressed genes (DEGs); green represents downregulated DEGs.

assists in the recognition of the mitochondrial proteins prior to intracellular transportation (42). Otherwise, dysfunction of mitochondria caused by permeability transition would lead to the apoptosis of muscle cells, which plays a principal role in the progression of COPD (43). Thus, the downregulation of *TSPO* expression observed suggests that it may have an important role in the loss of muscle force occurring in AECOPD.

TMD1 was another key downregulated gene, observed in module 3. Tropomodulin is a binding protein of tropomyosin, existing in the muscle cells, and is extracted from erythrocytes. It is necessary for many key biological functions including cell migration, differentiation and muscle contraction (44). The most important cause of the progression of AECOPD is oxygen deficiency resulting from several factors, such as the transformation of pulmonary blood vessel structures, manifested by hyperplasia and hypertrophy of pulmonary arterial muscle cells, leading to

the incassation of membranes and fibroblast proliferation (45). In the present study, *TMD1* was predicted to be regulated by miR-23a. It has been reported that the upregulation of miR-23a inhibits the development of B cells (46).

GOT1, displaying downregulated expression in AECOPD, was found in both modules 2 and 3 of the downregulated genes. *GOT1* encodes glutamic-oxaloacetic transaminase 1, which is a cytoplasmic form of glutamic-oxaloacetic transaminase that is involved in amino acid metabolism. De Palma *et al* have also reported the decreased expression of GOT1 in patients with Ullrich congenital muscular dystrophy compared to controls (47). However, there have been no reports regarding the role of GOT1 in AECOPD, to the best of our knowledge. However, it has been reported as a putative target gene of miR-9 by Thulin *et al* (48), which is consistent with our predictions. miR-9 is known to play a key role in the activation of monocytes

and/or macrophages during inflammatory responses (49). More importantly, it is also reported to play a role in Huntington's disease, a type of motor neuron disease (50). Furthermore, the abnormal expression of miR-9 alters motor neuron subtype differentiation as well as columnar development of spinal cords in chick embryos (51). Although the involvement of miR-9 in AECOPD has not been reported to date, to the best of our knowledge, it is possible that this miRNA may play a role in the loss of muscle force in AECOPD, and *GOT1* may also be involved in this process.

Finally, another two miRNAs, miR-15a and miR-16, were also found overexpressed in AECOPD patients in the present study. The pathway of Wnt signaling is known to be a promising target for mediating the development of COPD (52). A knockout of *Wnt2* gene would induce lung hypoplasia and pulmonary hemorrhage caused by the abnormal muscle cells (53). Notably, miR-15a and miR-16-1 are reported to inhibit Wnt signaling (54). Thus, we hypothesized that both miR-15a and miR-16 play key roles in preventing the progression of AECOPD.

In conclusion, the present study identified some DEGs, such as *NCL*, *GOT1*, *SOD2*, *KLF6*, *XRCC5*, *TSPO* and *TMOD1*, and several miRNAs (e.g., miR-1, miR-9 and miR-23a) which may be associated with the pathomechanism of AECOPD. Among them, *SOD2*, *KLF6* and *XRCC5* may be involved in AECOPD due to infection via immune system development. The present study provides in-depth knowledge of the pathogenesis underlying the loss of muscle force during an acute exacerbation of COPD, despite using non-experimental methods. Since the public microarray data used in this study comes from a small sample size - 4 male patients with acute COPD and 5 male patients with stable COPD - it is necessary to validate our findings using experimental methods in a Chinese population with a larger sample size.

References

- Vestbo J, Hurd SS, Agustí AG, Jones PW, Vogelmeier C, Anzueto A, Barnes PJ, Fabbri LM, Martinez FJ, Nishimura M, *et al*: Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 187: 347-365, 2013.
- Uzun S, Djamin R, Hoogsteden H, Aerts J and van der Eerden M: Acute exacerbations of chronic obstructive pulmonary disease. In: *Oncogenesis, Inflammation and Parasitic Tropical Diseases of the Lung*. Kayembe JM (ed). InTech, Chapter 4, pp77-98, 2013. <http://dx.doi.org/10.5772/54867>.
- Carrillo A, Ferrer M, Gonzalez-Diaz G, Lopez-Martinez A, Llamas N, Alcazar M, Capilla L and Torres A: Noninvasive ventilation in acute hypercapnic respiratory failure caused by obesity hypoventilation syndrome and chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 186: 1279-1285, 2012.
- Kim V, Rogers TJ and Criner GJ: New concepts in the pathobiology of chronic obstructive pulmonary disease. *Proc Am Thorac Soc* 5: 478-485, 2008.
- Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, Abraham J, Adair T, Aggarwal R, Ahn SY, *et al*: Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380: 2095-2128, 2012.
- Woodhead M, Blasi F, Ewig S, Garau J, Huchon G, Ieven M, Ortqvist A, Schaberg T, Torres A, van der Heijden G, *et al*: Joint Taskforce of the European Respiratory Society and European Society for Clinical Microbiology and Infectious Diseases: Guidelines for the management of adult lower respiratory tract infections - full version. *Clin Microbiol Infect* 17 (Suppl 6): E1-E59, 2011.
- Coventry PA, Gemmell I and Todd CJ: Psychosocial risk factors for hospital readmission in COPD patients on early discharge services: a cohort study. *BMC Pulm Med* 11: 49, 2011.
- Swallow EB, Reyes D, Hopkinson NS, Man WD, Porcher R, Cetti EJ, Moore AJ, Moxham J and Polkey MI: Quadriceps strength predicts mortality in patients with moderate to severe chronic obstructive pulmonary disease. *Thorax* 62: 115-120, 2007.
- Rabinovich RA, Bastos R, Ardite E, Llinàs L, Orozco-Levi M, Gea J, Vilaró J, Barberà JA, Rodríguez-Roisin R, Fernández-Checa JC and Roca J: Mitochondrial dysfunction in COPD patients with low body mass index. *Eur Respir J* 29: 643-650, 2007.
- Perotin JM, Dury S, Renois F, Deslee G, Wolak A, Duval V, De Champs C, Lebargy F and Andreoletti L: Detection of multiple viral and bacterial infections in acute exacerbation of chronic obstructive pulmonary disease: a pilot prospective study. *J Med Virol* 85: 866-873, 2013.
- Remels AH, Gosker HR, van der Velden J, Langen RC and Schols AM: Systemic inflammation and skeletal muscle dysfunction in chronic obstructive pulmonary disease: state of the art and novel insights in regulation of muscle plasticity. *Clin Chest Med* 28: 537-552, vi, 2007.
- Ezzie ME, Crawford M, Cho JH, Orellana R, Zhang S, Gelinas R, Batte K, Yu L, Nuovo G, Galas D, *et al*: Gene expression networks in COPD: microRNA and mRNA regulation. *Thorax* 67: 122-131, 2012.
- Van Pottelberge GR, Mestdagh P, Bracke KR, Thas O, van Durme YM, Joos GF, Vandesompele J and Brusselle GG: MicroRNA expression in induced sputum of smokers and patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 183: 898-906, 2011.
- Akbas F, Coskunpinar E, Aynaci E, Oltulu YM and Yildiz P: Analysis of serum micro-RNAs as potential biomarker in chronic obstructive pulmonary disease. *Exp Lung Res* 38: 286-294, 2012.
- Lewis A, Riddoch-Contreras J, Natanek SA, Donaldson A, Man WD, Moxham J, Hopkinson NS, Polkey MI and Kemp PR: Downregulation of the serum response factor/miR-1 axis in the quadriceps of patients with COPD. *Thorax* 67: 26-34, 2012.
- Crul T, Testelmans D, Spruit MA, Troosters T, Gosselink R, Geeraerts I, Decramer M and Gayan-Ramirez G: Gene expression profiling in vastus lateralis muscle during an acute exacerbation of COPD. *Cell Physiol Biochem* 25: 491-500, 2010.
- Barrett T, Troup DB, Wilhite SE, Ledoux P, Rudnev D, Evangelista C, Kim IF, Soboleva A, Tomashevsky M and Edgar R: NCBI GEO: Mining tens of millions of expression profiles - database and tools update. *Nucleic Acids Res* 35: D760-D765, 2007.
- Davis S and Meltzer PS: GEOquery: a bridge between the Gene Expression Omnibus (GEO) and BioConductor. *Bioinformatics* 14: 1846-1847, 2013.
- Gautier L, Cope L, Bolstad BM and Irizarry RA: affy - analysis of Affymetrix GeneChip data at the probe level. *Bioinformatics* 20: 307-315, 2004.
- Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, Ellis B, Gautier L, Ge Y, Gentry J, *et al*: Bioconductor: Open software development for computational biology and bioinformatics. *Genome Biol* 5: R80, 2004.
- Šmídl V and Quinn A: The variational Bayes method in signal processing. In: *Signals and Communication Technology*. Springer Berlin Heidelberg, 2006.
- Szekely GJ and Rizzo ML: Hierarchical clustering via joint between-within distances: extending Ward's minimum variance method. *J Classif* 22: 151-183, 2005.
- Mukherjee S, Chen Z and Gangopadhyay A: A privacy-preserving technique for Euclidean distance-based mining algorithms using Fourier-related transforms. *VLDB J* 15: 293-315, 2006.
- Dennis G Jr, Sherman BT, Hosack DA, Yang J, Gao W, Lane HC and Lempicki RA: DAVID: Database for annotation, visualization, and integrated discovery. *Genome Biol* 4: 3, 2003.
- Jensen LJ, Kuhn M, Stark M, Chaffron S, Creevey C, Muller J, Doerks T, Julien P, Roth A, Simonovic M, *et al*: STRING 8 - a global view on proteins and their functional interactions in 630 organisms. *Nucleic Acids Res* 37: D412-D416, 2009.
- Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B and Ideker T: Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 13: 2498-2504, 2003.

27. Fujita Y, Hayashida K, Nagai M, Inoue S, Matsumoto H, Okabe N, Reiprich TH, Sarazin CL and Takizawa M: Suzaku observation of the Ophiuchus galaxy cluster: one of the hottest cool core clusters. *Publ Astron Soc Jpn Nihon Tenmon Gakkai* 60: 1133-1142, 2008.
28. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES and Mesirov JP: Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci USA* 102: 15545-15550, 2005.
29. Plaisier CL, Pan M and Baliga NS: A miRNA-regulatory network explains how dysregulated miRNAs perturb oncogenic processes across diverse cancers. *Genome Res* 22: 2302-2314, 2012.
30. Wu X, Sun X, Chen C, Bai C and Wang X: Dynamic gene expressions of peripheral blood mononuclear cells in patients with acute exacerbation of chronic obstructive pulmonary disease: a preliminary study. *Crit Care* 18: 508, 2014.
31. Gabellini D, Green MR and Tupler R: Inappropriate gene activation in FSHD: a repressor complex binds a chromosomal repeat deleted in dystrophic muscle. *Cell* 110: 339-348, 2002.
32. van Deutekom JC, Wijnga C, van Tienhoven EA, Gruter AM, Hewitt JE, Padberg GW, van Ommen GJ, Hofker MH and Frants RR: FSHD associated DNA rearrangements are due to deletions of integral copies of a 3.2 kb tandemly repeated unit. *Hum Mol Genet* 2: 2037-2042, 1993.
33. Rao PK, Kumar RM, Farkhondeh M, Baskerville S and Lodish HF: Myogenic factors that regulate expression of muscle-specific microRNAs. *Proc Natl Acad Sci USA* 23: 8721-8726, 2006.
34. Puig-Vilanova E, Aguiló R, Rodríguez-Fuster A, Martínez-Llorens J, Gea J and Barreiro E: Epigenetic mechanisms in respiratory muscle dysfunction of patients with chronic obstructive pulmonary disease. *PLoS One* 9: e111514, 2014.
35. Barreiro E, de la Puente B, Minguella J, Corominas JM, Serrano S, Hussain SN and Gea J: Oxidative stress and respiratory muscle dysfunction in severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 171: 1116-1124, 2005.
36. Zelko IN, Mariani TJ and Folz RJ: Superoxide dismutase multigene family: A comparison of the CuZn-SOD (SOD1), Mn-SOD (SOD2), and EC-SOD (SOD3) gene structures, evolution, and expression. *Free Radic Biol Med* 33: 337-349, 2002.
37. Togliatto G, Trombetta A, Dentelli P, Cotogni P, Rosso A, Tschöp MH, Granata R, Ghigo E and Brizzi MF: Unacylated ghrelin promotes skeletal muscle regeneration following hindlimb ischemia via SOD-2-mediated miR-221/222 expression. *J Am Heart Assoc* 2: e000376, 2013.
38. Wilson SR, Joshi AD and Elferink CJ: The tumor suppressor Kruppel-like factor 6 is a novel aryl hydrocarbon receptor DNA binding partner. *J Pharmacol Exp Ther* 345: 419-429, 2013.
39. Mgbemena V, Segovia J, Chang TH and Bose S: KLF6 and iNOS regulates apoptosis during respiratory syncytial virus infection. *Cell Immunol* 283: 1-7, 2013.
40. Taccioli GE, Gottlieb TM, Blunt T, Priestley A, Demengeot J, Mizuta R, Lehmann AR, Alt FW, Jackson SP and Jeggo PA: Ku80: product of the XRCC5 gene and its role in DNA repair and V(D)J recombination. *Science* 265: 1442-1445, 1994.
41. Brody JS and Spira A: State of the art. Chronic obstructive pulmonary disease, inflammation, and lung cancer. *Proc Am Thorac Soc* 3: 535-537, 2006.
42. Becker T, Vögtle FN, Stojanovski D and Meisinger C: Sorting and assembly of mitochondrial outer membrane proteins. *Biochim Biophys Acta* 1777: 557-563, 2008.
43. Baraldo S, Turato G, Badin C, Bazzan E, Beghé B, Zuin R, Calabrese F, Casoni G, Maestrelli P, Papi A, *et al*: Neutrophilic infiltration within the airway smooth muscle in patients with COPD. *Thorax* 59: 308-312, 2004.
44. Fowler VM, Greenfield NJ and Moyer J: Tropomodulin contains two actin filament pointed end-capping domains. *J Biol Chem* 278: 40000-40009, 2003.
45. Peinado VI, Pizarro S and Barberà JA: Pulmonary vascular involvement in COPD. *Chest* 134: 808-814, 2008.
46. Kong KYI, Owens KS, Rogers JH, Mullenix J, Velu CS, Grimes HL and Dahl R: miR-23A microRNA cluster inhibits B-cell development. *Exp Hematol* 38: 629-640.e621, 2010.
47. De Palma S, Capitanio D, Vasso M, Braghetta P, Scotton C, Bonaldo P, Lochmüller H, Muntoni F, Ferlini A and Gelfi C: Muscle proteomics reveals novel insights into the pathophysiological mechanisms of collagen VI myopathies. *J Proteome Res* 13: 5022-5030, 2014.
48. Thulin P, Wei T, Werngren O, Cheung L, Fisher RM, Grandér D, Corcoran M and Ehrenborg E: MicroRNA-9 regulates the expression of peroxisome proliferator-activated receptor δ in human monocytes during the inflammatory response. *Int J Mol Med* 31: 1003-1010, 2013.
49. Bazzoni F, Rossato M, Fabbri M, Gaudiosi D, Mirolo M, Mori L, Tamassia N, Mantovani A, Cassatella MA and Locati M: Induction and regulatory function of miR-9 in human monocytes and neutrophils exposed to proinflammatory signals. *Proc Natl Acad Sci USA* 106: 5282-5287, 2009.
50. Packer AN, Xing Y, Harper SQ, Jones L and Davidson BL: The bifunctional microRNA miR-9/miR-9* regulates REST and CoREST and is downregulated in Huntington's disease. *J Neurosci* 28: 14341-14346, 2008.
51. Otaegi G, Pollock A, Hong J and Sun T: MicroRNA miR-9 modifies motor neuron columns by a tuning regulation of FoxP1 levels in developing spinal cords. *J Neurosci* 31: 809-818, 2011.
52. Reuter S, Beckert H and Taube C: Take the Wnt out of the inflammatory sails: modulatory effects of Wnt in airway diseases. *Lab Invest* 2: 177-185, 2015.
53. Goss AM, Tian Y, Cheng L, Yang J, Zhou D, Cohen ED and Morrissey EE: Wnt2 signaling is necessary and sufficient to activate the airway smooth muscle program in the lung by regulating myocardin/Mrtf-B and Fgf10 expression. *Dev Biol* 356: 541-552, 2011.
54. Chen CH, Dixon RA, Ke LY and Willerson JT: Vascular progenitor cells in diabetes mellitus: roles of Wnt signaling and negatively charged low-density lipoprotein. *Circ Res* 9: 1038-1040, 2009.