

JAK/STAT3 and Smad3 activities are required for the wound healing properties of *Periplaneta americana* extracts

QIN SONG¹, YUXIN XIE², QIHENG GOU³, XIAOQIANG GUO¹, QIAN YAO¹ and XIAOJUN GOU¹

¹College of Pharmacy and Bioengineering, Chengdu University, Chengdu, Sichuan 610106; ²Cancer Center, West China Hospital, Sichuan University; ³State Key Laboratory of Biotherapy/Collaborative Innovation Center of Biotherapy, West China Hospital, Sichuan University, Chengdu, Sichuan 610041, P.R. China

Received July 6, 2016; Accepted June 13, 2017

DOI: 10.3892/ijmm.2017.3040

Abstract. *Periplaneta americana* extracts (PAEs) play a crucial role in skin wound healing. However, their molecular effects and signaling pathways in regenerating tissues and cells are not clear. In this study, we refined the PAE from *Periplaneta americana* to investigate the mechanisms underlying skin wound healing. The human keratinocyte line HaCaT was selected and a mouse model of deep second-degree thermal burn was established for *in vitro* and *in vivo* studies, respectively. PAE treatment induced the proliferation and migration of HaCaT cells and wound healing in the burn model. Furthermore, the effects of PAE on wound healing were found to depend on the Janus-activated kinase/signal transducer and activator of transcription 3 (JAK/STAT3) pathway and Smad3 activities, according to western blot analysis and immunohistochemical (IHC) assays *in vitro* and *in vivo*. Pretreatment with a STAT3 inhibitor blocked the cell proliferation and migration induced by PAE. The results indicate the wound-healing function of PAE via enhanced JAK/STAT3 signaling and Smad3 activities. Our studies provide a theoretical basis underlying the role of PAE in cutaneous wound healing.

Introduction

Periplaneta americana (Linnaeus) (Blattodea: Blattidae), the largest insect member of the family Blattidae, is one of the most ancient insect groups with the strongest vitality. We explored its bioactive substances and the unique physiological

mechanism. *P. americana* has been used in traditional Chinese medicine (TCM) to eliminate stagnant blood, clear hematocoele, detoxify and promote urination and detumescence (1). *P. americana* has also been exploited as an alternative naturopathic remedy for impaired healing of ulcers, burn wounds, tuberculosis, ulcerative colitis, heart disease and cancer (2-6). The major focus of this study is on the promotion of wound healing.

In the early 1980s, W_{11-a12}, also known as Kangfuxin, was refined from the alcohol extract of *P. americana* (PAE), and was effective against thermal lesions, such as burns and scalds (1,7). PAE derived from *P. americana* dry worm contains peptides, polyols, epidermal growth factors, sticky sugar acids, amino acids and other active substances (8). Recent studies have explored the pharmacological effects and mechanisms of PAE in wound healing. Histopathological studies suggest that the necrotic tissue and inflammatory exudate induced by thermal lesions or chemical damages were significantly reduced after PAE treatment in rat and rabbit models (9). In addition, increased epithelial repair area, follicular regeneration, early fibrous tissue repair and rapid decrease of inflammatory cells were detected in the PAE-treated groups. PAE also promoted the synthesis and secretion of extracellular cell matrix (ECM) in the wound (10); activated cellular functions by accelerating the opening of macrophage ion channels (11); increased the number of neutrophils, improved the granulocyte chemotactic function and spontaneous movement, resulting in wound repair and clearance (12,13).

However, wound healing is a complex process of cell proliferation, migration, matrix synthesis and contraction, and involves various types of cells and regulatory mechanisms. Resident cells (keratinocytes, fibroblasts and endothelial cells) and inflammatory cells participate in wound healing (2,14). Evidence has revealed that several signaling pathways are associated with wound healing via triggering their target gene expression, such as the Janus-activated kinase/signal transducer and activator of transcription 3 (JAK/STAT3) signaling (15-18). In wound healing, cytokines contribute to activate STATs and the activated JAK/STAT3 pathway controls the proliferation and differentiation necessary for wound healing (19,20). Furthermore, through activation of JAK/STAT3 signaling cascades, the cytokine induces anti-apoptotic pathways and anti-microbial molecules to help prevent tissue damage and aid in their repair (21-23). In addition, a study demonstrated a critical role for STAT3 in

Correspondence to: Professor Xiaojun Gou, College of Pharmacy and Bioengineering, Chengdu University, Chengdu, Sichuan 610106, P.R. China
E-mail: 402052973@qq.com

Abbreviations: PAEs, *Periplaneta americana* extracts; JAK/STAT3, Janus-activated kinase/signal transducer and activator of transcription 3; IHC, immunohistochemical staining; TCM, traditional Chinese medicine; ECM, extracellular cell matrix

Key words: *Periplaneta americana* extracts, Janus-activated kinase, signal transducer and activator of transcription 3, Smad3, wound healing, signaling pathway

the migration but not proliferation of keratinocytes in wound healing (24). The pivotal roles of Smad3 signaling in cutaneous wound healing have been well documented (15,16). Smad3 binds with a Smad mediator (SMAD4) to form a complex, moving into the nucleus and regulates the expression of genes including those involved in keratinocyte migration, fibroblast infiltration and extracellular matrix construction (25,26). Additionally, Smad3 could balance the reepithelialization and fibrogenesis of the repaired tissues (27,28).

Previous studies have mainly focused on the effect of PAE on cellular repair (10-13). The resident biological functions and cellular signal transduction pathways underlying the healing effect of PAE are not fully established. In the present study, we investigated the biological function and mechanisms of PAE in human keratinocyte cells and rat skin injury models to facilitate the clinical application of PAE.

Materials and methods

Cell lines. The spontaneously immortalized human keratinocyte HaCaT cell line was obtained from the American Type Culture Collection (ATCC, Rockville, MD, USA) and was cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal calf serum (FCS) and 1% penicillin-streptomycin. Culture conditions were maintained constant at 37°C in a 5% CO₂ humidified atmosphere.

Preparation of PAE. *P. americana* was obtained from the Good Agricultural Practice (GAP) breeding base (Sichuan, China). *P. americana* (200 g) powder was extracted with 90% EtOH (1.2 liters) twice at 80°C. After solvent evaporation, the ethanol extract was recovered. The extract (20 g) was suspended in water (200 ml) at 80°C and filtered through a 0.22- μ m filter membrane in appropriate concentrations and stored at -20°C until use. HPLC-diode array detector (HPLC-DAD) was used for the study of *P. americana* extract. Diamonsil C18 (250 \times 4.6 mm; 5 μ m) was selected as the chromatography column. The optimized mobile phase consisted of solvent A (3% v/v methanol in water containing 0.07% v/v acetic acid) and solvent B (methanol). The following gradient was used to represent time (min)/mobile phase A (%)/mobile phase B (%): 0.0/100/0, 10/100/0, 20/70/30, 21/50/50, 35/0/100, at a flow rate of 0.6 ml/min at 25°C, and 254 nm in 10 μ l injection volume.

In vitro proliferation, migration and wound healing assays. Proliferation was determined using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. Cells were seeded in a volume of 200 μ l (2,000 cells/well) on 96-well plates after cultivation with or without PAE. The culture medium containing serum was replaced by MTT every 24 h. A final MTT concentration of 0.5 mg/ml was added to the wells followed by incubation for 4 h at 37°C. The supernatant was discarded and replaced with dimethyl sulfoxide (DMSO) (150 μ l/well). The optical densities (OD) were measured at 570 nm with a microplate reader (Bio-Rad, Hercules, CA, USA). The experiment was repeated in triplicate.

Cell migration was analyzed in a 24-well plate with Millicell Cell Culture Insert, using an 8- μ m pore size polycarbonate membrane (Millipore Corporation, Billerica, MA,

USA). After 48 h of cultivation in a medium with or without PAE (0.3125 mg/ml), the cells (2 \times 10⁴/well) were transferred into the upper chamber with the non-coated membrane and suspended in 200 μ l of serum-free DMEM. In the lower chamber, 600 μ l of the medium supplemented with 10% fetal bovine serum was added. After incubation for 24 h, the chambers were fixed with 4% paraformaldehyde for 20 min, and stained with hematoxylin for 15 min. Images were captured with an optical microscope.

The wound healing assay was performed by seeding the cell cultures on a 6-well plate and grown to 90% confluency, followed by 48 h of starvation in serum-free medium with or without PAE (0.3125 mg/ml). The culture medium was removed and the monolayers were scratched using a 200- μ l pipette to create a uniform cell-free wound area. Debris was removed by gently washing with sterile phosphate-buffered saline (PBS). Cell migration into the wound area was monitored and photographed at 0, 24 and 48 h using an optical microscope.

Western blot analysis and antibodies. Cells were lysed with radioimmunoprecipitation assay (RIPA) buffer and phosphatase inhibitors. The protein concentration was measured using the bicinchoninic acid (BCA)-protein quantification assay (Beyotime Biotechnology, Shanghai, China). Normalized lysates (30 μ g) were separated by electrophoresis on 8-12% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) gel and transferred to a PVDF membrane (Millipore Corporation). The membrane was blocked for 1 h at room temperature and incubated overnight at 4°C with primary antibody. After three washes with TBST, the membrane was incubated with horseradish peroxidase-(HRP)-conjugated IgG. Signals were visualized with enhanced chemiluminescence (ECL; Millipore Corporation). Primary antibodies against Smad3 (#9523), phospho-Stat3 (#9145), Stat3 (#12640), JAK1 (#3344), JAK2 (#3230) (dilution, 1:1,000; Cell Signaling Technology, Beverly, MA, USA), phospho-Smad3 (ab52903), phospho-JAK2 (ab32101) (dilution, 1:1,000; Abcam, Cambridge, MA, USA), phospho-JAK1 (sc-101716; dilution, 1:1,000; Santa Cruz Biotechnology, Inc., Dallas, TX, USA) were used.

Immunofluorescence staining. Each group of HaCaT cells was washed with PBS followed by fixation with 4% paraformaldehyde (pH 7.4) in 6-well plates. Cells were incubated with 0.5% Triton X-100 for 30 min at room temperature, and treated with 5% BSA for 1 h. Cells were incubated with the following primary antibodies: Smad3 (#9523; dilution, 1:100), phospho-Stat3 (#9145; dilution, 1:100), Stat3 (#12640; dilution, 1:500) (all from Cell Signaling Technology), and phospho-Smad3 (ab52903; dilution, 1:100; Abcam) overnight at 4°C. The cells were incubated with the corresponding fluorescent dye-conjugated secondary antibodies (#4412; dilution, 1:200; Cell Signaling Technology) at 37°C for 1 h, and protected from light. The cells were visualized via fluorescence microscopy.

Rat skin injury models. Healthy adult C57 male mice were purchased from the West China School of Preclinical and Forensic Medicine, Sichuan University, Sichuan, China. All the experiments were conducted according to the Guide for the Care and Use of Laboratory Animals at the Animal

Experimental Center of Sichuan University. We obtained ethical approval of the Medical Ethics Committee of Sichuan University with approval no. K2016033. Thermal injuries were created with a solid aluminum bar measuring 10 mm in diameter, and pre-heated in boiling water at a temperature of 100°C. The bar was maintained symmetrically in contact with the skin on the dorsal flank for 15 sec (29). The right skin of the dorsal flank was wiped with PAE (original fluid, 5 mg/ml), while the left was treated with normal saline as the untreated control. The daily treatment regimen lasted for 21 days. After injury, to minimize the suffering of the animals, each mouse was housed individually and fed with sterilized food and tap water to prevent infection. The wound healing rates were measured on days 0, 7, 14 and 21 after injury, and the complete wound healing time was calculated as follows: Healing rate = (original wound area - non-healing wound area)/original wound area (30). Mice were sacrificed after 3 weeks.

Immunohistochemistry. All the immunohistochemical assays were conducted following the manufacturer's instructions. Briefly, the skin tissues were fixed in formalin and embedded in paraffin. Consecutive paraffin sections (4- μ m) of the tissue samples were prepared and incubated overnight at 4°C with primary antibodies followed by incubation with peroxidase-labeled polymer conjugated to goat anti-rabbit immunoglobulins (EnVision/HRP; Dako, Glostrup, Denmark). Primary antibodies against Smad3 (#9523), phospho-Stat3 (#9145), Stat3 (#12640), JAK1 (#3344), JAK2 (#3230) (dilution, 1:1,000; all from Cell Signaling Technology), phospho-Smad3 (ab52903), phospho-JAK2 (ab32101) (dilution, 1:1,000; both from Abcam), phospho-JAK1 (sc-101716; dilution 1:1,000; Santa Cruz Biotechnology, Inc.) were used.

Statistical analysis. Statistical analyses were performed using SPSS 19.0 (SPSS, Inc., Chicago, IL, USA) or Prism 5.0 (GraphPad Software, La Jolla, CA, USA). Quantitative data were analyzed using a two-tailed Student's t-test, and one-way analysis of variance (ANOVA) followed by Dunnett's multiple comparison post-test. Differences were considered statistically significant at $P < 0.05$, $P < 0.05$ and $P < 0.01$.

Results

Chromatographic separation of PAE. PAE contained polyalcohols, amino acids, pyrimidines and proteoglycans. Analysis of PAE revealed the presence of uracil (retention time 7.626 min, 1.145 mg/g), hypoxanthine (retention time 11.707 min, 4.253 mg/g), and inosine (retention time 20.427 min, 8.156 mg/g) (Fig. 1).

PAE increases cell proliferation and migration of human keratinocytes in vitro. The MTT assay was used to confirm the proliferation of HaCaT cells in the presence of PAE at three time-points. The low (0.3125 mg/ml) dose of the extract induced optimal growth in HaCaT cells at 24, 48 and 72 h, respectively ($p < 0.05$) (Fig. 2A). Conversely, high doses of PAE (1.25 and 2.5 mg/ml) inhibited cell proliferation. Furthermore, the treatment with the extract caused obvious proliferation after 48 h of treatment. Transwell and wound healing assays were used to determine HaCaT cell migration in skin. Fig. 2B




Figure 1. Chromatographic separation of *Periplaneta americana* extract (PAE). (A) HPLC chromatogram of authentic standards tested. (B) HPLC chromatogram of PAE at 254 nm: peak 1: uracil (1.145 mg/g), peak 2: hypoxanthine (4.253 mg/g), and peak 3: inosine (8.156 mg/g). Identification was based on retention time, and UV spectra compared with commercial standards.

shows that compared with the control group, cell migration in the PAE-treated group increased significantly ($p < 0.01$). Furthermore, consistent with the results from the Transwell assay, data from the wound healing assay also showed significantly improved migration of wound closure in the PAE treatment group compared with the control group in the HaCaT cells ($p < 0.01$) (Fig. 2C).

PAE enhances the JAK/STAT3 signaling pathway and Smad3 activities in human keratinocytes. To better understand the signaling pathway or mechanism underlying the PAE-mediated regulation of HaCaT cells, the signaling molecules associated with proliferation and migration were screened. After PAE treatment, the expression levels of p-JAK1, p-JAK2 and its downstream molecule p-STAT3 were markedly upregulated in the HaCaT cells, while the total protein levels (JAK1, JAK2 and STAT3) were unchanged ($p < 0.01$) (Fig. 3A). In addition, phosphorylation of Smad3 (p-Smad3) was also significantly increased, while Smad3 was not affected ($p < 0.01$) (Fig. 3B). However, nuclear factor- κ B (NF- κ B)/p65 and β -catenin expression was not distinctly different between the PAE treatment and the control groups of HaCaT cells ($p > 0.05$) (Fig. 3C).

Effects of STAT3 inhibitor on PAE-mediated proliferation and migration of HaCaT cells. To further confirm that PAE increases cell proliferation and migration of human keratinocytes via JAK/STAT3 signaling, the inhibitor of STAT3




Figure 2. Effects of different doses of *Periplaneta americana* extract (PAE) on differentiation at different time-points and effects on migration in HaCaT cells. (A) Left: the survival fraction of HaCaT cells following treatment with different doses of the extract for 24 h (normalized to control group); middle: the survival fraction of HaCaT cells following treatment with different doses of the extract for 48 h (normalized to control group); right: the survival fraction of HaCaT cells following treatment with different doses of the extract for 72 h (normalized to control group). (B) HaCaT cells were treated with or without PAE (0.3125 mg/ml) for 48 h, and cells treated without PAE served as controls. HaCaT cells were plated on Matrigel-coated membranes for migration assays and incubated for 24 h. Migratory cells were counted in 10 random fields and expressed as the average number of cells/field. (C) Wound healing assay: images obtained at 0, 24 and 48 h after scratching. Data represent mean \pm SD. ** $p < 0.01$ vs. the control group.

was used to verify the effects of PAE on cell proliferation and migration. A STAT3 inhibitor, stattic, was used to block STAT3 activation. Phosphorylation of STAT3 was decreased after treatment with stattic in a dose-dependent manner in the HaCaT cells. However, the total expression of STAT3 was not changed. Furthermore, when cells were treated with PAE, after inhibition of STAT3 phosphorylation by stattic, the p-STAT3 levels remained unchanged (Fig. 4A and B). Furthermore, as shown in Fig. 4C and D, PAE-induced cell growth and migration were abrogated by pretreatment with stattic in the HaCaT cells. The result confirmed that the PAE extract promoted HaCaT cell proliferation and migration by activating the JAK/STAT3 signaling pathway.

PAE increases phosphorylation of STAT3 and Smad3 expression and alters their distribution. The distribution pattern of STAT3 and Smad3 was investigated using fluorescence

immunostaining. The results demonstrated that PAE treatment increased STAT3 phosphorylation in the cytoplasm followed by nuclear translocation (Fig. 5A). By contrast, the total expression of STAT3 was constant irrespective of PAE stimulation in the HaCaT cells (Fig. 5B). Furthermore, phosphorylation of Smad3 (p-Smad3) was also significantly increased in PAE-treated cells, while total Smad3 remained unchanged (Fig. 5C and D). The altered phosphorylation of STAT3 and Smad3 suggested activation of the JAK/STAT3 and Smad3 signaling cascade by PAE in HaCaT cells.

PAE promotes wound healing in vivo. To establish the *in vivo* effect of PAE on cutaneous wound healing, we established a mouse model of deep second-degree thermal burn (16). Skin lesions were not prominent initially, but eventually appeared and worsened. The results demonstrated that the burn wound in the PAE (original fluid, 50 mg/ml)-treated side (right




Figure 3. Effects of *Periplaneta americana* extract (PAE) on cell proliferation and migration-related signaling molecules in HaCaT cells. HaCaT cells were treated with or without PAE (0.3125 mg/ml) for 48 h, and cells treated without PAE served as the controls. (A) The protein expression of Janus-activated kinase/signal transducer and activator of transcription 3 (JAK/STAT3) signaling pathway, (B) Smad3 and its phosphorylation, (C) NF-κB/p65 and β-catenin were determined by western blotting. β-actin was used as a loading control. Data represent mean \pm SD. ** $p < 0.01$ vs. the control group. n.s. not significant.

side) showed significant fibroblast proliferation and fibrosis resulting in scar tissue formation compared with the control (left side) (Fig. 6A). Histopathological analysis revealed an increase in epithelial repair, follicle regeneration and additional fibrous tissues in the PAE-treated groups (Fig. 6B). The healing rates and times of skin wounds without and with PAE treatment were sequentially recorded and evaluated. The wound healing rates were measured on days 0, 7, 14 and 21 after burn, and the complete wound healing time was calculated. The healing rate of the treated group was significantly higher than that of the control group at different time-points ($p < 0.05$) (Table I and Fig. 6C). The results suggest that PAE accelerates the healing of burn wounds.

JAK/STAT3 signaling and Smad3 activities in PAE-mediated wound healing. To further confirm the signaling cascade and mechanism of PAE-induced regulation of wound healing, immunohistochemical (IHC) and western blot analysis were used to assay the JAK/STAT3 signaling and phosphorylation of Smad3 in burn wound specimens. Enhanced nuclear staining

of p-JAK1, p-JAK2, p-STAT3 and p-Smad3 was detected in the PAE treatment group compared with the control group. By contrast, the nuclear expression of JAK1, JAK2, STAT3 and Smad3 proteins showed no significant difference between the control wounds and the treated wounds (Fig. 7A). Consistent with the IHC findings, the western blot analysis further validated the increased expression of p-JAK1, p-JAK2, p-STAT3, p-Smad3, but not JAK1, JAK2, STAT3 and Smad3 proteins (Fig. 7B).

Discussion

The beneficial effects of *Periplaneta americana* extracts (PAEs) on skin wound healing have been adequately demonstrated, suggesting their potential role as bioactive agents for alternative treatment of skin wounds (1,2). However, the role of PAE in the regulation of target cells and the molecular mechanisms of wound healing in the skin are still unclear. In the present study, we extracted and purified PAE from *P. americana* to explore the underlying mechanism. We established skin wound healing models *in vitro* and *in vivo* to demonstrate that




Figure 4. Effects of STAT3 inhibitor on *Periplaneta americana* extract (PAE)-mediated signal transduction, proliferation and migration in HaCaT cells. (A) HaCaT cells were pretreated with different concentrations of the STAT3 inhibitor statistic (0, 1, 5 and 10 μM) for 20 min, and p-STAT3, total STAT3 and β -actin were detected by western blotting. (B) HaCaT cells were incubated in the medium containing PAE at the indicated concentrations for 48 h after pretreatment with 10 μM statistic for 20 min. The expression of p-STAT3, total STAT3 and β -actin was detected by western blotting. HaCaT cells were incubated with a medium containing PAE at the indicated concentrations for 48 h after pretreatment with statistic (10 μM) for 20 min. (C) Cell proliferation was determined by MTT and (D) migration was tested by Transwell assay. ** $p < 0.01$.

PAE promoted the proliferation and migration of human keratinocyte HaCaT cells and wound healing in a mouse model. The effects of PAE were mediated via the JAK/STAT3 and Smad3 signaling pathways *in vitro* and *in vivo*.

Wound healing comprises four distinct but overlapping phases: hemostasis, inflammation, proliferation and remodeling (2,31). Keratinocytes are a major skin component and an important determinant of wound healing efficiency (15,32,33). In this study, we found that PAE promoted the proliferation and migration of human immortalized keratinocyte HaCaT cells. To confirm the pharmacological effects of PAE and establish the optimal extraction process, the human immortalized keratinocyte line HaCaT was used as an *in vitro* wound healing model for the evaluation of the cellular and molecular effects of PAE. The results showed an increase in proliferation and migration of

keratinocytes after treatment with PAE (0.3125 mg/ml) for 48 h. In addition, PAE significantly increased the healing rates and time by enhancing the epithelial repair, follicle regeneration and fibrous tissue proliferation in cutaneous thermal burns *in vivo*.

Although the effects of PAE on wound healing have long been recognized, the underlying molecular mechanisms remain largely unknown. Studies suggest the role of numerous signaling pathways in cutaneous wound healing (15-18). In this study, the expression of JAK/STAT3 signaling and the activation of Smad3, NF- κB /p65 and β -catenin *in vitro* and in wound tissues *in vivo* were evaluated by western blot analysis and validated immunohistochemically. We found that the expression and nuclear translocation of p-JAK1, p-JAK2 and its downstream p-STAT3 were markedly upregulated in the HaCaT cells and wound tissues after PAE treatment. In addition, the levels of




Figure 5. Immunofluorescence analysis of p-STAT3/STAT3 and p-Smad3/Smad3 protein. Cells were treated with *Periplaneta americana* extract (PAE) for 48 h, followed by fixation and incubation to determine immunofluorescence. The antibody against (A and B) p-STAT3/STAT3 and (C and D) p-Smad3/Smad3 presented green fluorescence, whereas a blue fluorescent signal was produced by DAPI staining of the cell nuclei. Areas of overlap between the green and the blue fluorescence were observed in the merged images. The arrows in (A) indicate p-STAT3 expression in the nucleus. The arrows in (B) indicate STAT3 expression in the nucleus and cytoplasm. The arrows in (C) indicate p-Smad3 expression in the nucleus. The arrows in (D) indicate Smad3 expression in the nucleus and cytoplasm.

p-Smad3 expression were increased in the treated cells and tissues, suggesting the active role of these signaling pathways in PAE-mediated wound healing. However, NF- κ B and Wnt signaling appeared to be minimally activated irrespective of PAE treatment because of limited expression of NF- κ B/P65 and β -catenin upregulation, or nuclear translocation.

Table I. Healing rates at different time-points after burn (%).

Group	7 day	14 day	21 day
Control	28.17 \pm 7.03	58.73 \pm 4.43	87.72 \pm 2.88
Treatment	37.34 \pm 8.39 ^a	83.61 \pm 4.61 ^a	97.69 \pm 0.82 ^a

^aThe healing rate of the treatment group was compared with the control group at varied time-points, $p < 0.05$.




Figure 6. Effects of *Periplaneta americana* extract (PAE) on wound healing of skin thermal burn *in vivo*. In the mouse model, the right burn wound of the dorsal flank was wiped with PAE (original fluid), while the left was treated daily with normal saline. (A) Representative images of burn wounds were monitored on days 0, 7, 14 and 21. (B) Histological H&E staining of burnt tissues on days 0, 14 and 21 after burn injury, respectively. (magnification, x100 and x400). (C) The wound healing rates. ^a $p < 0.05$ vs. the control group.

STAT3 is a member of the latent transcription factor family that acts as a downstream effector of cytokine and growth factor receptor signaling. The canonical JAK/STAT signaling pathway involves the activation of JAK or growth factor receptor kinases, phosphorylation of STAT proteins,




Figure 7. Effects of *Periplaneta americana* extract (PAE) on Janus-activated kinase/signal transducer and activator of transcription 3 (JAK/STAT3) signaling and Smad3 activities in burn wound specimens *in vivo*. (A) Immunohistochemical (IHC) analysis of JAK/STAT3 signaling cascade, Smad3 and its phosphorylation in PAE-treated or control tissues. Representative images are shown (magnification, x400). (B) Western blotting of JAK/STAT3 signaling proteins, Smad3 and its phosphorylation using β -actin as a loading control. Data are expressed as mean \pm SD. The arrows in (A) indicate p-JAK1/JAK1, p-JAK2/JAK2, p-STAT3/STAT3, p-Smad3/Smad3 expression in the nucleus and cytoplasm, respectively.

their dimerization and translocation into the nucleus whereas STATs act as transcription factors with pleiotropic downstream effects (34). The pivotal roles of JAK/STAT3 signaling and TGF- β /Smad3 in cutaneous wound healing are well known (15,35-37). The proliferation and differentiation of keratinocytes during wound healing are regulated by cytokines and chemokines, which are secreted by resident and inflammatory cells and activate the transcription factor

STAT3 (38,39). Moreover, the specific ablation of STAT3 in the follicular and interfollicular keratinocytes resulted in impaired wound healing (24). Our results suggested that the enhanced phosphorylation and activation of JAK/STAT3 and Smad3 by the active ingredients in PAE accelerates wound healing efficiency. However, the composition of active ingredients and the downstream factors need to be defined to elucidate the mechanism of PAE.

Consequently, PAE promoted wound healing *in vitro* and *in vivo* experimental models, which were strongly correlated with the activation of JAK/STAT3 and Smad3 signaling. Pretreatment with statin inhibited HaCaT cell proliferation and migration induced by PAE. Furthermore, the enhanced activities of JAK/STAT3 and Smad3 pathways may represent major molecular targets of PAE, in accelerated wound healing.

Acknowledgements

This study was supported by the National Natural Science Foundation of China (no. 81403194).

References

- Luo TS, Gao MT, Ma FF, Liu GM and Zhang CG: Research advances in pharmacological action and clinical application of *Periplaneta americana*. *Agric Sci Technol* 13: 888-892, 2012.
- Chen XH, Ran XZ, Sun RS, Shi CM, Su Y, Guo CH and Cheng TM: Protective effect of an extract from *Periplaneta americana* on hematopoiesis in irradiated rats. *Int J Radiat Biol* 85: 607-613, 2009.
- Yan MC, Yang H and Yang RM: Observation of the clinical effects of W_{11-a12}. *Yunnan Med* 3: 138-140, 1987.
- Pan XL: The observation and nurse care with Kangfuxin local medicine - change in treating diabetes acromelic gangrene. *J Nurse Train* 11: 26-27, 1996.
- Wang HL and Liu J: Clinical observation of patients burned at III degree in small proportion of total surface area treated with Kangfuxin. *J Binzhou Med Coll* 22: 368, 1999.
- Lin Q, Cao D, Yang YQ, *et al*: Study on action of Kangfuxin solution on experimental gastric ulcer. *Chin Trad Pat Med* 23: 122-124, 2001.
- Du GM and Li WL: The clinical and laboratory studies about effects of Kangfuxin on superoxide dismutase and cell immune function of old persons. *Yunnan J Chin Trad Med* 10: 32-33, 1989.
- Man HX, Huang L, Na KG, *et al*: Research progress of chemical component and biological activity in medicinal *Periplaneta americana*. *Anti-infect Pharmacol* 11: 403-407, 2014.
- Wang ZY, Huang XH, Xie YK, Chen S and Wang S: Influence of Kangfuxin liquid on the wound healing of experimental animal burns and scalds. *J Tradit Chin Med* 52: 1316-1317, 2011.
- Shu CX, Cheng TM and Yan GM: Effects of whole body irradiation on several components of skin wound extracellular matrix and the repair-promoting action of W_{11-a12}. *Chin J Traumatol* 11: 604-607, 2001.
- Shu CX, Ye BL, Cheng TM and Xiao JS: Effects of irradiation and W_{11-a12} on anion-selective channel of mouse peritoneal macrophage. *Acta Acade Med Milita Tertiae* 23: 290-292, 2001.
- Chen XH, Cheng TM and Ai GP: Effects of systemic irradiation and W_{11-a12} on neutrophils in wounds. *Acta Acade Med Milita Tertiae* 23: 287-289, 2001.
- Chen XH, Sun RS, Cheng TM, *et al*: Neutrophil apoptosis in wound of irradiated rats and effects of W_{11-a12} in accelerating wound healing. *Chin J Clin Rehabil* 9: 108-110, 2005.
- Amadeu TP, Seabra AB, de Oliveira MG and Monte-Alto-Costa A: Nitric oxide donor improves healing if applied on inflammatory and proliferative phase. *J Surg Res* 149: 84-93, 2008.
- Li PN, Li H, Zhong LX, Sun Y, Yu LJ, Wu ML, Zhang LL, Kong QY, Wang SY and Lv DC: Molecular events underlying maggot extract promoted rat *in vivo* and human *in vitro* skin wound healing. *Wound Repair Regen* 23: 65-73, 2015.
- Pakyari M, Farrokhi A, Maharlooee MK and Ghahary A: Critical role of transforming growth factor beta in different phases of wound healing. *Adv Wound Care (New Rochelle)* 2: 215-224, 2013.
- Ren X, Ge M, Qin X, Xu P, Zhu P, Dang Y, Gu J and Ye X: S100a8/NF- κ B signal pathway is involved in the 800-nm diode laser-induced skin collagen remodeling. *Lasers Med Sci* 31: 673-678, 2016.
- Shi Y, Shu B, Yang R, Xu Y, Xing B, Liu J, Chen L, Qi S, Liu X, Wang P, *et al*: Wnt and Notch signaling pathway involved in wound healing by targeting c-Myc and Hes1 separately. *Stem Cell Res Ther* 6: 120, 2015.
- Tokumaru S, Sayama K, Yamasaki K, Shirakata Y, Hanakawa Y, Yahata Y, Dai X, Tohyama M, Yang L, Yoshimura A, *et al*: SOCS3/CIS3 negative regulation of STAT3 in HGF-induced keratinocyte migration. *Biochem Biophys Res Commun* 327: 100-105, 2005.
- Yasukawa H, Ohishi M, Mori H, Murakami M, Chinen T, Aki D, Hanada T, Takeda K, Akira S, Hoshijima M, *et al*: IL-6 induces an anti-inflammatory response in the absence of SOCS3 in macrophages. *Nat Immunol* 4: 551-556, 2003.
- Lejeune D, Dumoutier L, Constantinescu S, Kruijer W, Schuringa JJ and Renauld JC: Interleukin-22 (IL-22) activates the JAK/STAT, ERK, JNK, and p38 MAP kinase pathways in a rat hepatoma cell line. Pathways that are shared with and distinct from IL-10. *J Biol Chem* 277: 33676-33682, 2002.
- Wolk K, Witte E, Witte K, Warszawska K and Sabat R: Biology of interleukin-22. *Semin Immunopathol* 32: 17-31, 2010.
- Yu R, Ding Y, Zhu L, Qu Y, Zhang C, Liu L and Chen L: IL-22 mediates the oral mucosal wound healing via STAT3 in keratinocytes. *Arch Oral Biol* 72: 14-20, 2016.
- Sano S, Itami S, Takeda K, Tarutani M, Yamaguchi Y, Miura H, Yoshikawa K, Akira S and Takeda J: Keratinocyte-specific ablation of Stat3 exhibits impaired skin remodeling, but does not affect skin morphogenesis. *EMBO J* 18: 4657-4668, 1999.
- Penn JW, Grobbelaar AO and Rolfe KJ: The role of the TGF- β family in wound healing, burns and scarring: A review. *Int J Burns Trauma* 2: 18-28, 2012.
- Hong HJ, Jin SE, Park JS, Ahn WS and Kim CK: Accelerated wound healing by smad3 antisense oligonucleotides-impregnated chitosan/alginate polyelectrolyte complex. *Biomaterials* 29: 4831-4837, 2008.
- Biernacka A, Dobaczewski M and Frangogiannis NG: TGF- β signaling in fibrosis. *Growth Factors* 29: 196-202, 2011.
- Werner S, Krieg T and Smola H: Keratinocyte-fibroblast interactions in wound healing. *J Invest Dermatol* 127: 998-1008, 2007.
- Tavares Pereira DS, Lima-Ribeiro MH, de Pontes-Filho NT, Carneiro-Leão AM and Correia MT: Development of animal model for studying deep second-degree thermal burns. *J Biomed Biotechnol* 2012: 460841, 2012.
- Zhang J, La X, Fan L, Li P, Yu Y, Huang Y, Ding J and Xing Y: Immunosuppressive effects of mesenchymal stem cell transplantation in rat burn models. *Int J Clin Exp Pathol* 8: 5129-5136, 2015.
- Diegelmann RF and Evans MC: Wound healing: An overview of acute, fibrotic and delayed healing. *Front Biosci* 9: 283-289, 2004.
- Hata S, Okamura K, Hatta M, Ishikawa H and Yamazaki J: Proteolytic and non-proteolytic activation of keratinocyte-derived latent TGF- β 1 induces fibroblast differentiation in a wound-healing model using rat skin. *J Pharmacol Sci* 124: 230-243, 2014.
- Yamaoka H, Sumiyoshi H, Higashi K, Nakao S, Minakawa K, Sumida K, Saito K, Ikoma N, Mabuchi T, Ozawa A, *et al*: A novel small compound accelerates dermal wound healing by modifying infiltration, proliferation and migration of distinct cellular components in mice. *J Dermatol Sci* 74: 204-213, 2014.
- Levy DE and Darnell JE Jr: Stats: Transcriptional control and biological impact. *Nat Rev Mol Cell Biol* 3: 651-662, 2002.
- Zhu BM, Ishida Y, Robinson GW, Pacher-Zavisin M, Yoshimura A, Murphy PM and Hennighausen L: SOCS3 negatively regulates the gp130-STAT3 pathway in mouse skin wound healing. *J Invest Dermatol* 128: 1821-1829, 2008.
- Murray PJ: The JAK-STAT signaling pathway: Input and output integration. *J Immunol* 178: 2623-2629, 2007.
- Tokumaru S, Sayama K, Shirakata Y, Komatsuzawa H, Ouhara K, Hanakawa Y, Yahata Y, Dai X, Tohyama M, Nagai H, *et al*: Induction of keratinocyte migration via transactivation of the epidermal growth factor receptor by the antimicrobial peptide LL-37. *J Immunol* 175: 4662-4668, 2005.
- Avitabile S, Odorisio T, Madonna S, Eyerich S, Guerra L, Eyerich K, Zambruno G, Cavani A and Cianfarani F: Interleukin-22 promotes wound repair in diabetes by improving keratinocyte pro-healing functions. *J Invest Dermatol* 135: 2862-2870, 2015.
- Nelson AM, Katseff AS, Ratliff TS and Garza LA: Interleukin 6 and STAT3 regulate p63 isoform expression in keratinocytes during regeneration. *Exp Dermatol* 25: 155-157, 2016.