

# Genome-wide DNA methylome alterations in acute coronary syndrome

DANDAN LI<sup>1\*</sup>, JING YAN<sup>1,2\*</sup>, YUNLONG YUAN<sup>1\*</sup>, CHENG WANG<sup>1</sup>, JIA WU<sup>1</sup>,  
QINGWEN CHEN<sup>1</sup>, JIAXI SONG<sup>1</sup> and JUNJUN WANG<sup>1</sup>

<sup>1</sup>Department of Clinical Laboratory, Jinling Hospital, Medical School of Nanjing University;

<sup>2</sup>Jingling Clinical Medical College of Nanjing Medical University, Nanjing, Jiangsu 210002, P.R. China

Received January 23, 2017; Accepted October 17, 2017

DOI: 10.3892/ijmm.2017.3220

**Abstract.** Acute coronary syndrome (ACS) is a common disease with high mortality and morbidity rates. The methylation status of blood DNA may serve as a potential early diagnosis and prevention biomarker for numerous diseases. The present study was designed to explore novel genome-wide aberrant DNA methylation patterns associated with ACS. The Infinium HumanMethylation450 assay was used to examine genome-wide DNA methylation profiles in 3 pairs of ACS and control group samples. Epigenome-wide DNA methylation, genomic distribution, Gene Ontology (GO) term and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analyses were performed. The results were confirmed using methylation-specific polymerase chain reaction (MSP) and Sequenom MassARRAY analyses in ACS, stable coronary artery disease (SCAD) and control samples. A total of 11,342 differentially methylated (DM) 5'-C-phosphate-G-3' (CpG) sites were identified, including 8,865 hypomethylated and 2,477 hypermethylated CpG sites in the ACS group compared with the control samples. They varied in frequency across genomic compartments, but were particularly notable in gene bodies and shores. The results of GO term and KEGG pathway

enrichment analyses revealed that the methylated genes were associated with certain biological processes and pathways. Despite the considerable variability in methylation data, the candidate selected possessed significant methylation alteration in mothers against decapentaplegic homolog 3 (SMAD3) transcription start site 155 (Chr1:67356838-Chr1:67356942). MSP analysis from 81 ACS samples, 74 SCAD samples and 53 healthy samples, and Sequenom MassARRAY analysis, confirmed that differential CpG methylation of SMAD3 was significantly corrected with the reference results of the HumanMethylation450 array. The data identified an ACS-specific DNA methylation profile with a large number of novel DM CpG sites, some of which may serve as candidate markers for the early diagnosis of ACS.

## Introduction

Acute coronary syndrome (ACS) is a type of cardiovascular disease (CVD) that occurs as a result of an acute obstruction in the coronary arteries, and it represents a leading cause of mortality and disability worldwide (1-3). A previous epidemiological survey revealed that ~50% of cardiovascular mortalities are due to ACS (4). Therefore, it is necessary to clarify the underlying biological mechanisms of ACS to enable early diagnosis and treatment of the disease.

The traditional diagnosis of ACS is achieved by a combination of electrocardiogram (ECG) results and the detection of specific serum markers (5). However, the sensitivity of ECGs is generally <50% and the application of serum markers, including creatine kinase isoenzyme MB (CK-MB), cardiac troponin (cTn)I and cTnT, also has limitations (6). Despite the high specificity of cTnT and cTnI, they may only be detected 6-12 h following myocardial ischemia, which is insufficient for early diagnosis and treatment of ACS prior to myocardial injury (7). The development of novel, sensitive and accurate ACS biomarkers are urgently required to enable the early diagnosis and treatment of ACS.

Epigenetics are defined as stable, heritable alterations in gene expression, which are not due to changes in the DNA sequence (8). The role of epigenetics has mainly been evaluated in relation to cancer (9), but selected previous studies have explored epigenetic involvement in CVD development and progression (10-13). DNA methylation-induced genetic

*Correspondence to:* Professor Junjun Wang or Dr Jiayi Song, Department of Clinical Laboratory, Jinling Hospital, Medical School of Nanjing University, 305 East Zhongshan Road, Nanjing, Jiangsu 210002, P.R. China  
E-mail: wangjunjun9202@163.com  
E-mail: strive1225@126.com

\*Contributed equally

**Abbreviations:** ACS, acute coronary syndrome; SCAD, stable coronary artery disease; GO, gene ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes; DM, differentially methylated; MSP, methylation-specific polymerase chain reaction; ECG, electrocardiogram; CK-MB, creatine kinase isoenzyme MB; cTn, cardiac troponin; CVD, cardiovascular disease; TSS, transcription start site; UTR, untranslated region; CAD, coronary artery disease

**Key words:** DNA methylation, genome-wide, alterations, acute coronary syndrome, Infinium HumanMethylation450 assay

repression is a major type of epigenetic modification (14). DNA methylation occurs at the C5 position of cytosine by the covalent modification of a methyl group, which is most commonly found in 5'-C-phosphate-G-3' (CpG) dinucleotides (15). Alterations in genome-wide and site-specific DNA methylation have been described in CVD, although their specific functions are still largely uninvestigated (16,17). Altered DNA methylation (hyper methylation or hypomethylation) is one of the early events in the pathogenesis of ACS (18,19). Hypermethylation of specific promoter regions of crucial genes, such as transcription factor genes, can repress the expression of these genes and alter biological functions, which in turn increases the risk of ACS (20). Furthermore, it has been reported that numerous risk factors, including hypertension (21,22), diabetes (23-26), atherosclerosis (27-30) and smoking (31,32), associated with the development of ACS may cause epigenetic alterations. These previous studies suggested that the detection of DNA methylation alterations may be a promising tool for early ACS prediction and diagnosis. However, systematic analyses of the changes to genome-wide DNA methylation in patients with ACS are still lacking. In the present study, the genome-wide aberrant DNA methylation patterns predominantly associated with ACS were explored. Additionally, the differentially methylated (DM) regions associated with ACS pathogenesis were characterized, and novel DNA methylation biomarkers potentially beneficial to the early diagnosis of ACS were identified.

## Materials and methods

**Study subjects.** The present study recruited 81 patients with ACS, 74 patients with stable coronary artery disease (SCAD) and 53 control subjects. The patients with CAD in the present study were admitted to the Department of Cardiology of Jinling Hospital (Nanjing, China) between January, 2014 and December, 2016, who were undergoing clinically necessary coronary angiography. Angiograms of all patients with CAD revealed  $\geq 50\%$  stenosis in a minimum of one coronary artery. The 81 patients with ACS included patients with acute myocardial infarction and unstable angina, with Braunwald classification II or III who exhibited a positive cardiac biomarker result (cTnI  $\geq 0.090$  ng/ml or cTnT  $\geq 0.014$  ng/ml), acute ischemic-type chest pain (lasting for  $>15$  min, duration from symptom onset to emergency admission within 72 h) and characteristic ECGs. The 74 patients with SCAD had a normal ECG and documented normal left ventricular contractility, with the exception of possible minor nonspecific ST-T features, and they had a  $\geq 1$  year history without any cardiac events or procedures suggestive of ACS. The exclusion criteria for the patients with CAD included mild disease identified on angiography (a stenosis of 10-50% of the luminal diameter in all three coronary arteries), prior coronary revascularization and the presence of renal disease. The 53 control subjects were recruited to the study from routine health examinations at Jinling Hospital over the same period. The control patients had a normal physical examination, ECG and laboratory tests, and were did not show signs of other diseases, including hyperlipidemia, hypertension and diabetes mellitus, or any clinically evident signs of atherosclerosis. The clinicopathological features of all the patients and controls are listed in Table I. The present study was approved by the Ethics Committee of

Jinling Hospital and all the recruited subjects provided written informed consent prior to the study.

**Blood sampling.** Blood samples were collected from the patients with 81 ACS and 74 patients with SCAD in tubes containing EDTA immediately following their hospitalization. The 53 control samples were drawn from the elbow vein of a paired healthy individual on an empty stomach. The samples were stored at  $-80^{\circ}\text{C}$  prior to analysis.

**DNA isolation and sodium bisulfite modification.** Genomic DNA was isolated from 100  $\mu\text{l}$  blood using the DNeasy Blood and Tissue kit (Qiagen, Inc., Valencia, CA, USA) according to the manufacturer's protocol. The quality and quantity of the extracted genomic DNA was analyzed with a NanoDrop<sup>TM</sup> 2000 Spectrophotometer. Bisulfite modification of 1  $\mu\text{g}$  DNA was conducted using an EpiTect Bisulfite kit (Qiagen, Inc.) according to the manufacturer's protocol.

**Infinium HumanMethylation450 assay.** A total of 3 patients with ACS were randomly selected and paired with 3 control subjects for the Infinium HumanMethylation450 assay (Illumina, Inc., San Diego, CA, USA). The Infinium HumanMethylation450 assay was performed according to the manufacturer's standard protocol. Processed methylation chips were scanned using an iScan reader (Illumina, Inc.). Paired control samples were processed on the same chip and all samples were processed at the same time to avoid chip-to-chip variation. The assays were conducted in the Beijing Genomics Institute (Beijing, China).

**Methylation-specific semi-quantitative polymerase chain reaction (MSP) analysis.** In the present study, MSP was developed for the detection of mothers against decapentaplegic homolog 3 (SMAD3; Chr1:67356838-Chr1:67356942) methylation in 81 patients with ACS, 74 patients with SCAD and 53 healthy individuals using two pairs of primers, one was specific for the methylated alleles while the other was specific for the unmethylated alleles. Primer sequences are listed in Table II. The thermocycling conditions were as follows: Complete denaturation of the DNA at  $95^{\circ}\text{C}$  for 5 min; 35 cycles of denaturation at  $95^{\circ}\text{C}$  for 30 sec, annealing at  $58^{\circ}\text{C}$  for 30 sec, extension at  $72^{\circ}\text{C}$  for 45 sec; and a final extension at  $72^{\circ}\text{C}$  for 10 min. The MSP products were assessed by electrophoresis in 2% agarose gels with 0.5  $\mu\text{g}/\text{ml}$  ethidium bromide and analyzed using a UVC1-1100 SmartView Pro 1100 Imaging system with version 1.0.0.2 software (Major Science, Saratoga, CA, USA). Images demonstrating the different methylation patterns were captured.

**Sequenom MassARRAY.** Primer pairs were designed using EpiDesigner (epidesigner.com) and are presented in Table II. MassArray analysis was performed in triplicate using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MassARRAY Analyzer 4; Sequenom, San Diego, CA, USA) according to manufacturer's protocol in 8 ACS patients, 8 SCAD patients and 8 healthy individuals randomly from each group. The experiments were conducted at the Beijing Genomics Institute.

**Differential methylation analysis.** The Infinium HumanMethylation450 assay data were normalized using

Table I. Clinical and biochemical characteristics of the participants included in the present study.

Variable	Group		
	Control (n=53)	SCAD (n=74)	ACS (n=81)
Age, years (mean $\pm$ range)	64.68 $\pm$ 8.18	68.99 $\pm$ 13.62	72.72 $\pm$ 12.93
Sex, n=m/f	34/19	47/27	42/39
History of hypertension, n (%)	0 (0)	58 (78)	61 (75)
History of diabetes, n (%)	0 (0)	15 (20)	37 (46)
TC, mmol/l	4.43 $\pm$ 0.47	4.16 $\pm$ 0.95	4.50 $\pm$ 0.73
TG, mmol/l	1.01 $\pm$ 0.33	1.48 $\pm$ 0.97 <sup>a</sup>	1.66 $\pm$ 0.57 <sup>b</sup>
HDL-C, mmol/l	1.39 $\pm$ 0.26	1.07 $\pm$ 0.30 <sup>b</sup>	0.94 $\pm$ 0.16 <sup>b</sup>
LDL-C, mmol/l	2.70 $\pm$ 0.58	2.74 $\pm$ 0.52	2.80 $\pm$ 0.33
Cardiac troponin I, ng/ml	-	0.03 $\pm$ 0.01	6.61 $\pm$ 17.31 <sup>c</sup>
Cardiac troponin T, ng/ml	-	0.015 $\pm$ 0.03	0.758 $\pm$ 1.95 <sup>d</sup>
Creatine kinase-MB, U/l	-	10.24 $\pm$ 7.44	45.57 $\pm$ 80.27 <sup>d</sup>

<sup>a</sup>P<0.05 and <sup>b</sup>P<0.001 vs. the control group; <sup>c</sup>P<0.05 and <sup>d</sup>P<0.01 vs. the SCAD group. HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; TC, total cholesterol; TG, triglyceride; SCAD, stable coronary artery disease; ACS, acute coronary syndrome; m, male; f, female.

Table II. Primer sequences for the methylation analysis of mothers against decapentaplegic homolog 3 (Chr1:67356838-Chr1:67356942).

Method	Primer	Sequence (5'-3')	Amplicon size (bp)
MSP	UF	TTTAGATATTGGTTAGTTGGTTGT	141
	UR	ACACATTAAAAAACTCCACAACCT	
	MF	AGATATCGGTTAGTCGGTTGC	141
	MR	CATTAAAAAACTCCGCGACTT	
Sequenom	F	AGGAAGAGAGGTTTTGTAATTTTTTATTTATTTGAGGG	352
MassARRAY	R	CAGTAATACGACTCACTATAGGGAGAAGGCTATTTTATTTCC AAAACAATAAACTCACA	

UF, unmethylated forward; UR, unmethylated reverse; MF, methylated forward; MR, methylated reverse; F, forward; R, reverse.

internal controls (normalization control probe pairs, which were designed to target the same region within housekeeping genes and had no underlying CpG sites in the probe) with GenomeStudio software (version 2011.1; Illumina, Inc.). For quality control, methylation measures of P>0.05 and samples with a CpG coverage <95% were removed. The methylation levels of CpG sites were calculated as the  $\beta$  score [ $\beta$  score = intensity of the methylated allele/(intensity of the unmethylated allele + intensity of the methylated allele + 100)]. Limma software (version 3.24.15; bioconductor.org/packages/release/bioc/html/limma.html) was used to perform paired t-tests followed by Benjamini and Hochberg correction for multiple testing in order to identify DM CpG sites in the ACS group vs. the control group. Sites with a Benjamini and Hochberg-corrected value of P<0.05 and a  $\beta$  score  $\geq$ 0.1 were considered to be significantly statistically different. A volcano plot was used to display the mean DNA

methylation differences for the DM CpG sites. Hierarchical clustering of the methylation data was performed when the  $\beta$  score was  $\geq$ 0.2.

**Functional annotation of DM CpG sites.** The enrichment analysis of the methylated regions for Gene Ontology (GO) (33) terms and Kyoto Encyclopedia of Genes and Genomes (KEGG) (34) pathways was conducted using the Genomic Regions Enrichment of Annotations Tool (35). Binomial statistics with false discovery rate (FDR) correction were used to identify significantly enriched ontologies. The most significant and replicable corresponding genes were selected to be candidate markers for ASC detection. These genes were chosen from the results of the differential methylation tests at the site, gene region and gene island region levels, and had a  $\beta$  score difference between the ACS and control samples of >0.3.




Figure 1. Density plots of the distribution of the  $\beta$  scores. The  $\beta$  score distribution for the three pairs of samples following normalization, as visualized in density plots.

Table III. DM CpG sites and genes in patients with acute coronary syndrome compared with the control group.

CpG site methylation	No. of CpG sites	No. of genes
Loci detected	484,352	42,343
DM	11,342	5,073
Hypermethylated	2,477	1,746
Hypomethylated	8,865	5,071

CpG, 5'-C-phosphate-G-3'; DM, differentially methylated.

**Statistical analysis.** Statistical analyses were performed using SPSS software (version 20.0; SPSS, Inc., Chicago, IL, USA). Normally distributed values are expressed as the mean  $\pm$  standard deviation. A one-sample Kolmogorov-Smirnov test was used to evaluate the normality of distribution of the variables. The significance of differences between variants among groups was analyzed by one-way analysis of variance, and the differences between groups were subsequently determined using a post hoc Fisher's least significant difference test. The differences in cTnI and CK-MB between two groups were analyzed using the Student's t-test. The Kruskal-Wallis H test was performed for non-parametric data.  $P < 0.05$  was considered to indicate a statistically significant difference.

## Results

**Normalization of methylation data.** For the methylated sites identified by Illumina HumanMethylation450 assay, the original data of the 3 patients with ACS and 3 paired controls were normalized to the internal controls to eliminate the possibility of system error, and the  $\beta$  scores were subsequently calculated. The  $\beta$  scores were generated based on the intensity of the methylated and unmethylated probes, and were used as an indicator of the degree of methylation at each CpG site. Subsequently, a built-in detection score filter was used prior to further calculations. This only left values with significantly higher mean signal intensities from multiple probes for a given CpG locus compared with those of the negative control in the same set of chip data. The average number of loci detected for the three pairs of ACS and control samples was 484,352 covering 42,343 genes (Table III). The density plots of the  $\beta$  score for the 6 samples exhibited a bimodal distribution (Fig. 1). These results indicate that the hybridization and amplification conditions were uniform for all samples.

**Epigenome-wide DNA methylation analysis.** Using Benjamini and Hochberg correction for multiple comparisons, a total of 11,342 (2.34%) DM CpG sites covering 5,073 genes were identified in patients with ACS compared with the control patients (Table III). Among the 11,342 DM CpG sites, 8,865 (78.2%) covering 5,071 genes were significantly hypomethylated, while 2,477 (21.8%) covering 1,746 genes were significantly hypermethylated in the ACS samples (Table III).






Figure 2. Volcano plot of the significant differential DNA methylation analysis. All 11,342 sites among the 3 paired acute coronary syndrome and control samples were presented in the volcano plot. Red markers indicate the 11,342 differentially methylated CpG sites between the ACS and the control samples. CpG, 5'-C-phosphate-G-3'; ACS, acute coronary syndrome.




Figure 3. Hierarchical cluster analysis of the significantly DM CpG sites. A hierarchical cluster analysis was performed on the 1,206 significantly differentially methylated CpG sites. DM, differentially methylated; ACS, acute coronary syndrome; CpG, 5'-C-phosphate-G-3'.

A volcano plot of the data exhibited the mean DNA methylation differences for all CpG sites (Fig. 2). The frequencies of all significant CpG sites between the ACS and control groups were analyzed (Table IV). A total of 9.04% (224) of the hypermethylated CpG sites and 11.8% (982) of the hypomethylated CpG sites had an absolute ACS/control group methylation difference  $\geq 20\%$ . Fig. 3 presents a hierarchical cluster analysis of the 1,206 CpG sites that distinguished the ACS samples from the control samples.

*Genomic distribution of DM CpG sites.* Significant DM CpG sites were analyzed based on their region and island region level throughout the genome (Table V). There were 6 gene-based regions, including the transcription start site (TSS)1500, TSS200 (within 200 base pairs of a TSS), 5'-untranslated region (5'-UTR), 1stExon, body and 3'-UTR. In addition, there were 5 CpG island-based regions, including the island, N-Shore, S-Shore, N-Shelf and S-Shelf sites. The gene and island regions were used to calculate the methyla-

Table IV. Frequency distribution of all significantly differentially methylated CpG sites between acute coronary syndrome and control samples by methylation status.

ACS vs. control group methylation $\beta$ score difference (%)	Hypermethylated CpG sites, n (%)	Cumulative (%)	Hypomethylated CpG sites, n (%)	Cumulative (%)
$\geq 50$	24 (0.97)	0.97	16 (0.18)	0.18
$\geq 40$ -<50	32 (1.29)	2.26	22 (0.25)	0.43
$\geq 30$ -<40	35 (1.41)	3.67	40 (0.45)	0.88
$\geq 20$ -<30	133 (5.37)	9.04	904 (10.20)	11.08
$\geq 10$ -<20	2,253 (90.96)	100	7,883 (88.92)	100
Total	2,477	-	8,865	-

CpG, 5'-C-phosphate-G-3'; ACS, acute coronary syndrome.

Table V. Statistics of DM CpG sites identified by site, region and island region level.

Test	Region	No. of DM CpG sites	No. of genes
CpG site-level	Whole genome	11,342	5,073
Region-level gene based	TSS1500	861	349
	TSS200	421	182
	5'-UTR	515	246
	1st Exon	73	31
	Body	2,662	1,250
	3'-UTR	277	130
Region-level island based	Island	1224	462
	N-Shore	1593	644
	S-Shore	1247	513
	N-Shelf	751	372
	S-Shelf	685	331

UTR, untranslated region; CpG, 5'-C-phosphate-G-3'; DM, differentially methylated; TSS, transcription start site; N, north; S, south.

tion index. All the given region definitions were based on the original Illumina methylation annotation for the Human Methylation450 BeadChip array. The gene-based variant of the region-level test resulted in the identification of 4,809 significant DM CpG sites, which covered 2,188 genes. This number contained all unique genes, which resulted from six different, separately analyzed gene categories: TSS1500 (349 genes), TSS200 (182 genes), 5'-UTR (246 genes), 1stExon (31 genes), body (1,250 genes) and 3'-UTR (130 genes). Similarly, the CpG island-based variant of the region-level test resulted in 1,224 DM CpG sites located in islands, and 1,593 and 1,247 DM CpG sites distributed inside the N-Shore and S-Shore, respectively, while 751 and 685 DM CpG sites were located in the N-Shelf and S-Shelf, respectively. In total, 5,500 unique DM CpG sites covering 2,322 genes were identified among the five CpG-related regions.

Table VI. Distribution of the genomic region of significantly differentially methylated CpG sites in the acute coronary syndrome compared with the control samples.

Genomic region of CpG sites	All CpG sites, n (%)	Hypermethylated CpG sites, n (%)	Hypomethylated CpG sites, n (%)
TSS1500	861 (17.90)	150 (15.17)	711 (18.61)
TSS200	421 (8.75)	68 (6.88)	353 (9.24)
5'-UTR	515 (10.71)	99 (10.01)	416 (10.89)
1st Exon	73 (1.52)	17 (1.72)	56 (1.47)
Body	2,662 (55.36)	597 (60.36)	2,065 (54.06)
3'-UTR	277 (5.76)	58 (5.86)	219 (5.73)
Total	4,809	989	3,820

CpG, 5'-C-phosphate-G-3'; UTR, untranslated region; TSS, transcription start site.

The genomic regions of the significantly hypermethylated or hypomethylated CpG sites were distributed similarly (Table VI). The largest portion of hypermethylated CpGs (60.36%) were located in the body of the genes and subsequently decreased in other categories (15.17% in TSS1500, 6.88% in TSS200, 10.01% in the 5'-UTR, 1.72% in the 1stExon and 5.86% in the 3'-UTR). In addition, over half (54.06%) of the significantly hypomethylated CpG sites were located in the body, while the rest were distributed in the TSS1500 (18.16%), TSS200 (9.24%), 5'-UTR (10.89%), 1stExon (1.47%) and 3'-UTR (5.73%) regions. Furthermore, the distribution analysis of DM CpG sites in island regions revealed that 18.5% (190) of the significantly hypermethylated CpG sites were in CpG island regions, with 47.81% (491) in CpG shores and 33.69% (346) in shelf regions. As for the significantly hypomethylated CpG sites, 23.12% (1,034) were located in islands, while the majority were identified in shore (52.52%) or shelf (24.36%) regions (Table VII), suggesting they serve a potential role in genomic instability.

*GO term and KEGG pathway enrichment analyses of the DM CpG sites.* To identify the functions and pathways possibly

Table VII. Distribution of island region of significantly differentially methylated CpG sites in ACS vs. control samples.

Region level island-based GpG sites	All GpG sites	Hypermethylated CpG sites, no. (%)	Hypomethylated CpG sites, no. (%)
Island	1,224 (22.25)	190 (18.50)	1,034 (23.12)
N-Shore	1,593 (28.96)	304 (29.60)	1,289 (28.82)
S-Shore	1,247 (22.68)	187 (18.21)	1,060 (23.70)
N-Shelf	751 (13.65)	166 (16.16)	585 (13.08)
S-Shelf	685 (12.46)	180 (17.53)	505 (11.28)
Total	5,500	1,027	4,473

CpG, 5'-C-phosphate-G-3'; N, north; S, south.




Figure 4. Top 10 terms in each GO category ranked according to their statistical significance. The top terms within the biological process, cellular component and molecular function categories are presented. GO, Gene Ontology.

influenced by the genes nearest the DM CpG sites in patients with ACS compared with the controls, GO term and KEGG pathway enrichment analyses were performed (Fig. 4). There were 767 functional categories of DM genes ( $FDR < 0.05$ ), the largest portion of the functional terms comprised of biological processes (603), while the rest were cellular components (76) or molecular functions (88). When the biological processes were used for categorization, it was observed that the most enriched groups included immune system processes, the regulation of responses to stimuli and cell activation. Categorization by cellular components indicated that proteins encoded by the analyzed genes were mainly associated with the cytoplasm. The molecular function category demonstrated significant enrichment for phospholipid, protein and lipid

binding activities. Fig. 4 presents the top 10 terms of the three GO categories ranked by statistical significance.

KEGG pathway analysis revealed a significant enrichment of genes associated with the hematopoietic cell lineage and endocytosis (Table VIII and Fig. 5). Among them, the immune system, and the transport and catabolism pathways attracted were ranked highest in terms of significance.

#### *Selection of significant genes as possible diagnostic markers.*

The selection criteria used to select potential diagnostic marker genes included CpG sites that were methylated in the site, gene and gene island region, and a  $\beta$  score difference between the ACS and control samples of  $>0.3$ . Following filtering, a list of 19 hypermethylated and 17 hypomethylated CpG sites were




Figure 5. Top 20 Kyoto Encyclopedia of Genes and Genomes pathways ranked according to their statistical significance.




Figure 6. Electrophoresis of the TSS1500 region of mothers against decapentaplegic homolog 3 (Chr1:67356838-Chr1:67356942) following methylation-specific polymerase chain reaction analysis. The M primers and U primers amplified the 141 bp products. M, methylated; U, unmethylated.

selected that matched these criteria (Table IX). These methylated CpG sites were mapped to 37 known genes.

**CpG methylation validation.** The methylation state of the representative TSS1500 region (SMAD3; Chr1:67356838-Chr1:67356942) on the basis of DM magnitude was detected by MSP (Figs. 6 and 7A) and Sequenom MassARRAY (Fig. 7B). These results were compared with the results from the HumanMethylation450 array. The frequency of semi-methylated sites (unmethylated/

methylated status) in patients with ACS was notably higher than that in the patients with SCAD and the healthy donors (Fig. 7A). The mean methylation level was identified as being significantly lower in patients with ACS compared with the SCAD and healthy control groups (Fig. 7B).

## Discussion

To the best of our knowledge, the present study presents the first genome-wide DNA methylation profiling of 3 paired ACS and normal human blood samples, using a high-throughput DNA methylation microarray covering >450,000 CpG sites on the human genome. The DM CpGs identified were enriched in genes associated with immune system processes, the regulation of responses to stimuli and cell activation, which suggests that they are of potential mechanistic and therapeutic relevance.

The impact of epigenetic mechanisms in cardiovascular pathophysiology is emerging as a key participant in the interface between genotype and phenotype variability. Several previous studies have focused on the role of DNA methylation in the development of atherosclerosis and CVD. Hiltunen *et al* (27) reported that a global hypomethylation of DNA was present in advanced human, mouse and rabbit atherosclerotic lesions. Notably, the results of Kim *et al* (36) indicated that elevated peripheral blood leukocyte DNA methylation was positively associated with the prevalence of CVD predisposing conditions and obesity in Chinese Singaporeans. A recent study also revealed that lower global DNA methylation is associated with



Table VIII. KEGG pathway enrichment analysis of the hypermethylated and hypomethylated 5'-C-phosphate-G-3' sites between the acute coronary syndrome and control samples by methylation status.

KEGG ID no.	Pathway name	Pathway class	No. of genes in pathway	P-value	FDR
hsa04640	Hematopoietic cell lineage	Immune system	43	2.13x10 <sup>-7</sup>	4.72x10 <sup>-5</sup>
hsa04144	Endocytosis	Transport and catabolism	83	3.63x10 <sup>-7</sup>	4.72x10 <sup>-5</sup>
hsa04940	Type I diabetes mellitus	Endocrine and metabolic diseases	25	1.55x10 <sup>-7</sup>	1.35x10 <sup>-4</sup>
hsa05330	Allograft rejection	Immune diseases	22	2.90x10 <sup>-6</sup>	1.89x10 <sup>-4</sup>
hsa04510	Focal adhesion	Cell communication	80	1.24x10 <sup>-5</sup>	5.51x10 <sup>-4</sup>
hsa05320	Autoimmune thyroid disease	Immune diseases	24	1.27x10 <sup>-5</sup>	5.51x10 <sup>-4</sup>
hsa05145	Toxoplasmosis	Infectious diseases: Parasitic	50	1.51x10 <sup>-5</sup>	5.63x10 <sup>-4</sup>
hsa04514	Cell adhesion molecules	Signaling molecules and interaction	58	2.12x10 <sup>-5</sup>	6.90x10 <sup>-4</sup>
hsa04066	HIF-1 signaling pathway	Signal transduction	47	2.54x10 <sup>-5</sup>	7.17x10 <sup>-4</sup>
hsa04650	Natural killer cell mediated cytotoxicity	Immune system	52	2.76x10 <sup>-5</sup>	7.17x10 <sup>-4</sup>
hsa04145	Phagosome	Transport and catabolism	60	3.41x10 <sup>-5</sup>	7.83x10 <sup>-4</sup>
hsa04380	Osteoclast differentiation	Development	54	3.61x10 <sup>-5</sup>	7.83x10 <sup>-4</sup>
hsa05332	Graft vs. host disease	Immune diseases	21	4.81x10 <sup>-5</sup>	9.63x10 <sup>-4</sup>
hsa04660	T cell receptor signaling pathway	Immune system	46	5.94x10 <sup>-5</sup>	1.10x10 <sup>-3</sup>
hsa05200	Pathways in cancer	Cancers: Overview	113	7.67x10 <sup>-5</sup>	1.33x10 <sup>-3</sup>
hsa05146	Amoebiasis	Infectious diseases: Parasitic	46	1.03x10 <sup>-4</sup>	1.67x10 <sup>-3</sup>
hsa04070	Phosphatidylinositol signaling system	Signal transduction	35	2.66x10 <sup>-4</sup>	4.03x10 <sup>-3</sup>
hsa05310	Asthma	Immune diseases	16	3.06x10 <sup>-4</sup>	4.03x10 <sup>-3</sup>
hsa05416	Viral myocarditis	Cardiovascular diseases	31	3.10x10 <sup>-4</sup>	4.03x10 <sup>-3</sup>
hsa04151	PI3K-Akt signaling pathway	Signal transduction	112	3.10x10 <sup>-4</sup>	4.03x10 <sup>-3</sup>
hsa05221	Acute myeloid leukemia	Cancers: Specific types	27	3.34x10 <sup>-4</sup>	4.14x10 <sup>-3</sup>
hsa05212	Pancreatic cancer	Cancers: Specific types	30	4.07x10 <sup>-4</sup>	4.82x10 <sup>-3</sup>
hsa04670	Leukocyte transendothelial migration	Immune system	46	5.73x10 <sup>-4</sup>	6.28x10 <sup>-3</sup>
hsa04630	Jak-STAT signaling pathway	Signal transduction	53	5.80x10 <sup>-4</sup>	6.28x10 <sup>-3</sup>
hsa05166	HTLV-I infection	Infectious diseases: Viral	89	7.26x10 <sup>-4</sup>	7.55x10 <sup>-3</sup>
hsa05032	Morphine addiction	Substance dependence	38	8.19x10 <sup>-4</sup>	8.01x10 <sup>-3</sup>
hsa05223	Non-small cell lung cancer	Cancers: Specific types	25	8.32x10 <sup>-4</sup>	8.01x10 <sup>-3</sup>
hsa04062	Chemokine signaling pathway	Immune system	67	9.16x10 <sup>-4</sup>	8.32x10 <sup>-3</sup>
hsa04150	mTOR signaling pathway	Signal transduction	27	9.29x10 <sup>-4</sup>	8.32x10 <sup>-3</sup>
hsa04662	B cell receptor signaling pathway	Immune system	32	1.06x10 <sup>-3</sup>	9.16x10 <sup>-3</sup>
hsa04915	Estrogen signaling pathway	Endocrine system	39	1.30x10 <sup>-3</sup>	1.05x10 <sup>-2</sup>
hsa04612	Antigen processing and presentation	Immune system	29	1.34x10 <sup>-3</sup>	1.05x10 <sup>-2</sup>
hsa04664	Fc epsilon RI signaling pathway	Immune system	30	1.36x10 <sup>-3</sup>	1.05x10 <sup>-2</sup>
hsa04060	Cytokine-cytokine receptor interaction	Signaling molecules and interaction	83	1.37x10 <sup>-3</sup>	1.05x10 <sup>-2</sup>
hsa05222	Small cell lung cancer	Cancers: Specific types	35	1.75x10 <sup>-3</sup>	1.30x10 <sup>-2</sup>
hsa00562	Inositol phosphate metabolism	Carbohydrate metabolism	26	2.97x10 <sup>-3</sup>	2.15x10 <sup>-2</sup>
hsa05140	Leishmaniasis	Infectious diseases: Parasitic	28	3.90x10 <sup>-3</sup>	2.74x10 <sup>-2</sup>
hsa04666	Fc gamma R-mediated phagocytosis	Immune system	35	5.30x10 <sup>-3</sup>	3.62x10 <sup>-2</sup>
hsa04360	Axon guidance	Development	47	5.58x10 <sup>-3</sup>	3.72x10 <sup>-2</sup>
hsa04725	Cholinergic synapse	Nervous system	41	7.36x10 <sup>-3</sup>	4.78x10 <sup>-2</sup>

KEGG, Kyoto Encyclopedia of Genes and Genomes; FDR, false discovery rate.

higher cardiovascular risk in postmenopausal women (37). These studies led to the use of DNA methylation as a novel biomarker for CAD. However, the majority of their results were obtained from animal experiments or gene-specific

DNA methylation, which lack of the systematic analysis of genome-wide changes to DNA methylation in patients with ACS. Zaina *et al* (38) identified an atherosclerosis-specific DNA methylation profile that highlighted the contribution of

Table IX. List of the top 19 hypermethylated and 17 hypomethylated corresponding genes selected as possible diagnostic markers.

## A, Hypermethylated corresponding genes

CpG site	ACS vs. control group methylation $\beta$ score difference	Gene affected	P-value	FDR
cg03951394	0.31859	DEAF1	4.61x10 <sup>-2</sup>	0.43464
cg11478607	0.33696	GSTT1	1.43x10 <sup>-3</sup>	0.24422
cg01387905	0.34720	ADAMTS2	1.35x10 <sup>-6</sup>	0.03278
cg13829625	0.35709	SIN3B	1.13x10 <sup>-6</sup>	0.03160
cg18424635	0.35947	HLA-DRB5	2.83x10 <sup>-2</sup>	0.39069
cg26951705	0.38792	ZNF787	4.81x10 <sup>-2</sup>	0.43861
cg00243527	0.39404	LIF	4.99x10 <sup>-2</sup>	0.44222
cg23782583	0.40248	JAKMIP3	4.53x10 <sup>-2</sup>	0.43270
cg00248861	0.40500	WHSC2	3.01x10 <sup>-2</sup>	0.39564
cg17005068	0.41609	GSTT1	1.49x10 <sup>-5</sup>	0.14324
cg10662395	0.46909	HCN2	3.48x10 <sup>-3</sup>	0.24464
cg19235645	0.47402	FAM101A	4.16x10 <sup>-2</sup>	0.42548
cg21829038	0.48534	LRRK1	3.54x10 <sup>-2</sup>	0.41016
cg17713488	0.49270	COL5A3	7.66x10 <sup>-5</sup>	0.24422
cg14594459	0.50806	PKHD1L1	4.34x10 <sup>-2</sup>	0.42957
cg18443741	0.79680	PYROXD1	2.48x10 <sup>-5</sup>	0.18747
cg19577958	0.54856	SERPINB9	4.46x10 <sup>-2</sup>	0.43166
cg04234412	0.64833	LOC391322	3.49x10 <sup>-7</sup>	0.01914
cg16836675	0.68159	TOM1L1	4.18x10 <sup>-3</sup>	0.24951

## B, Hypomethylated corresponding genes

cg20592836	-0.59591	TP53INP2	1.33x10 <sup>-2</sup>	0.32577
cg03549208	-0.55680	ICA1L	3.62x10 <sup>-2</sup>	0.41253
cg27141807	-0.53542	SPSB4	2.27x10 <sup>-2</sup>	0.37233
cg24851651	-0.48666	CCS	1.31x10 <sup>-2</sup>	0.32488
cg06180910	-0.43609	GSTT1	1.11x10 <sup>-4</sup>	0.24422
cg10864200	-0.41318	PCGF3	1.12x10 <sup>-2</sup>	0.31160
cg24875593	-0.40991	PDXK	5.26x10 <sup>-3</sup>	0.26054
cg04613734	-0.39972	TRPS1	4.07x10 <sup>-2</sup>	0.42315
cg19961153	-0.38234	ALPPL2	4.02x10 <sup>-2</sup>	0.42219
cg05141159	-0.37248	DEXI	3.71x10 <sup>-2</sup>	0.41476
cg06049774	-0.34306	PLAC9	9.33x10 <sup>-3</sup>	0.29687
cg23231631	-0.33541	GABRB1	3.02x10 <sup>-6</sup>	0.05052
cg23731272	-0.32429	SMAD3	5.37x10 <sup>-3</sup>	0.26137
cg08253809	-0.32072	C14orf119	3.87x10 <sup>-2</sup>	0.41857
cg01655658	-0.31142	HLA-L	4.82x10 <sup>-7</sup>	0.02123
cg13120108	-0.31114	PRKCH	4.08x10 <sup>-2</sup>	0.42353
cg20433858	-0.30584	OR2L13	3.36x10 <sup>-3</sup>	0.24464

CpG, 5'-C-phosphate-G-3'; ACS, acute coronary syndrome; FDR, false discovery rate.

different genes and pathways to the disorder; however, this study did not take into account the characteristics of acute onset in patients with atherosclerotic CVD.

The present study analyzed the changes to genome-wide DNA methylation in patients with ACS using the Illumina HumanMethylation450 assay. A total of 11,342 CpG sites across the whole genome demonstrated statistically significant DM in the ACS samples compared with the control samples. The majority

of the DM CpG sites were hypomethylated (8,865; 78.2%) rather than hypermethylated (2,477; 21.8%), which was consistent with the results of a previous methylation study in atherosclerotic lesions and CAD (28). In addition, global hypomethylation, particularly in repeated sequences, serves an important role in increasing chromosomal instability and may lead to ACS (39). This suggests that alteration of the DNA methylation profile, particularly hypomethylation, is an important process in the




Figure 7. Representative measurements of the SMAD3 TSS1500 methylation status. (A) The TSS1500 methylation frequency of SMAD3 in 81 patients with ACS, 74 patients with SCAD and 53 healthy individuals was measured using methylation-specific semi-quantitative polymerase chain reaction analysis. (B) The DNA methylation level of SMAD3 TSS1500 using Sequenom MassARRAY in 8 paired ACS, SCAD and healthy individuals was measured. SMAD3, mothers against decapentaplegic homolog 3; MM, fully methylated; MU, methylated/unmethylated or semi-methylated; ACS, acute coronary syndrome; SCAD, stable coronary artery disease.

pathogenesis of ACS. Sharma *et al* (40) proposed that genomic DNA methylation in patients with CAD is significantly altered, as higher genomic DNA methylation changes were observed in patients with CAD than in controls. Kim *et al* (36) examined the association between peripheral blood leukocyte global genomic DNA methylation and the prevalence of CVD (myocardial infarction and stroke) or its risk factors (hypertension and diabetes) to verify the value of DNA methylation as a CVD risk biomarker. However, Kim *et al* (36) only compared differences in the overall methylation status between normal subjects and patients with coronary heart disease; further details on the differences in DNA methylation sites were not mentioned.

The genomic distribution of significant CpG sites revealed that significantly DM CpG sites were more frequently located at the body (55.3%) and shore (51.6%) sites. It has been previously reported that methylation at the CpG island shore regions regulates promoter activity (41), and that differential DNA methylation in gene bodies and intergenic loci indicates more complex mechanistic implications, as they are considered to promote transcriptional elongation, and to regulate gene splicing and enhancer activities (42).

The GO term and KEGG pathway enrichment analyses results demonstrated a significant enrichment of the corresponding methylated genes associated with the immune system processes, regulation of responses to stimuli and cell activation, which are all associated with the hematopoietic cell lineage and endocytosis signaling pathways. The proteins encoded by the corresponding methylated genes were mainly associated with the cytoplasm, and were enriched in the molecular functions of phospholipid, protein and lipid binding activities. These results are highly consistent with data from a recent genome-wide systematic pathway analysis of body mass index, which provided insights into the genetics of CVD and its risk factors (43). The results of the present study revealed a significant enrichment to immune system processes in the ACS samples, which may be due to the altered maturation status and possible immune exhaustion of immune cells in the peripheral blood of patients with ACS (44). Additionally, the neutrophil/lymphocyte ratio (NLR) has been identified to be independently associated with the severity and 3-year outcomes of CAD: Patients with higher NLR values had more advanced obstructive CAD and a poorer prognosis (45). CVD

is often accompanied by disorders of lipid metabolism, and previous studies have demonstrated that elevated concentrations of oxidized lipoprotein were associated with the presence and severity of ACS (46,47). In the present study, the network analysis for GO term and KEGG pathway enrichment of the DM CpG sites revealed significant enrichment change in patients with ACS, which indicated that the methylation alterations at these sites were biologically meaningful.

In the present study, 19 hypermethylated genes (DEAF1, GSTT1, ADAMTS2, SIN3B, HLA-DRB5, ZNF787, LIF, JAKMIP3, WHSC2, HCN2, FAM101A, LRRK1, COL5A3, PKHD1L1, PYROXD1, SERPINB9, LOC391322 and TOM1L1) and 17 hypomethylated genes (TP53INP2, ICA1L, SPSB4, CCS, GSTT1, PCGF3, PDXK, TRPS1, ALPPL2, DEX1, PLAC9, GABRB1, SMAD3, C14orf119, HLA-L, PRKCH and OR2L13) demonstrated a potential association with ACS. These genes may thus represent potential novel candidate biomarkers for the early diagnosis of ACS. Three DM genes were identified that have been deeply studied in CVD: SMAD3, SERPINB9 and PRKCH. SMAD3, a key contributor to the transforming growth factor- $\beta$  pathway signaling, has been reported to serve an essential role in CAD (48-51). Meta-analyses of genome-wide association studies have identified that an intronic single nucleotide polymorphism (SNP) in SMAD3 is associated with protection from CAD (52). SERPINB9 is most commonly known for its role in the control of granzyme B, which can be applicable for prevention of ACS and atherosclerotic plaque formation in patients with left ventricular remodeling following acute myocardial infarction (53). However, SERPINB9 also serves a role in suppressing interleukin-1 $\beta$  maturation through inhibiting caspase-1 (54). Genome-wide association and molecular mechanism studies have revealed that the SNPs rs2230500 (55) and rs2230500 (56,57) in PRKCH were significantly associated with the risk of ischemic stroke. However, whether SMAD3, SERPINB9 or PRKCH methylation is associated with the pathogenesis of ACS is currently unknown. The validation experiments performed for SMAD3 and TSS1500 (Chr1:67356838-Chr1:67356942) in the present study indicated that the methylation degree was negatively associated with the pathogenesis of ACS, and this was significantly corrected by the reference results of the Illumina HumanMethylation450 assay.

In conclusion, the present study characterized the genome-wide DNA methylation patterns occurring in ACS, and identified significantly differentiated CpG sites and genes that were correlated with ACS. Additionally, GO term and KEGG pathway enrichment analyses provided information on the processes and pathways that the methylated CpG sites and the participating genes affected. These results may help clarify the role of aberrant methylation in ACS pathogenesis, and provide the basis for future research into potentially abnormally methylated genes as biomarkers for the prevention and early diagnosis of ACS.

## Acknowledgements

The present study was supported by the National Natural Science Foundation of China (grant nos. 81271904, 81401742, 81572073 and 81572074), the Special-Funded Program on National Key Scientific Instruments and Equipment Development of China (grant no. 2012YQ030261) and Jinling Hospital Foundation (grant no. 2017060).

## References

- Falk E, Nakano M, Bentzon JF, Finn AV and Virmani R: Update on acute coronary syndromes: The pathologists' view. *Eur Heart J* 34: 719-728, 2013.
- Corcoran D, Grant P and Berry C: Risk stratification in non-ST elevation acute coronary syndromes: Risk scores, biomarkers and clinical judgment. *Int J Cardiol* 8: 131-137, 2015.
- O'Gara PT, Kushner FG, Ascheim DD, Casey DE Jr, MK, de Lemos JA, Ettinger SM, Fang JC, Fesmire FM, Franklin BA, *et al*: American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation* 127: e362-e425, 2013.
- Redfors B, Råmunddal T, Angerås O, Dworeck C, Haraldsson I, Ioanes D, Petursson P, Libungan B, Odenstedt J, Stewart J, *et al*: Angiographic findings and survival in patients undergoing coronary angiography due to sudden cardiac arrest in western Sweden. *Resuscitation* 90: 13-20, 2015.
- Hofer TP, Frankenberger M, Mages J, Lang R, Meyer P, Hoffmann R, Colige A and Ziegler-Heitbrock L: Tissue-specific induction of ADAMTS2 in monocytes and macrophages by glucocorticoids. *J Mol Med (Berl)* 86: 323-332, 2008.
- Tomson T, Surges R, Delamont R, Haywood S and Hesdorffer DC: Who to target in sudden unexpected death in epilepsy prevention and how? Risk factors, biomarkers, and intervention study designs. *Epilepsia* 57 (Suppl 1): 4-16, 2016.
- Mosleh W, Abdel-Qadir H and Farkouh M: Biomarkers in the emergency workup of chest pain: Uses, limitations, and future. *Cleve Clin J Med* 80: 589-598, 2013.
- Wu SS, Lin X, Yuan LQ and Liao EY: The role of epigenetics in arterial calcification. *BioMed Res Int* 2015: 320849, 2015.
- Heyn H and Esteller M: DNA methylation profiling in the clinic: Applications and challenges. *Nat Rev Genet* 13: 679-692, 2012.
- Handy DE, Castro R and Loscalzo J: Epigenetic modifications: Basic mechanisms and role in cardiovascular disease. *Circulation* 123: 2145-2156, 2011.
- Wallace RG, Twomey LC, Custaud MA, Moyna N, Cummins PM, Mangone M and Murphy RP: Potential diagnostic and prognostic biomarkers of epigenetic drift within the cardiovascular compartment. *BioMed Res Int* 2016: 2465763, 2016.
- Loscalzo J and Handy DE: Epigenetic modifications: Basic mechanisms and role in cardiovascular disease (2013 Grover Conference series). *Pulm Circ* 4: 169-174, 2014.
- Turgeon PJ, Sukumar AN and Marsden PA: Epigenetics of cardiovascular disease - A new 'Beat' in coronary artery disease. *Med Epigenet* 2: 37-52, 2014.
- Kandi V and Vadakedath S: Effect of DNA methylation in various diseases and the probable protective role of nutrition: A mini-review. *Cureus* 7: e309, 2015.
- Razin A, Webb C, Szyf M, Yisraeli J, Rosenthal A, Naveh-Mani T, Sciaky-Gallili N and Cedar H: Variations in DNA methylation during mouse cell differentiation in vivo and in vitro. *Proc Natl Acad Sci USA* 81: 2275-2279, 1984.
- Afzali M, Nakhaee A, Tabatabaei SP, Tirgar-Fakheri K and Hashemi M: Aberrant promoter methylation profile of Niemann-pick type C1 gene in cardiovascular disease. *Iran Biomed J* 17: 77-83, 2013.
- Nazarenko MS, Markov AV, Lebedev IN, Freidin MB, Sleptcov AA, Koroleva IA, Frolov AV, Popov VA, Barbarash OL and Puzyrev VP: A comparison of genome-wide DNA methylation patterns between different vascular tissues from patients with coronary heart disease. *PLoS One* 10: e0122601, 2015.
- Li Y, Liu Y, Strickland FM and Richardson B: Age-dependent decreases in DNA methyltransferase levels and low transmethylation micronutrient levels synergize to promote overexpression of genes implicated in autoimmunity and acute coronary syndromes. *Exp Gerontol* 45: 312-322, 2010.
- Kim JM, Stewart R, Kang HJ, Bae KY, Kim SW, Shin IS, Hong YJ, Ahn Y, Jeong MH and Yoon JS: BDNF methylation and depressive disorder in acute coronary syndrome: The K-DEPACS and EsDEPACS studies. *Psychoneuroendocrinology* 62: 159-165, 2015.
- Lü CX, Xu RD, Cao M, Wang G, Yan FQ, Shang SS, Wu XF, Ruan L, Quan XQ and Zhang CT: FOXP3 demethylation as a means of identifying quantitative defects in regulatory T cells in acute coronary syndrome. *Atherosclerosis* 229: 263-270, 2013.
- Zhang LN, Liu PP, Wang L, Yuan F, Xu L, Xin Y, Fei LJ, Zhong QL, Huang Y, Xu L, *et al*: Lower ADD1 gene promoter DNA methylation increases the risk of essential hypertension. *PLoS One* 8: e63455, 2013.
- Kim JD, Lee A, Choi J, Park Y, Kang H, Chang W, Lee MS and Kim J: Epigenetic modulation as a therapeutic approach for pulmonary arterial hypertension. *Exp Mol Med* 47: e175, 2015.
- Wang Z, Zheng Y, Hou C, Yang L, Li X, Lin J, Huang G, Lu Q, Wang CY and Zhou Z: DNA methylation impairs TLR9 induced Foxp3 expression by attenuating IRF-7 binding activity in fulminant type 1 diabetes. *J Autoimmun* 41: 50-59, 2013.
- Pasquier J, Hoarau-Véhot J, Fakhro K, Rafii A and Abi Khalil C: Epigenetics and cardiovascular disease in diabetes. *Curr Diab Rep* 15: 108, 2015.
- Babu M, Durga Devi T, Mäkinen P, Kaikkonen M, Lesch HP, Junttila S, Laiho A, Ghimire B, Gyenesei A and Ylä-Herttuala S: Differential promoter methylation of macrophage genes is associated with impaired vascular growth in ischemic muscles of hyperlipidemic and type 2 diabetic mice: Genome-wide promoter methylation study. *Circ Res* 117: 289-299, 2015.
- Yip L, Fuhlbrigge R, Taylor C, Creusot RJ, Nishikawa-Matsumura T, Whiting CC, Schartner JM, Akter R, von Herrath M and Fathman CG: Inflammation and hyperglycemia mediate Deaf1 splicing in the pancreatic lymph nodes via distinct pathways during type 1 diabetes. *Diabetes* 64: 604-617, 2015.
- Hiltunen MO, Turunen MP, Häkkinen TP, Rutanen J, Hedman M, Mäkinen K, Turunen AM, Aalto-Setälä K and Ylä-Herttuala S: DNA hypomethylation and methyltransferase expression in atherosclerotic lesions. *Vasc Med* 7: 5-11, 2002.
- Castillo-Díaz SA, Garay-Sevilla ME, Hernández-González MA, Solís-Martínez MO and Zaina S: Extensive demethylation of normally hypermethylated CpG islands occurs in human atherosclerotic arteries. *Int J Mol Med* 26: 691-700, 2010.
- Jiang YZ, Manduchi E, Stoeckert CJ Jr and Davies PF: Arterial endothelial methylome: Differential DNA methylation in atherosusceptible disturbed flow regions in vivo. *BMC Genomics* 16: 506, 2015.
- Eden A, Gaudet F, Waghmare A and Jaenisch R: Chromosomal instability and tumors promoted by DNA hypomethylation. *Science* 300: 455, 2003.
- Virani S, Rentschler KM, Nishijo M, Ruangyuttikarn W, Swaddiwudhipong W, Basu N and Rozek LS: DNA methylation is differentially associated with environmental cadmium exposure based on sex and smoking status. *Chemosphere* 145: 284-290, 2016.
- Zhu X, Li J, Deng S, Yu K, Liu X, Deng Q, Sun H, Zhang X, He M, Guo H, *et al*: Genome-wide analysis of DNA methylation and cigarette smoking in a Chinese population. *Environ Health Perspect* 124: 966-973, 2016.
- Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, *et al*: The Gene Ontology Consortium: Gene ontology: Tool for the unification of biology. *Nat Genet* 25: 25-29, 2000.
- Kanehisa M and Goto S: KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res* 28: 27-30, 2000.



35. McLean CY, Bristor D, Hiller M, Clarke SL, Schaar BT, Lowe CB, Wenger AM and Bejerano G: GREAT improves functional interpretation of cis-regulatory regions. *Nat Biotechnol* 28: 495-501, 2010.
36. Kim M, Long TI, Arakawa K, Wang R, Yu MC and Laird PW: DNA methylation as a biomarker for cardiovascular disease risk. *PLoS One* 5: e9692, 2010.
37. Ramos RB, Fabris V, Lecke SB, Maturana MA and Spritzer PM: Association between global leukocyte DNA methylation and cardiovascular risk in postmenopausal women. *BMC Med Genet* 17: 71, 2016.
38. Zaina S, Heyn H, Carmona FJ, Varol N, Sayols S, Condom E, Ramírez-Ruz J, Gomez A, Gonçalves I, Moran S, *et al*: DNA methylation map of human atherosclerosis. *Circ Cardiovasc Genet* 7: 692-700, 2014.
39. Li J, Chen L, Du L and Li M: Cage the firefly luciferin! - a strategy for developing bioluminescent probes. *Chem Soc Rev* 42: 662-676, 2013.
40. Sharma P, Kumar J, Garg G, Kumar A, Patowary A, Karthikeyan G, Ramakrishnan L, Brahmachari V and Sengupta S: Detection of altered global DNA methylation in coronary artery disease patients. *DNA Cell Biol* 27: 357-365, 2008.
41. Bockmühl Y, Patchev AV, Madejska A, Hoffmann A, Sousa JC, Sousa N, Holsboer F, Almeida OF and Spengler D: Methylation at the CpG island shore region upregulates Nr3c1 promoter activity after early-life stress. *Epigenetics* 10: 247-257, 2015.
42. Jones PA: Functions of DNA methylation: Islands, start sites, gene bodies and beyond. *Nat Rev Genet* 13: 484-492, 2012.
43. Zhao X, Gu J, Li M, Xi J, Sun W, Song G and Liu G: Pathway analysis of body mass index genome-wide association study highlights risk pathways in cardiovascular disease. *Sci Rep* 5: 13025, 2015.
44. Zidar DA, Mudd JC, Juchnowski S, Lopes JP, Sparks S, Park SS, Ishikawa M, Osborne R, Washam JB, Chan C, *et al*: Altered maturation status and possible immune exhaustion of CD8 T lymphocytes in the peripheral blood of patients presenting with acute coronary syndromes. *Arterioscler Thromb Vasc Biol* 36: 389-397, 2016.
45. Arbel Y, Finkelstein A, Halkin A, Birati EY, Revivo M, Zuzut M, Shevach A, Berliner S, Herz I, Keren G, *et al*: Neutrophil/lymphocyte ratio is related to the severity of coronary artery disease and clinical outcome in patients undergoing angiography. *Atherosclerosis* 225: 456-460, 2012.
46. Wang JJ, Zhang CN, Meng Y, Han AZ, Gong JB and Li K: Elevated concentrations of oxidized lipoprotein(a) are associated with the presence and severity of acute coronary syndromes. *Clin Chim Acta* 408: 79-82, 2009.
47. Wang JJ, Han AZ, Meng Y, Gong JB, Zhang CN, Li K and Liu YX: Measurement of oxidized lipoprotein (a) in patients with acute coronary syndromes and stable coronary artery disease by 2 ELISAs: Using different capture antibody against oxidized lipoprotein (a) or oxidized LDL. *Clin Biochem* 43: 571-575, 2010.
48. Lee AM, Shimizu C, Oharaseki T, Takahashi K, Daniels LB, Kahn A, Adamson R, Dembitsky W, Gordon JB and Burns JC: Role of TGF- $\beta$  signaling in remodeling of noncoronary artery aneurysms in kawasaki disease. *Pediatr Dev Pathol* 18: 310-317, 2015.
49. Turner AW, Nikpay M, Silva A, Lau P, Martinuk A, Linseman TA, Soubeyrand S and McPherson R: Functional interaction between COL4A1/COL4A2 and SMAD3 risk loci for coronary artery disease. *Atherosclerosis* 242: 543-552, 2015.
50. Peng Q, Deng Y, Yang X, Leng X, Yang Y and Liu H: Genetic variants of ADAM17 are implicated in the pathological process of Kawasaki disease and secondary coronary artery lesions via the TGF- $\beta$ /SMAD3 signaling pathway. *Eur J Pediatr* 175: 705-713, 2016.
51. Lusis AJ: Cardiovascular disease genes come together. *Atherosclerosis* 242: 630-631, 2015.
52. Turner AW, Martinuk A, Silva A, Lau P, Nikpay M, Eriksson P, Folkersen L, Perisic L, Hedin U, Soubeyrand S, *et al*: Functional analysis of a novel genome-wide association study signal in SMAD3 that confers protection from coronary artery disease. *Arterioscler Thromb Vasc Biol* 36: 972-983, 2016.
53. Saito Y, Kondo H and Hojo Y: Granzyme B as a novel factor involved in cardiovascular diseases. *J Cardiol* 57: 141-147, 2011.
54. van der Burgh R, Meeldijk J, Jongeneel L, Frenkel J, Bovenschen N, van Gijn M and Boes M: Reduced serpinB9-mediated caspase-1 inhibition can contribute to autoinflammatory disease. *Oncotarget* 7: 19265-19271, 2016.
55. Hata J, Kubo M and Kiyohara Y: Genome-wide association study for ischemic stroke based on the Hisayama study. *Nihon Eiseigaku Zasshi* 66: 47-52, 2011 (In Japanese).
56. Wu L, Xi B, Hou D, Zhao X, Liu J, Cheng H, Zhou X, Shen Y, Wang X and Mi J: The SNP (rs2230500) in PRKCH decreases the risk of carotid intima-media thickness in a Chinese young adult population. *PLoS One* 7: e40606, 2012.
57. Wu L, Shen Y, Liu X, Ma X, Xi B, Mi J, Lindpaintner K, Tan X and Wang X: The 1425G/A SNP in PRKCH is associated with ischemic stroke and cerebral hemorrhage in a Chinese population. *Stroke* 40: 2973-2976, 2009.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.