

Identification of aberrantly expressed long non-coding RNAs in postmenopausal osteoporosis

QI FEI*, XIAODONG BAI*, JISHENG LIN, HAI MENG, YONG YANG and AI GUO

Department of Orthopedics, Beijing Friendship Hospital, Capital Medical University, Beijing 100050, P.R. China

Received October 19, 2017; Accepted February 14, 2018

DOI: 10.3892/ijmm.2018.3575

Abstract. Postmenopausal osteoporosis (PMOP) is a common skeletal disorder in postmenopausal women. The present study aimed to identify the key long non-coding RNAs (lncRNAs) in PMOP through RNA sequencing. RNA sequencing was performed to obtain the expression profile of lncRNAs and mRNAs in blood samples of patients with PMOP and normal controls (NCs). Following the identification of differentially expressed mRNAs (DEmRNAs) and differentially expressed lncRNAs (DElncRNAs), the DElncRNA-DEmRNA co-expression network was constructed. A search was performed for the DEGs transcribed within a 100-kb window upstream or downstream of DElncRNAs, which served as nearby DEmRNAs of DElncRNAs. Functional annotation of the DEmRNAs co-expressed with DElncRNAs was performed. The GSE56815 dataset was used to verify the expression of selected DEmRNAs and DElncRNAs. Three blood samples from patients with PMOP and two blood samples from NCs were used for RNA sequencing. Compared with the NC group, a total of 185 DEmRNAs and 51 DElncRNAs were obtained in PMOP. A total of 3,057 co-expression DElncRNA-DEmRNA pairs and 97 DElncRNA-nearby DEmRNA pairs were obtained. Six DEmRNAs [diacylglycerol O-acyltransferase 2, potassium voltage-gated channel subfamily S member 1, peptidase inhibitor 3, secretory leukocyte peptidase inhibitor, galectin-related protein and alkaline phosphatase, liver/bone/kidney (ALPL)] were nearby co-expressed genes of four DElncRNAs, including LOC105376834, LOC101929866, LOC105374771 and LOC100506113. Three PMOP-associated DEmRNAs, including ALPL, suppressor of cytokine signaling 3 and adrenomedullin, were co-expressed with the hub

DElncRNAs (LINC00963, LOC105378415, LOC105377067, HCG27, LOC101928143 and LINC01094) of the positively and negatively co-expressed DElncRNA-DEmRNA interaction network. The expression of selected DEmRNAs and DElncRNAs was consistent with the RNA-sequencing results. In conclusion, the present study identified the key DEmRNAs and DElncRNAs in PMOP, which may provide clues for understanding the mechanism and developing novel biomarkers for PMOP.

Introduction

Osteoporosis is a systemic skeletal disorder characterized by a reduction in bone mineral density (BMD) and disrupted bone architecture, which results in a higher risk of bone fractures (1,2). It is reported that ~50% of postmenopausal women suffer from osteoporosis, which is defined as postmenopausal osteoporosis (PMOP) (3,4). The basic pathogenesis of PMOP involves an imbalance between bone resorption by osteoclasts and bone formation by osteoblasts, which is mainly induced by decreased estrogen. As PMOP is a chronic disease, it imposes a significant financial burden on postmenopausal women (4,5). Therefore, it is crucial to uncover the mechanism and develop accurate diagnostic biomarkers of PMOP.

Previous studies have indicated that osteoporosis and BMD are heritable (6,7), and >60 susceptible loci have been found to be associated with osteoporosis and BMD (6). Among these, polymorphisms of several genes have been found to be involved in PMOP, including tumor necrosis factor (TNF)- α , interleukin (IL)10, osteoprotegerin, estrogen receptor 1 gene, estrogen receptor α , cannabinoid receptor 2, vitamin D receptor gene and LDL receptor related protein 5 (8-14).

Long non-coding RNAs (lncRNAs) are a set of non-coding RNAs containing >200 nucleotides. There has been increased interest focused on lncRNAs, which have been found to be involved in diseases, including cancer and osteoporosis by regulating their target genes at the transcriptional, post-transcriptional and epigenetic levels (15,16). An lncRNA, DANCER was found to be involved in PMOP by regulating TNF- α and IL6 (16). LncRNA MEG3 can suppress the osteogenic differentiation of bone marrow mesenchymal stem cells induced by PMOP (17). However, reports of lncRNAs in PMOP remain limited.

In the present study, the lncRNA and mRNA expression profile of blood samples from patients with PMOP and

Correspondence to: Dr Qi Fei or Dr Ai Guo, Department of Orthopedics, Beijing Friendship Hospital, Capital Medical University, 95 Yong'an Road, Xicheng, Beijing 100050, P.R. China
E-mail: spinefei@126.com
E-mail: guoaiortho@126.com

*Contributed equally

Key words: long non-coding RNA, RNA sequencing, postmenopausal osteoporosis, co-expression

normal controls (NCs) were identified by high-throughput RNA-sequencing. To the best of our knowledge, the present study is the first to obtain the lncRNA expression profiles of PMOP by RNA sequencing. Based on the identified differentially expressed lncRNAs (DElncRNAs) and differentially expressed mRNAs (DEmRNAs) in PMOP, compared with NC, the DElncRNAs-DEmRNAs co-expression network was constructed. The potential roles of these DElncRNAs were further examined according to the functional annotation of their co-expressed DEmRNAs. These findings may provide clues for understanding the pathogenesis and novel insight for developing diagnostic biomarkers of PMOP.

Materials and methods

Patients and samples. From April 2016 to March 2017, three women with PMOP and two healthy women from Beijing Friendship Hospital were enrolled in the present study. The inclusion criteria of patients with PMOP were as follows: i) Postmenopausal women who were diagnosed with osteoporosis. Osteoporosis was defined by the World Health Organization criteria of a BMD T-score of -2.5 standard deviations below the average for a young adult at peak bone density in the femoral neck, total hip, or L1-L4; ii) clinically symptomatic postmenopausal women with painful vertebral fractures verified by X-ray and MRI within the last 6 months, who returned for further examination and treatment. The patient characteristics are listed in Table I. All individuals provided written informed consent for use of their samples in the present study. The present study was approved by the Ethics Committee of Beijing Friendship Hospital, Capital Medical University (Beijing, China; 2017-P2-084-01). From every participant, a 2.5 ml peripheral whole blood was collected in PAXgene® RNA blood tubes (PreAnalytiX GmbH, Hombrechtikon, Switzerland) and stored at -80°C prior to processing.

RNA isolation and sequencing. RNA isolation was performed using the PAXgene blood RNA kit (PreAnalytiX GmbH) according to the manufacturer's protocol. The concentration and purity of RNA were assessed using a Nanodrop ND-2000 spectrophotometer (Thermo Fisher Scientific, Inc., Waltham, MA, USA). The integrity of RNA was assessed using a 2% agarose gel. A RIN value was obtained using an Agilent 2100 bioanalyzer. The criteria for cDNA library construction were as follows: i) Total RNA $>5\ \mu\text{g}$; ii) concentration of RNA $\geq 200\ \text{ng/ml}$; iii) OD260/280 value 1.8-2.2.

Following removal of the ribosomal RNA using the Ribo-Zero Magnetic kit (EpiCentre, Madison, WI, USA), the RNA was purified and fragmented into 200-500-base pair fragments. The RNA fragments were primed with random hexamer primers and the first cDNA strand was synthesized, with the second cDNA strand synthesized with dUTP instead of dTTP. The blunt ends of double-stranded DNA were produced from cohesive ends of double-stranded DNA using End Repair Enzyme mix (New England BioLabs, Inc., Ipswich, MA, USA). Subsequently, 3' end adenylation and adapter ligation were performed. When the second digested cDNA strand was digested using the UNG enzyme (Illumina, Inc., San Diego, CA, USA), polymerase chain reaction (PCR)

was performed with PCR Primer Cocktail (Illumina, Inc.) and PCR Master Mix (Illumina, Inc.) to amplify the libraries. The following thermocycling conditions were used for the PCR: Initial denaturation at 98°C for 30 sec; 15 cycles of 98°C for 10 sec, 65°C for 30 sec and 72°C for 30 sec, followed by a final extension step of 72°C for 5 min. Certified Low Range Ultra Agarose (Bio-Rad Laboratories, Inc., Hercules, CA, USA) was used to purify the libraries, and the libraries were quantified using Picogreen double-stranded DNA quantitation kit (Molecular Probes; Thermo Fisher Scientific, Inc.) on a TBS380 fluorometer (Promega Corporation, Madison, WI, USA). The qualified libraries were amplified on cBot to generate the cluster on the flowcell using TruSeq PE Cluster kit V3-cBot-HS (Illumina, Inc.) according to the manufacturer's protocol. Sequencing was performed on the Illumina HiSeq Xten platform (Illumina, Inc.).

Quality control of raw sequence and mapping of clean reads. The FASTQ sequence data were obtained from the RNA-seq data using Base Calling V0.11.4 (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). To obtain the high quality clean data, the low quality reads including adaptor sequences, sequences with a quality score <20 , and sequences with an N base rate of raw reads $>10\%$ were removed by using Cutadapt V1.9.1 (<https://cutadapt.readthedocs.io/en/stable/>). With TopHat release 2.1.1 (<http://tophat.cbcb.umd.edu/>) and Ensemble gene annotation, clean reads were aligned with the human reference genome, Ensemble GRCh38.p7 (ftp://ftp.ncbi.nlm.nih.gov/genomes/Homo_sapiens). The expression of mRNAs and lncRNAs was determined and outputted using Cuffquant V2.2.1 (<http://cufflinks.cbcb.umd.edu/>).

Identification of DEmRNAs and DElncRNAs in PMOP compared with NC. Fragments per Kilobase of exon per million fragments mapped (FPKM) was used to determine the transcription abundance of lncRNAs and mRNAs. The FPKMs of lncRNAs and mRNAs were calculated using Cuffdiff (<http://cole-trapnell-lab.github.io/cufflinks/cuffdiff/index.html>). A paired t-test was performed to obtain the DEmRNAs and DElncRNAs in PMOP compared with NC. The thresholds of the DEmRNAs and DElncRNAs was $P < 0.05$.

DElncRNA-DEmRNA co-expression network. To further examine the potential roles of DElncRNAs and DEmRNAs in PMOP the DElncRNA-DEmRNA co-expression network was constructed. Firstly, the Pearson's correlation coefficient (PCC) between the expression levels of each DElncRNA-DEmRNA pair in the PMOP and the NC group were calculated. Secondly, DElncRNA-DEmRNA pairs with an absolute value of PCC ≥ 0.90 and $P < 0.05$ were defined as co-expressed DElncRNA-DEmRNA pairs. Those co-expressed DElncRNA-DEmRNA pairs in which the expression level of DEmRNAs was positively correlated with the expression level of DElncRNAs in PMOP were defined as positively co-expressed DElncRNA-DEmRNA pairs. Co-expressed DElncRNA-DEmRNA pairs in which the expression level of DEmRNAs was negatively correlated with the expression level of DElncRNAs in PMOP were defined as negatively

Table I. Patient characteristics.

Characteristic	Case 1	Case 2	Case 3	Control 1	Control 2
Age (years)	81	68	79	67	68
Sex	Female	Female	Female	Female	Female
BMI (kg/m ²)	23.1	15.6	15.6	28.6	22.5
BMD-T score	-3	-4.2	-3.7	1.5	0.4
History of smoking	No	No	No	No	No
History of alcohol intake	No	No	No	No	No
History of coffee or carbonated drink intake	No	No	No	No	No
Family history of matrilineal family	No	No	No	No	No
Lack of physical activity	No	No	No	No	No
Bone metabolism-associated disease	No	No	No	No	No
Bone metabolism-associated drugs	No	No	No	No	No

BMI, body mass index; BMD, bone mineral density.

co-expressed DElncRNA-DEmRNA pairs. The positively and negatively co-expressed DElncRNA-DEmRNA networks were visualized using Cytoscape 3.1 (<http://cytoscape.org/>).

Nearby DEmRNAs of the DElncRNAs. In order to identify the targeted DEmRNAs of DElncRNAs by *cis*-regulatory effects, a search was performed for the DEmRNAs transcribed within a 100-kb window upstream or downstream of DElncRNAs, which served as nearby *cis*-targeted DEmRNAs of DElncRNAs.

Functional annotation of DEmRNAs co-expressed lncRNAs. Functional annotation, including Gene Ontology (GO) function and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analyses of the DEmRNAs co-expressed with DElncRNAs was performed using the GeneCoDis3 tool (<http://genecodis.cnb.csic.es/analysis>). A hypergeometric test was used to obtain the P-value. The false discovery rate (FDR; corrected P-value) of <0.05 was set as the cut-off for significant GO terms and KEGG pathways.

Validation in the Gene Expression Omnibus (GEO) dataset. The GSE56815 dataset was obtained from the GEO (<https://www.ncbi.nlm.nih.gov/geo/>), which consisted of 20 postmenopausal women with low hip BMD (case group) and 20 postmenopausal women with high hip BMD (normal group). All 40 females were Caucasian. The expression pattern of selected DElncRNAs and DEmRNAs was verified using the GSE56815 dataset. GSE7158 was another dataset obtained from GEO, which consisted of 12 women with a low peak bone mass (case group) and 14 women with a high peak bone mass (normal group). GSE7158 was also used to validate the expression pattern of selected DElncRNAs.

Results

RNA-sequencing data. Total RNA extracted from each of the blood samples met the criteria for cDNA library construction

and RNA-sequencing. Following trimming of the raw reads, 6.8×10^7 , 6.8×10^7 and 6.7×10^7 clean reads were obtained from the three respective blood samples from patients with postmenopausal osteoporosis; 6.8×10^7 and 6.7×10^7 clean reads were obtained from the two respective NCs. All of the clean reads were aligned to the human reference genome (GRCh38.p7) and the mapped ratio of all samples was >80%.

DEmRNAs and DElncRNAs in PMOP. A total of 185 significantly DEmRNAs (184 upregulated DEmRNAs and one downregulated DEmRNAs) were obtained with $P < 0.05$. The top 30 significant DEmRNAs are listed in Table II. A total of 51 significantly DElncRNAs (25 upregulated DElncRNAs and 26 downregulated DElncRNAs) were obtained with $P < 0.05$ (Table III). LOC105372321 was the most markedly upregulated DElncRNA and LOC105374771 was the most markedly downregulated DElncRNA in PMOP, compared with NC. NOD-like receptor family pyrin domain containing 6 was the most significantly upregulated DEmRNA and PAGE family member 2B was the only downregulated DEmRNA. Furthermore, these DElncRNAs were distributed in all chromosomes (chr.), with the exception of chr. 5, 15, 17, 18, 21 and sex chr. X and Y, whereas the DEmRNAs were widely distributed in all chromosomes, except sex chr.Y (Fig. 1).

DElncRNA-DEmRNA co-expression network. Based on the expression levels of DElncRNAs and DEmRNAs, the PCC describing the co-expression association between 185 DElncRNAs and 51 DEmRNAs, was calculated. A total of 3,057 DElncRNA-DEmRNA co-expression pairs were obtained with an absolute value of $PCC \geq 0.90$ and $P < 0.05$. Among these, a total of 2,756 lncRNA-mRNA pairs were identified as being positively co-expressed, whereas 301 lncRNA-mRNA pairs were negatively co-expressed. The positively co-expressed DElncRNA-DEmRNA network (Fig. 2A) consisted of 215 nodes and 2,756 edges, and its hub lncRNAs were LOC105378415 (degree=159), LOC105377067 (degree=157), HCG27 (degree=157),

Table II. Top 30 significantly DEGs in patients with postmenopausal osteoporosis compared with normal controls.

DEG	Locus	log2 (fold-change)	Regulation	P-value
FOLR3	chr11:72135709-72139892	-4.03	Down	5.00x10 ⁻⁰⁵
PI3	chr20:45174898-45176544	-2.88	Down	5.00x10 ⁻⁰⁵
KRT23	chr17:40922695-40987135	-2.76	Down	5.00x10 ⁻⁰⁵
CD177	chr19:43353658-43366075	-2.57	Down	5.00x10 ⁻⁰⁵
REM2	chr14:22883164-22887680	-2.15	Down	5.00x10 ⁻⁰⁵
NLRP6	chr11:278364-285942	-2.14	Down	5.00x10 ⁻⁰⁵
MANSC1	chr12:12329262-12350541	-2.06	Down	1.00x10 ⁻⁰⁴
LRG1	chr19:4537214-4540024	-2.14	Down	1.50x10 ⁻⁰⁴
HCAR2	chr12:122695781-122710104	-1.89	Down	2.00x10 ⁻⁰⁴
ADM	chr11:10304979-10307402	-1.67	Down	2.00x10 ⁻⁰⁴
MAK	chr6:10762722-10838954	-2.07	Down	2.50x10 ⁻⁰⁴
ABCG1	chr21:42199688-42304389	-1.98	Down	4.50x10 ⁻⁰⁴
LGALS1	chr2:64454192-64461383	-1.69	Down	5.00x10 ⁻⁰⁴
BTNL8	chr5:180899076-180952166	-1.89	Down	5.50x10 ⁻⁰⁴
CCNJL	chr5:160251651-160339592	-1.84	Down	6.50x10 ⁻⁰⁴
HIP1	chr7:75533297-75738976	-1.51	Down	6.50x10 ⁻⁰⁴
SOCS3	chr17:78356776-78360079	-1.85	Down	8.00x10 ⁻⁰⁴
PADI2	chr1:17066760-17119453	-1.85	Down	8.00x10 ⁻⁰⁴
TGFA	chr2:70402845-70554015	-1.60	Down	1.05x10 ⁻⁰³
KIAA0226L	chr13:46341999-46390042	-1.35	Down	1.30x10 ⁻⁰³
OSM	chr22:30262827-30266843	-1.65	Down	1.40x10 ⁻⁰³
CRISPLD2	chr16:84819980-84909510	-1.44	Down	1.55x10 ⁻⁰³
KCNS1	chr20:45091213-45101112	-1.37	Down	1.55x10 ⁻⁰³
IL1B	chr2:112829757-112836842	-1.52	Down	1.70x10 ⁻⁰³
KAZN	chr1:13893386-15220480	-2.41	Down	2.10x10 ⁻⁰³
HLX	chr1:220832762-220885059	-1.31	Down	2.20x10 ⁻⁰³
ABCA1	chr9:104781001-104939096	-2.05	Down	2.25x10 ⁻⁰³
HSPA1A	chr6:31815513-31817942	-1.45	Down	2.35x10 ⁻⁰³
KREMEN1	chr22:29058671-29168333	-1.52	Down	2.45x10 ⁻⁰³

DEG, differentially expressed genes.

LOC101928143 (degree=154) and LINC00963 (degree=154). The negatively co-expressed DElncRNA-DEmRNA network (Fig. 2B) consisted of 175 nodes and 301 edges, and its hub lncRNAs were LINC01094 (degree=135) and LOC105371455 (degree=85).

Nearby DEmRNAs of DElncRNAs. A total of 97 DElncRNAs nearby DEmRNA pairs were obtained. LOC101928595, LOC101929866 and HCG27 had 13, 8 and 7 nearby DEmRNAs, respectively, and were the top three DElncRNAs with the most nearby DEmRNAs (Table IV). DElncRNAs nearby DEmRNA pairs in which the expression levels of DEmRNAs were co-expressed with DElncRNAs are listed in Table V.

Functional annotation. Based on the GO enrichment analysis of DEmRNAs co-expressed with DElncRNAs, inflammatory response (FDR=3.14E-5), anti-apoptosis (FDR=5.44E-05), plasma membrane (FDR=1.48E-11), integral to membrane

(FDR=6.06E-10), protein binding (FDR=1.39E-06), and receptor activity (FDR=8.92E-06) were the most significantly enriched GO terms in PMOP (Fig. 3A-C). Hematopoietic cell lineage (FDR=0.000244565), Osteoclast differentiation (FDR=0.000438367) and Cytokine-cytokine receptor interaction (FDR=0.00212347) were the most significantly enriched KEGG pathways in PMOP (Fig. 3D).

Validation in the GEO dataset. The expression patterns of selected DElncRNAs (LINC00963, LOC105376834, LOC101929866, LOC105374771 and LOC100506113) and DEmRNAs [alkaline phosphatase, liver/bone/kidney (ALPL), suppressor of cytokine signaling 3 (SOCS3), secretory leukocyte peptidase inhibitor (SLPI) and CD177] were verified using the GSE56815 dataset. As shown in Fig. 4A-D, SOCS3, SLPI and CD177 were upregulated in PMOP, which was consistent with the RNA-sequencing results. ALPL was downregulated in PMOP, which was inconsistent with the RNA-sequencing results. However, only one of these five

Table III. Significantly differentially expressed lncRNAs in patients with postmenopausal osteoporosis compared with normal controls.

LncRNA	Locus	Regulation	log2 (fold change)	P-value
LOC105374771	chr2:64390955-64425399	Down	-2.57	5.00x10 ⁻⁰⁵
LOC105372321	chr19:21444103-21464331	Up	3.58	5.00x10 ⁻⁰⁵
PSMD5-AS1	chr9:120843041-120854373	Down	-1.70	1.00x10 ⁻⁰⁴
PAX8-AS1	chr2:113215996-113278950	Down	-2.41	7.00x10 ⁻⁰⁴
LOC105372578	chr20:24919978-24932985	Down	-2.70	1.65x10 ⁻⁰³
LINC00570	chr2:11393980-11403077	Up	2.20	1.95x10 ⁻⁰³
LOC105369213	chr16:81739026-81777351	Down	-1.82	5.00x10 ⁻⁰⁵
LOC105378020	chr6:137943074-137957648	Up	inf	3.75x10 ⁻⁰³
SNHG5	chr6:85677006-85678733	Up	1.83	4.15x10 ⁻⁰³
LOC105378415	chr10:88061829-88104391	Down	-1.37	4.80x10 ⁻⁰³
LOC105374150	chr3:148439991-148465791	Up	1.76	0.01
LINC00282	chr13:51804681-51845150	Down	-1.69	0.01
LOC102724231	chr3:44421131-44424025	Down	-1.46	0.01
LINC00211	chr2:37826246-37875863	Down	-1.92	0.01
LOC101929638	chr22:29180622-29205834	Up	1.77	0.01
JHDM1D-AS1	chr7:140177260-140179640	Up	1.77	0.01
LOC105370449	chr14:34551436-34557529	Up	2.89	0.01
LOC105372881	chr1:207365821-207373252	Down	-1.14	0.01
LOC105373262	chr1:244230505-244325182	Down	-1.75	0.01
LOC105371455	chr1:157225405-157283617	Up	1.68	0.01
LOC100507487	chr4:128428015-128519398	Up	3.25	0.01
LOC101929866	chr20:45178476-45191638	Down	-1.37	0.01
LINC00963	chr9:129488659-129513686	Down	-1.08	0.02
LOC101928143	chr14:73460934-73463642	Down	-1.16	0.02
LOC399715	chr10:6326543-6335982	Down	-1.37	0.02
LOC105373730	chr2:165821976-165848198	Up	1.43	0.02
LOC100507639	chr4:141321123-141332617	Up	1.45	0.02
LOC105374768	chr2:64299870-64344064	Down	-1.34	0.02
LOC100506159	chr12:9936578-9943495	Down	-1.91	0.02
LOC105376834	chr1:21585689-21591187	Down	-2.23	0.02
LINC01094	chr4:78645993-78684501	Up	1.29	0.02
LOC105378085	chr6:159586906-159604657	Down	-1.36	0.02
LOC105377067	chr3:46130889-46190381	Down	-1.26	0.02
LOC105369823	chr12:69624414-69699416	Up	2.14	0.02
LOC101929422	chr14:101120762-101123545	Up	2.50	0.02
LOC105375328	chr7:64944845-64950665	Up	1.73	0.02
LINC01271	chr20:50292719-50321342	Down	-2.13	0.03
LOC102723828	chr4:31997378-32155406	Up	inf	0.03
LOC105377384	chr4:116344095-116355205	Up	2.43	0.03
LOC100506113	chr11:75801640-75814797	Down	-1.26	0.03
LOC105377782	chr8:2199669-2206204	Up	1.20	0.03
HCG27	chr6:31197759-31203968	Down	-0.90	0.03
LINC01137	chr1:37454878-37474443	Down	-1.84	0.03
LOC105374546	chr4:26859623-26860599	Up	3.84	0.03
LOC105376995	chr20:62533992-62536728	Up	1.78	0.03
LOC105374852	chr2:88016353-88021354	Up	1.72	0.04
LOC105378701	chr1:47172216-47177080	Up	2.11	0.04
LOC101928595	chr16:30096429-30113557	Down	-0.98	0.04
LOC105372991	chr22:30447958-30472047	Up	1.33	0.04
LOC105374769	chr2:64299870-64344064	Down	-3.63	0.05
GAS5	chr1:173863247-173867987	Up	1.03	0.05

LncRNA, long non-coding RNA; Inf, infinite.

Table IV. Nearby DE mRNAs of DE lncRNAs in postmenopausal osteoporosis.

Count	DE lncRNA	lncRNA location	mRNA	mRNA location
1	LOC105376834	chr1:21585689-21591187	ALPL	chr1:21508981-21578412
4	LINC01137	chr1:37454878-37474443	ZC3H12A	chr1:37474517-37484377
			SNIP1	chr1:37531436-37554344
			DNALI1	chr1:37556918-37595985
			GNL2	chr1:37556918-37595985
2	LOC105378701	chr1:47172216-47177080	CYP4Z1	chr1:47067487-47118320
			CYP4A22	chr1:47137424-47149738
1	LOC105371455	chr1:157225405-157283617	ETV3	chr1:157121190-157138591
4	GAS5	chr1:173863247-173867987	CENPL	chr1:173799549-173824639
			ZBTB37	chr1:173868094-173891122
			SERPINC1	chr1:173903803-173917378
			RC3H1	chr1:173931083-173993072
2	LOC105372881	chr1:207365821-207373252	CD55	chr1:207321471-207360966
			CR2	chr1:207454299-207489895
1	LOC105373262	chr1:244230505-244325182	C1orf100	chr1:244352062-244389896
1	LINC00570	chr2:11393980-11403077	E2F6	chr2:11444374-11466177
1	LOC105374771	chr2:64390955-64425399	LGALS1	chr2:64454192-64461383
2	LOC105374852	chr2:88016353-88021354	RGPD2	chr2:87748086-87992864
			SMYD1	chr2:88067779-88113384
1	LOC105373730	chr2:165821976-165848198	GALNT3	chr2:165747802-165796352
1	LOC102724231		TOPAZ1	chr3:44241885-44338010
2	LOC105377067	chr3:46130889-46190381	XCR1	chr3:46016989-46086803
			CCR1	chr3:46201708-46208341
1	LOC105374546	chr4:26859623-26860599	STIM2	chr4:26860690-27025381
1	LINC01094	chr4:78645993-78684501	ANXA3	chr4:78551587-78610451
1	LOC100507639	chr4:141321123-141332617	ZNF330	chr4:141220293-141234697
7	HCG27	chr6:31197759-31203968	C6orf15	chr6:31111222-31112555
			PSORS1C1	chr6:31114830-31140092
			CDSN	chr6:31114830-31140092
			PSORS1C2	chr6:31114830-31140092
			CCHCR1	chr6:31142438-31158238
			POU5F1	chr6:31164336-31170693
			HLA-C	chr6:31268748-31272136
1	LOC105378020	chr6:137943074-137957648	TNFAIP3	chr6:137823668-137883314
1	LOC105378085	chr6:159586906-159604657	SOD2	chr6:159679063-159789703
4	LOC105375328	chr7:64944845-64950665	ZNF138	chr7:64794387-64853800
			LOC441239	chr7:64882492-64937316
			ZNF117	chr7:64974451-65006746
			ERV3-1	chr7:64974451-65006746
2	LINC00963	chr9:129488659-129513686	NTMT1	chr9:129608883-129642169
			C9orf50	chr9:129608883-129642169
2	LOC399715	chr10:6326543-6335982	PFKFB3	chr10:6144801-6254648
			PRKCQ	chr10:6393037-6585361
2	LOC105378415	chr10:88061829-88104391	PTEN	chr10:87863437-87975287
			RNLS	chr10:88131897-88583860
2	LOC100506113	chr11:75801640-75814797	MOGAT2	chr11:75701595-75732958
			DGAT2	chr11:75768732-75801536
4	LOC100506159	chr12:9936578-9943495	KLRF2	chr12:9881488-9932430
			CLEC2A	chr12:9881488-9932430
			CLEC12B	chr12:10006137-10030606
			CLEC9A	chr12:10030676-10066030
2	LOC105369823	chr12:69624414-69699416	LRRC10	chr12:69608563-69611162
			BEST3	chr12:69624414-69699416

Table IV. Continued.

Count	DElncRNA	lncRNA location	mRNA	mRNA location
4	LINC00282	chr13:51804681-51845150	WDFY2	chr13:51584193-51804206
			DHRS12	chr13:51584193-51804206
			CCDC70	chr13:51861980-51866236
			ATP7B	chr13:51932668-52012130
3	LOC101928143	chr14:73460934-73463642	NUMB	chr14:73275209-73458580
			HEATR4	chr14:73478483-73634418
			C14orf169	chr14:73478483-73634418
13	LOC101928595	chr16:30096429-30113557	INO80E	chr16:29996208-30023280
			DOC2A	chr16:29996208-30023280
			C16orf92	chr16:30023333-30053026
			FAM57B	chr16:30023333-30053026
			ALDOA	chr16:30053089-30070420
			PPP4C	chr16:30075975-30085377
			TBX6	chr16:30085792-30091919
			GDPD3	chr16:30096429-30113557
			MAPK3	chr16:30114104-30123309
			CORO1A	chr16:30183392-30189076
			BOLA2B	chr16:30192929-30206927
			SLX1A	chr16:30192929-30206927
			SULT1A3	chr16:30192929-30206927
2	LOC105369213	chr16:81739026-81777351	CMIP	chr16:81445169-81711762
			PLCG2	chr16:81779257-81962693
3	LOC105372578	chr20:24919978-24932985	CST7	chr20:24949229-24959928
			APMAP	chr20:24962924-24992974
			ACSS1	chr20:25006229-25058182
8	LOC101929866	chr20:45178476-45191638	STK4	chr20:44966473-45079977
			KCNS1	chr20:45091213-45101112
			WFDC5	chr20:45109451-45116321
			WFDC12	chr20:45123425-45124465
			PI3	chr20:45174898-45176544
			SEMG1	chr20:45206963-45209773
			SEMG2	chr20:45221368-45224458
			SLPI	chr20:45230820-45290352
1	LINC01271	chr20:50292719-50321342	LINC01272	chr20:50267467-50279795
2	LOC105376995	chr20:62533992-62536728	GATA5	chr20:62463496-62475970
			MIR1-1HG	chr20:62543069-62570764
3	LOC101929638	chr22:29180622-29205834	KREMEN1	chr22:29058671-29168333
			RHBDD3	chr22:29259854-29300525
			EWSR1	chr22:29259854-29300525
5	LOC105372991	chr22:30447958-30472047	SF3A1	chr22:30331987-30378655
			SEC14L2	chr22:30396940-30436501
			SEC14L3	chr22:30447958-30472047
			SEC14L6	chr22:30522796-30546717
			GAL3ST1	chr22:30554634-30574588

DElncRNAs, differentially expressed long non-coding RNAs.

DElncRNAs, LINC00963, was detected in GSE56815, which may be due to the restriction of the microarray. LINC00963 was downregulated in PMO, which showed the same pattern with that in the RNA-sequencing results (Fig. 4E).

The expression pattern of six DElncRNAs (PSMD5-AS1, PAX8-AS1, JHDM1D-AS1, LINC00963, LOC100506113 and HCG27) was validated by GSE7158. Five DElncRNAs (PSMD5-AS1, PAX8-AS1, LINC00963, LOC100506113 and




Figure 1. Circos plots of the distribution of DElncRNAs and DEmrRNAs in PMOP on chromosomes. The outer layer cycle represents the chromosome map of the human genome hg19. The red inner layer and blue inner layer represent the distribution of DEmrRNAs and DElncRNAs in PMOP on different chromosomes, respectively. DE, differentially expressed; lncRNAs, long non-coding RNAs; PMOP, post-menopausal osteoporosis.

Table V. DElncRNA-nearby DEmrRNA pairs in which DEmrRNAs are co-expressed with DElncRNAs.

DEmrRNA	DElncRNA	PCC	P-value
DGAT2	LOC100506113	9.77×10^{-01}	4.05×10^{-03}
KCNS1	LOC101929866	9.02×10^{-01}	3.64×10^{-02}
PI3	LOC101929866	9.65×10^{-01}	7.89×10^{-03}
SLPI	LOC101929866	9.48×10^{-01}	1.41×10^{-02}
LGALS1	LOC105374771	9.93×10^{-01}	6.56×10^{-04}
ALPL	LOC105376834	9.60×10^{-01}	9.51×10^{-03}

DE, differentially expressed; lncRNA, long non-coding RNA; PCC, Pearson's correlation coefficient; DGAT2, diacylglycerol O-acyltransferase 2; KCNS1, potassium voltage-gated channel subfamily S member 1; PI3, peptidase inhibitor 3; SLPI, secretory leukocyte peptidase inhibitor; LGALS1, galectin-related protein; ALPL, alkaline phosphatase, liver/bone/kidney.

HCG27) were downregulated, whereas JHDM1D-AS1 was upregulated, in PMOP compared with NC (Fig. 5), which was the same pattern found in the RNA-sequencing results.

Discussion

Although the function of the majority of lncRNAs remains to be elucidated, previous studies have indicated that lncRNAs may be involved in the pathogenesis of PMOP. Identifying the key DElncRNAs in PMOP not only provides novel clues for understanding the function of lncRNAs, but also contributes to developing novel biomarkers of PMOP.

In the present study, the landscape of lncRNAs in PMOP was obtained and a total of 51 DElncRNAs in PMOP were identified. With the exception of LINC00963 and GAS5, no previous study has reported on the function of the remaining 49 DElncRNAs. In addition, the present study is the first, to the best of our knowledge, to show that these 51 DElncRNAs may be associated with PMOP.

LINC00963 is reported to be involved in cell viability, motility and invasiveness in prostate cancer cells by affecting the expression of epidermal growth factor receptor (18). In the present study, LINC00963 was a significantly downregulated lncRNA in PMOP. Whether LINC00963 is involved in PMOP by regulating the viability, motility and invasiveness of osteoclasts and osteoblast requires further investigation.

Although the functions of lncRNAs remain to be fully elucidated, previous studies have indicated that lncRNAs are




Figure 2. DElncRNA-DEmRNA co-expression network. The PCC between the expression levels of each DElncRNA-DEmRNA pair in PMOP and NC groups were calculated. (A) DElncRNA-DEmRNA pairs with $PCC \geq 0.90$ and $P < 0.05$ were considered to be positively co-expressed DElncRNA-DEmRNA pairs and (B) DElncRNA-DEmRNA pairs with $PCC \leq -0.90$ and $P < 0.05$ were considered to be negatively co-expressed DElncRNA-DEmRNA pairs. The rhombi and the ellipses represent DElncRNAs and DEmRNAs in PMOP, respectively. The blue and red colors represent downregulation and upregulation in PMOP, respectively. DE, differentially expressed; lncRNAs, long non-coding RNAs; PMOP, post-menopausal osteoporosis; PCC, Pearson's correlation coefficient; NC, normal control.

important in regulating the expression levels of genes and proteins, and are involved in a variety of biochemical processes and diseases (19-21). To date, calculating the correlation coefficients between the expression levels of lncRNAs and genes has been the most popular approach to identify potential target genes of lncRNAs (22,23). Accumulated evidence has indicated that lncRNA-mRNA co-expression analysis can be used to examine the biological functions of lncRNAs in various diseases by examining their co-expressed mRNAs (24-26). In addition, several lncRNA-gene pairs have been validated by *in vitro* experiments (27).

In the present study, LINC00963 was a hub lncRNA of the positively co-expressed DElncRNA-DEmRNA network. Among its 154 co-expressed DEmRNAs, SOCS3 and adrenomedullin (ADM) were two of the top 20 DEmRNAs in PMOP. ADM is a 52-amino acid peptide with several biological functions. Previous studies have demonstrated that ADM is closely associated with regulating bone formation (28). The expression of ADM has been detected in chondrocytes and osteoblasts (29). ADM can promote growth of chondrocytes and osteoblasts *in vitro* (29). Additionally, apoptotic cell death in serum-starved osteoblasts can be reduced by ADM (30).




Figure 3. Significantly enriched GO terms and KEGG pathways of DEmRNAs co-expressed with DElncRNAs in PMOP. GO and KEGG pathway enrichment analyses of DEmRNAs co-expressed with DElncRNAs was performed using the online GeneCoDis3 tool (<http://genecodis.cnb.csic.es/analysis>). A P-value was obtained using a hypergeometric test. FDR (corrected P-value) < 0.05 was set as the cut-off for significant GO terms and KEGG pathways. The y-axis shows GO terms or KEGG pathways and the x-axis presents counts of DEmRNAs in PMOP enriched in GO terms or KEGG pathways. The color scale represented -log FDR (A) BP; (B) MF; (C) CC; (D) KEGG pathways. GO, Gene Ontology; KEGG, Kyoto Encyclopedia of Genes and Genomes; DE, differentially expressed; lncRNAs, long non-coding RNAs; PMOP, post-menopausal osteoporosis; BP, biological process; MF, molecular function; CC, cellular component; FDR, false discovery rate.




Figure 4. Validation of selected DElncRNAs and DEmRNAs in GSE56815. The x-axes show PMOP (case) and normal control groups, and the y-axes show the log₂ transformation to the intensities. Student's t-test was used to calculate the significant differences in DElncRNAs and DEmRNAs between PMOP and normal control groups. (A) alkaline phosphatase, liver/bone/kidney; (B) suppressor of cytokine signaling 3; (C) CD177; (D) secretory leukocyte peptidase inhibitor; (E) LINC00963. DE, differentially expressed; lncRNAs, long non-coding RNAs; PMOP, post-menopausal osteoporosis.

In the present study, the expression of ADM was significantly downregulated in the blood samples of patients with PMOP. It was hypothesized that downregulated ADM may be involved in PMOP through reducing the bone formation induced by the reduced proliferation of osteoblasts. The SOCS family includes cytokine-inducible negative regulators of cytokine signaling. As a member of the SOCS family, SOCS3 can be regulated by various cytokines (31). Previous studies have reported that increased SOCS3 elevated transforming growth factor- β , TNF- α and RANK ligand (RANKL)-induced osteoclast formation, and promoted precursors to the osteoclast lineage through the inhibition of specific anti-osteoclastic Janus kinase/signal transducer and activator of transcription signals (31). In addition, increased SOCS3 is closely associated with inflammation-induced bone loss (32). SOCS3 is also involved in RANKL-mediated dendritic cell-derived osteoclastogenesis by regulating associated cytokine signaling (32). Diabetes-associated inflammation-induced alveolar bone loss can also be regulated by SOCS3. In the present

study, downregulated SOCS3 was detected in blood samples of patients with PMOP, which suggested that SOCS3 may also be a regulator of PMOP. It was hypothesized that LINC00963-ADM and LINC00963-SOCS3 interactions may be key in PMOP.

Another lncRNA, GAS5, has been reported to regulate apoptosis in prostate cancer, breast cancer, renal cell carcinoma and gastric cancer (33-36). In the present study, GAS5 was a significantly upregulated DElncRNA in PMOP, which had four nearby DEmRNAs (centromere protein L, zinc finger and BTB domain containing 37, serpin family C member 1, and ring finger and CCCH-type domains 1). It was hypothesized that GAS5 may be involved in PMOP by regulating apoptosis and these four genes.

lncRNAs have also been shown to regulate gene expression in *cis*. LOC105376834-ALPL and LOC101929866-SLPI were two DElncRNA-DEmRNA co-expression pairs in PMOP. In addition, ALPL and SLPI were nearby DEmRNAs of LOC105376834 and LOC101929866, respectively. It was




Figure 5. Validation of selected DElncRNAs in GSE7158. The x-axes show PMOP (case) and normal control groups, and the y-axes show relative expression levels. Student's t-test was used to calculate the significant differences in DElncRNAs between the PMOP and normal control groups. (A) PSMD5-AS1; (B) PAX8-AS1; (C) JHDM1D-AS1; (D) LINC00963; (E) LOC100506113; (F) HCG27. * $P < 0.05$. DE, differentially expressed; lncRNAs, long non-coding RNAs; PMOP, post-menopausal osteoporosis.

hypothesized that LOC105376834 and LOC101929866 may regulate the expression of ALPL and SLPI by a *cis*-effect, in which the DElncRNAs were also co-expressed with DElncRNAs. ALPL is an osteoblast marker and is reported to be closely associated with the development of osteoporosis (37,38). Downregulated ALPL can reflect decreased activity of osteoblasts, bone formation and extracellular matrix mineralization (37). Previous studies have detected downregulated ALPL in bone tissue samples of patients with PMOP and ovariectomized mice, a model of postmenopausal osteoporosis (37,39,40). In the present study, the downregulation of ALPL was detected in blood samples from patients with PMOP, which confirmed the importance of ALPL in PMOP and may serve as a diagnostic marker of PMOP. As ALPL was a nearby co-expressed DEmRNA of LOC105376834, it was hypothesized that LOC105376834 may be involved in PMOP by *cis*-regulating the expression of ALPL.

SLPI encodes a serine protease inhibitor, which protects epithelial tissues from serine proteases. Additionally, SLPI is an anti-inflammatory mediator (41). SLPI can contribute to wound healing by decreasing the excessive inflammatory response, elevating keratinocyte proliferation and increasing collagen deposition by suppressing the activity of protease (42). To the best of our knowledge, the association between SLPI and PMOP has not been reported previously. A significant downregulation of SLPI was detected in patients with PMOP in the present study. As accumulated evidence has indicated that various inflammatory conditions are involved with osteoporosis (43), the present study hypothesized that SLPI may be involved in PMOP by regulating the inflammatory condition. In addition, estrogen treatment has been shown to increase the expression of SPLI in alveolar epithelial cells in ovariectomized mice (44). The same result was found in the rat uterus following treatment with estrogen (45). It was hypothesized that reduced estrogen may be involved in PMOP by regulating SLPI. SLPI was the nearby co-expressed DEmRNA of LOC101929866, which suggested that LOC101929866 may

be associated with PMOP. The other two nearby co-expressed DEmRNAs (potassium voltage-gated channel subfamily S member 1, and peptidase inhibitor 3) of LOC101929866 may also be involved in PMOP.

CD177 was the third significant DEmRNA in PMOP, which may also be an estrogen-associated gene. CD177 encodes a glycosyl-phosphatidylinositol-linked cell surface glycoprotein associated with neutrophil activation. Although there was no previous report on the association between CD177 and PMOP, a low expression of CD177 was found to be involved in clonal myeloid disorders, particularly myelodysplasia (46). A significantly upregulated level CD177 was previously detected in breast cancer cells following treatment with estrogen receptors- β agonists (47), which suggested that CD177 was closely associated with estrogen. It was hypothesized that reduced CD177 may also be involved in PMOP by regulating estrogen. The precise role of CD177 in PMOP requires further investigation.

Besides LOC105376834 and LOC10192986, LOC105374771 and LOC100506113 were two downregulated DElncRNAs in PMOP, which had nearby co-expressed DEmRNAs. Therefore, LOC100506113 and LOC105374771 may be involved in PMOP by regulating the expression of diacylglycerol O-acyltransferase 2 and LGALS1, respectively. In addition, LOC105374771 was the most markedly downregulated lncRNA, which was co-expressed with 130 DEmRNAs, including ALPL, SOCS3, ADM, CD177 and SLPI. LOC105374771 may affect the pathogenesis of PMOP by regulating the expression of these DEmRNAs.

Besides LINC00963, the other hub lncRNAs of the positively and negatively co-expressed DElncRNAs-DEmRNAs network were LOC105378415, LOC105377067, HCG27, LOC101928143 and LINC01094. Three PMOP-associated DEmRNAs, including ALPL, SOCS3 and ADM, were common co-expressed DEmRNAs of these hub DElncRNAs, which indicated the importance of these DElncRNAs in PMOP.

As hematopoietic cell lineage and osteoclast differentiation are two well-known pathways in PMOP. DEmRNAs enriched in these two pathways and their co-expressed DElncRNAs may be involved in PMOP by regulating hematopoietic cell lineage or osteoclast differentiation.

In conclusion, the present study identified five DEmRNAs (ALPL, SOCS3, ADM, SLPI and CD177) co-expressed with DElncRNAs, which may be involved in PMOP. DElncRNAs in PMOP, including LINC00963, LOC105376834, LOC101929866, LOC105374771 and LOC100506113, may be involved in the pathogenesis of PMOP by regulating the expression of their nearby and co-expressed DEmRNAs and the pathway of osteoclast differentiation. The results of the present study may provide a foundation for future investigations of lncRNAs in PMOP and contribute in developing novel diagnostic biomarkers and drug design for PMOP. However, the sample size for RNA sequencing in the present study was small, and the difference in body mass index between the PMOP and NC groups may have affected the results of RNA-sequencing, which were limitations of the study. Although the validation based on GSE56815 and GSE7158 suggested that the RNA-sequencing results were generally reliable, investigations with a larger sample size are required to confirm this conclusion. In addition, further experiments are required to address the biological significance of key lncRNAs and genes in PMOP.

Acknowledgements

Not applicable.

Funding

This study was supported by the Beijing Health System High Level Health Technical Personnel Training Project (grant no. 2015-3-009).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

QF and AG were responsible for conception and design of the experiments. QF and XDB performed the experiments. XDB and JSL analyzed the data. HM and YY supplied reagents, materials and analysis tools. All named authors wrote this manuscript and have agreed to the publication of this manuscript, and it does not infringe on any copyright or property rights.

Ethics approval and consent to participate

All individuals provided written informed consent for use of their samples in the present study. The present study was approved by the Ethics Committee of Beijing Friendship Hospital, Capital Medical University (Beijing, China; 2017-P2-084-01).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Zhang Y, Liu H, Zhang C, Zhang T, Zhang B, Li L, Chen G, Fu D and Wang K: Endochondral ossification pathway genes and postmenopausal osteoporosis: Association and specific allele related serum bone sialoprotein levels in Han Chinese. *Sci Rep* 5: 16783, 2015.
2. Ma M, Chen X, Lu L, Yuan F, Zeng W, Luo S, Yin F and Cai J: Identification of crucial genes related to postmenopausal osteoporosis using gene expression profiling. *Aging Clin Exp Res* 28: 1067-1074, 2016.
3. Taguchi A, Ohtsuka M, Nakamoto T, Naito K, Tsuda M, Kudo Y, Motoyama E, Suei Y and Tanimoto K: Identification of post-menopausal women at risk of osteoporosis by trained general dental practitioners using panoramic radiographs. *Dentomaxillofac Radiol* 36: 149-154, 2007.
4. Sasser AC, Taylor M, Birnbaum HG, Schoenfeld MJ, Oster EF and Rousculp M: Assessing the economic impact of chronic conditions in postmenopausal women. *Expert Opin Pharmacother* 6: 1803-1814, 2005.
5. Sasser AC, Rousculp MD, Birnbaum HG, Oster EF, Lufkin E and Mallet D: Economic burden of osteoporosis, breast cancer, and cardiovascular disease among postmenopausal women in an employed population. *Womens Health Issues* 15: 97-108, 2005.
6. Richards JB, Zheng HF and Spector TD: Genetics of osteoporosis from genome-wide association studies: Advances and challenges. *Nat Rev Genet* 13: 576-588, 2012.
7. Liu YJ, Zhang L, Papasian CJ and Deng HW: Genome-wide association studies for osteoporosis: A 2013 update. *J Bone Metab* 21: 99-116, 2014.
8. Kotrych D, Dziedzic V, Safranow K, Sroczynski T, Staniszewska M, Juzyszyn Z and Pawlik A: TNF- α and IL10 gene polymorphisms in women with postmenopausal osteoporosis. *Eur J Obstet Gynecol Reprod Biol* 199: 92-95, 2016.
9. Shang DP, Lian HY, Fu DP, Wu J, Hou SS and Lu JM: Relationship between estrogen receptor 1 gene polymorphisms and postmenopausal osteoporosis of the spine in Chinese women. *Genet Mol Res* 15, 2016.
10. Zhang C, Ma J, Chen G, Fu D, Li L and Li M: Evaluation of common variants in CNR2 gene for bone mineral density and osteoporosis susceptibility in postmenopausal women of Han Chinese. *Osteoporos Int* 26: 2803-2810, 2015.
11. Luo D, Liu Y, Zhou Y, Chen Z, Yang L, Liu Y, Xu Q, Xu H, Kuang H, Huang Q, *et al*: Association between dietary phytoestrogen intake and bone mineral density varied with estrogen receptor alpha gene polymorphisms in southern Chinese postmenopausal women. *Food Funct* 6: 1977-1983, 2015.
12. Singh M, Singh P, Singh S, Juneja PK and Kaur T: Vitamin D receptor (VDR) gene polymorphism influences the risk of osteoporosis in postmenopausal women of Northwest India. *Arch Osteoporos* 8: 147, 2013.
13. Kitjaroenanth A, Hananantachai H, Phonrat B, Preutthipan S and Tungtrongchitr R: Low density lipoprotein receptor-related protein 5 gene polymorphisms and osteoporosis in Thai menopausal women. *J Negat Results Biomed* 15: 16, 2016.
14. Cvijetic S, Grazio S, Kosovic P, Uremovic M, Nemic T and Bobic J: Osteoporosis and polymorphisms of osteoprotegerin gene in postmenopausal women-a pilot study. *Reumatologia* 54: 10-13, 2016.
15. Terashima M, Tange S, Ishimura A and Suzuki T: MEG3 Long Noncoding RNA contributes to the epigenetic regulation of epithelial-mesenchymal transition in lung cancer cell lines. *J Biol Chem* 292: 82-99, 2017.
16. Rajpathak SN, Vellarikkal SK, Patowary A, Scaria V, Sivasubbu S and Deobagkar DD: Human 45,X fibroblast transcriptome reveals distinct differentially expressed genes including long noncoding RNAs potentially associated with the pathophysiology of Turner syndrome. *PLoS One* 9: e100076, 2014.
17. Wang Q, Li Y, Zhang Y, Ma L, Lin L, Meng J, Jiang L, Wang L, Zhou P and Zhang Y: LncRNA MEG3 inhibited osteogenic differentiation of bone marrow mesenchymal stem cells from postmenopausal osteoporosis by targeting miR-133a-3p. *Biomed Pharmacother* 89: 1178-1186, 2017.

18. Wang L, Han S, Jin G, Zhou X, Li M, Ying X, Wang L, Wu H and Zhu Q: Linc00963: A novel, long non-coding RNA involved in the transition of prostate cancer from androgen-dependence to androgen-independence. *Int J Oncol* 44: 2041-4449, 2014.
19. Mercer TR, Dinger ME and Mattick JS: Long non-coding RNAs: Insights into functions. *Nat Rev Genet* 10: 155-159, 2009.
20. Costa FF: Non-coding RNAs: New players in eukaryotic biology. *Gene* 357: 83-94, 2005.
21. Fitzgerald KA and Caffrey DR: Long noncoding RNAs in innate and adaptive immunity. *Curr Opin Immunol* 26: 140-146, 2014.
22. Liao Q, Liu C, Yuan X, Kang S, Miao R, Xiao H, Zhao G, Luo H, Bu D, Zhao H, *et al*: Large-scale prediction of long non-coding RNA functions in a coding-non-coding gene co-expression network. *Nucleic Acids Res* 39: 3864-3878, 2011.
23. Dong R, Jia D, Xue P, Cui X, Li K, Zheng S, He X and Dong K: Genome-wide analysis of long noncoding RNA (lncRNA) expression in hepatoblastoma tissues. *PLoS One* 9: e85599, 2014.
24. Wang P, Fu H, Cui J and Chen X: Differential lncRNA-mRNA co-expression network analysis revealing the potential regulatory roles of lncRNAs in myocardial infarction. *Mol Med Rep* 13: 1195-1203, 2016.
25. Fang M, Zhang P, Zhao Y and Liu X: Bioinformatics and co-expression network analysis of differentially expressed lncRNAs and mRNAs in hippocampus of APP/PS1 transgenic mice with Alzheimer disease. *Am J Transl Res* 9: 1381-1391, 2017.
26. Yang L, Yi K, Wang H, Zhao Y and Xi M: Comprehensive analysis of lncRNAs microarray profile and mRNA-lncRNA co-expression in oncogenic HPV-positive cervical cancer cell lines. *Oncotarget* 7: 49917-49929, 2016.
27. Wan ZY, Song F, Sun Z, Chen YF, Zhang WL, Samartzis D, Ma CJ, Che L, Liu X, Ali MA, *et al*: Aberrantly expressed long noncoding RNAs in human intervertebral disc degeneration: A microarray related study. *Arthritis Res Ther* 16: 465, 2014.
28. Martínez-Herrero S, Larrayoz IM, Ochoa-Callejero L, Fernández LJ, Allueva A, Ochoa I and Martínez A: Prevention of bone loss in a model of postmenopausal osteoporosis through adrenomedullin inhibition. *Front Physiol* 7: 280, 2016.
29. Montuenga LM, Martínez A, Miller MJ, Unsworth EJ and Cuttitta F: Expression of adrenomedullin and its receptor during embryogenesis suggests autocrine or paracrine modes of action. *Endocrinology* 138: 440-451, 1997.
30. Uzan B, Villemain A, Garel JM and Cressent M: Adrenomedullin is anti-apoptotic in osteoblasts through CGRP1 receptors and MEK-ERK pathway. *J Cell Physiol* 215: 122-128, 2008.
31. Sun GJ, Guo T, Chen Y, Xu B, Guo JH and Zhao JN: Significant pathways detection in osteoporosis based on the bibliometric network. *Eur Rev Med Pharmacol Sci* 17: 1-7, 2013.
32. Zhang X, Alnaeeli M, Singh B and Teng YT: Involvement of SOCS3 in regulation of CD11c⁺ dendritic cell-derived osteoclastogenesis and severe alveolar bone loss. *Infect Immun* 77: 2000-2009, 2009.
33. Pickard MR, Mourtada-Maarabouni M and Williams GT: Long non-coding RNA GAS5 regulates apoptosis in prostate cancer cell lines. *Biochim Biophys Acta* 1832: 1613-1623, 2013.
34. Sun M, Jin F, Xia R, Kong R, Li JH, Xu TP, Liu YW, Zhang EB, Liu XH and De W: Decreased expression of long noncoding RNA GAS5 indicates a poor prognosis and promotes cell proliferation in gastric cancer. *BMC Cancer* 14: 319, 2014.
35. Qiao HP, Gao WS, Huo JX and Yang ZS: Long non-coding RNA GAS5 functions as a tumor suppressor in renal cell carcinoma. *Asian Pac J Cancer Prev* 14: 1077-1082, 2013.
36. Mourtada-Maarabouni M, Pickard MR, Hedge VL, Farzaneh F and Williams GT: GAS5, a non-protein-coding RNA, controls apoptosis and is downregulated in breast cancer. *Oncogene* 28: 195-208, 2009.
37. Balla B, Kosa JP, Kiss J, Borsy A, Podani J, Takács I, Lazáry A, Nagy Z, Bácsi K, Speer G, *et al*: Different gene expression patterns in the bone tissue of aging postmenopausal osteoporotic and non-osteoporotic women. *Calcif Tissue Int* 82: 12-26, 2008.
38. Rodríguez JP, Montecinos L, Ríos S, Reyes P and Martínez J: Mesenchymal stem cells from osteoporotic patients produce a type I collagen-deficient extracellular matrix favoring adipogenic differentiation. *J Cell Biochem* 79: 557-565, 2000.
39. Hsiao HB, Lin H, Wu JB and Lin WC: Kinsenoside prevents ovariectomy-induced bone loss and suppresses osteoclastogenesis by regulating classical NF- κ B pathways. *Osteoporos Int* 24: 1663-1676, 2013.
40. de Castro LF, Lozano D, Portal-Núñez S, Maycas M, De la Fuente M, Caeiro JR and Esbrit P: Comparison of the skeletal effects induced by daily administration of PTHrP (1-36) and PTHrP (107-139) to ovariectomized mice. *J Cell Physiol* 227: 1752-1760, 2012.
41. McKiernan PJ, McElvaney NG and Greene CM: SLPI and inflammatory lung disease in females. *Biochem Soc Trans* 39: 1421-1426, 2011.
42. Park JJ, Bae CS, Choi BD, Jeong SJ, Wang G, Lim DS, Kim BO, Cho YS, Kim SJ and Jeong MJ: Induction of secretory leukocyte protease inhibitor (SLPI) in estradiol valerate (EV) induced polycystic ovary. *Arch Pharm Res* 34: 1389-1397, 2011.
43. Al-Daghri NM, Aziz I, Yakout S, Aljohani NJ, Al-Saleh Y, Amer OE, Sheshah E, Younis GZ and Al-Badr FB: Inflammation as a contributing factor among postmenopausal Saudi women with osteoporosis. *Medicine (Baltimore)* 96: e5780, 2017.
44. Draijer C, Hylkema MN, Boersma CE, Klok PA, Robbe P, Timens W, Postma DS, Greene CM and Melgert BN: Sexual maturation protects against development of lung inflammation through estrogen. *Am J Physiol Lung Cell Mol Physiol* 310: L166-L174, 2016.
45. Chen DJ, Xu X, Cheon YP, Bagchi MK and Bagchi IC: Estrogen induces expression of secretory leukocyte protease inhibitor in rat uterus. *Biol Reprod* 71: 508-514, 2004.
46. Meyerson HJ, Osei E, Schweitzer K, Blidaru G, Edinger A and Balog A: CD177 expression on neutrophils: In search of a clonal assay for myeloid neoplasia by flow cytometry. *Am J Clin Pathol* 140: 658-669, 2013.
47. Lattrich C, Schüler S, Häring J, Skrzypczak M, Ortmann O and Treeck O: Effects of a combined treatment with tamoxifen and estrogen receptor β agonists on human breast cancer cell lines. *Arch Gynecol Obstet* 289: 163-171, 2014.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.