Thromboxane A2 receptor antagonist SQ29548 attenuates SH-SY5Y neuroblastoma cell impairments induced by oxidative stress

GAOYU CAI^{1*}, AIJUAN YAN^{2*}, NINGZHEN FU³ and YI FU¹

¹Department of Neurology, Rui Jin Hospital, Shanghai Jiao Tong University, Shanghai 200025;

²Department of Neurology, Xin Hua Hospital, Shanghai Jiao Tong University, Shanghai 200082;

³Department of Pancreatic Surgery, Rui Jin College of Clinical Medicine, Rui Jin Hospital, Shanghai Jiao Tong University, Shanghai 200025, P.R. China

Received September 28, 2017; Accepted March 21, 2018

DOI: 10.3892/ijmm.2018.3589

Abstract. Thromboxane A2 receptor (TXA2R) serves a vital role in numerous neurological disorders. Our previous study indicated that SQ29548, an antagonist of TXA2R, attenuated the induced neuron damage in cerebral infarction animals; however, the underlying mechanism remains unknown. Certain studies revealed a new role of TXA2R in the regulation of oxidative stress, which is one of the basic pathological processes in neurological disorders. Thus, the present study attempted to examine whether the inhibition of TXA2R with SQ29548 helped to protect the nerve cells against oxidative stress. SQ29548 was utilized as a TXA2R antagonist, and relevant assays were performed to detect the cell viability, cellular reactive oxygen species (ROS) level, cell apoptosis, expression levels of superoxide dismutase-2 (SOD2), catalase and caspases, and activation of mitogen-activated protein kinase (MAPK) pathways. It was observed that hydrogen peroxide (H₂O₂) dose-dependently reduced the viability of SH-SY5Y cells. In addition, H₂O₂ raised the level of ROS in cells, inhibited the expression levels of SOD2 and catalase, and potentially enhanced cell apoptosis and the expression of caspases via activating the MAPK pathways. Pretreatment with SQ29548 not only rescued the viability of SH-SY5Y cells, but also ameliorated the intracellular ROS level and the expression levels of SOD2 and catalase. Furthermore, it decreased the cell apoptosis and the expression of caspases, possibly via the inhibition of MAPK pathways. In conclusion,

SQ29548, an antagonist of TXA2R, improved the antioxidant capacities of SH-SY5Y cells and reduced the cell apoptosis through the inhibition of MAPK pathways.

Introduction

Thromboxane A2 receptor (TXA2R), a member of the G protein-coupled receptor family (1), is broadly distributed in platelets (2), as well as epithelial (3), smooth muscle (4), glial and nerve cells in the brain (5). TXA2R is regarded as a traditional coagulation and inflammation-associated receptor, which is also closely associated with neurological disorders. A previous study of our research group indicated that mutation on TXA2R influenced the morbidity of cerebral infarctions in humans, which may be associated with the changes of coagulation functions (6). Also, it was observed that SQ29548, an antagonist of TXA2R, was able to attenuate the cerebral ischemia-reperfusion damage in mice and improve their behavioral scores (7), thus revealing the potential therapeutic effect of TXA2R antagonists. However, the underlying mechanisms involved remain unclear. A number of studies revealed a new role for TXA2R in regulating the cellular oxidative stress and the consequent apoptotic process (8-10). TXA2R antagonists are reported to ameliorate organic oxidative stress and protect the function of organs (9), which provided a new clue for the treatment of oxidative stress-associated nervous system diseases.

Oxidative stress is an important contributor in the pathogenesis and pathological processes of several neurological disorders (11,12). In general, the cellular reactive oxygen species (ROS) production and the antioxidative defense were observed to be in a dynamic balance. For instance, antioxidative enzymes (13), including superoxide dismutase (SOD) and catalase, efficiently eliminated the ROS produced by cellular respiration. Cerebral infarction (CI) or other neurological disorders lead to an excess of ROS production, which distorts the balance between ROS production and antioxidant defenses. The excessive ROS production results in cellular damage and apoptosis via attacking the lipid, protein and DNA of cells, and activating signaling pathways, such as the

Correspondence to: Dr Yi Fu, Department of Neurology, Rui Jin Hospital, School of Medicine, Shanghai Jiao Tong University, 197 Rui Jin Er Road, Shanghai 200025, P.R. China
E-mail: fuyiki@sina.com

*Contributed equally

Key words: antagonists, neurons, neuroprotection, oxidative stress, thromboxane A2 receptor

mitogen-activated protein kinase (MAPK) pathways (14,15). Numerous studies have proven that the elevation of cellular antioxidative stress capacity was beneficial for the limitation of ROS generation, acceleration of ROS elimination, alleviation of ROS damage and inhibition of MAPK signaling pathways to reduce cell death (13,16). Compared with other organs (17), brain tissue is sensitive to oxidative stress damage due to its characteristics of high energy consumption and weak antioxidative stress capacity. Therefore, the identification of a novel effective neuroprotective agent is essential in the prevention and treatment of oxidative stress-associated neurological disorders.

In view of the association between TXA2R and oxidative stress, increasing attention has concentrated on the role of TXA2R and its antagonists in the treatment of oxidative stress-associated neurological disorders and the underlying molecular mechanisms. In the present study, an *in vitro* oxidative stress model was established in order to investigate whether the TXA2R antagonist SQ29548 regulated the cellular oxidative stress response in nerve cells and determined the mechanisms involved.

Materials and methods

Reagents. Hydrogen peroxide (H_2O_2 ; cat. no. 323381; used for cell experiments), dimethyl sulfoxide (DMSO; cat. no. D2650), poly-D-lysine hydrobromide (cat. no. P6407), nicotinamide adenine dinucleotide (NADH; cat. no. N4505), sodium pyruvate (cat. no. P4562) and Triton X-100 (cat. no. T8787) were purchased from Sigma-Aldrich (Merck KGaA, Darmstadt, Germany). SQ29548 (cat. no. 19025) was obtained from Cayman Chemical Company (Ann Arbor, MI, USA). Bovine serum albumin (BSA; cat. no. 36102ES) was from Yeasen Biotechnology Co., Ltd. (Shanghai, China). Water-soluble tetrazolium salt (WST-1; cat. no. C0035), 2',7'-dichlorofluorescein diacetate (DCFH-DA; cat. no. S0033) and DAPI (cat. no. C1002) were purchased from Beyotime Institute of Biotechnology (Shanghai, China). ApoScreen Annexin V Apoptosis-phycoerythrin (PE) kit (cat. no. 10010-09) was obtained from Southern Biotech (Birmingham, AL, USA). Terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL; cat. no. TUN002) staining reagents were purchased from Mai bio (Shanghai, China). An antibody against TXA2R was from Santa Cruz Biotechnology, Inc. (Dallas, TX, USA; cat. no. sc-30036). Antibodies against c-Jun N-terminal kinases (JNK; cat. no. 9926), extracellular signal-regulated kinase (ERK; cat. no. 9926), p38 (cat. no. 9926), phosphorylated (p)-JNK (cat. no. 9910), p-ERK (cat. no. 9910), p-p38 (cat. no. 9910), cleaved caspase3 (cat. no. 9661), caspase3 (cat. no. 9662), cleaved caspase9 (cat. no. 9501), caspase9 (cat. no. 9502) and β -tubulin (cat. no. 15115), as well as the anti-rabbit horseradish peroxidase-conjugated secondary antibody (cat. no. 7074), were purchased from Cell Signaling Technology, Inc. (Danvers, MA, USA). Antibodies for SOD2 (cat. no. ab13534) and catalase (cat. no. ab16731) were from Abcam (Cambridge, MA, USA). An antibody for NeuN (cat. no. MAB377) was from EMD Millipore (Billerica, MA, USA). The Alexa-594 antibody (cat. no. A21203; Invitrogen), TRIzol reagent (cat. no. 15596026) and BCA protein detection kit (cat. no. 23225) were purchased from Thermo Fisher

Scientific, Inc. (Waltham, MA, USA). Paraformaldehyde (PFA; cat. no. 80096618), potassium phosphate (cat. no. 10017518), 30% hydrogen peroxide (cat. no. 10011208; used for TUNEL staining) from Sinopharm Chemical Reagent Co., Ltd. (Shanghai, China). The primers used in this study were synthesised by Sangon Biotech Co., Ltd. (Shanghai, China).

Cell culture and oxidative stress model. SH-SY5Y cells were acquired from the Cell Bank of Shanghai Institute of Biochemistry and Cell Biology, Chinese Academy of Sciences (Shanghai, China). SH-SY5Y cells were incubated in the Dulbecco's modified Eagle's medium (DMEM; HyClone; GE Healthcare, Logan, UT, USA) supplemented with 10% fetal bovine serum (FBS; HyClone; GE Healthcare) and 1% penicillin-streptomycin (Invitrogen; Thermo Fisher Scientific, Inc.) at 37°C in a 5% CO_2 incubator. When the cell confluence reached 80%, cells were collected and transferred into 96-well plates at a cell density of 2×10^4 cells/well for WST-1 assay, while 24-well plates (1×10^5 cells/well) were used for DCFH-DA, TUNEL and lactate dehydrogenase (LDH) assays, and 6-well plates (3×10^5 cells/well) were used for western blot analysis, reverse transcription-quantitative polymerase chain reaction (RT-qPCR) and flow cytometry. Subsequently, the cells were incubated in DMEM without sodium pyruvate with 10% FBS and 1% penicillin-streptomycin for 24 h. All the plates used were pretreated with poly-D-lysine hydrobromide solution (0.1 mg/ml) for 24 h and then washed three times with phosphate-buffered saline (PBS; HyClone; GE Healthcare). A total of 24 h after the cells transferred to the plates, DMSO was added to the control group and SQ29548 (dissolved in DMSO) in different concentrations were added to the treatment groups. After 2-h treatment, the cells were washed three times with PBS. Next, the cells were treated with or without H_2O_2 in different concentrations in the medium for 1 h, followed by washing three times with PBS. Finally, DMEM (without sodium pyruvate) with 10% FBS and 1% penicillin-streptomycin was used for subsequent culture and experiments.

WST-1 assay. A WST-1 assay was utilized to detect the cell viability, according to the protocol provided by the manufacturer. After transferring the cells to the plates for 24 h, the cells were treated with SQ29548 (0, 0.1, 0.5, 1.0 and 10 μ M) for 2 h or treated with H_2O_2 (0, 0.05, 0.1, 0.2, 0.5 and 1.0 mM) for 1 h or treated with H_2O_2 (0, 0.1 mM) for 1 h following pretreatment with SQ29548 (0, 0.1, 0.5, 1.0 and 10 μ M) for 2 h. Subsequent to the incubation for 24 h, 10 μ l WST-1 solution was added to the SH-SY5Y cells. After 1 h of incubation at 37°C, a microplate reader (Bio-Rad Laboratories, Inc., Hercules, CA, USA) was used to measure the absorbance of each well at 450 nm.

LDH assay. An LDH assay was conducted as reported previously (18). The cells were treated with H_2O_2 (0, 0.1 mM) for 1 h after pretreated with SQ29548 (0, 0.1, 1.0 μ M) for 2 h. After 24 h of incubation, 150 μ l culture medium of post-treated SH-SY5Y cells was collected and added into 96-well plates. Next, 150 μ l reaction buffer containing 500 mM potassium phosphate, 7.5 mM sodium pyruvate and 1.5 mM NADH was added successively and the absorbance of each well at 340 nm was measured over 30 sec with the microplate reader.

Intracellular ROS level assay. The intracellular ROS level was detected with DCFH-DA reagents. The cells were treated with H_2O_2 (0 or 0.1 mM) for 1 h following pretreatment with SQ29548 (0, 0.1, 1.0 μ M) for 2 h. Treated SH-SY5Y cells were washed three times with PBS after 1-h incubation. Next, 200 μ l DMEM (without sodium pyruvate, serum and antibiotics) with 10 μ M DCFH-DA was added into each well. After 20 min of incubation at 37°C, DMEM (without sodium pyruvate, serum and antibiotics) was used to wash the cells for three times. The plates were then observed with an inverted fluorescence microscope (Leica Microsystems GmbH, Wetzlar, Germany) at an excitation wavelength of 488 nm and an emission wavelength of 525 nm. All the data were subjected to fluorescence intensity analysis with ImageJ software, version 1.14 (National Institutes of Health, Bethesda, MA, USA).

Annexin V-PE and 7-aminoactinomycin D (7-AAD) double-staining assay. Flow cytometry was performed as reported previously (19). The cells were treated with H_2O_2 (0, 0.1 mM) for 1 h following pretreatment with SQ29548 (0, 0.1, 1.0 μ M) for 2 h. Treated SH-SY5Y cells were digested after 1-h incubation, and 10 μ l of which was then used for cell counting with Fuchs-Rosenthal counting chambers. Subsequently, the cells were centrifuged at 74 x g for 4 min at 4°C. Centrifuged cells were resuspended with cold 1X Annexin binding buffer to obtain a density of a 1×10^6 cells/ml. A total of 10 μ l Annexin V-PE was added to 100 μ l cell suspension and the cell suspension was then incubated in the dark for 15 min at 4°C. After the incubation, 380 μ l cold 1X Annexin binding buffer and 10 μ l 7-AAD were added to the cell suspension at 4°C, and the cell suspension was immediately subjected to apoptosis assay by flow cytometry (Bio-Rad Laboratories, Inc.).

TUNEL staining. The cells were treated with H_2O_2 (0, 0.1 mM) for 1 h following pretreatment with SQ29548 (0, 0.1, 1.0 μ M) for 2 h. After 1 h of incubation, the treated SH-SY5Y cells were fixed with 4% PFA for 10 min at 4°C and sealed with methanol containing 3% H_2O_2 , and 0.3% Triton X-100 was then used for membrane rupture. Next, 50 μ l *In Situ* Cell Death reagent was added to each slide, and then cells were incubated without light exposure for 60 min. Subsequent to 150 μ l DAPI staining (1:300) for 3 min at room temperature, slides were sealed and observed with a confocal fluorescence microscope (Leica Microsystems GmbH).

RT-qPCR. The cells were treated with H_2O_2 (0, 0.1 mM) for 1 h following pretreatment with SQ29548 (0, 0.1 μ M) for 2 h. After 12 h of incubation, total RNA was extracted from the treated SH-SY5Y cells with TRIzol reagent and was reverse transcribed into cDNA with PrimeScript RT reagent kit (Takara Bio, Inc., Otsu, Japan), as previously described (20). qPCR was subsequently implemented on an Applied Biosystems 7500 Real-Time PCR system (Thermo Fisher Scientific, Inc.) using a SYBR Green kit (Takara Bio, Inc.) under the following conditions: Denaturation at 95°C for 10 sec, followed by 40 cycles at 95°C for 5 sec and at 60°C for 30 sec. The primers used in qPCR were as follows: SOD2, 5'-GGTGGTCATATCAATCATAG-3' (forward) and 5'-AGTGAATAAGGTTTGTGT-3' (reverse); catalase, 5'-TTAATCCATTCGATCTCACC-3' (forward) and 5'-GGCGGTGAGTGTCAGGATAG-3' (reverse); GAPDH,

5'-TCTTCTTTTTCGTCGCCAG-3' (forward) and 5'-GGG GGCAGAGATGATGACC-3' (reverse). GAPDH served as the loading control. All the data were analyzed with the comparative threshold cycle (Cq) method (21).

Western blot analysis. The cells were treated with H_2O_2 (0, 0.1 mM) for 1 h following pretreatment with SQ29548 (0, 0.1 μ M) for 2 h. Protein was extracted with standard lysis buffer (Cell Signaling Technology, Inc.). Following purification, a BCA protein detection kit was used for protein concentration measurement. Equal amounts of extracted protein (30 μ g) were separated with 10% SDS-PAGE. Subsequently the proteins were transferred to nitrocellulose membranes at 300 mA for 90 min. After 1 h of incubation in 5% BSA for blocking at room temperature, membranes were incubated at 4°C overnight with primary antibodies diluted at 1:1,000. Subsequent to washing three times with TBST, anti-rabbit horseradish peroxidase-conjugated secondary antibody diluted at 1:5,000 was added for 1-h incubation at room temperature. Following further washing with TBST for three times, the specific bands were visualized with an enhanced chemiluminescence reagent (Thermo Fisher Scientific, Inc.) in a gel imaging system (Tanon Science & Technology Co., Ltd., Shanghai, China). ImageJ software was applied for gray analysis of the data.

Statistical analysis. All the data were analyzed with GraphPad Prism software, version 5.0 (GraphPad Software, Inc., La Jolla, CA, USA). Comparison between two groups was performed with an unpaired t-test, while comparison among several groups was performed with one-way analysis of variance, followed by a Student-Newman-Keuls test. All the data are expressed as the mean $\pm$ standard error of the mean. $P < 0.05$ was regarded as an indicator of a statistically significant difference.

Results

SQ29548 attenuates the decrease in the viability of SH-SY5Y cells induced by H_2O_2 . Previous studies have reported (5,22) the expression of TXA2R on the membrane of nerve cells. In the present study, the expression of TXA2R on the surface of SH-SY5Y cells was confirmed by staining with TXA2R antibody and DAPI (Fig. 1A). Subsequently, to evaluate the function of SQ29548 and H_2O_2 on SH-SY5Y cell viability, a WST-1 assay was applied after cells were treated with different SQ29548 and H_2O_2 concentrations. It was observed that, compared with the untreated control group, treatment with DMSO or with 0.1, 0.5, 1.0 and 10 μ M SQ29548 resulted in cell viability of 98.86 ± 4.55 , 99.15 ± 2.90 , 95.97 ± 2.55 , 100.20 ± 1.86 and $95.33 \pm 4.03\%$, respectively (Fig. 1B). No statistically significant difference was observed among these groups ($P > 0.05$), indicating that SQ29548 did not evidently affect the SH-SY5Y cell viability. However, treatment with 0.05, 0.1, 0.2, 0.5 and 1.0 mM H_2O_2 reduced the SH-SY5Y cell viability to 83.01 ± 1.52 , 31.91 ± 1.85 , 28.16 ± 1.44 , 25.32 ± 1.44 and $23.30 \pm 1.65\%$, respectively, compared with the untreated control group (Fig. 1C). Considering other relevant findings (17,23), the concentration of 0.1 mM H_2O_2 was selected for subsequent experiments.

Figure 1. TXA2R existed on the surface of SH-SY5Y cells and SQ29548 inhibited the decrease of cell viability in SH-SY5Y cells induced by H₂O₂. (A) TXA2R was distributed on the surface of SH-SY5Y cells. A WST-1 assay was used to examine the changes in cell viability subsequent to treatment with (B) SQ29548, (C) H₂O₂ and (D) SQ29548 and H₂O₂ combined. (E) The LDH release of SH-SY5Y cells treated with SQ29548 and H₂O₂. Data are expressed as the mean ± standard error of the mean. n=3. *P<0.05, **P<0.01 and ***P<0.001. TXA2R, thromboxane A2 receptor; H₂O₂, hydrogen peroxide; DMSO, dimethyl sulfoxide; n.s., no significant difference.

Figure 2. SQ29548 reduced the ROS level in SH-SY5Y cells treated with H₂O₂. (A) DCFH-DA was performed to evaluate the intracellular ROS level. (B) Comparisons of DCF fluorescence intensity quantization among all the groups. The results are shown as the mean ± standard error of the mean (n=3). *P<0.05 and **P<0.01. ROS, reactive oxygen species; H₂O₂, hydrogen peroxide; DCFH-DA, 2',7'-dichlorofluorescein diacetate; DMSO, dimethyl sulfoxide; n.s., no significant difference.

Figure 3. SQ29548 ameliorated the decreases in SOD2 and catalase levels induced by H₂O₂ in SH-SY5Y cells. (A) Western blot analysis evaluated the protein expression levels of SOD2 and catalase. Gray value quantification of (B) SOD2 and (C) catalase expression levels. (D) SOD2 and (E) catalase mRNA expression levels are shown. Results are expressed as the mean $\pm$ standard error of the mean (n=3). *P<0.05, **P<0.01 and ***P<0.001. SOD2, superoxide dismutase-2; H₂O₂, hydrogen peroxide; DMSO, dimethyl sulfoxide; n.s., no significant difference.

An oxidative stress model was then established in SH-SY5Y cells by H₂O₂ treatment (0.1 mM). H₂O₂-induced cells that were also pretreated with 0.1, 0.5, 1.0 and 10 μ M SQ29548 for 2 h exhibited improved viability of 44.29 $\pm$ 1.40, 43.28 $\pm$ 1.07, 42.32 $\pm$ 2.03 and 40.27 $\pm$ 1.29%, respectively. By contrast, the cell viability of the H₂O₂-treated only group was 32.27 $\pm$ 1.54%, compared with that of the control group (Fig. 1D). The results indicated the protective effect of SQ29548 on cell viability. Furthermore, this protective effect was independent of the concentration of SQ29548.

The LDH assay demonstrated that, in the H₂O₂-treated only group, the LDH release was 3.30 $\pm$ 0.06-fold that of the control group; by contrast, 0.1 and 1.0 μ M SQ29548 significantly decreased the LDH release to 2.57 $\pm$ 0.09-fold and 2.74 $\pm$ 0.14-fold that of the control, respectively (Fig. 1E). Therefore, the protective effect of SQ29548 on cell viability was further confirmed.

SQ29548 reduces the intracellular ROS level in SH-SY5Y cells induced by H₂O₂. A DCFH-DA assay was performed to evaluate the effect of SQ29548 on the intracellular ROS level of SH-SY5Y cells (Fig. 2A). H₂O₂ increased the SH-SY5Y intracellular ROS level to 4.95 $\pm$ 0.97-fold that of the control group (Fig. 2B). However, pretreatment with 0.1 and 1.0 μ M SQ29548 decreased the intracellular ROS level to 2.01 $\pm$ 0.10-fold and 1.69 $\pm$ 0.16-fold, respectively, of the control group level (Fig. 2B). Thus, SQ29548 reduced the ROS generation in SH-SY5Y cells induced by H₂O₂.

SQ29548 ameliorates the decrease in SOD2 and catalase levels in SH-SY5Y cells induced by H₂O₂. To determine whether the change of ROS level was mediated by the regulatory effect of SQ29548 on intracellular antioxidant proteins, the intracellular SOD2 and catalase protein and mRNA levels were evaluated. Upon treatment with H₂O₂, the protein levels

of SOD2 (Fig. 3A and B) and catalase (Fig. 3A and C) were significantly inhibited, while their mRNA levels were also decreased to 56.05 $\pm$ 7.95% (Fig. 3D) and 45.83 $\pm$ 9.70% (Fig. 3E) of the control group levels, respectively. By contrast, pretreatment with SQ29548 significantly ameliorated the H₂O₂-induced decrease in the protein levels of SOD2 and catalase (Fig. 3B and C), while the mRNA levels were rescued to 85.70 $\pm$ 5.75% (Fig. 3D) and 80.17 $\pm$ 7.02% (Fig. 3E) of the control group levels, respectively. Therefore, SQ29548 appeared to ameliorate the H₂O₂-induced decrease in the expression levels of SOD2 and catalase in SH-SY5Y cells, and to improve the antioxidative stress capacities of SH-SY5Y cells.

SQ29548 reduces the apoptosis of SH-SY5Y cells induced by H₂O₂. Oxidative stress may result in cell damage and consequently induce cell apoptosis. To investigate the impact of SQ29548 on SH-SY5Y cell apoptosis, Annexin V-PE/7-AAD double-staining assay (Fig. 4A-C) and TUNEL staining (Fig. 4D and E) were selected to estimate the cell apoptosis. The Annexin V-PE/7-AAD double-staining assay revealed that H₂O₂ increased the ratio of early-stage and late-stage apoptosis in SH-SY5Y cells to 26.23 $\pm$ 2.46 and 15.4 $\pm$ 1.89%, as compared with the ratios of 5.82 $\pm$ 1.31 and 2.07 $\pm$ 1.07% in the control group, respectively (Fig. 4B and C). However, pretreatment with 0.1 and 1.0 μ M SQ29548 significantly abated the early-stage apoptosis cell ratio to 18.18 $\pm$ 1.56 and 16.33 $\pm$ 1.50%, respectively (Fig. 4B), while the late-stage apoptosis cell ratio was at 8.85 $\pm$ 0.57 and 8.58 $\pm$ 1.05%, respectively (Fig. 4C). No significant difference was observed between the 0.1 and 1.0 μ M SQ29548 groups (P>0.05).

TUNEL staining was also applied as a detector of cell apoptosis. Compared with the control group, H₂O₂ markedly raised the ratio of TUNEL-positive cells to 15.21 $\pm$ 2.71% from 0.90 $\pm$ 0.1%. Pretreatment with 0.1 and 1.0 μ M SQ29548 significantly reduced the ratio of TUNEL-positive cells to 9.19 $\pm$ 0.82

Figure 4. SQ29548 reduced the cell apoptosis of SH-SY5Y cells induced by H_2O_2 . (A) Flow cytometry analysis using Annexin V-PE/7-AAD double staining. Normal cells are observed in Q3 quadrant, early-stage apoptotic cells in Q4, late-stage apoptotic cells in Q2 and cell debris in Q1. Cell ratio of (B) early-stage and (C) late-stage apoptotic cells among all the groups. (D) TUNEL staining of the apoptotic cells. (E) Percentage of TUNEL-positive cells. Results are expressed as the mean $\pm$ standard error of the mean (n=3). * P <0.05 and ** P <0.01. H_2O_2 , hydrogen peroxide; DMSO, dimethyl sulfoxide; PE, phycoerythrin; 7-AAD, 7-aminoactinomycin D; TUNEL, terminal deoxynucleotidyl transferase dUTP nick end labeling; n.s., no significant difference.

Figure 5. SQ29548 inhibited the expression levels of cleaved caspase9 and cleaved caspase3 in SH-SY5Y cells induced by H_2O_2 . (A) Western blot analysis was performed to evaluate the expression levels of cleaved caspase9, caspase9, cleaved caspase3 and caspase3. Gray value quantification of (B) cleaved caspase9/total caspase9, (C) cleaved caspase3/total caspase3, (D) caspase9 and (E) caspase3. No significant difference was observed in the expression of total caspase9 and total caspase3 among the different groups. Results are expressed as the mean $\pm$ standard error of the mean (n=3). * P <0.05 and ** P <0.01. H_2O_2 , hydrogen peroxide; DMSO, dimethyl sulfoxide.

Figure 6. SQ29548 inhibited the activation of ERK and p38 signaling pathways in SH-SY5Y cells induced by H₂O₂. (A) Western blot analysis results on the expression levels of p-JNK, JNK, p-ERK, ERK, p-p38 and p38. (B) p-JNK/total JNK, (C) p-ERK/total ERK and (D) p-p38/total p38. No significant difference was observed in the expression levels of total JNK, ERK and p38 among the groups. Results are expressed as the mean ± standard error of the mean (n=3). *P<0.05, **P<0.01 and ***P<0.001. JNK, c-Jun N-terminal kinases; ERK, extracellular signal-regulated kinase; p-, phosphorylated; H₂O₂, hydrogen peroxide; DMSO, dimethyl sulfoxide; n.s., no significant difference.

and 8.00±1.00%, respectively (Fig. 4D and E), indicating that SQ29548 reduced the cell apoptosis induced by H₂O₂.

SQ29548 inhibits cleaved caspase expression in SH-SY5Y cells induced by H₂O₂. Caspases are typical apoptosis-associated proteins, whose cleaved products promote cell apoptosis. The present study attempted to determine how SQ29548 affected the expression of caspases and their cleaved products in SH-SY5Y cells induced by H₂O₂. The results demonstrated that no significant differences existed in the expression levels of caspase3 and caspase9 among all the groups (P>0.05), while H₂O₂ markedly elevated the expression levels of cleaved caspase9 and cleaved caspase3 (Fig. 5A). However, SQ29548-pretreated cells exhibited significantly inhibited expression levels of cleaved caspase9 (Fig. 5B) and cleaved caspase3 (Fig. 5C). Thus, SQ29548 possibly inhibited the expression of cleaved caspases.

SQ29548 inhibits ERK and p38 signaling pathway activation in SH-SY5Y cells induced by H₂O₂. MAPK pathways participate in a number of cellular physiological processes and regulate cell apoptosis induced by oxidative stress. In order to examine how SQ29548 interfered in MAPK pathways in SH-SY5Y cells induced by H₂O₂, the activation of JNK, ERK and p38 signaling pathways was examined (Fig. 6A). It was observed that H₂O₂ evidently activated JNK (Fig. 6B), ERK (Fig. 6C) and p38 (Fig. 6D) signaling pathways in SH-SY5Y cells, while SQ29548 pretreatment inhibited the activation of ERK (Fig. 6C) and p38 (Fig. 6D) signaling pathways. Therefore, SQ29548 may protect SH-SY5Y cells against oxidative stress via the inhibition of ERK and p38 signaling pathway activation.

Discussion

The results of the present study suggested that SQ29548, one of the TXA2R antagonists, was able to improve the antioxidative stress capacities of cells, reduce intracellular ROS levels and attenuate nerve cell apoptosis via the inhibition of ERK and p38 signaling pathway activation.

TXA2R, which is mainly distributed in platelets, has been suggested to be an important membrane receptor in the coagulation and inflammation processes in humans (6). In recent years, it has been reported that TXA2R also exists in nerve cells (5), and participates in the regulation of a series of physiological processes in peripheral and central nervous cells (5,24,25). Our previous study observed that the mutation of TXA2R affected the coagulation system function and further affected the onset of CI (6). Upon the application of SQ29548, an antagonist of TXA2R, ischemia-reperfusion damage of nerve cells and neurological function defects were attenuated in mice, although the underlying mechanism remains unclear (7). Furthermore, recent studies revealed that TXA2R is also involved in other physiological processes. For instance, activated TXA2R was found to inhibit growth factor expression levels in vascular endothelial cells, which further inhibited angiogenesis and vascular development (26). TXA2R was also closely associated with the cellular oxidative stress response. It has been observed that TXA2R participated in the regulation of cellular oxidative stress (8) and induced cell apoptosis (10), while TXA2R antagonists helped to reduce oxidative stress (9). These aforementioned studies indicated that, in certain neurological disorders, TXA2R may participate in the regulation of oxidative stress and the subsequent apoptotic process of nervous cells. Therefore, it is inferred that

TXA2R antagonists may protect nervous cells through the regulation of cellular oxidative stress.

SH-SY5Y cells were selected in the present study as the *in vitro* human nerve cell model, while SQ29548 was used as a TXA2R antagonist and H_2O_2 was used to simulate oxidative stress *in vitro*. SH-SY5Y, a neuron cell line, is widely used in studies investigating neurological disorders (27,28) and oxidative stress in nerve cells (17,28). SQ29548 has been proposed to be a specific, reliable and safe TXA2R antagonist (29), and is broadly utilized in studies on TXA2R (7,25,30,31). H_2O_2 , as one of main ROS products in cellular oxidative stress (32), is typically used to establish an *in vitro* oxidative stress cell model (23,33). In the current study, SH-SY5Y cells were initially cultured in DMEM; however, due to the well-known role of sodium pyruvate as a H_2O_2 scavenger and neuroprotective agent (34), the medium was altered to sodium pyruvate-negative DMEM following the cell plating, in order to minimize the confounder effects. The results revealed that different doses of SQ29548 had no significant effect on the viability of SH-SY5Y cells. However, different concentrations of H_2O_2 significantly influence the cell viability, with even a low concentration of H_2O_2 exhibiting a strong cytotoxic function. In addition, the present study observed that different doses of SQ29548 effectively protected against the H_2O_2 -induced inhibition of SH-SY5Y cell viability, and this protective effect was independent of the dose used.

The DCFH-DA experiment conducted in the current study verified that SQ29548 reduced intracellular ROS levels in SH-SY5Y cells treated with H_2O_2 , which may be associated with the regulation of intracellular antioxidant protein production following TXA2R inhibition. Intracellular antioxidant proteins, including SOD2 and catalase, serve to eliminate excessive ROS in cells (13). Generally, intracellular superoxide anion would be dismutated to H_2O_2 by SODs, particularly SOD2, in mitochondria; subsequently, catalase catalyzes H_2O_2 to H_2O and O_2 molecules. However, excessive oxidative stress response not only exceeds the elimination capacity of the antioxidant system, but also impairs the antioxidant system and inhibits the expression of antioxidant proteins (35,36). The present study demonstrated decreased protein and mRNA levels of SOD2 and catalase in SH-SY5Y cells treated with H_2O_2 . However, SQ29548 treatment rescued these decreased levels, protected the antioxidant system and improved the antioxidative stress capacities of SH-SY5Y cells. SQ29548 also significantly reduced the H_2O_2 -induced cell apoptosis and inhibited the expression levels of cleaved caspase9 and cleaved caspase3. Caspases are apoptosis-associated proteins (37,38), whose cleaved products promote cell apoptosis (38). In addition, the MAPK signaling pathways, which possess numerous roles in cell physiological processes, also participate in the cell apoptosis process induced by oxidative stress (39,40). For instance, it has been reported that, as a member of the MAPK family, the activation of the ERK signaling pathway induced by H_2O_2 caused SH-SY5Y cell apoptosis, and the inhibition of this pathway improved the cell survival (41). Furthermore, the inhibition of p38 pathway, another MAPK pathway, was reported to be beneficial against cell apoptosis induced by H_2O_2 (42). The two aforementioned phenomena may be associated with the ERK (43) and p38 (44) signaling pathways, which participated in the cell apoptosis process and regulated the

expression of apoptosis-associated proteins, such as caspases. Similarly, the present study indicated that SQ29548 regulated the ERK and p38 signaling pathway activation in SH-SY5Y cells treated with H_2O_2 , suggesting that SQ29548 possibly inhibited SH-SY5Y cell apoptosis and the expression of intracellular apoptosis-associated proteins through interfering with the ERK and p38 signaling pathways.

Oxidative stress is a vital pathological process in CI and other neurological disorders (45). Excessive cerebral oxidative stress severely impairs the neurons, thus, it is important to investigate how to protect nerve cells affected by oxidative stress response. In fact, TXA2R is a known therapeutic target in various diseases, and SQ29548 has been reported to protect against ischemia-reperfusion-induced intestinal impairment (46). There are also certain studies indicating that SQ29548 was able to treat the microvascular blood-brain barrier disruption caused by hyperglycemia and TXA2 (47). However, investigation of the association between TXA2R and oxidative stress in nerve cells has not been previously conducted. The present study further verified the potential of TXA2R as a novel therapeutic target in oxidative stress-associated neurological disorders, such as CI. Among oxidative stress studies, drugs such as edaravone (48) and esculetin (49) were demonstrated to be able to protect cells under oxidative stress. Nevertheless, the current study found that the TXA2R antagonist SQ29548 functioned similarly to these drugs.

However, certain issues remain to be solved. For instance, it is unclear whether extending SQ29548 treatment time amplifies the protective effect, whether the long-term inhibition of TXA2R affected the growth and development of neurons, and whether the nerve cell protection function of SQ29548 existed in other disease models. In future studies, we will investigate other *in vivo* and *in vitro* models, extend the SQ29548 treatment time and discuss the function of TXA2R antagonists in other neurological disease models. Such investigations are essential in clarifying the therapeutic mechanism of CI and other neurological disorders.

In conclusion, the present study demonstrated that SQ29548, an antagonist of TXA2R, was able to improve the antioxidative stress capacities of SH-SY5Y cells, and reduce cell apoptosis via the inhibition of ERK and p38 signaling pathways. This provided a new basis for the therapy of oxidative stress-associated neurological disorders.

Acknowledgements

Not applicable.

Funding

The present study was supported by key basic research grants from the Science and Technology Commission of Shanghai Municipality (grant no. 16JC1400500) and the Science and Technology Commission of Shanghai Municipality (grant no. 12ZR1418600).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

GC and AY designed and performed the experiments, analyzed the data and wrote the paper. GC, AY, NF and YF reviewed and edited the manuscript. All authors read and approved the manuscript.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Miggin SM and Kinsella BT: Expression and tissue distribution of the mRNAs encoding the human thromboxane A₂ receptor (TP) alpha and beta isoforms. *Biochim Biophys Acta* 1425: 543-559, 1998.
- Kassouf N, Ambily A, Watson S, Hassock S, Authi HS, Srivastava S, Watson SP and Authi KS: Phosphatidylinositol-3,4,5-trisphosphate stimulates Ca²⁺ elevation and Akt phosphorylation to constitute a major mechanism of thromboxane A₂ formation in human platelets. *Cell Signal* 27: 1488-1498, 2015.
- Bode M and Mackman N: Regulation of tissue factor gene expression in monocytes and endothelial cells: Thromboxane A₂ as a new player. *Vascul Pharmacol* 62: 57-62, 2014.
- Dagher OK, Jaffa MA, Habib A, Ziyadeh FN, Luttrell LM and Jaffa AA: Abstract 133: Mechanistic insights into bradykinin and thromboxane receptors heterodimerization in vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol* 36: A133, 2016.
- Sumimoto S, Muramatsu R and Yamashita T: Thromboxane A₂ stimulates neurite outgrowth in cerebral cortical neurons via mitogen activated protein kinase signaling. *Brain Res* 1594: 46-51, 2015.
- Shao J, Fu Y, Yang W, Yan J, Zhao J, Chen S and Xia W: Thromboxane A₂ receptor polymorphism in association with cerebral infarction and its regulation on platelet function. *Curr Neurovasc Res* 12: 15-24, 2015.
- Yan A, Zhang T, Yang X, Shao J, Fu N, Shen F, Fu Y and Xia W: Thromboxane A₂ receptor antagonist SQ29548 reduces ischemic stroke-induced microglia/macrophages activation and enrichment, and ameliorates brain injury. *Sci Rep* 6: 35885, 2016.
- Zhang M, Dong Y, Xu J, Xie Z, Wu Y, Song P, Guzman M, Wu J and Zou MH: Thromboxane receptor activates the AMP-activated protein kinase in vascular smooth muscle cells via hydrogen peroxide. *Circ Res* 102: 328-337, 2008.
- Xu S, Jiang B, Maitland KA, Bayat H, Gu J, Nadler JL, Corda S, Lavielle G, Verbeuren TJ, Zuccollo A and Cohen RA: The thromboxane receptor antagonist S18886 attenuates renal oxidant stress and proteinuria in diabetic apolipoprotein E-deficient mice. *Diabetes* 55: 110-119, 2006.
- Touchberry CD, Elmore CJ, Srinivas S and Wacker MJ: Thromboxane A₂ mediates apoptosis in cardiomyocytes via IP₃. *FASEB J* 27: 1122-1128, 2013.
- Rezzani R, Rodella LF and Bonomini F: Metabolic syndrome, aging and involvement of oxidative stress. *Aging Dis* 6: 109-120, 2015.
- Salim S: Oxidative stress and the central nervous system. *J Pharmacol Exp Ther* 360: 201-205, 2016.
- Birben E, Sahiner UM, Sackesen C, Erzurum S and Kalayci O: Oxidative stress and antioxidant defense. *World Allergy Organ J* 5: 9-19, 2012.
- Manzanero S, Santoro T and Arumugam TV: Neuronal oxidative stress in acute ischemic stroke: Sources and contribution to cell injury. *Neurochem Int* 62: 712-718, 2013.
- Schieber M and Chandel NS: ROS function in redox signaling and oxidative stress. *Curr Biol* 24: R453-R462, 2014.
- Pisoschi AM and Pop A: The role of antioxidants in the chemistry of oxidative stress: A review. *Eur J Med Chem* 97: 55-74, 2015.
- Jiang XW, Bai JP, Zhang Q, Hu XL, Tian X, Zhu J, Liu J, Meng WH and Zhao QC: Caffeoylquinic acid derivatives protect SH-SY5Y neuroblastoma cells from hydrogen peroxide-induced injury through modulating oxidative status. *Cell Mol Neurobiol* 37: 499-509, 2017.
- Shao J, Liu T, Xie QR, Zhang T, Yu H, Wang B, Ying W, Mruk DD, Silvestrini B, Cheng CY and Xia W: Adjudin attenuates lipopolysaccharide (LPS)- and ischemia-induced microglial activation. *J Neuroimmunol* 254: 83-90, 2013.
- Xu C, Hu Y, Hou L, Ju J, Li X, Du N, Guan X, Liu Z, Zhang T, Qin W, *et al*: β -Blocker carvedilol protects cardiomyocytes against oxidative stress-induced apoptosis by up-regulating miR-133 expression. *J Mol Cell Cardiol* 75: 111-121, 2014.
- Yan A, Cai G, Xia W and Fu Y: Thromboxane A₂ receptor antagonist SQ29548 suppresses the LPS-induced release of inflammatory cytokines in BV2 microglia cells via suppressing MAPK and NF- κ B signaling pathways. *Mol Med Rep* 16: 2491-2496, 2017.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2^{- $\Delta\Delta$ CT} method. *Methods* 25: 402-408, 2001.
- Wacker MJ, Tevis O, Hanke J, Howard T, Gilbert W and Orr JA: Characterization of thromboxane A₂ receptor and TRPV1 mRNA expression in cultured sensory neurons. *Neurosci Lett* 515: 12-17, 2012.
- Qian D, Li Z, Zhang Y, Huang Y, Wu Q, Ru G, Chen M and Wang B: Response of mouse zygotes treated with mild hydrogen peroxide as a model to reveal novel mechanisms of oxidative stress-induced injury in early embryos. *Oxid Med Cell Longev* 2016: 1521428, 2016.
- Cudd TA and Wood CE: Thromboxane A₂ receptor antagonism prevents hormonal and cardiovascular responses to mineral acid infusion. *Am J Physiol* 267: R1235-R1240, 1994.
- Rebel AA, Urquhart SA, Puig KL, Ghatak A, Brose SA, Golovko MY and Combs CK: Brain changes associated with thromboxane receptor antagonist SQ 29,548 treatment in a mouse model. *J Neurosci Res* 93: 1279-1292, 2015.
- Benndorf RA, Schwedhelm E, Gnann A, Taheri R, Kom G, Didé M, Steenpass A, Ergun S and Böger RH: Isoprostanes inhibit vascular endothelial growth factor-induced endothelial cell migration, tube formation, and cardiac vessel sprouting in vitro, as well as angiogenesis in vivo via activation of the thromboxane A₂ receptor: A potential link between oxidative stress and impaired angiogenesis. *Circ Res* 103: 1037-1046, 2008.
- Lopes FM, Schröder R, da Frota ML Jr, Zanotto-Filho A, Müller CB, Pires AS, Meurer RT, Colpo GD, Gelain DP, Kapczinski F, *et al*: Comparison between proliferative and neuron-like SH-SY5Y cells as an in vitro model for Parkinson disease studies. *Brain Res* 1337: 85-94, 2010.
- Cho GW, Koh SH, Kim MH, Yoo AR, Noh MY, Oh S and Kim SH: The neuroprotective effect of erythropoietin-transduced human mesenchymal stromal cells in an animal model of ischemic stroke. *Brain Res* 1353: 1-13, 2010.
- Chakraborty R, Bhullar RP, Dakshinamurti S, Hwa J and Chelikani P: Inverse agonism of SQ 29,548 and ramatroban on thromboxane A₂ receptor. *PLoS One* 9: e85937, 2014.
- Yang W, Yan A, Zhang T, Shao J, Liu T, Yang X, Xia W and Fu Y: Thromboxane A₂ receptor stimulation enhances microglial interleukin-1 β and NO biosynthesis mediated by the activation of ERK pathway. *Front Aging Neurosci* 8: 8, 2016.
- Yun DH, Song HY, Lee MJ, Kim MR, Kim MY, Lee JS and Kim JH: Thromboxane A₂ modulates migration, proliferation, and differentiation of adipose tissue-derived mesenchymal stem cells. *Exp Mol Med* 41: 17-24, 2009.
- Kumar KH and Khanum F: Hydroalcoholic extract of cyperus rotundus ameliorates H₂O₂-induced human neuronal cell damage via its anti-oxidative and anti-apoptotic machinery. *Cell Mol Neurobiol* 33: 5-17, 2013.
- Huang YY, Nagata K, Tedford CE, McCarthy T and Hamblin MR: Low-level laser therapy (LLLT) reduces oxidative stress in primary cortical neurons in vitro. *J Biophotonics* 6: 829-838, 2013.

34. Jagtap JC, Chandele A, Chopde BA and Shastry P: Sodium pyruvate protects against H₂O₂ mediated apoptosis in human neuroblastoma cell line-SK-N-MC. *J Chem Neuroanat* 26: 109-118, 2003.
35. Chen H, Yoshioka H, Kim GS, Jung JE, Okami N, Sakata H, Maier CM, Narasimhan P, Goeders CE and Chan PH: Oxidative stress in ischemic brain damage: Mechanisms of cell death and potential molecular targets for neuroprotection. *Antioxid Redox Signal* 14: 1505-1517, 2011.
36. Miguel F, Augusto AC and Gurgueira SA: Effect of acute vs chronic H₂O₂-induced oxidative stress on antioxidant enzyme activities. *Free Radic Res* 43: 340-347, 2009.
37. D'Amelio M, Cavallucci V and Cecconi F: Neuronal caspase-3 signaling: Not only cell death. *Cell Death Differ* 17: 1104-1114, 2010.
38. Brentnall M, Rodriguez-Menocal L, De Guevara RL, Cepero E and Boise LH: Caspase-9, caspase-3 and caspase-7 have distinct roles during intrinsic apoptosis. *BMC Cell Biol* 14: 32, 2013.
39. Hsieh CJ, Kuo PL, Hsu YC, Huang YF, Tsai EM and Hsu YL: Arctigenin, a dietary phytoestrogen, induces apoptosis of estrogen receptor-negative breast cancer cells through the ROS/p38 MAPK pathway and epigenetic regulation. *Free Radic Biol Med* 67: 159-170, 2014.
40. Sui X, Kong N, Ye L, Han W, Zhou J, Zhang Q, He C and Pan H: p38 and JNK MAPK pathways control the balance of apoptosis and autophagy in response to chemotherapeutic agents. *Cancer Lett* 344: 174-179, 2014.
41. Ruffels J, Griffin M and Dickenson JM: Activation of ERK1/2, JNK and PKB by hydrogen peroxide in human SH-SY5Y neuroblastoma cells: Role of ERK1/2 in H₂O₂-induced cell death. *Eur J Pharmacol* 483: 163-173, 2004.
42. Filomeni G, Piccirillo S, Rotilio G and Ciriolo MR: p38^{MAPK} and ERK1/2 dictate cell death/survival response to different pro-oxidant stimuli via p53 and Nrf2 in neuroblastoma cells SH-SY5Y. *Biochem Pharmacol* 83: 1349-1357, 2012.
43. Zhuang S, Yan Y, Daubert RA, Han J and Schnellmann RG: ERK promotes hydrogen peroxide-induced apoptosis through caspase-3 activation and inhibition of Akt in renal epithelial cells. *Am J Physiol Renal Physiol* 292: F440-F447, 2007.
44. Zhuang S, Demirs JT and Kochevar IE: p38 mitogen-activated protein kinase mediates bid cleavage, mitochondrial dysfunction, and caspase-3 activation during apoptosis induced by singlet oxygen but not by hydrogen peroxide. *J Biol Chem* 275: 25939-25948, 2000.
45. Liu J, Yang L, Tian H and Ma Q: Cathepsin D is involved in the oxygen and glucose deprivation/reperfusion-induced apoptosis of astrocytes. *Int J Mol Med* 38: 1257-1263, 2016.
46. Wen SH, Ling YH, Liu WF, Qiu YX, Li YS, Wu Y, Shen JT, Xia ZY and Liu KX: Role of 15-F₂t-isoprostane in intestinal injury induced by intestinal ischemia/reperfusion in rats. *Free Radic Res* 48: 907-918, 2014.
47. Zhao Z, Hu J, Gao X, Liang H, Yu H, Liu S and Liu Z: Hyperglycemia via activation of thromboxane A₂ receptor impairs the integrity and function of blood-brain barrier in microvascular endothelial cells. *Oncotarget* 8: 30030-30038, 2017.
48. Zhao ZY, Luan P, Huang SX, Xiao SH, Zhao J, Zhang B, Gu BB, Pi RB and Liu J: Edaravone protects HT22 neurons from H₂O₂-induced apoptosis by inhibiting the MAPK signaling pathway. *CNS Neurosci Ther* 19: 163-169, 2013.
49. Han MH, Park C, Lee DS, Hong SH, Choi IW, Kim GY, Choi SH, Shim JH, Chae JI, Yoo YH and Choi YH: Cytoprotective effects of esculetin against oxidative stress are associated with the upregulation of Nrf2-mediated NQO1 expression via the activation of the ERK pathway. *Int J Mol Med* 39: 380-386, 2017.