

Mitochondria-targeted antioxidant therapy for an animal model of PCOS-IR

YU DING^{1*}, ZHAOCHANG JIANG^{2*}, BOHOU XIA³, LIZONG ZHANG⁴, CAIJUAN ZHANG⁵ and JIANHANG LENG¹

¹Central Laboratory, Hangzhou First People's Hospital; ²Department of Pathology, Second Affiliated Hospital, Zhejiang University, School of Medicine, Hangzhou, Zhejiang 310006; ³Department of Pharmacy, Hunan Chinese Medical University, Changsha, Hunan 410208; ⁴Department of Experimental Animal Center, Zhejiang Chinese Medical University, Hangzhou, Zhejiang 310053; ⁵Department of Gynecological and Obstetrics, Hangzhou First People's Hospital, Zhejiang University, School of Medicine, Hangzhou, Zhejiang 310006, P.R. China

Received June 16, 2018; Accepted October 19, 2018

DOI: 10.3892/ijmm.2018.3977

Abstract. Polycystic ovary syndrome (PCOS) is a common endocrine disorder with unknown etiology and unsatisfactory clinical treatment. Considering the ethical limitations of studies involving humans, animal models that reflect features of PCOS and insulin resistance (IR) are crucial resources in investigating this syndrome. Our previous study showed that mitochondrial dysfunction resulted from pathogenic mutations of mitochondrial DNA (mtDNA), and that oxidative stress had an active role in the phenotypic manifestation of PCOS-IR. Therefore, it was hypothesized that limiting oxidative stress and mitochondrial damage may be useful and effective for the clinical treatment of PCOS-IR. For this purpose, the present study examined the therapeutic effects of the mitochondria-targeted antioxidant MitoQ₁₀ for PCOS-IR. Furthermore, the histopathology was used to analysis the ovarian morphological changes. The endocrine and reproductive related parameters were analyzed by ELISA approach. A PCOS-IR model was successfully established by subcutaneous injection of rats with testosterone propionate and feeding a high-fat diet. The 30 female Sprague-Dawley rats were then divided into three groups, comprising a control (n=10), animal model (PCOS-IR, n=10) and MitoQ₁₀ treatment (n=10) group. It was found that MitoQ₁₀ significantly improved the IR condition and reversed the endocrine and reproductive conditions of PCOS. In addition, the impaired mitochondrial functions were improved following MitoQ₁₀ administration. Notably, western blot results suggested that this antioxidant

reduced the expression levels of apoptosis-related proteins cytochrome *c* and B-cell lymphoma-2 (Bcl-2)-associated X protein, whereas the anti-apoptotic protein Bcl-extra large was increased following MitoQ₁₀ treatment. Taken together, the data indicated that the MitoQ₁₀ may have a beneficial favorable therapeutic effect on animals with PCOS-IR, most likely via the protection of mitochondrial functions and regulation of programmed cell death-related proteins.

Introduction

PCOS is a common endocrine and metabolic disease affecting ~10% of women of reproductive age (1). Clinically, PCOS may manifest a variety of phenotypes, including chronic anovulation, hyperandrogenism, insulin resistance (IR) and infertility. Women with PCOS are also more prone to developing diabetes, coronary heart disease and metabolic syndrome (2). Therefore, the management of PCOS-IR remains a significant clinical challenge in gynecology.

Oxidative stress (OS), resulting from an imbalance between radicals and antioxidant defense, has been found to be a main pathophysiological mechanism in various human diseases (3). This occurs due to the overproduction of specific molecules (4), including reactive oxygen species (ROS), which are generated from nitric oxide (NO) and malondialdehyde (MDA) and can damage all components of the cell, including proteins, lipids and DNA (5). Increasing evidence indicates that OS may be associated with IR and obesity, and may also contribute to PCOS and its metabolic associations (6,7).

Mitochondria produce the majority of the body's cellular energy via oxidative phosphorylation, releasing ROS as a toxic byproduct (8). The increased ROS production may have negative consequences; for example, enhancing the OS in tissues and organisms, disrupting the natural genetic code and inducing the mitochondrial-mediated apoptosis (9). Our previous studies indicated that mtDNA pathogenic mutations led to mitochondrial dysfunctions and, together with OS, contributed to the progression of PCOS-IR (10,11). Considering the importance of mitochondria in energy production and ROS generation, it was hypothesized that mitochondria-targeted

Correspondence to: Dr Yu Ding, Central Laboratory, Hangzhou First People's Hospital, Zhejiang University, School of Medicine, 261 Huansha Road, Hangzhou, Zhejiang 310006, P.R. China
E-mail: dingyu.zj@gmail.com

*Contributed equally

Key words: mitochondrial, MitoQ₁₀, antioxidant therapy, animal model, polycystic ovary syndrome and insulin resistance

antioxidants may offer potential as a novel, potential useful drug to prevent the pathogenesis of PCOS-IR.

MitoQ₁₀ is one of these mitochondria-targeted antioxidants, which directly binds to human mitochondria. At the molecular level, it consists of a triphenylphosphonium cation (TPP⁺)² attached to a ubiquinone moiety (Fig. 1) (12). Previous studies have suggested that MitoQ₁₀ can accelerate the absorbance of CoQ₁₀ into mitochondria in a mouse model of Alzheimer's disease, and this antioxidant is considered to combat ROS generation in mitochondria (13,14). *In vitro* and *in vivo* experiments have indicated that MitoQ₁₀ has antioxidant effects (13-15). In addition, MitoQ₁₀ had been found to be a novel therapeutic intervention in human autoimmune disease (15) and cardiovascular disease (16,17). However, whether MitoQ₁₀ has a therapeutic effect on PCOS-IR remains to be elucidated.

In the present study, a rat model manifesting the features of PCOS-IR was first generated; this was followed by investigation of the effects of MitoQ₁₀ on the rats and further discussion of the possible molecular mechanism.

Materials and methods

Animals. A total of 30 female Sprague-Dawley rats (SPF grade, 3 weeks old, 230-250 g) [SCXK (Hu) 2013-0016] were provided by the Experimental Animal Center, Zhejiang Chinese Medical University (Hangzhou, China). The rats were housed with four to six animals per cage under standard laboratory conditions (25°C clean environment with 50% humidity, 12-h light/dark cycle), all animals had free access to regular food and tap water. All experiments were performed in strict accordance with the Animal Care and Use Committee of Zhejiang Chinese Medical University.

Generation of the PCOS-IR rat model. The 30 rats were divided into three groups for the following stage of treatment, which were as follows: i) Control group (n=10): Mice received a normal rodent diet and injection with olive oil at a volume equal to injections in the experimental groups; ii) PCOS-IR model group (n=10): 100 µg of testosterone propionate (Sigma-Aldrich; Merck KGaA, Darmstadt, Germany) was subcutaneously injected into animals, which was dissolved in 0.05 ml commercial corn oil. The rats were also fed a high-fat diet, consisting of 54.2% standard diet, 16.8% lard, 15% sucrose, 9% casein, 1% minerals, 1% vitamins and 3% malt dextrin. Following 8 consecutive weeks of treatment, the mice were sacrificed and the vaginal cytology was analyzed using the methylene blue staining method by LEICA DM1000 light microscope (Leica Microsystems GmbH, Wetzlar, Germany); iii) MitoQ₁₀ treatment group (n=10): Following establishment of the PCOS-IR animal model, 10 PCOS-IR rats were given clean drinking water which contained 500 µmol/l MitoQ₁₀ (Sigma-Aldrich; Merck KGaA) for 8 consecutive weeks. The solutions for MitoQ₁₀ were reformulated fresh every 3 days, and were stored at 4°C in a dark room.

Detection of endocrine-related parameters. The orbital venous blood of each animal in the control, PCOS-IR and MitoQ₁₀ treatment groups were collected following overnight fasting. The serum levels of fasting plasma glucose (FPG) and fasting




Figure 1. Structure and accumulation of MitoQ₁₀. (A) Chemical structure of MitoQ₁₀. (B) MitoQ₁₀ accumulates (5-10 fold) within the cell driven by the plasma membrane potential, and accumulates (100-500 fold) within the mitochondria, driven by the mitochondrial membrane potential.

insulin (FINS) were then analyzed. In addition, the levels of hormones in the rats, including testosterone (T), lactate dehydrogenase (LH), follicle-stimulating hormone (FSH), FPG and FINS, were determined using ELISA kits (Elabscience, Wuhan, China), according to the protocols provided by the manufacturer. A total of three samples were selected from each group for analysis. The HOMA-IR was determined as follows: $\text{HOMA-IR} = \text{FPG (mmol/l)} \times \text{FINS (mU/l)} / 22.5$; IR was considered present if the value of HOMA-IR was >2.8 (18).

Histopathology. All rats were bilaterally ovariectomized following treatment. The animals were anesthetized with isoflurane, and the ovaries were removed through dorsolateral incisions. The ovaries were subsequently underwent 10% formalin-fixed and paraffin-embedded, incubated overnight at 4°C, stained with hematoxylin for 5 min and eosin for 3 min under room temperature (25°C); two pathologists evaluated the morphological changes between different groups using the LEICA DM1000 light microscope (Leica Microsystems GmbH).

Measurement of the levels of OS-associated biomarkers. To examine the expression levels of OS-related biomarkers, the contents of MDA, total antioxidant capacity (T-AOC), and the activity of superoxide dismutase (SOD) and glutathione (GSH) in tissues were analyzed using the kits provided by Nanjing Jiancheng Bioengineering Institute (Nanjing, China) according to the manufacturer's protocol.




Figure 2. Features of the PCOS-IR rat model. MitoQ₁₀ improved the morphology of ovarian follicles and increases the granulosa cells of the PCOS-IR rat model. Magnification, x200. PCOS-IR, polycystic ovary syndrome and insulin resistance.




Figure 3. Serum levels of T, LH, FSH and LH/FSH in rats in the Control, PCOS-IR and MitoQ₁₀ treatment groups. ^{##}P<0.05, compared with the control; ^{**}P<0.05, compared with the PCOS-IR group. PCOS-IR, polycystic ovary syndrome and insulin resistance; T, testosterone; LH, lactate dehydrogenase; FSH, follicle-stimulating hormone.

ATP analysis. The ATP concentrations in the ovarian tissues were analyzed using the ATP assay kit, according to the manufacturer's protocols. The absorbance was then measured at 636 nm (Nanjing Jiancheng Bioengineering Institute).

Isolation of mitochondria. The isolation of mitochondria was performed at 4°C by different centrifugation (700 x g for 10 min; 3,000 x g for 15 min) steps using a mitochondria isolation kit (Baosai Biosciences, Inc., Beijing, China). The final pellet of mitochondria was resuspended in buffer, stored on ice, stored for 4°C and used for experiments within 4 h (19). The BCA assay was used to determine the final protein concentration.

Analysis of mitochondrial membrane potential (MMP) and ROS. MMP was determined using the JC-1 kit (cat. no. T4069, Sigma-Aldrich; Merck KGaA), according to our previous study (11). In brief, 50 µg of the purified rat mitochondria protein was mixed with 5 µg/ml JC-1 staining for 10 min at 37°C; subsequently, a microscope reader (Tecan Group, Inc., Ltd., Mannendorf, Switzerland) was used to record the green and red fluorescence of JC-1 at 530 and 590 nm, respectively (20). Qualification of the ROS levels were assessed using 20 µM of ROS-sensitive dye 2',7'-dichlorodihydrofluorescein diacetate (H₂DCFDA), which cleaves to become DCF (cat. no. D6883; Sigma-Aldrich; Merck KGaA) (21).

Western blot analysis. For western blot analysis, the total proteins from the fresh rat ovarian tissues were extracted based on the protocol originally proposed by Dignam *et al* (22) with modifications. The 50 µg protein samples were boiled for 5 min in Laemmli buffer and then separated by 12% SDS polyacrylamide gel at 180 V. Subsequently, the separated proteins were transferred onto a polyvinylidene difluoride membrane at 400 mA for 2 h (EMD Millipore, Billerica, MA, USA). Following incubation of the PVDF membrane with blocking solution, the protein bands were mixed with special anti-Bax (cat. no. 34260; 1:500; Signalway Antibody LLC, College Park, MD, USA), anti-Bcl-xL (cat. no. 21061; 1:500; Signalway Antibody LLC), anti-Cyto C (cat. no. ab33484; 1:500; Abcam) and anti-Bcl-2 (cat. no. ab59348; 1:500; Abcam) antibodies overnight at 4°C. The blots were then washed and incubated with horseradish peroxidase (HRP)-conjugated secondary antibodies for 1 h at room temperature, the second antibodies were as follows: Anti-rabbit IgG, HRP-linked antibody (cat. no. 7074; 1:500; Signalway Antibody LLC) and anti-mouse IgG, HRP-linked antibody (cat. no. 7076; 1:500; Signalway Antibody LLC). The immune complexes were visualized using an enhanced chemiluminescence kit and analyzed with the ImageJ[®] LAS 4000 mini program (National Institutes of Health, Bethesda, MD, USA).

Statistical analysis. All values are expressed as the mean ± standard deviation. The statistical analyses were performed using SPSS 19.0 software (IBM SPSS, Armonk,




Figure 4. Serum concentrations of FPG, FIN and the HOMA-IR in the Control, PCOS-IR and MitoQ₁₀ treatment groups. ^{##}P<0.05, compared with the control; ^{**}P<0.05, compared with the PCOS-IR group. PCOS-IR, polycystic ovary syndrome and insulin resistance; FPG, fasting plasma glucose; FIN, fasting insulin; HOMA-IR.

NY, USA). The results were analyzed with one-way analysis of variance followed by Tukey's post hoc test for multiple comparisons. P<0.05 was considered to indicate a statistically significant difference.

Results

Reproductive and metabolic features of the PCOS-IR rat model. Following 8 consecutive weeks under treatment with testosterone propionate, together with a high fat diet, the PCOS-IR mouse model was successfully established. As presented in Fig. 2, the PCOS-IR animals exhibited vaginal keratosis and abnormal estrous cycle. In addition, on measuring endocrine parameters, it was noted that all animals exhibited IR as the HOMA-IR was >2.8.

Recovery of endocrine status following MitoQ₁₀ treatment. As presented in Figs. 3 and 4, the rats exposed to testosterone propionate had significantly increased T, LH, LH/FSH, FIN and HOMA-IR levels, compared with those in the control group (all P<0.01). Following MitoQ₁₀ treatment, the levels of these parameters were decreased significantly, compared with those in the controls (all P<0.01). However, no statistically significant differences were observed in the levels of FPG or FSH prior to and following MitoQ₁₀ treatment.

MitoQ₁₀ decreases OS. To determine whether MitoQ₁₀ reversed the OS of PCOS-IR rats, the expression levels of MDA, T-AOC, SOD and GSH were analyzed in the rat tissues. As presented

in Fig. 5, the levels of T-AOC, SOD and GSH, which reflected the antioxidant capacity in rats, were enhanced in the MitoQ₁₀ treatment group, compared with those in the PCOS-IR rat model (all P<0.01). Consistent with this finding, the level of MDA, an OS-related biomarker, was significantly decreased following MitoQ₁₀ administration (P<0.01).

MitoQ₁₀ protects against mitochondrial damage. To determine whether the administration of MitoQ₁₀ improved the mitochondrial function, the levels of ATP, MMP and ROS were analyzed in the Control, PCOS-IR and MitoQ₁₀ treatment groups. As shown in Fig. 6, it was found that the antioxidant significantly protected against mitochondrial damage (all P<0.05).

Detection of the expression of Cyto C, Bcl-xL, Bcl-2 and Bax. The present study further evaluated the expression levels of the apoptotic and anti-apoptotic proteins, Cyto C, Bcl-xL, Bcl-2 and Bax, in the different groups to examine whether the MitoQ₁₀ had a therapeutic effect. As presented in Figs. 7 and 8, it was found that, compared with the control group, the PCOS-IR rats exhibited higher levels of Cyto C and Bax and a higher ratio of Bax to Bcl-2, whereas the level of Bcl-xL was decreased. However, following MitoQ₁₀ treatment, the levels of Cyto C, Bax and the ratio of Bax to Bcl-2 were decreased and the level of Bcl-xL was increased, compared with levels in the PCOS-IR group. However, no statistically significant difference was observed in the level of Bcl-2 between these groups.




Figure 5. Analysis of oxidative stress-associated biomarkers in the Control, PCOS-IR and MitoQ₁₀ treatment groups. ##P<0.05, compared with the control; **P<0.05, compared with the PCOS-IR group. PCOS-IR, polycystic ovary syndrome and insulin resistance; MDA, malondialdehyde; T-AOC, total antioxidant capacity; SOD, superoxide dismutase; GSH, glutathione.

Discussion

In the present study, the therapeutic effects of mitochondria-targeted antioxidant MitoQ₁₀ on an animal model of PCOS-IR were investigated. Accumulating evidence has demonstrated that the elevated production of ROS has an active role in the progression and pathogenesis of PCOS (23,24). Natural antioxidants, including vitamin C and vitamin E, cannot scavenge sufficient ROS as they cannot directly bind to mitochondria, highlighting the importance of developing novel targeted therapies. MitoQ₁₀ can solve this problem; this antioxidant is composed of ubiquinone moieties attached to a TPP moiety by a 10-carbon alkyl chain. Lipophilic cations can readily move through phospholipid bilayers, therefore, MitoQ₁₀ can be taken up by mitochondria and be effective in combating ROS (25,26). Once within the mitochondria, almost all of the accumulated MitoQ₁₀ resides in the matrix surface of the inner-membrane and prevents OS and mitochondrial dysfunction (27).

In the present study, to examine the therapeutic effects of MitoQ₁₀, an animal model of PCOS-IR was generated via the subcutaneous injection of testosterone propionate, together with a high-fat diet. In general, this animal model exhibits the clinical and biochemical characterizations of women with PCOS, including abnormal estrous cycles, markedly increased total body weight, polycystic ovaries and variable levels of T, LH, FSH, FIN and FPG in the serum of rats. Of the 20 rats injected, 100% exhibited the IR features. Following

treatment with MitoQ₁₀, the morphological characteristic of polycystic ovaries and the plasma concentrations of FIN in the PCOS group were improved. These data suggested that MitoQ₁₀ exerted a therapeutic effect against the development of PCOS-IR.

MDA, T-AOC, SOD and GSH are regarded as common markers to evaluate OS levels. Of these, MDA is the product of lipid oxidation, whereas T-AOC indicates the ability of all antioxidants in different foods to clean harmful free radicals in blood and cells (28). SOD is an enzyme which can catalyze superoxide radicals into ordinary molecular oxygen or hydrogen peroxide, and GSH is another important antioxidant that can protect cells against ROS. As presented in Fig. 5, compared with those in the control group, the rats in the PCOS-IR group had higher MDA levels and lower T-AOC, SOD and GSH levels, suggesting increased OS in PCOS-IR. However, the animals treated with MitoQ₁₀ exhibited decreased OS (all P<0.05).

Increasing evidence suggests that the overproduction of ROS may activate stress signals to the mitochondria and be important in a number of biological processes; this suggests that mitochondrial dysfunction, together with the impaired antioxidant system, may be involved in the pathophysiological mechanism and progression of PCOS-IR (29,30). There are two general pathways of apoptosis based on different apoptotic stimuli, termed the death receptor pathway and the mitochondrial-mediated pathway. In particular, increased ROS production destroys mitochondrial function, which may




Figure 6. Analysis of mitochondrial function in the Control, PCOS-IR and MitoQ₁₀ treatment groups. [#] $P < 0.05$, compared with the controls; ^{**} $P < 0.05$, compared with the PCOS-IR group. PCOS-IR, polycystic ovary syndrome and insulin resistance; MMP, mitochondrial membrane potential; ROS, reactive oxygen species.

subsequently lead to the loss of MMP and release of Cyto C (31,32). Therefore, the present study measured the ROS levels, the level of MMP and the expression of apoptosis-related proteins to confirm whether MitoQ₁₀ induced apoptosis via the mitochondrial pathway. The data revealed that the PCOS-IR rats treated with MitoQ₁₀ exhibited recovery of mitochondrial functions, including increased ATP and MMP levels; by contrast, the ROS level was significantly decreased. This supported the hypothesis that MitoQ₁₀ can reduce the cellular OS resulting from the mitochondrial dysfunction responsible for PCOS-IR.

Bcl-2 and Bcl-xL are anti-apoptotic members of the Bcl-2 family, which function as a 'life/death switch' that integrates diverse inter- and intracellular cues to determine whether or not the stress apoptotic pathway is activated (33,34). In response to apoptotic stimuli, the balance and interactions of anti-apoptotic and pro-apoptotic Bcl proteins influence the activation of downstream pro-apoptotic proteins Bcl-2 homologous antagonist/killer (Bak) and Bax (35). Of note, Bak and Bax are two nuclear-encoded proteins present in higher eukaryotes that are able to pierce the mitochondrial outer membrane to mediate cell death by apoptosis (36). Once the mitochondrial-mediated cell death pathway is activated, the conformations of Bax and Bak are altered, MMP is decreased and pro-apoptotic proteins are released from the intermembrane space into the cytosol (37,38). Following release into the cytoplasm, Cyto C stimulates apoptosome formation followed by the activation of caspase-9. Therefore, the release of Cyto C is an early event in the pathway of mitochondrial-mediated




Figure 7. Relative expression of Cyt C, Bcl-xL, Bcl-2 and Bax in the ovaries of the Control, PCOS-IR and MitoQ₁₀ treatment groups. PCOS-IR, polycystic ovary syndrome and insulin resistance; Cyt C, cytochrome c; Bcl-2, B-cell lymphoma 2; Bcl-xL, Bcl-extra large; Bax, Bcl-2-associated X protein.

apoptosis. As presented in Figs. 7 and 8, the present study found that MitoQ₁₀ significantly increased the expression of Bcl-xL,




Figure 8. Qualification of the expression levels of Cyt C, Bcl-xL, Bcl-2 and Bax in the ovaries of the Control, PCOS-IR and MitoQ₁₀ treatment groups. ##P<0.05, compared with the control; **P<0.05, compared with the PCOS-IR group. PCOS-IR, polycystic ovary syndrome and insulin resistance; Cyt C, cytochrome c; Bcl-2, B-cell lymphoma 2; Bcl-xL, Bcl-extra large; Bax, Bcl-2-associated X protein.

whereas the levels of total Cyt C and Bax decreased following MitoQ₁₀ treatment. However, no difference was observed in the expression level of Bcl-2 prior to and following MitoQ₁₀ administration, suggesting that MitoQ₁₀ may not affect the downregulation of Bcl-2. A range of stimuli induce apoptosis by releasing Cyt C from the mitochondria into the cytoplasm, where it activates caspases; the mechanisms by which these stimuli cause Cyt C release from mitochondria remain to be fully elucidated, however, some or all may involve increased mitochondrial OS (39). MitoQ₁₀ can assist in elucidating the role of oxidative damage in apoptosis, subsequently preventing apoptotic cell death.

Although the detection of Cyt C was not based on the total protein, it was noted that detecting total Cyt C protein has been reported in several studies; for example, Eleftheriadis *et al* (40) demonstrated that serum Cyt C levels may be a biomarker for mitochondrial and cellular damage. In addition, Li *et al* (41) measured Cyt C levels via an optical microfiber, and a study by Radhakrishnan *et al* (42) used an ELISA approach to determine plasma Cyt C levels. Overall, although the qualification of total Cyt C levels cannot reflect programmed cell death, it is only the way for reflecting the early event of apoptosis.

Based on the above observations, the present study hypothesized that the possible molecular mechanism underlying MitoQ₁₀ in the treatment of PCOS-IR rats may be as follows: Administration of MitoQ₁₀ may bind to mitochondria and scavenge ROS, indicating its antioxidant effects. Subsequently, decreased ROS levels protect the pancreatic cells from death, most likely from mitochondrial-mediated apoptosis in the form of increasing the expression levels of Bcl-xL, decreasing the expression levels of Cyt C and Bax, and recovering mitochondrial functions, with improved ATP and MMP levels. As a result, the decreased OS may prevent the pancreatic cells from death or apoptosis. Finally, the endocrine and reproductive

conditions of PCOS-IR recovered following MitoQ₁₀ treatment. Therefore, this antioxidant may be a potential agent for patients with PCOS-IR in the future.

Acknowledgements

The authors would like to thank the members of their lab for discussion. The authors are grateful to Dr Qi Liu from Shaoxing People's Hospital, Zhejiang University for critical reading of this manuscript.

Funding

This study was supported by the grants from the Hangzhou Bureau of Science and Technology (grant no. 20150633B16), the Ministry of Public Health from Zhejiang Province (grant nos. 2013KYA158 and 2018ZH019) and the Natural Science Foundation of Zhejiang Province (grant nos. LY14H270008 and LY15H280007).

Availability of data and materials

The datasets used and/or analyzed during the present study are available from the corresponding author on reasonable request.

Authors' contributions

YD designed the experiments, ZJ performed the histopathological analysis, BX performed the statistical analysis, LZ performed the animal experiments, CZ and JL analyzed the data. YD wrote the manuscript. All authors discussed the results and implications and commented on the manuscript at all stages. All authors read and approved the final manuscript.

Ethics approval and consent to participate

All experiments were approved by the Animal Care and Use Committee of Zhejiang Chinese Medical University.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- McCartney CR and Marshall JC: Clinical Practice. Clinical practice. Polycystic ovary syndrome. *N Engl J Med* 375: 54-64, 2016.
- Ali AT: Polycystic ovary syndrome and metabolic syndrome. *Ceska Gynekol* 80: 279-289, 2015.
- Kalyanaraman B: Teaching the basics of redox biology to medical and graduate students: Oxidants, antioxidants and disease mechanisms. *Redox Biol* 1: 244-257, 2013.
- Liu X, Zhu Q, Zhang M, Yin T, Xu R, Xiao W, Wu J, Deng B, Gao X, Gong W, *et al*: Isoliquiritigenin ameliorates acute pancreatitis in mice via inhibition of oxidative stress and modulation of the Nrf2/HO-1 pathway. *Oxid Med Cell Longev* 2018: 7161592, 2018.
- Dalle-Donne I, Rossi R, Colombo R, Giustarini D and Milzani A: Biomarkers of oxidative damage in human disease. *Clin Chem* 52: 601-623, 2006.
- Escobar-Morreale HF, Luque-Ramírez M and San Millán JL: The molecular-genetic basis of functional hyperandrogenism and the polycystic ovary syndrome. *Endocr Rev* 26: 251-282, 2005.
- Zuo T, Zhu M and Xu W: Roles of oxidative stress in polycystic ovary syndrome and cancers. *Oxid Med Cell Longev* 2016: 8589318, 2016.
- Picard M, Wallace DC and Burelle Y: The rise of mitochondria in medicine. *Mitochondrion* 30: 105-116, 2016.
- Böttger EC and Schacht J: The mitochondrion: A perpetrator of acquired hearing loss. *Hear Res* 303: 12-19, 2013.
- Ding Y, Xia BH, Zhang CJ and Zhuo GC: Mitochondrial tRNA^{Leu}(UUR) C3275T, tRNA^{Gln} T4363C and tRNA^{Lys} A8343G mutations may be associated with PCOS and metabolic syndrome. *Gene* 642: 299-306, 2018.
- Ding Y, Xia BH, Zhang CJ and Zhuo GC: Mutations in mitochondrial tRNA genes may be related to insulin resistance in women with polycystic ovary syndrome. *Am J Transl Res* 9: 2984-2996, 2017.
- Kelso GF, Porteous CM, Coulter CV, Hughes G, Porteous WK, Ledgerwood EC, Smith RA and Murphy MP: Selective targeting of a redox-active ubiquinone to mitochondria within cells: antioxidant and antiapoptotic properties. *J Biol Chem* 276: 4588-4596, 2001.
- Smith RA and Murphy MP: Animal and human studies with the mitochondria-targeted antioxidant MitoQ. *Ann NY Acad Sci* 1201: 96-103, 2010.
- Manczak M, Mao P, Calkins MJ, Cornea A, Reddy AP, Murphy MP, Szeto HH, Park B and Reddy PH: Mitochondria-targeted antioxidants protect against amyloid-beta toxicity in Alzheimer's disease neurons. *J Alzheimers Dis* 20 (Suppl 2): S609-S631, 2010.
- Mao P, Manczak M, Shirendeb UP and Reddy PH: MitoQ, a mitochondria-targeted antioxidant, delays disease progression and alleviates pathogenesis in an experimental autoimmune encephalomyelitis mouse model of multiple sclerosis. *Biochim Biophys Acta* 1832: 2322-2331, 2013.
- Graham D, Huynh NN, Hamilton CA, Beattie E, Smith RA, Cochemé HM, Murphy MP and Dominiczak AF: Mitochondria-targeted antioxidant MitoQ₁₀ improves endothelial function and attenuates cardiac hypertrophy. *Hypertension* 54: 322-328, 2009.
- McLachlan J, Beattie E, Murphy MP, Koh-Tan CH, Olson E, Beattie W, Dominiczak AF, Nicklin SA and Graham D: Combined therapeutic benefit of mitochondria-targeted antioxidant, MitoQ₁₀, and angiotensin receptor blocker, losartan, on cardiovascular function. *J Hypertens* 32: 555-564, 2014.
- Keskin M, Kurtoglu S, Kendirci M, Atabek ME and Yazici C: Homeostasis model assessment is more reliable than the fasting glucose/insulin ratio and quantitative insulin sensitivity check index for assessing insulin resistance among obese children and adolescents. *Pediatrics* 115: e500-e503, 2005.
- Yarian CS, Toroser D and Sohal RS: Aconitase is the main functional target of aging in the citric acid cycle of kidney mitochondria from mice. *Mech Ageing Dev* 127: 79-84, 2006.
- Yuan Y, Chen Y, Zhang P, Huang S, Zhu C, Ding G, Liu B, Yang T and Zhang A: Mitochondrial dysfunction accounts for aldosterone-induced epithelial-to-mesenchymal transition of renal proximal tubular epithelial cells. *Free Radic Biol Med* 53: 30-43, 2012.
- Raza H, John A and Howarth FC: Increased oxidative stress and mitochondrial dysfunction in Zucker diabetic rat liver and brain. *Cell Physiol Biochem* 35:1241-1251, 2015.
- Dignam JD, Lebovitz RM and Roeder RG: Accurate transcription initiation by RNA polymerase II in a soluble extract from isolated mammalian nuclei. *Nucleic Acids Res* 11: 1475-1489, 1983.
- Victor VM, Rovira-Llopis S, Bañuls C, Diaz-Morales N, Martinez de Marañon A, Rios-Navarro C, Alvarez A, Gomez M, Rocha M and Hernández-Mijares A: Insulin resistance in PCOS patients enhances oxidative stress and leukocyte adhesion: Role of myeloperoxidase. *PLoS One* 11: e0151960, 2016.
- Serrano Mujica L, Bridi A, Della Mèa R, Rissi VB, Guarda N, Moresco RN, Premeaor MO, Antoniazzi AQ, Gonçalves PBD and Comim FV: Oxidative stress and metabolic markers in pre- and postnatal polycystic ovary syndrome rat protocols. *J Inflamm Res* 11: 193-202, 2018.
- Ross MF, Kelso GF, Blaikie FH, James AM, Cochemé HM, Filipovska A, Da Ros T, Hurd TR, Smith RA and Murphy MP: Lipophilic triphenylphosphonium cations as tools in mitochondrial bioenergetics and free radical biology. *Biochemistry (Mosc)* 70: 222-230, 2005.
- Firsov AM, Kotova EA, Orlov VN, Antonenko YN and Skulachev VP: A mitochondria-targeted antioxidant can inhibit peroxidase activity of cytochrome c by detachment of the protein from liposomes. *FEBS Lett* 590: 2836-2843, 2016.
- James AM, Cochemé HM and Murphy MP: Mitochondria-targeted redox probes as tools in the study of oxidative damage and ageing. *Mech Ageing Dev* 126: 982-986, 2005.
- Yilmaz N, Inal HA, Gorkem U, Sargin Oruc A, Yilmaz S and Turkkan A: Follicular fluid total antioxidant capacity levels in PCOS. *J Obstet Gynaecol* 36: 654-657, 2016.
- Abu Bakar MH, Sarmidi MR, Tan JS and Mohamad Rosdi MN: Celastrol attenuates mitochondrial dysfunction and inflammation in palmitate-mediated insulin resistance in C3A hepatocytes. *Eur J Pharmacol* 799: 73-83, 2017.
- Hafizi Abu Bakar M, Kian Kai C, Wan Hassan WN, Sarmidi MR, Yaakob H and Zaman Huri H: Mitochondrial dysfunction as a central event for mechanisms underlying insulin resistance: The roles of long chain fatty acids. *Diabetes Metab Res Rev* 31: 453-475, 2015.
- Yan Y, Su X, Liang Y, Zhang J, Shi C, Lu Y, Gu L and Fu L: Emodin azide methyl anthraquinone derivative triggers mitochondrial-dependent cell apoptosis involving in caspase-8-mediated Bid cleavage. *Mol Cancer Ther* 7: 1688-1697, 2008.
- Wang XH, Jia DZ, Liang YJ, Yan SL, Ding Y, Chen LM, Shi Z, Zeng MS, Liu GF and Fu LW: Lgf-YL-9 induces apoptosis in human epidermoid carcinoma KB cells and multidrug resistant KB200 cells via reactive oxygen species-independent mitochondrial pathway. *Cancer Lett* 249: 256-270, 2007.
- Kluck RM, Bossy-Wetzel E, Green DR and Newmeyer DD: The release of cytochrome c from mitochondria: a primary site for Bcl-2 regulation of apoptosis. *Science* 275: 1132-1136, 1997.
- Yang J, Liu X, Bhalla K, Kim CN, Ibrado AM, Cai J, Peng TI, Jones DP and Wang X: Prevention of apoptosis by Bcl-2: Release of cytochrome c from mitochondria blocked. *Science* 275: 1129-1132, 1997.
- Nechushtan A, Smith CL, Lamensdorf I, Yoon SH and Youle RJ: Bax and Bak coalesce into novel mitochondria-associated clusters during apoptosis. *J Cell Biol* 153: 1265-1276, 2001.
- Susin SA, Lorenzo HK, Zamzami N, Marzo I, Snow BE, Brothers GM, Mangion J, Jacotot E, Costantini P, Loeffler M, *et al*: Molecular characterization of mitochondrial apoptosis-inducing factor. *Nature* 397: 441-446, 1999.

37. Zhang JY, Yi T, Liu J, Zhao ZZ and Chen HB: Quercetin induces apoptosis via the mitochondrial pathway in KB and KBv200 cells. *J Agric Food Chem* 61: 2188-2195, 2013.
38. Westphal D, Dewson G, Czabotar PE and Kluck RM: Molecular biology of Bax and Bak activation and action. *Biochim Biophys Acta* 1813: 521-531, 2011.
39. Hampton MB and Orrenius S: Dual regulation of caspase activity by hydrogen peroxide: Implications for apoptosis. *FEBS Lett* 414: 552-556, 1997.
40. Eleftheriadis T, Pissas G, Liakopoulos V and Stefanidis I: Cytochrome c as a potentially clinical useful marker of mitochondrial and cellular damage. *Front Immunol* 7: 279, 2016.
41. Li H, Huang Y, Chen C, Xiao A, Hou G, Huang Y, Feng X and Guan BO: Real-time cellular cytochrome c monitoring through an optical microfiber: Enabled by a silver-decorated graphene nanointerface. *Adv Sci (Weinh)* 5: 1701074, 2018.
42. Radhakrishnan J, Origenes R, Littlejohn G, Nikolich S, Choi E, Smite S, Lamoureux L, Baetiong A, Shah M and Gazmuri RJ: Plasma cytochrome c detection using a highly sensitive electrochemiluminescence enzyme-linked immunosorbent assay. *Biomark Insights* 12: 1177271917746972, 2017.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.