p57^{KIP2}-mediated inhibition of human trophoblast apoptosis and promotion of invasion *in vitro*

GUO-QIAN HE^{1,2}, GUANG-YU LIU^{1,3}, WEN-MING XU^{1,4}, HUI-JUAN LIAO^{1,4}, XING-HUI LIU^{1,3} and GUO-LIN HE^{1,3}

¹Key Laboratory of Birth Defects and Related Diseases of Women and Children (Sichuan University), Ministry of Education, Departments of ²Pediatrics and ³Obstetrics and Gynecology; ⁴Joint Laboratory of Reproductive Medicine, West China Second University Hospital, Sichuan University, Chengdu, Sichuan 610041, P.R. China

Received February 11, 2018; Accepted April 19, 2019

DOI: 10.3892/ijmm.2019.4175

Abstract. Placental hypoxia serves a role in the early stages of normal pregnancy and is involved in the pathophysiology of preeclampsia. Previously, it was suggested that p57kinase inhibitory protein (KIP)2 regulates the cell cycle during embryogenesis and apoptosis. Recent evidence has indicated that p57KIP2 is increased in preeclamptic placentas and absence of p57KIP2 induces preeclampsia-type symptoms in rats. However, effects of p57^{KIP2} on apoptosis under hypoxic conditions remain to be elucidated. In the present study, HTR-8/SVneo trophoblasts were cultured under hypoxic conditions ($2\% O_2$). Knockdown using small interfering (si)RNA and overexpression of p57KIP2 were utilized to explore the biological function of p57^{KIP2} in apoptosis and cell function in vitro. Furthermore, expression of p57^{KIP2} and apoptosis were evaluated by western blotting, flow cytometry and TUNEL assays, and the response of trophoblasts to hypoxia and the role of p57KIP2 in trophoblast migration and invasion was assessed. The role of p57KIP2 in the JNK signaling pathway in HTR-8/SVneo trophoblasts was further studies. In vitro, protein expression of p57KIP2 was increased in HTR-8/SVneo cells exposed to 2% O2. Exogenous p57KIP2 overexpression significantly decreased the expression of pro-apoptosis proteins, including p53, Bax and cleaved caspase3, under hypoxic conditions for 24 h. In addition, knockdown of p57^{KIP2} increased the response to apoptosis following hypoxia for 24 h. The present study revealed that overexpression of p57^{KIP2} decreased the levels of phosphorylated-JNK. JNK inhibitor treatment combined with the overexpression of p57KIP2 significantly decreased the levels of apoptosis and

Correspondence to: Dr Guo-Lin He, Key Laboratory of Birth Defects and Related Diseases of Women and Children (Sichuan University), Ministry of Education, West China Second University Hospital, Sichuan University, 20 Renmin South Road, Chengdu, Sichuan 610041, P.R. China E mail: xiaopi0500@163.com

E-mail: xiaopi0509@163.com

increased cell invasion and migration. Taken together, p57^{KIP2} knockdown significantly increased apoptosis in HTR-8/SVneo cells exposed to 2% O₂, whereas overexpression of p57^{KIP2} had opposite effects, mediated by the JNK/stress activated protein kinase (SAPK) signaling pathway. The results indicated that hypoxia-induced expression of p57^{KIP2} promoted trophoblast migration and invasion by mediating the JNK/SAPK signaling pathway, which is crucial during placentation. These results may provide a novel molecular mechanism to understand the involvement of p57^{KIP2} in the pathogenesis of preeclampsia.

Introduction

In the early stages of pregnancy (<10 weeks), trophoblasts exist in a hypoxic environment $(1-2\% O_2)$ and hypoxia serves an important role in normal trophoblast differentiation and development (1-4). In addition, hypoxia has been shown to directly induce trophoblast apoptosis and represents a significant pathological factor that contributes to the deficient placentation observed in the early onset of preeclampsia (PE) (5-8). PE is one of the most common pathological complications of pregnancy and is a leading cause of maternal and perinatal morbidity and mortality (9). As for the pathogenesis of PE, it is commonly believed that inadequate migration and the reduced invasive ability of placental extravillous trophoblasts (EVTs) may contribute to PE development (10). A recent study with PE placental explants reported that increased apoptosis and lack of proliferation stimulation were observed in trophoblasts (11). These studies suggest that placental hypoxia is critical for trophoblast physiology and pathology, and apoptosis may be involved in diseases of placental origin. Gaining greater understanding of these mechanisms and pathophysiology in the first trimester of pregnancy will be beneficial for PE treatment and therapy approaches.

p57 [also known as cyclin-dependent kinase inhibitor 1C or kinase inhibitory protein (KIP)-2; p57^{KIP2}] is ubiquitously expressed in human placental cells (12,13). p57^{KIP2} is considered to be a master regulator of the cell cycle during embryogenesis (14-17). Pregnant mice with p57^{KIP2} deficiency presented with symptoms similar to PE (10). Several studies have shown the proapoptotic role of p57^{KIP2} in the pathogenesis of different cancer cell lines (18-21). An antiapoptotic activity of p57^{KIP2}

Key words: p57 kinase inhibitory protein 2, apoptosis, preeclampsia, JNK

has also been observed during embryogenesis (19,22-24). A previous study reported that p57^{KIP2} was significantly increased in all cell types of preeclamptic human placentas with the exception of syncytiotrophoblasts (12). A decrease in p57KIP2 expression was implicated with abnormal trophoblast proliferation (25). However, the role of p57^{KIP2} in trophoblast function and apoptosis remains unclear. In the present study, p57KIP2 expression in trophoblasts (HTR-8/SVneo) under hypoxic conditions was evaluated and attempts were made to further investigate the effect of p57^{KIP2} on migration, invasion and apoptosis in response to hypoxia. The results showed that overexpression of p57KIP2 decreased apoptosis and increased cell invasion and migration in HTR-8/SVneo cells exposed to 2% O₂. This may be mediated by the JNK/stress activated protein kinase (SAPK) signaling pathway. The present study may provide a novel molecular mechanism to understand the involvement of p57KIP2 in the pathogenesis of preeclampsia.

Materials and methods

Cell culture and treatment. The human cell line HTR-8/SVneo was provided by Professor Wenming Xu (Sichuan University, Chengdu, China) (26), derived from short-lived primary EVTs, was used in the present study. Cells were cultured in RPMI-1640 medium (Invitrogen; Thermo Fisher Scientific, Inc.) supplemented with 10% fetal bovine serum (FBS; Gibco; Thermo Fisher Scientific, Inc.), 100 U/ml penicillin and 100 μ g/ml streptomycin. Cells were plated in 6-wells plates (2x10⁵ cells/well). After 24 h in non-hypoxic atmosphere (37°C; 5% CO₂; 21% O₂), cells were transferred to an incubator with 2% O₂ and 5% CO₂ atmosphere for 0, 12, 24, 48 or 72 h at 37°C.

Plasmid constructs and transfection. Small interfering (si)RNA that mediated the knockdown of human p57KIP2 (si-p57) and the nonspecific siRNA control (si-ctrl) were obtained from Thermo Fisher Scientific, Inc. The present study used the following oligos: si-p57, forward, 5'-GGACCUUCCCAGUACUAG UTT-3' and reverse, 5'-ACUAGUACUGGGAAGGUCCTT-3'; and si-ctrl, forward, 5'-AAGGGACTTCCTGTAACAAT GCA-3' and reverse, 5'-CTGGAACGGTGAAGGTGACA-3'. To ectopically express p57KIP2, the synthetic human p57KIP2 sequence (950 bp) was cloned into the plasmid vector. The overexpression plasmids of p57KIP2 (pcDNA3.1-p57) and empty vector plasmid as the control (pcDNA3.1) were purchased from ProteinTech Group, Inc. Cells (2x10⁵) were cultured in 6-well plates overnight prior to transfection. At 70% confluence, cells were transiently transfected with pcDNA3.1 empty vector, pcDNA3.1-p57, si-p57 or si-ctrl at 50 nmol/l. Cell transfection was performed using Lipofectamine 3000™ (Thermo Fisher Scientific, Inc.) according to the manufacturer's instruction. Following transfection for 24 h, cells were incubated under hypoxic conditions for 24 h and then harvested to test the knockdown or overexpression efficiency via RT-qPCR and western blotting, respectively.

Drug treatment. The JNK inhibitor SP600125 was purchased from Sigma-Aldrich (Merck KGaA) and dissolved in DMSO (final concentration, 10 μ M). Transfected HTR-8/SVneo cells (2x10⁵) were incubated with SP600125 for 30 min prior to and throughout hypoxia culturing for 24 h. Cells were

then harvested for subsequent experiments. All cells were compared with a control group, which received an equal volume of DMSO.

Cell migration and invasion assays. Cell migration and invasion was determined by Transwell assay as described previously (27). For the migration assay, cells were resuspended in RPMI-1640 medium containing 1% FBS and placed into the upper well of a Transwell chamber (pore diameter, $8 \,\mu\text{m}$) at 2.5x10⁵ cells/well. Medium containing 10% FBS was placed into the lower wells as a chemoattractant. Cells were incubated under hypoxic conditions for 24 h. Invasion assays were performed as the migration assays, but the filters were coated with growth factor-induced Matrigel (BD Biosciences). Following incubation, cells on the upper surface of the membrane were collected with a sterile cotton swab. Cells that migrated to the lower surface were fixed and stained with 0.1% crystal violet for 10 min at room temperature. The number of migrated and invaded cells was examined via digital light microscopy (magnification, x20). A total of ten visual fields were counted for each sample.

Flow cytometry. HTR-8/SVneo cells were harvested after 24 h hypoxia using trypsin without EDTA, washed with PBS, resuspended in 1 ml binding buffer (eBioscienceTM Annexin V-FITC Apoptosis kit; #BMS500FI-100; Invitrogen; Thermo Fisher Scientific, Inc.) and stained for 15 min with FITC-Annexin V and propidium iodide in the dark at room temperature. The analysis of the cells was conducted via flow cytometry (FACScan; BD Biosciences) equipped with CellQuest V6.0 (BD Biosciences). Cells were sorted into living, necrotic, early apoptotic and late apoptotic cells. The relative ratio of early to late apoptotic cells was counted for further comparison. This assay was repeated >3 times.

TUNEL assays. Apoptosis was validated using the *In Situ* Cell Death Detection kit (Roche Diagnostics). Cells $(5x10^4)$ on coverslips were fixed with 4% PFA for 30 min at room temperature and washed twice with PBS (pH 7.0). Cells were then treated with 3% H₂O₂ for 10 min at room temperature, 0.1% Triton X-100 for 2 min at 4°C, incubated with TUNEL reaction mixture at 37°C for 1 h and then stained with DAPI (1:100) at room temperature for 10 min. Images were taken using a fluorescent inverted microscope (magnification, x40; Nikon Corporation). The percentage of TUNEL-positive cells to apoptotic nuclei was determined in three independent samples and quantified using ImageJ 1.8.0 (National Instituted of Health). The apoptotic index (AI) was calculated as follows: AI = (number of apoptotic cells/total number of cells) x100%.

Western blot analysis. Cells were lysed in cold radioimmunoprecipitation assay lysis buffer (Beyotime Institute of Biotechnology) containing protease inhibitor cocktails. Bicinchoninic acid assays were used to quantify protein concentrations. Equal amounts of protein (80 μ g) were separated on 10% SDS-PAGE gels and transferred to PVDF membranes. Membranes were incubated with rabbit monoclonal antibodies, including anti-p57^{KIP2} (1:700; #2557; Cell Signaling Technology, Inc.), anti-p53 (1:1,000; #ab183544; Abcam), anti-Bax (1:2,000; #50599-2-Ig; ProteinTech Group, Inc.),



Figure 1. Hypoxia induces expression of $p57^{KIP2}$. HTR-8/SVneo cells were subjected to hypoxia for 0, 12, 24, 48 and 72 h. Protein expression was assessed by western blot and (A) images and (B) quantification data are presented for $p57^{KIP2}$; β -actin was used as internal control. *P<0.05 vs. 0 h. KIP, kinase inhibitory protein 2.



Figure 2. $p57^{KIP2}$ affects the expression of apoptosis-associated proteins at hypoxia for 24 h. HTR-8/SVneo cells were transfected with pcDNA3.1-p57, si-p57 or respective controls, followed by hypoxia treatment for 24 h. (A) mRNA expression of $p57^{KIP2}$ in the transfected cells, determined by RT-qPCR. Western blot (B) images and (C) quantification determined in the transfected cells for $p57^{KIP2}$, p53, Bax, cleaved caspase3 and Bcl-2; β -actin was used as internal control. [#]P<0.05 vs. pcDNA3.1; [&]P<0.05 vs. si-ctrl. si-ctrl, siRNA control; si-p57, siRNA targeting $p57^{KIP2}$; KIP, kinase inhibitory protein 2.

284



Figure 3. p57^{KIP2} affects cell apoptosis as determined by flow cytometry and TUNEL assays. HTR-8/SVneo cells were transfected with pcDNA3.1-p57, si-p57 or respective controls, followed by hypoxia treatment for 24 h. Flow cytometry data for the (A) pcDNA3.1, (B) pcDNA3.1-p57, (C) si-ctrl and (D) si-p57 transfected cells exposed to 2% O₂ for 24 h. Quantitative analysis of (E) early and (F) late apoptosis rates determined by flow cytometry. (G) TUNEL-positive cells (white arrow) following hypoxia 24 h; scale bar, 10 μ m. (H) Quantitative analysis of the TUNEL data. [#]P<0.05 vs. pcDNA3.1; [&]P<0.05 vs. si-ctrl. si-ctrl, siRNA control; si-p57, siRNA targeting p57^{KIP2}; KIP, kinase inhibitory protein 2.



Figure 4. JNK signaling pathway inhibitor and overexpression of $p57^{KIP2}$ decrease apoptosis-associated proteins under hypoxic conditions. HTR-8/SVneo cells were transfected with pcDNA3.1-p57 or pcDNA3.1, treated with JNK inhibitor or DMSO, or a combination of both, followed by hypoxia for 24 h. Western blot (A) images and (B) quantification p-JNK, JNK, p53, Bcl-2, Bax and cleaved caspase3 levels; β -actin was used as internal control. $^{e}P<0.05$ vs. pcDNA3.1; $^{s}P<0.05$ vs. pcDNA3.1+DMSO. KIP, kinase inhibitory protein 2; SP, JNK inhibitor SP600125; p-, phosphorylated.

anti-Bcl-2 (1:1,000; #12789-1-AP; ProteinTech Group, Inc.), anti-caspase3 (1:1,000; #9662; Cell Signaling Technology, Inc.), anti-cleaved caspase3 (1:1,000; #9664P; Cell Signaling Technology, Inc.), anti-JNK (1:1,000; #9252; Cell Signaling Technology, Inc.), anti-phosphorylated (p-)Thr183/Tyr18-JNK (p-JNK; 1:1,000; #4671; Cell Signaling Technology, Inc.) and anti-β-actin (1:10,000; #AF0003; Zhengneng Biotechnology, Inc.) overnight at 4°C. Membranes were washed with TBST and incubated with a secondary horseradish peroxidase-conjugated anti-rabbit or anti-mouse antibody (1:10,000; #A0208 or #A0208; Zhengneng Biotechnology, Inc.) at room temperature for 1 h followed by enhanced chemiluminescence assay for



Figure 5. Cleaved caspase3 is affected by p57^{KIP2} under hypoxia. HTR-8/SVneo cells were transfected with pcDNA3.1-p57 or pcDNA3.1, treated with JNK inhibitor or DMSO, or a combination of both and p57^{KIP2} knockdown was achieved by transfection with si-p57; all procedures were followed by hypoxia for 24 h. Immunofluorescence images with cleaved caspase3 (red; white arrow) and nuclear stain (DAPI; blue) for (A) pcDNA3.1, (B) pcDNA3.1-p57, (C) si-ctrl and (D) si-p57 transfected cell, for (E) DMSO and (F) SP treated cells, and for (G) pcDNA3.1 transfected and DMSO and (H) pcDNA3.1-p57 transfected and SP treated cells; scale bar, 10 μ m. (I) Quantified optical density results for cleaved caspase3 analysis. *P<0.05 vs. pcDNA3.1; *P<0.05 vs. si-ctrl; *P<0.05 vs. DMSO; *P<0.05 vs. pcDNA3.1+DMSO. si-ctrl, siRNA control; si-p58, siRNA targeting p57^{KIP2}; KIP, kinase inhibitory protein 2; SP, JNK inhibitor SP600125.

visualization (EMD Millipore). β -actin was used as a loading control. The optical density of the bands was quantitatively analyzed using ImageJ 1.8.0 (National Institutes of Health).

RT-qPCR. SYBR-Green-based RT-qPCR (Invitrogen; Thermo Fisher Scientific, Inc.) was used to examine changes in mRNA levels of $p57^{KIP2}$. Total RNA was isolated from the cells using TRIzol[®] reagent (Invitrogen; Thermo Fisher Scientific, Inc.). PCR was performed with the ABI Prism 7500 Sequence Detection system (Thermo Fisher Scientific, Inc.). The reaction was subjected to 95°C for 60 sec followed by 40 cycles of 95°C for 15 sec and 60°C for 60 sec. Primers for $p57^{KIP2}$ were obtained from Thermo Fisher Scientific, Inc. β -actin was used as the internal control. All reactions were performed in triplicate. The primer sequences were used as follows: $p57^{KIP2}$, forward, 5'-TTCTCAGGCGCTGATCTCTT-3' and reverse,

5'-AGCTGCACTCGGGGATTT-3'; and β -actin, forward, 5'-AAGGGACTTCCTGTAACAATGCA-3' and reverse, 5'-CTGGAACGGTGAAGGTGACA-3'. All results were normalized to β -actin. Relative expression was calculated using the 2^{- $\Delta\Delta$ Cq}} method (27).

Immunofluorescence. Cells (5x10⁴) were grown on a coverslip and fixed with 4% PFA at room temperature for 30 min. Cells were probed with rabbit anti-cleaved caspase3 at 4°C overnight. Alexa Fluor 594 goat anti-rabbit IgG-FITC secondary antibody (1:500; Invitrogen; Thermo Fisher Scientific, Inc.) was added for 2 h at room temperature. DAPI (1:100) was used to stain the cell nuclei for 10 min at room temperature. Cells were mounted with ProLong[®] Gold antifade reagent (Invitrogen; Thermo Fisher Scientific, Inc.). Images were obtained by an inverted fluorescence microscope (magnification, x60; Nikon



Figure 6. $p57^{KIP2}$ and JNK inhibitor affect cell migration under hypoxic conditions. HTR-8/SVneo cells were transfected with pcDNA3.1-p57 or pcDNA3.1, treated with JNK inhibitor or DMSO, or a combination of both and $p57^{KIP2}$ knockdown was achieved by transfection with si-p57; all procedures were followed by hypoxia for 24 h. Microscopy images if the Transwell migration assay for (A) pcDNA3.1, (B) pcDNA3.1-p57, (C) si-ctrl and (D) si-p57 transfected cell, for (E) DMSO and (F) SP treated cells, and for (G) pcDNA3.1 transfected and DMSO and (H) pcDNA3.1-p57 transfected and SP treated cells; magnification, x20; scale bar, 10 μ m. (I) Number of migrated cells. [#]P<0.05 vs. pcDNA3.1; ^{*}P<0.05 vs. si-ctrl; ^{\$}P<0.05 vs. DMSO; ^{*}P<0.05 vs. pcDNA3.1+DMSO. si-ctrl, siRNA control; si-p58, siRNA targeting p57^{KIP2}; KIP, kinase inhibitory protein 2; SP, JNK inhibitor SP600125.

Corporation) and analyzed using Image-Pro Plus 6.0 (Media Cybernetics, Inc.) and Adobe Photoshop CS5 (Adobe Systems, Inc.).

Statistical analysis. All data are expressed as the mean \pm standard deviation. Comparisons between two groups were analyzed by unpaired Student's t-tests. One-way ANOVA followed by Tukey's test was used to assess differences among >2 groups. Data were analyzed using GraphPad Prism 5 (GraphPad Software, Inc.). P<0.05 was considered to indicate a statistically significant difference.

Results

Hypoxia treatment induces $p75^{KIP2}$ expression. To investigate the effect of hypoxia on the expression of $p57^{KIP2}$ in trophoblasts, HTR-8/SVneo cells were incubated with 2% O₂. It was observed that the expression of $p57^{KIP2}$ was significantly increased when exposed to hypoxia for 24-72 h compared with the cells at 0 h (Fig. 1). In the following experiments, hypoxia treatment for 24 h was chosen to elucidate the mechanism.

Overexpression and downregulation of $p57^{KIP2}$ in human trophoblasts. HTR-8/SVneo cells were transfected with the pcDNA3.1-p57, si-p57 or respective controls for 24 h. Following transfection, cells were exposed to hypoxia for further 24 h. RT-qPCR results demonstrated that $p57^{KIP2}$ was significantly overexpressed in the pcDNA3.1-p57 transfected cells compared with the empty vector control and $p57^{KIP2}$ was significantly downregulated in the presence of si-p57 compared with si-ctrl (Fig. 2A). In addition, the western blotting results of $p57^{KIP2}$ were consistent with the RT-qPCR findings (Fig. 2B).

p57^{KIP2} expression affects cell apoptosis under hypoxic conditions. Western blot analysis, flow cytometry and TUNEL assays 288



Figure 7. $p57^{KIP2}$ and JNK inhibitor affect cell invasion under hypoxic conditions. HTR-8/SVneo cells were transfected with pcDNA3.1-p57 or pcDNA3.1, treated with JNK inhibitor or DMSO, or a combination of both and $p57^{KIP2}$ knockdown was achieved by transfection with si-p57; all procedures were followed by hypoxia for 24 h. Microscopy images if the Transwell invasion assay for (A) pcDNA3.1, (B) pcDNA3.1-p57, (C) si-ctrl and (D) si-p57 transfected cell, for (E) DMSO and (F) SP treated cells, and for (G) pcDNA3.1 transfected and DMSO and (H) pcDNA3.1-p57 transfected and SP treated cells; magnification, x20; scale bar, 10 μ m. (I) Number of invaded cells. [#]P<0.05 vs. pcDNA3.1; ^{*}P<0.05 vs. si-ctrl; ^{\$}P<0.05 vs. pcDNA3.1+DMSO. si-ctrl, siRNA control; si-p58, siRNA targeting $p57^{KIP2}$; KIP, kinase inhibitory protein 2; SP, JNK inhibitor SP600125.

were used to obtain comprehensive information regarding the impact of p57^{KIP2} on cell apoptosis. As shown in Fig. 2B, overexpression of p57^{KIP2} increased the protein levels of the antiapoptosis protein Bcl-2 and inhibited expression of the proapoptosis proteins p53, Bax and cleaved caspase3 compared with the empty vector control. The analysis further indicated that expression levels of Bcl-2 decreased and expression of p53, Bax and cleaved caspase3 increased in cells with p57^{KIP2} knockdown compared with the si-ctrl transfected cells. In HTR-8/SVneo cells transfected with si-p57 a significant increase in apoptosis was further observed compared with the si-ctrl cells and a decrease in apoptosis was observed in the p57^{KIP2} overexpressing cells compared with the empty vector control (Fig. 3A-D). Using TUNEL, a similar rate of cell apoptosis was determined for the different samples (Fig. 3E and F). p57^{KIP2} inhibits the JNK signaling pathway to affect apoptosis under hypoxia. The present study next explored the potential mechanisms underlying the regulation of p57^{KIP2} on apoptosis under hypoxic conditions. It has been reported that p57^{KIP2} interacts with JNK/SAPK through its QT domain, mediating a variety of cellular activities, including cell differentiation and survival (28). As shown in Fig. 4, there was a decreasing trend in p-JNK levels after 24 h of hypoxia in p57^{KIP2} overexpressing cells compared with the empty vector control. SP600125, a JNK inhibitor, decreased the ratio of p-JNK/JNK at 24 h compared with the DMSO-treated control cells. Inhibition of JNK further decreased the expression of Bax and cleaved caspase3, and increased the expression of Bcl-2 compared with the DMSO control. Overexpression of p57^{KIP2} in combination with JNK inhibitor treatment markedly decreased the ratio of p-JNK/JNK and expression of Bax and cleaved caspase3, but increased Bcl-2 expression compared with the empty vector plus DMSO-treated control. The effect of overexpression of p57KIP2 in combination with JNK inhibitor treatment on Bax and Bcl-2 appeared stronger compared with the overexpression of p57^{KIP2} or JNK inhibitor treatment alone. The inhibition of JNK had no effect on p53 expression levels. Cleaved caspase3 proteolytically cleaves and activates other caspases and is thought to serve an important role in apoptosis (29). As shown in Fig. 5, immunofluorescence results suggested that cleaved caspase3 was mainly located in the cytoplasm and cell nucleus following 24 h of hypoxia. Quantification of the data showed that cleaved caspase3 was significantly increased following si-p57 transfection and 24 h hypoxia compared with the si-ctrl (Fig. 5I). Overexpression of p57^{KIP2} and inhibition of the JNK pathway significantly decreased the optical density of cleaved caspase3 at 24 h hypoxia compared with the respective controls. It is suggested that p57^{KIP2} may regulate the JNK signaling pathway to inhibit apoptosis during hypoxic injury.

 $p57^{KIP2}$ expression affects cell migration and invasion under hypoxic conditions. HTR-8/SVneo cells have been extensively used to study trophoblast migration and invasion (30). Hypoxia markedly affects trophoblast biological processes, including cell migration and invasion (27). To determine the function of $p57^{KIP2}$ in trophoblast migration, HTR8/SVneo cells were transfected with pcDNA3.1-p57, si-p57, JNK inhibitor or respective controls and exposed to 2% O₂ for 24 h. It was revealed that HTR-8/SVneo cells overexpressing $p57^{KIP2}$ exhibited significantly increased migration and invasion abilities compared with the empty vector control (Figs. 6 and 7). $p57^{KIP2}$ knockdown significantly decreased cell migration and invasion following 24 h of hypoxia compared with the si-ctrl cells. This indicated that $p57^{KIP2}$ enhanced migration and invasion abilities of trophoblasts under hypoxia.

As shown in Fig. 6F, the JNK inhibitor significantly increased HTR-8/SVneo cell migration and invasion at 24 h of hypoxia compared with the DMSO treated control. Overexpression of p57^{KIP2} combined with JNK inhibitor treatment significantly increased HTR-8/SVneo cell migration and invasion compared with the empty vector transfected and DMSO treated cells. The results indicated that under hypoxic conditions p57^{KIP2} affected HTR-8/SVneo cell migration and invasion through the JNK signaling pathway.

Discussion

As a member of the CIP/KIP family, $p57^{KIP2}$ is considered to be a master regulator of the cell cycle during embryogenesis, serving a role in cell cycle control and regulating the induction of apoptosis (31,32). It has been shown that $p57^{KIP2}$ has a minor proapoptotic effect on its own, sensitizing cells to apoptosis (17). A previous study reported that $p57^{KIP2}$ expression enhanced apoptosis in HeLa cells and showed that $p57^{KIP2}$ was a target of caspase activity (21). Other studies have reported an increase in apoptosis and altered differentiation during mouse development in a $p57^{KIP2}$ knockout model (17,33). Previously, it has been shown that $p57^{KIP2}$ serves a role in antiapoptosis regulation, suggesting that whether it promotes or inhibits apoptosis is mainly cellular context-dependent (24). In agreement with these previous studies, the present study revealed that overexpression of p57^{KIP2} increased expression of the antiapoptotic protein Bcl-2 following hypoxia and silencing p57^{KIP2} induced the expression of the proapoptosis-associated proteins p53, Bax and cleaved caspase3. It appeared that in the HTR-8/SVneo hypoxia model, p57^{KIP2} acted as an antiapoptotic molecule.

The family of mammalian mitogen-activated protein kinases includes several subgroups, such as extracellular signal-regulated kinase and JNK/SAPK (34). Accordingly, the JNK/SAPK signaling pathway is implicated in the control of cell growth, transformation, survival and death. It has been shown that p57KIP2 interacts with and inhibits the kinase activity of JNK/SAPK through its QT domain (29,34). The present study provided evidence showing that p57^{KIP2} affected the phosphorylation of JNK under hypoxic conditions. Furthermore, p57^{KIP2} modulated apoptosis by negatively regulating the JNK signaling pathway. In addition, overexpression of p57KIP2 or treatment with the JNK inhibitor increased cell migration and invasion under hypoxic conditions. Treatment of the p57KIP2 overexpressing cells with the JNK inhibitor further affected cell apoptosis, migration and invasion. These results showed that p57^{KIP2} functioned as an inhibitory protein of JNK and affected HTR-8/SVneo migration and invasion.

In conclusion, the results of the present study indicated that $p57^{KIP2}$ served a protective role in apoptosis and increased cell invasion and migration through the JNK signaling pathway under hypoxic conditions. These results suggested that at hypoxia, the $p57^{KIP2}$ -JNK network may serve an important role in regulating HTR-8/SVneo apoptosis and function.

Acknowledgements

The authors would like to thank Mrs. Yan Chen (West China Second University Hospital, Sichuan University, China) for her help advice on flow cytometry.

Funding

The current work was supported by Science and Technology Support Projects in Sichuan Province (grant nos. 2013SZ0004, 2014JY0158 and 2019YFS0411).

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Authors' contributions

GQH, WMX and GLH conceived and designed the study. GQH, GYL and HJL performed the experiments. XHL performed the analysis of data. GHQ and GLH wrote the manuscript. GLH revised the article. All authors read and approved the final manuscript.

Ethics approval and consent for participation

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Wang K, Chen Y, Ferguson SD and Leach RE: MTA1 and MTA3 regulate HIF1a expression in hypoxia-treated human trophoblast cell line HTR8/Svneo. Med J Obstet Gynecol 1: pii: 1017 2013.
- Leslie K, Whitley GS, Herse F, Dechend R, Ashton SV, Laing K, Thilaganathan B and Cartwright JE: Increased apoptosis, altered oxygen signaling, and antioxidant defenses in first-trimester pregnancies with high-resistance uterine artery blood flow. Am J Pathol 185: 2731-2741, 2015.
- Tuuli MG, Longtine MS and Nelson DM: Review: Oxygen and trophoblast biology-a source of controversy. Placenta (32 Suppl 2): S109-S118, 2011.
- Highet AR, Khoda SM, Buckberry S, Leemaqz S, Bianco-Miotto T, Harrington E, Ricciardelli C and Roberts CT: Hypoxia induced HIF-1/HIF-2 activity alters trophoblast transcriptional regulation and promotes invasion. Eur J Cell Biol 94: 589-602, 2015.
- Sharma S, Norris WE and Kalkunte S: Beyond the threshold: An etiological bridge between hypoxia and immunity in preeclampsia. J Reprod Immunol 85: 112-116, 2010.
- 6. Zou Y, Yu X, Lu J, Jiang Z, Zuo Q, Fan M, Huang S and Sun L: Decorin-mediated inhibition of human trophoblast cells proliferation, migration, and invasion and promotion of apoptosis in vitro. Biomed Res Int 2015: 201629, 2015.
- Zhang D, Liu H, Zeng J, Miao X, Huang W, Chen H, Huang Y, Li Y and Ye D: Glucocorticoid exposure in early placentation induces preeclampsia in rats via interfering trophoblast development. Gen Comp Endocrinol 225: 61-70, 2016.
- 8. Heazell AE, Buttle HR, Baker PN and Crocker IP: Altered expression of regulators of caspase activity within trophoblast of normal pregnancies and pregnancies complicated by preeclampsia. Reprod Sci 15: 1034-1043, 2008.
- Saito S and Nakashima A: A review of the mechanism for poor placentation in early-onset preeclampsia: The role of autophagy in trophoblast invasion and vascular remodeling. J Reprod Immunol 101-102: 80-88, 2014.
- Kanayama N, Takahashi K, Matsuura T, Sugimura M, Kobayashi T, Moniwa N, Tomita M and Nakayama K: Deficiency in p57Kip2 expression induces preeclampsia-like symptoms in mice. Mol Hum Reprod 8: 1129-1135, 2002.
- Kadyrov M, Schmitz C, Black S, Kaufmann P and Huppertz B: Pre-eclampsia and maternal anaemia display reduced apoptosis and opposite invasive phenotypes of extravillous trophoblast. Placenta 24: 540-548, 2003.
- Unek G, Ozmen A, Mendilcioglu I, Simsek M and Korgun ET: The expression of cell cycle related proteins PCNA, Ki67, p27 and p57 in normal and preeclamptic human placentas. Tissue Cell 46: 198-205, 2014.
- Unek G, Ozmen A, Kipmen-Korgun D and Korgun ET: Immunolocalization of PCNA, Ki67, p27 and p57 in normal and dexamethasone-induced intrauterine growth restriction placental development in rat. Acta Histochem 114: 31-40, 2012.
- Lee MH, Reynisdóttir I and Massagué J: Cloning of p57KIP2, a cyclin-dependent kinase inhibitor with unique domain structure and tissue distribution. Genes Dev 9: 639-649, 1995.
- and tissue distribution. Genes Dev 9: 639-649, 1995.
 15. Kamura T, Hara T, Kotoshiba S, Yada M, Ishida N, Imaki H, Hatakeyama S, Nakayama K and Nakayama KI: Degradation of p57Kip2 mediated by SCFSkp2-dependent ubiquitylation. Proc Natl Acad Sci USA 100: 10231-10236, 2003.
- Lu Z and Hunter T: Ubiquitylation and proteasomal degradation of the p21(Cip1), p27(Kip1) and p57(Kip2) CDK inhibitors. Cell Cycle 9: 2342-2352, 2010.

- Pateras IS, Apostolopoulou K, Niforou K, Kotsinas A and Gorgoulis VG: p57KIP2: 'Kip'ing the cell under control. Mol Cancer Res 7: 1902-1919, 2009.
- Kavanagh E, Vlachos P, Emourgeon V, Rodhe J and Joseph B: p57(KIP2) control of actin cytoskeleton dynamics is responsible for its mitochondrial pro-apoptotic effect. Cell Death Dis 3: e311, 2012.
- Yan Y, Frisén J, Lee MH, Massagué J and Barbacid M: Ablation of the CDK inhibitor p57Kip2 results in increased apoptosis and delayed differentiation during mouse development. Genes Dev 11: 973-983, 1997.
- Zhang P, Wong C, Liu D, Finegold M, Harper JW and Elledge SJ: p21(CIP1) and p57(KIP2) control muscle differentiation at the myogenin step. Genes Dev 13: 213-224, 1999.
- Samuelsson MK, Pazirandeh A and Okret S: A pro-apoptotic effect of the CDK inhibitor p57(Kip2) on staurosporine-induced apoptosis in HeLa cells. Biochem Biophys Res Commun 296: 702-709, 2002.
- 22. Hsu S, Yu FS, Lewis J, Singh B, Borke J, Osaki T, Athar M and Schuster G: Induction of p57 is required for cell survival when exposed to green tea polyphenols. Anticancer Res 22: 4115-4120, 2002.
- 23. Zhang P, Liégeois NJ, Wong C, Finegold M, Hou H, Thompson JC, Silverman A, Harper JW, DePinho RA and Elledge SJ: Altered cell differentiation and proliferation in mice lacking p57KIP2 indicates a role in Beckwith-Wiedemann syndrome. Nature 387: 151-158, 1997.
- 24. Rossi MN and Antonangeli F: Cellular response upon stress: p57 contribution to the final outcome. Mediators Inflamm 2015: 259325, 2015.
- 25. Takahashi K, Kobayashi T and Kanayama N: p57(Kip2) regulates the proper development of labyrinthine and spongiotrophoblasts. Mol Hum Reprod 6: 1019-1025, 2000.
- 26. Yu Q, Qiu Y, Wang X, Tang J, Liu Y, Mei L, Li M, Yang M, Tang L, Gao H, *et al*: Efficient siRNA transfer to knockdown a placenta specific lncRNA using RGD-modified nano-liposome: A new preeclampsia-like mouse model. Int J Pharm 546: 115-124, 2018.
- 27. Zhang Z, Li P, Wang Y and Yan H: Hypoxia-induced expression of CXCR4 favors trophoblast cell migration and invasion via the activation of HIF-1α. Int J Mol Med 42: 1508-1516, 2018.
- 28. Chang TS, Kim MJ, Ryoo K, Park J, Eom SJ, Shim J, Nakayama KI, Nakayama K, Tomita M, Takahashi K, *et al*: p57KIP2 modulates stress-activated signaling by inhibiting c-Jun NH2-terminal kinase/stress-activated protein Kinase. J Biol Chem 278: 48092-48098, 2003.
- 29. Choudhary GS, Al-Harbi S and Almasan A: Caspase-3 activation is a critical determinant of genotoxic stress-induced apoptosis. Methods Mol Biol 1219: 1-9, 2015.
- 30. Yang Y, Zhang J, Gong Y, Liu X, Bai Y, Xu W and Zhou R: Increased expression of prostasin contributes to early-onset severe preeclampsia through inhibiting trophoblast invasion. J Perinatol 35: 16-22, 2015.
- Ma J, Li J, Yang S, Huang K, Dong X, Sui C and Zhang H: P57 and cyclin G1 express differentially in proliferative phase endometrium and early pregnancy decidua. Int J Clin Exp Med 8: 5144-5149, 2015.
- 32. Korgun ET, Celik-Ozenci C, Acar N, Cayli S, Desoye G and Demir R: Location of cell cycle regulators cyclin B1, cyclin A, PCNA, Ki67 and cell cycle inhibitors p21, p27 and p57 in human first trimester placenta and deciduas. Histochem Cell Biol 125: 615-624, 2006.
- 33. Matsuoka S, Edwards MC, Bai C, Parker S, Zhang P, Baldini A, Harper JW and Elledge SJ: p57KIP2, a structurally distinct member of the p21CIP1 Cdk inhibitor family, is a candidate tumor suppressor gene. Genes Dev 9: 650-662, 1995.
- 34. Minden A and Karin M: Regulation and function of the JNK subgroup of MAP kinases. Biochim Biophys Acta 1333: F85-F104, 1997.