

miR-146a-5p suppresses ATP-binding cassette subfamily G member 1 dysregulation in patients with refractory *Mycoplasma pneumoniae* via interleukin 1 receptor-associated kinase 1 downregulation

HU-NIAN LI*, XU ZHAO*, YONG-JIU ZHA, FANG DU, JIE LIU and LIANG SUN

Emergency and Critical Care Center, Renmin Hospital, Hubei University of Medicine, Shiyan, Hubei 442000, P.R. China

Received September 24, 2018; Accepted May 15, 2019

DOI: 10.3892/ijmm.2019.4380

Abstract. In the present study, we examined the function of microRNA (miR)-146a-5p in patients with refractory *Mycoplasma pneumoniae* pneumonia. In brief, the expression of miR-146a-5p was reduced in patients with refractory *Mycoplasma pneumoniae* pneumonia. Downregulation of miR-146a-5p reduced inflammation in an *in vitro* model of refractory *Mycoplasma pneumoniae* pneumonia, whilst overexpression of miR-146a-5p promoted inflammation. Downregulation of miR-146a-5p induced the protein expression of ATP-binding cassette subfamily G member 1 (ABCG1) and interleukin 1 receptor-associated kinase 1 (IRAK-1), while suppressed expression was observed of the aforementioned proteins following overexpression of miR-146a-5p in an *in vitro* model of refractory *Mycoplasma pneumoniae* pneumonia. The administration of small interfering RNA against RXR or IRAK-1 attenuated the effects of miR-146a-5p on inflammation in an *in vitro* model of refractory *Mycoplasma pneumoniae* pneumonia. Collectively, these results suggested that miR-146a-5p reduced ABCG1 expression in refractory *Mycoplasma pneumoniae* pneumonia via downregulation of IRAK-1.

Introduction

Mycoplasma pneumoniae pneumonia (MPP), a common acute respiratory tract infectious disease in children that has a variety of clinical manifestations, is caused by *Mycoplasma*

pneumoniae (MP) infection, and can lead to multi-organ and multi-system damage (1). In recent years, the cases of refractory MPP (RMPP) are increasing annually. Additionally, the risk of extra-pulmonary complications is elevated, which can severely affect the health of patients (1). MPP affects the quality of life of children and their families; therefore clinical investigations into this disease is required (2,3).

MicroRNA (miRNA/miR) can regulate gene expression by activating or suppressing gene transcription, serving a key role in physiological development, as well as the development of disease (4). Additionally, miRNA not only exerts a crucial part in cell differentiation and organ development (5), but can also serve as a molecular marker for various physiological and pathological states (5). Numerous studies indicated that miRNAs are closely associated with the genesis, development and prognosis of pulmonary infection (6,7); however, no intensive studies have been performed, yet further investigation is required (6). On the contrary, research into miRNAs have provided notable insight into the molecular mechanism, clinical diagnosis and treatment for pulmonary infectious disease (6). Osei *et al* (8) revealed that decreased levels of miR-146a-5p in chronic obstructive pulmonary disease-associated fibroblasts may induce a more pronounced pro-inflammatory phenotype. Pradhan *et al* reported that miRNAs interfere with translation of their target gene and regulate a variety of biological actions exerted by these target genes (9).

Member 1 of human transporter subfamily (ABCA1) and ATP-binding cassette subfamily G member 1 (ABCG1) belong to the ATP binding cassette transporter (ABC) superfamily, whose main function is to promote the outflow of intracellular free cholesterol (10). The ABCA1- and ABCG1-regulated cholesterol outflow from macrophages is key step in preventing and reversing foam cell formation (10).

The ABCA1- and ABCG1-regulated cholesterol outflow from macrophages serves a key role in scavenging excessive cholesterol in tissues, including vascular walls (11,12). Therefore, dysfunctions in ABCA1 and ABCG1 may lead to excessive cholesterol accumulation in macrophages, forming foam cells, which subsequently invade the vascular wall and promote MPP genesis and development (11,12).

Interleukin (IL)-1 receptor-associated kinases (IRAKs) are the members with similar composition and structure of

Correspondence to: Dr Liang Sun, Emergency and Critical Care Center, Renmin Hospital, Hubei University of Medicine, 37 Chaoyang Middle Road, Shiyan, Hubei 442000, P.R. China
Email: slzaaz222009@126.com

*Contributed equally

Key words: microRNA-146a-5p, refractory *Mycoplasma pneumoniae* pneumonia, ATP-binding cassette subfamily G member 1, interleukin 1 receptor-associated kinase 1

serine-threonine; four members have been reported as of yet, including IRAK-1, IRAK-2, IRAK-M and IRAK-4 (13). Among them, only IRAK-1 and IRAK-4 possess kinase activity, while IRAK-4 is considered as an essential factor required to activate the Toll-IL receptor and myeloid differentiation primary response 88 (MyD88)-dependent pathway (13,14). Following the phosphorylation processes in the aforementioned pathways, IRAKs dissociate from MyD88, and bind with tumor necrosis factor receptor (TNF) associated factor 6 (TRAF6) to form an IRAK1-TRAF6 complex (14). Subsequently, nuclear factor- κ B (NF- κ B) and transcription factor activated activator protein-1 are activated, while the release of pro-inflammatory cytokines, including IL-6, IL-1 and TNF- α is promoted, inducing the downstream cascade of inflammatory reactions, resulting in tissue inflammatory injury (15). Li *et al* (16) revealed that miR-146a-5p antagonized advanced glycation end products (AGEs)- and *Porphyromonas gingivalis* (P.g)-LPS-induced ABCA1 and ABCG1 dysregulation in macrophages via IRAK-1 downregulation (16). In the present study, the function of miR-146a-5p in patients with refractory MPP was investigated.

Materials and methods

Patients with MPP. Children diagnosed (male, n=10; female, n=10) with MPP were enrolled from Renmin Hospital. The age range was 1 month to 12 years. The peripheral blood of all patients was collected, and patients underwent chest radiography and *M. pneumonia* tests, including specific IgM in by ELISA. The exclusion criteria for the enrollment of patients were: i) Those with congenital heart diseases, hereditary metabolic diseases, neurological disorders, bronchopulmonary dysplasia, and immunodeficiency; and ii) patients co-infected with other pathogens. The present study was approved by the Ethics Committee of Renmin Hospital, Hubei University of Medicine. Written informed consent obtained from the family of patients.

Quantification of miRNA level. Total RNA was extracted from lung tissue or cell samples using TRIzol reagent (Thermo Fisher Scientific, Inc.) and cDNA was obtained using a RevertAid First Strand cDNA Synthesis kit (Thermo Fisher Scientific, Inc.) at 42°C for 60 min and 82°C for 10 sec. 500 ng RNA was labeled and using an equimolar concentration of Cyanine-3-CTP-labelled universal mouse reference (Stratagene). miRNA analysis was performed on an ABI 7500 real-time PCR system (Applied Biosystems; Thermo Fisher Scientific, Inc) using TaqMan Universal PCR Master Mix (Thermo Fisher Scientific, Inc.). miRNA expression levels were calculated using an endogenous control with the $2^{-\Delta\Delta C_q}$ method (17). PCR amplification conditions were as follows: Initial denaturation at 95°C for 10 min, 40 cycles of 95°C for 30 sec, 60°C for 30 sec, 72°C for 30 sec. Sequence information: miR-146a-5p: Forward, 5'-GGGGTGAGAACTGAATTCAT-3' and reverse, 5'-CAGTGCGTGTCTGGAGT-3'; U6: Forward, 5'-GCTTCGGCAGCACATATACTAAAAT-3' and reverse, 5'-CGCTTCACGAATTTGCGTGTTCAT-3'. Data were examined using Genespring 12.6.1 (Agilent Technologies) and Agilent Feature Extraction Software (version A.10.7.3.1).

Gene chip analysis. The total RNA was extracted using mirVana™ miRNA Isolation kit (Ambion; Thermo Fisher

Scientific, Inc.). The Agilent Human miRNA array (8*60K, V16.0, Agilent Technologies, Inc.) and total RNA was labeled by Cy-3 using miRNA Complete Labeling and Hyb kit (5190-0456, Agilent Technologies, Inc.). Data was performed by Agilent Microarray Scanner (G2565CA, Agilent Technologies, Inc.) and normalized by Quantile algorithm with Gene Spring Software 12.6 (Agilent Technologies, Inc.).

MPP model, and hematoxylin and eosin staining. Ethical approval was obtained from Renmin Hospital, Hubei University of Medicine. Male C57BL/6 mice (4-5 g, 1 week old, n=20) were purchased from the Animal laboratory of Renmin Hospital. C57BL/6 mice of model group were injected with 2 mg/kg of lipopolysaccharide (LPS, intraperitoneally) for 12 h under anesthesia with 50 mg/kg sodium pentobarbital as reported previously by Fang *et al* (18). C57BL/6 mice of sham group were injected with 50 mg/kg sodium pentobarbital. Then, mice were sacrificed via decollation whilst anesthetized. Lung tissue samples were collected and fixed with 4% paraformaldehyde for 24 h at room temperature or stored at -80°C.

Lung tissue samples fixed with paraformaldehyde were paraffin-embedded. The samples were cut into 10- μ m sections using a paraffin slicing machine (RM2235; Leica Microsystems GmbH), stained with hematoxylin and eosin for 10 min at room temperature, and observed under a light microscope (magnification, x100, BH3-MJL; Olympus Corporation).

Cell culture. A549 cells was maintained in Dulbecco's Modified Eagle's medium (Gibco; Thermo Fisher Scientific, Inc.) supplemented with 10% fetal bovine serum (Gibco; Thermo Fisher Scientific, Inc.). 100 ng/ml miR-146a-5p (5'-CCGATATATATCCTCACTT-3'), anti-miR-146a-5p (5'-ATGGGCTATATAGGAGTGAACC-3') and negative mimics (5'-TTCTCCGAACGTGTCACGT-3'), small interfering RNA (si)-IRAK-1 (sc-35704, Santa Cruz Biotechnology, Inc.) or si-RXR (sc-36447, Santa Cruz Biotechnology, Inc.) were transfected into cells using Lipofectamine® 2000 for 48 h. After 48 h, cells were treated with 100 ng/ml of LPS for 6 h at 37°C as reported by Guo and Cheng (19).

ELISA kits. Serum samples were centrifuged at 2,000 x g for 20 min at 4°C and cell samples were centrifuged at 1,000 x g for 10 min at 4°C. Serum samples were used to measure the levels of TNF- α (cat. no. H052), IL-1 β (cat. no. H002), IL-6 (cat. no. H007) and IL-18 (H015) using ELISA kits (Nanjing Jiancheng Institute of Bioengineering). Cell samples were lysed with radioimmunoprecipitation assay (RIPA; Beyotime Institute of Biotechnology) and used to measure TNF- α , IL-1 β , IL-6 and IL-18 levels using ELISA kits.

Luciferase reporter assay. The network of signaling pathway components revealed that IRAK-1 may targets of miR-146a-5p be important using http://www.targetscan.org/vert_71/. IRAK-1-expressing pc-DNA-3.1 plasmids and anti-miR-146a-5p were co-transfected into cells using Lipofectamine 2000 for 48 h. Reporter activity levels were quantified 48 h after transfection using a Dual-Luciferase Reporter Assay kit (Beyotime Institute of Biotechnology). Measurements were obtained using a Lumat LB 9507




Figure 1. Expression of miR-146a-5p in patients or mice with refractory MPP. (A) TNF- α , (B) IL-1 β , (C) IL-6 and (D) IL-18 levels. Lung tissues from mice were analyzed using (E) H&E staining (magnification, x100), (F) gene chip for miRNA expression, and (G and H) reverse transcription-quantitative polymerase chain reaction for miR-146a-5p expression in mice with refractory MPP and patients with refractory MPP. **P<0.01 vs. sham group. Control, control normal group; MPP, *Mycoplasma pneumoniae* pneumonia; miR, microRNA; IL, interleukin; Sham, sham control group; TNF- α , tumor necrosis factor- α .

instrument (Berthold Technologies). Normalization was performed via comparisons with *Renilla* luciferase activity.

Western blotting analysis. Cells were harvested and lysed in RIPA buffer. Protein content was measured using a BCA assay (Beyotime Institute of Biotechnology). Protein (50 μ g) was separated via 10% SDS-PAGE and transferred to polyvinylidene difluoride membranes, which were washed in Tris-buffered saline with Tween-20 (TBST) for three times and blocked with 5% non-fat in TBST for 1 h at room temperature. Subsequently, the membranes were incubated with antibodies against ABCG1 (ab218528, 1:1,000, Abcam), IRAK-1 (sc-515512, 1:1,000, Santa Cruz Biotechnology, Inc.), retinoid X

receptor (RXR, sc-46659, 1:1,000, Santa Cruz Biotechnology, Inc.), liver X receptor (LXR, sc-271064, 1:1,000, Santa Cruz Biotechnology, Inc.) and GAPDH (sc-69778, 1:5,000, Santa Cruz Biotechnology, Inc.) at 4°C overnight. The membrane was washed three times with TBST for 15 min and incubated with a horseradish peroxidase-conjugated secondary antibody (sc-2004, 1:5,000, Santa Cruz Biotechnology, Inc.) at 37°C for 1 h. Proteins were detected by Super Signal chemiluminescent reagent (Thermo Fisher Scientific, Inc.) and measured using FlowJo 7.6.1 (FlowJo, LLC).

Immunofluorescence (IF). Cells were washed with PBS and fixed with 4% paraformaldehyde for 20 min at room




Figure 2. Anti-miR-146a-5p regulates inflammation in an *in vitro* model of refractory *Mycoplasma pneumoniae* pneumonia. Reverse transcription-quantitative polymerase chain reaction for (A) miR-146a-5p, (B) TNF- α , (C) IL-1 β , (D) IL-6, (E) IL-18, (F) COX-2 and (G) iNOS levels. ** $P < 0.01$ vs. negative group. anti-146a-5p, downregulation of miR-146a-5p expression group; COX-2, cyclooxygenase-2; IL, interleukin; iNOS, inducible nitric oxide synthase; miR, microRNA; negative, negative mimics group; TNF- α , tumor necrosis factor- α .

temperature. Cells were blocked with 5% bovine serum albumin (Beyotime Institute of Biotechnology) and 0.1% Triton X-100 for 1 h and incubated with anti-IRAK-1 (Santa Cruz Biotechnology, Inc.) at 4°C overnight. Cells were washed with PBS and incubated with goat anti-rabbit IgG-CFL 555 (Santa Cruz Biotechnology, Inc.) for 1 h at room temperature. Cells were washed with PBS and stained with DAPI for 15 min at room temperature. Then, cells were washed with PBS and observed under a light microscope (magnification, x100, BH3-MJL; Olympus Corporation).

Statistical analysis. Data are presented as the mean \pm standard deviation. Statistical significance between groups was determined by a Student's t-test or one-way analysis of variance followed by a Tukey's post-hoc test. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Expression of miR-146a-5p in patients or mice with refractory MPP. To investigate the effects of miR-146a-5p in a mouse

model with refractory MPP, we first analyzed changes in the expression of miR-146a-5p in a mouse model with refractory MPP. The results revealed that the expression serum levels of TNF- α , IL-1 β , IL-6 and IL-18 were significantly increased in MPP mice, compared with sham mice (Fig. 1A-D). H&E staining showed that pulmonary alveoli were smaller in the MPP mouse model, compared with sham mice (Fig. 1E). In addition, the expression of miR-146a-5p was significantly reduced in mice with refractory MPP, compared with the sham group (Fig. 1F and G). Consistently, miR-146a-5p expression was significantly downregulated in patients with refractory MPP compared with the control normal group (Fig. 1H). Therefore, miR-146a-5p may serve a role in the inflammation associated with refractory MPP.

miR-146a-5p regulates inflammation in an *in vitro* model of refractory MPP. Then, downregulation of miR-146a-5p significantly increased the levels of TNF- α , IL-1 β , IL-6 and IL-18 in an *in vitro* model of refractory MPP, compared with negative control group (Fig. 2). However, overexpression of miR-146a-5p significantly reduced the levels of TNF- α ,





Figure 3. miR-146a-5p regulates inflammation in an *in vitro* model of refractory *Mycoplasma pneumoniae* pneumonia. Reverse transcription-quantitative polymerase chain reaction for (A) miR-146a-5p expression. ELISA of (B) TNF- α , (C) IL-1 β , (D) IL-6, (E) IL-18, (F) COX-2 and (G) iNOS levels. **P<0.01 vs. negative group. COX-2, cyclooxygenase-2; IL, interleukin; iNOS, inducible nitric oxide synthase; miR, microRNA; negative, negative mimics group; TNF- α , tumor necrosis factor- α .

IL-1 β , IL-6 and IL-18 in an *in vitro* model of refractory MPP compared with the negative control group (Fig. 3). These results suggested that miR-146a-5p was involved in the inflammation associated with MPP *in vitro*.

miR-146a-5p regulates the protein expression of ABCG1/IRAK-1 in an in vitro model of refractory MPP. To confirm the mechanism of miR-146a-5p in inflammation in an *in vitro* model of refractory MPP, miR-146a-5p and anti-miR-146a-5p mimics were respectively transfected into cells. Gene chip analysis revealed that the protein expression levels of ABCG1 and IRAK-1 were increased in an *in vitro* model of refractory MPP following downregulation of miR-146a-5p, compared with the negative group (Fig. 4A). The 3-untranslated region of IRAK-1 is complementary to the seed sequence of anti-miR-146a-5p; the luciferase activity levels were significantly increased in an *in vitro* model of refractory MPP following downregulation of miR-146a-5p, compared with the negative control group (Fig. 4B and C). The network of signaling pathway components revealed that ABCG1 and IRAK-1 may be important in the development of MPP (Fig. 4D)

using http://www.targetscan.org/vert_71/. These results of IF analysis revealed that IRAK-1 protein expression was induced in an *in vitro* model of refractory MPP following downregulation of miR-146-5p compared with the negative control group (Fig. 4E). Upregulation of miR-146a-5p significantly reduced the protein expression of IRAK-1, RXR, LXR, p65 and ABCG1 in an *in vitro* model of refractory MPP in comparison with the negative control group (Fig. 5). However, downregulation of miR-146a-5p significantly increased the protein expression of IRAK-1, RXR, LXR, p65 and ABCG1 in an *in vitro* model of refractory MPP, compared with the negative control group (Fig. 6).

Small interfering RNA (si)-IRAK-1 attenuates the effects of miR-146a-5p on inflammation and ABCG1 in an in vitro model of refractory MPP. To evaluate the function of IRAK-1 in the effects of miRNA-146a-5p on inflammation in an *in vitro* model of infantile pneumonia, si-IRAK-1 was used to downregulate the protein expression of IRAK-1; the expression levels of RXR, LXR, p65 and ABCG1 were significantly downregulated in an *in vitro* model of infantile pneumonia, compared with the group transfected with anti-miR-146a-5p



si-RXR attenuates the effects of miR-146a-5p on inflammation and ABCG1 in an in vitro model of refractory Mycoplasma Pneumoniae pneumonia. Finally, si-RXR was used to suppress the protein expression of RXR, LXR, p65 and ABCG1 in an *in vitro* model of refractory MPP

following downregulation of miR-146a-5p. Compared with the anti-miR-146a-5p transfected group, silencing of RXR expression significantly suppressed the expression of the aforementioned proteins (Fig. 8A-E). In addition, downregulation of RXR significantly attenuated the effects of anti-miR-146a-5p on the levels of TNF- α , IL-1 β , IL-6 and IL-18 in an *in vitro* model of refractory MPP (Fig. 8F-K). Overall, the results demonstrated that miR-146a-5p may be critical for inflammation and the expression of ABCG1 in an *in vitro* model of refractory MPP.




Figure 4. Continued. (E) Immunofluorescence for IRAK-1 protein expression (magnification, x200). miR, microRNA; negative, negative mimics group; anti-146a-5p, downregulation of miR-146a-5p expression group; ABCG1, ATP-binding cassette subfamily G member 1; CHK1, checkpoint kinase 1; hsa, *homo sapiens*; IRAK-1, interleukin 1 receptor-associated kinase 1; NEDD1, neural precursor cell expressed, developmentally downregulated 1; PDK1, phosphoinositide-dependent kinase 1; PI3K, phosphoinositide 3-kinase; TRAF6, tumor necrosis factor associated factor 6; Tpl2, mitogen-activated protein kinase kinase kinase 8; UTR, untranslated region; ZEB1, zinc finger E-box-binding homeobox 1.




Figure 5. MicroRNA-146a-5p regulates ABCG1/IRAK-1 protein expression in an *in vitro* model of refractory *Mycoplasma pneumoniae* pneumonia. (A-E) Densitometry analysis of IRAK-1, RXR, LXR, p65 and ABCG1 protein expression following western blotting. (F) Western blot gel. ** $P < 0.01$ vs. negative group. ABCG1, ATP-binding cassette subfamily G member 1; IRAK-1, interleukin 1 receptor-associated kinase 1; LXR, liver X receptor; RXR, retinoid X receptor; negative, negative mimics group; 146a-5p, overexpression of microRNA-146a-5p expression group.

Discussion

RMPP is a type of MPP, which cannot be alleviated with macrolide antibiotics; instead, RMPP rapidly develops into a

form of intra-pulmonary disease, thereby potentially inducing various extra-pulmonary complications as a life-threatening condition (19). Therefore, early diagnosis and treatment are problems to overcome, yet the precise pathogenesis and




Figure 6. MicroRNA-146a-5p regulates ABCG1/IRAK-1 protein expression in an *in vitro* model of refractory *Mycoplasma pneumoniae* pneumonia. (A-E) Densitometry analysis IRAK-1, RXR, LXR, p65 and ABCG1 protein expression following western blotting. (F) Western blot gel. ** $P < 0.01$ vs. negative group. ABCG1, ATP-binding cassette subfamily G member 1; IRAK-1, interleukin 1 receptor-associated kinase 1; LXR, liver X receptor; RXR, retinoid X receptor; negative, negative mimics group; anti-146a-5p, downregulation of microRNA-146a-5p expression group.

regulatory factors of MP infection remain unclear (20). Previously, the role of pattern recognition receptors in MP infection has attracted wide attention (20). Our findings demonstrated that miR-146a-5p expression was inhibited in patients with refractory MPP. Osei *et al* (8) showed that decreased levels of miR-146a-5p in chronic obstructive pulmonary disease-associated fibroblasts may induce a more pronounced pro-inflammatory phenotype. Ramkaran *et al* (21) reported miR-146a as a target for lowering inflammation in coronary artery disease patients. These results indicated miR-146a as a potential target for regulating inflammation in MPP. Additionally, we used LPS to induce refractory MPP in a mouse model. LPS can exhibit a broad range of effects and may not recapitulate the conditions of mycoplasma infection as a whole; thus, this poses as a the limitation of the present study. We aim to generate a more precise model in the future to validate our findings.

miRNAs serve a key role in regulating gene expression, and its role has attracted increasing attention (22). Numerous miRNAs are involved in inducing an intra-pulmonary inflammatory response, which is closely associated with tumors (22). Epidemiological analysis suggests that, ~1/4 of tumors are induced by chronic infection and other chronic inflammatory

responses, while active measures to control the inflammatory response may suppress tumor genesis (23). However, the relationship between miRNAs and the intra-pulmonary inflammatory response, and its role require further investigation (24), which may provide novel insight into the differential expression of miRNAs in the blood to diagnose and treat lung disease (22). Our findings demonstrated that downregulation of miR-146a-5p increased the levels of TNF- α , IL-1 β , IL-6 and IL-18 in an *in vitro* model of refractory MPP. Wu *et al* (25) reported that miR-146a-5p inhibits TNF- α -induced adipogenesis via targeting insulin receptors in primary porcine adipocytes. In the present study, A549 cells were treated with LPS to generate an *in vitro* model of *Mycoplasma pneumoniae* infection; however, as of the aforementioned limitations, additional models are required to validate our findings.

Accumulating evidence has revealed that, IRAK-1/4 serves critical roles in regulating the Toll-like receptor (TLR) pathway (26). IRAK1 phosphorylation by IRAK4 is the earliest activation step in the TLR and MyD88-dependent pathway, which is closely related to the formation of an early receptor complex and activation of downstream signaling molecules (26). Of note, the expression of downstream cytokines associated with the TLR signaling pathway is severely




Figure 7. Si-IRAK-1 reverses the effects of anti-miR-146a-5p on inflammation and ABCG1 in an *in vitro* model of refractory *Mycoplasma pneumoniae* pneumonia. (A-E) Densitometry analysis of IRAK-1, RXR, LXR, p65 and ABCG1 protein expression following western blotting. (F) Western blot gel. (G) TNF- α , (H) IL-1 β , (I) IL-6, (J) IL-18, (K) COX-2 and (L) iNOS levels. **P<0.01 vs. negative group; ##P<0.01 vs. anti-146a-5p group. ABCG1, ATP-binding cassette subfamily G member 1; COX-2, cyclooxygenase-2; IL, interleukin; iNOS, inducible nitric oxide synthase; IRAK-1, interleukin 1 receptor-associated kinase 1; LXR, liver X receptor; RXR, retinoid X receptor; negative, negative mimics group; anti-146a-5p, downregulation of microRNA-146a-5p expression group; Si, small interfering RNA; TNF- α , tumor necrosis factor- α .




Figure 8. Small interfering-RXR reverses the effects of miR-146a-5p on inflammation and ABCG1 in an *in vitro* model of refractory *Mycoplasma pneumoniae* pneumonia. (A-D) Densitometry analysis of RXR, LXR, p65 and ABCG1 protein expression following western blotting. (E) Western blot gel. (F) TNF-α, (G) IL-1β, (H) IL-6, (I) IL-18, (J) COX-2 and (K) iNOS levels. **P<0.01 vs. negative group; ##P<0.01 vs. anti-146a-5p group. ABCG1, ATP-binding cassette subfamily G member 1; COX-2, cyclooxygenase-2; IL, interleukin; iNOS, inducible nitric oxide synthase; miR, microRNA; IRAK-1, interleukin 1 receptor-associated kinase; LXR, liver X receptor; RXR, retinoid X receptor; negative, negative mimics group; anti-146a-5p, downregulation of miR-146a-5p expression group; TNF-α, tumor necrosis factor-α.




Figure 9. miR-146a-5p reduces ABCG1 dysregulation in refractory *Mycoplasma pneumoniae* pneumonia via downregulation of IRAK-1. ABCG1, ATP-binding cassette subfamily G member 1; IRAK-1, interleukin 1 receptor-associated kinase; LXR, liver X receptor; miR, microRNA; RXR, retinoid X receptor.

suppressed in IRAK4^{-/-} knockout mice (11). In addition, research on IRAKs, which play key regulatory roles in the TLR signaling pathway, has focused on peripheral cells and IRAK-1/4-induced inflammation (27). Our findings revealed that downregulation of miR-146a-5p induced IRAK-1, RXR, LXR, p65 and ABCG1 protein expression in an *in vitro* model of refractory MPP. Lo *et al* (28) reported that miR-146a-5p mediates high glucose-induced endothelial inflammation via targeting IRAK-1 expression.

ABCG1 belongs to the adenosine triphosphate binding cassette subfamily G family of proteins, which can regulate outflow of the intracellular free cholesterol and prevent the formation of foam cells (29). ABCG1 is also highly expressed in endothelial cells, and its effect on promoting cholesterol outflow from endothelial cells serves a key role in protecting

the normal endothelial cell function (30). Upregulation of ABCG1 expression can reduce TNF- α -induced vascular endothelial cell injury, the mechanism of action of which may be related to its suppression on TNF- α -induced vascular endothelial cell inflammation (10). Our results indicate that si-IRAK-1 and si-RXR reduced the effects of miR-146a-5p on inflammation and ABCG1 in an *in vitro* model of refractory MPP. Li *et al* (16) showed that miR-146a-5p antagonized AGEs- and P.g-LPS-induced ABCA1 and ABCG1 dysregulation in macrophages via IRAK-1 downregulation (16).

In conclusion, the present study demonstrated that miR-146a-5p expression in refractory MPP was reduced. miR-146a-5p was proposed to attenuate inflammation and ABCG1 expression in refractory MPP via the IRAK-1/RXR/LXR signaling pathway (Fig. 9). We proposed a novel anti-inflammatory role of miR-146a-5p in refractory MPP, suggesting that miR-146a-5p may be a potential therapeutic target for the treatment of refractory MPP.

Acknowledgements

Not applicable.

Funding

No funding was received.

Availability of data and materials

The analyzed data sets generated during the study are available from the corresponding author on reasonable request.

Authors' contributions

LS made substantial contributions to the design of the study; HNL, XZ, YJZ, FD and JL performed the experiments. LS and HNL analyzed the data, and LS wrote the manuscript. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The present study was approved by Renmin Hospital, Hubei University of Medicine.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that have no competing interests.

References

- Jeong J, Kang I, Kim S, Park KH, Park C and Chae C: Comparison of 3 vaccination strategies against porcine reproductive and respiratory syndrome virus, mycoplasma hyopneumoniae, and porcine circovirus type 2 on a 3 pathogen challenge model. *Can J Vet Res* 82: 39-47, 2018.
- Zhang Y, Zhou Y, Li S, Yang D, Wu X and Chen Z: The clinical characteristics and predictors of refractory mycoplasma pneumoniae pneumonia in children. *PLoS One* 11: e0156465, 2016.
- Marrie TJ, Beecroft M, Herman-Gnjidic Z and Poulin-Costello M: Symptom resolution in patients with mycoplasma pneumoniae pneumonia. *Can Respir J* 11: 573-577, 2004.
- Zhao J, Chen C, Guo M, Tao Y, Cui P, Zhou Y, Qin N, Zheng J, Zhang J and Xu L: MicroRNA-7 deficiency ameliorates the pathologies of acute lung injury through elevating KLF4. *Front Immunol* 7: 389, 2016.
- Alipour SD, Adcock IM, Garssen J, Mortaz E, Varahram M, Mirsaeidi M and Velayati A: The roles of miRNAs as potential biomarkers in lung diseases. *Eur J Pharmacol* 791: 395-404, 2016.
- Jadideslam G, Ansarin K, Sakhinia E, Alipour S, Pouremamali F and Khabbazi A: The microRNA-326: Autoimmune diseases, diagnostic biomarker, and therapeutic target. *J Cell Physiol* 233: 9209-9222, 2018.
- Yan X, Li W, Yang L, Dong W, Chen W, Mao Y, Xu P, Li D, Yuan H and Li YH: MiR-135a protects vascular endothelial cells against ventilator-induced lung injury by inhibiting PHLPP2 to activate PI3K/Akt pathway. *Cell Physiol Biochem* 48: 1245-1258, 2018.
- Osei ET, Florez-Sampedro L, Tasena H, Faiz A, Noordhoek JA, Timens W, Postma DS, Hackett TL, Heijink IH and Brandsma CA: miR-146a-5p plays an essential role in the aberrant epithelial-fibroblast cross-talk in COPD. *Eur Respir J* 49: 1602538, 2017.
- Pradhan AK, Emdad L, Das SK, Sarkar D and Fisher PB: The enigma of miRNA regulation in cancer. *Adv Cancer Res* 135: 25-52, 2017.
- Cao XJ, Zhang MJ, Zhang LL, Yu K, Xiang Y, Ding X, Fan J, Li JC and Wang QS: TLR4 mediates high-fat diet induced physiological changes in mice via attenuating PPARgamma/ABCG1 signaling pathway. *Biochem Biophys Res Commun* 503: 1356-1363, 2018.
- Guo L, Chen CH, Zhang LL, Cao XJ, Ma QL, Deng P, Zhu G, Gao CY, Li BH, Pi Y, *et al*: IRAK1 mediates TLR4-induced ABCA1 downregulation and lipid accumulation in VSMCs. *Cell Death Dis* 6: e1949, 2015.
- Ma G, Huang X, Bi Y, Ren D, Xu F, Sun Q, Zhang R, Hu J, Niu W, Guo Z, *et al*: Association study between ABCB1, ABCB6 and ABCG1 polymorphisms and major depressive disorder in the Chinese han population. *Psychiatry Res* 270: 1170-1171, 2018.
- Li YW, Zhao F, Mo ZQ, Luo XC, Li AX and Dan XM: Characterization, expression, and functional study of IRAK-1 from grouper, *epinephelus coioides*. *Fish Shellfish Immunol* 56: 374-381, 2016.
- Zeng KW, Zhang T, Fu H, Liu GX and Wang XM: Schisandrin B exerts anti-neuroinflammatory activity by inhibiting the toll-like receptor 4-dependent MyD88/IKK/NF- κ B signaling pathway in lipopolysaccharide-induced microglia. *Eur J Pharmacol* 692: 29-37, 2012.
- Lee KH, Jeong J, Woo J, Lee CH and Yoo CG: Globular adiponectin exerts a pro-inflammatory effect via I κ B/NF- κ B pathway activation and anti-inflammatory effect by IRAK-1 downregulation. *Mol Cells* 41: 762-770, 2018.
- Li X, Ji Z, Li S, Sun YN, Liu J, Liu Y, Tian W, Zhou YT and Shang XM: miR-146a-5p antagonized AGEs- and P.g-LPS-induced ABCA1 and ABCG1 dysregulation in macrophages via IRAK-1 downregulation. *Inflammation* 38: 1761-1768, 2015.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
- Fang X, Liu X, Meng C, Fu Y, Wang X, Li B, Tu F, Zhao F and Ren S: Breed-linked polymorphisms of porcine toll-like receptor 2 (TLR2) and TLR4 and the primary investigation on their relationship with prevention against mycoplasma pneumoniae and bacterial LPS challenge. *Immunogenetics* 65: 829-834, 2013.
- Guo J and Cheng Y: MicroRNA-1247 inhibits lipopolysaccharides-induced acute pneumonia in A549 cells via targeting CC chemokine ligand 16. *Biomed Pharmacother* 104: 60-68, 2018.
- Vranckx K, Maes D, Marchioro SB, Villarreal I, Chiers K, Pasmans F and Haesebrouck F: Vaccination reduces macrophage infiltration in bronchus-associated lymphoid tissue in pigs infected with a highly virulent mycoplasma hyopneumoniae strain. *BMC Vet Res* 8: 24, 2012.
- Ramkaran P, Khan S, Phulukdaree A, Moodley D and Chuturgoon AA: miR-146a polymorphism influences levels of miR-146a, IRAK-1, and TRAF-6 in young patients with coronary artery disease. *Cell Biochem Biophys* 68: 259-266, 2014.
- Chen X, Cheng JY and Yin J: Predicting microRNA-disease associations using bipartite local models and hubness-aware regression. *RNA Biol* 15: 1192-1205, 2018.

23. Yang Q, Zhang D, Li Y, Li Y and Li Y: Paclitaxel alleviated liver injury of septic mice by alleviating inflammatory response via microRNA-27a/TAB3/NF- κ B signaling pathway. *Biomed Pharmacother* 97: 1424-1433, 2018.
24. Liu H, He Y, Jiang Z, Shen S, Mei J and Tang M: Prodigiosin alleviates pulmonary fibrosis through inhibiting miRNA-410 and TGF- β 1/ADAMTS-1 signaling pathway. *Cell Physiol Biochem* 49: 501-511, 2018.
25. Wu D, Xi QY, Cheng X, Dong T, Zhu XT, Shu G, Wang LN, Jiang QY and Zhang YL: miR-146a-5p inhibits TNF- α -induced adipogenesis via targeting insulin receptor in primary porcine adipocytes. *J Lipid Res* 57: 1360-1372, 2016.
26. Vollmer S, Strickson S, Zhang T, Gray N, Lee KL, Rao VR and Cohen P: The mechanism of activation of IRAK1 and IRAK4 by interleukin-1 and toll-like receptor agonists. *Biochem J* 474: 2027-2038, 2017.
27. Ma H, Zhang S, Xu Y, Zhang R and Zhang X: Analysis of differentially expressed microRNA of TNF- α -stimulated mesenchymal stem cells and exosomes from their culture supernatant. *Arch Med Sci* 14: 1102-1111, 2018.
28. Lo WY, Peng CT and Wang HJ: MicroRNA-146a-5p mediates high glucose-induced endothelial inflammation via targeting Interleukin-1 receptor-associated kinase 1 expression. *Front Physiol* 8: 551, 2017.
29. Chen S, Villeneuve L, Jonker D, Couture F, Laverdière I, Cecchin E, Innocenti F, Toffoli G, Lévesque E and Guillemette C: ABCC5 and ABCG1 polymorphisms predict irinotecan-induced severe toxicity in metastatic colorectal cancer patients. *Pharmacogenet Genomics* 25: 573-583, 2015.
30. Aleidi SM, Howe V, Sharpe LJ, Yang A, Rao G, Brown AJ and Gelissen IC: The E3 ubiquitin ligases, HUWE1 and NEDD4-1, are involved in the post-translational regulation of the ABCG1 and ABCG4 lipid transporters. *J Biol Chem* 290: 24604-24613, 2015.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.