

Ablation of CXCR4 expression in cardiomyocytes exacerbates isoproterenol-induced cell death and heart failure

MIN CHENG¹, CAN CHEN¹, KUNWU YU¹, XIAO LV², QIUTANG ZENG¹, NIANGUO DONG³ and FENG ZHU¹

Departments of ¹Cardiology, ²Orthopedics and ³Surgery, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei 430022, P.R. China

Received July 11, 2022; Accepted November 21, 2022

DOI: 10.3892/ijmm.2022.5216

Abstract. CXCR4 is a seven-transmembrane-spanning Gi-coupled receptor for the SDF-1 chemokine and plays a critical role in cardiovascular development and post-injury repair. However, the specific role of CXCR4 in cardiomyocytes is incompletely understood. It was hypothesized that CXCR4 activation in cardiomyocytes antagonizes β -adrenoceptor/Gs signaling-induced cardiac dysfunction. Cardiomyocyte-specific CXCR4 knockout (CXCR4-^{CM}KO) mice were generated by crossing CXCR4^{fl/fl} and MHC-Cre^{+/+} mice. Their cardiac structure and function in the basal state are equivalent to that of the control MHC-Cre^{+/+} littermates until at least 4 months old. However, following continuous subcutaneous administration of isoproterenol (Iso) via an osmotic mini-pump, the ventricular myocardial contractility, dilation, cardiomyocyte apoptosis, and interstitial fibrosis are worse in CXCR4-^{CM}KO mice than in MHC-Cre^{+/+} littermates. In the cultured H9C2 cardiomyocytes, SDF-1 treatment markedly attenuated Iso-induced apoptosis and reduction in phospho-Akt, and this protective effect was lost by knockdown of CXCR4 or by co-treatment with Gi inhibitors. In conclusion, CXCR4 promotes cardiomyocyte survival and heart function during β -adrenergic stress.

Introduction

The C-X-C chemokine receptor type 4 (CXCR4) is a G_{α_i} protein-coupled receptor, and its only identified ligand is the CXC chemokine stromal-derived factor-1 (SDF-1). The SDF-1/CXCR4 axis is essential for organogenesis during embryonic development, including hematopoiesis, cardiac

septum formation, and neuronal development; genetic deletion of CXCR4 or SDF-1 leads to almost identical developmental defects and embryonic lethality (1-3). In the adult animals, previous evidence has shown that SDF-1/CXCR4 axis plays an indispensable role in the maintenance of stem/progenitor cells in the bone marrow niche, recruitment of stem/progenitor cells to sites of injury, ischemic neovascularization and tissue repair (4-10). These findings have led to the notion of targeting SDF-1/CXCR4 axis for therapy of ischemic heart disease (10-14).

The role of CXCR4 signaling in cardiomyocyte biology, however, is not well understood. Recent evidence suggested that in the developing heart, SDF-1/CXCR4 signaling is crucial not only for epicardia-guided coronary vessel formation but also for the expansion of the second heart field (6,15). In the adult, CXCR4 is expressed in cardiomyocytes across mice, rats and humans (16). Notably, SDF-1 treatment of isolated mouse papillary muscles or rat cardiomyocytes attenuates β -adrenergic agonist Isoproterenol (Iso)-induced calcium mobilization and contractility (17), and cardiac CXCR4 knockout mice develop progressive cardiomyopathy (18). Importantly, CXCR4 is expressed at high levels in the cardiomyocytes of patients with end-stage heart failure (19), the first cause of death in the developed countries.

Heart failure is featured by reduced cardiac pump function unable to maintain the systemic demand of blood supply, for which the impaired β -adrenergic receptor (β -AR) signaling plays a central role (20). β -ARs, predominantly β_1 (80%) and β_2 (20%) ARs in a normal mammalian heart, primarily couple G_{α_s} , although β_2 has been shown to also couple G_{α_i} . Upon binding to ligands (catecholamine hormone epinephrine and neurotransmitter norepinephrine) or agonist (isoproterenol), β -ARs activate G_{α_s} GTPase, leading to adenylyl cyclase activation, cAMP production, and signaling events including protein kinase A (PKA) activation and phosphorylation (activation) of proteins involved in cellular calcium mobilization and contraction (21). While acute activation of β -ARs increase heart rate (chronotropy) and contractility (inotropy) to augment cardiac performance, over or chronic activation results in cardiomyocyte death, adverse cardiac remodeling, and progressive deterioration of function (22-26). Thus, β blockers are one of most widely used treatment and have been shown to increase the long-term survival of patients with chronic heart failure (27,28). Furthermore, evidence over last

Correspondence to: Dr Min Cheng or Dr Feng Zhu, Department of Cardiology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, 1377 Jiefang Avenue, Wuhan, Hubei 430022, P.R. China
E-mail: min_cheng@hust.edu.cn
E-mail: zhufeng@hust.edu.cn

Key words: CXCR4, cardiomyocytes, Gs stress, cardiac remodeling, apoptosis, adrenergic, heart failure

three decades suggested that β_1 AR not β_2 AR plays a causal role in cardiomyocyte apoptosis and heart failure (26,29-31). Mice lacking β_1 and β_2 ARs or overexpressing β_2 ARs have essentially normal basal cardiac function (32,33), whereas mice overexpressing β_1 AR, $G\alpha_s$, or active-PKA display a similar cardiac phenotype, including initial increased responsiveness to catecholamines and hypertrophy followed by cardiomyocyte death, fibrosis, cardiac dilation and failure (34-36).

In the present study, the role of cardiomyocyte CXCR4 in Iso-induced heart failure in cardiomyocyte-specific CXCR4 knockout (^{CM}KO) mice was investigated. It was found that these mice display normal heart structure and function in the basal state but are more susceptible to Iso-induced cardiomyocyte apoptosis, fibrosis and heart failure, and that SDF-1/CXCR4 signaling attenuates Iso-induced cardiomyocyte death in a $G\alpha_i$ dependent manner.

Materials and methods

Mice. Male, 8-10 weeks old mice on C57BL/6J background with body weights between 25 and 27 g were used. CXCR4 $^{fl/fl}$ mice and α -MHC-Cre mice were obtained from Jax lab and bred to generate cardiomyocyte-specific CXCR4 knockout (MHC-Cre $^{+}$; CXCR4 $^{fl/fl}$ or ^{CM}KO) mice. All animal experiments in the present study were approved (IACUC approval no. 3005) by the Animal Care and Use Committee of Huazhong University of Science and Technology (Wuhan, China) and performed in compliance with the 'Guide for the Care and Use of Laboratory Animals' (NIH publication) and relevant ethical regulations for animal testing and research. Animal numbers for the experiments were pre-calculated based on the reported similar studies from the authors and other laboratories with 80% power (37,38). The basic characteristics of our model animals were reported, and physiological and biochemical measurements were performed in double-blinded fashion. After Iso or Saline minipump implantation, the animal health and behaviour were monitored daily for the first 3 days, then every other day until they were euthanized at day 14. Over the course of the study, no experimental animal experienced conditions needed for euthanasia before the end of experiments. For euthanasia, the mice were placed in a designated chamber with continuous introduction of 100% CO $_2$ at a fill rate of 50% of the chamber volume per min and for a duration of 15 min; the death was verified by loss of consciousness, head sinking, and disappearance of muscle tone. Then the mice were subjected to a secondary method, either vital organ removal (for 30 mice used in histological analyses) or cervical dislocation (for 30 mice used in echocardiographic analyses), to ensure death. For colony maintenance, mice were fed *ad libitum* and maintained at 22-24°C ambient temperature and 40-60% humidity under a 12:12-h light: dark cycle.

Chemicals and bioagents. All chemicals and bioagents used in the present study, including Iso, AMD3100, Gi inhibitor Pertussis toxin (all from Sigma-Aldrich; Merck KGaA), SDF-1 (R&D Systems, Inc.) were purchased from commercial sources and authenticated by the providers. Their concentrations, Iso (100 μ M), SDF-1 (500 ng/ml), and AMD3100 (100 ng/ml), used on H9C2 cells were based on pilot studies by the authors and consistent with a similar study previously reported (39).

Continuous Iso infusion model. ^{CM}KO and Cre mice received Iso (45 mg/kg/day) or saline (control) continuously via Alzet mini-osmotic pumps (model 2004; Durect Corporation) for up to 28 days. Pumps were prepared and subcutaneously implanted in the mice by following the manufacturer's instructions. The surgery was performed under anesthesia by intraperitoneal injection of sterile pentobarbital sodium (50 mg/kg body weight, Sigma-Aldrich; Merck KGaA) and post-operative care was performed by following our approved animal study protocol. For pain management, Metacam was subcutaneously injected (1 mg/kg) at the end of surgery and continued twice daily for 2 days. The heart function was monitored via echocardiography at day 0, 3, 7, and 14. Then the mice were euthanized, and the cardiac tissues were harvested for histological analyses.

Echocardiography. Cardiac function was recorded and analyzed using Vevo 770 high-resolution ultrasound system (VisualSonics, Inc.) as previously described (10). Mice were lightly anesthetized via 2% isoflurane inhalation and secured with surgical tapes to the platform in supine position. Heart rates were recorded. Data were collected after three days of training. Parasternal long-axis view, short-axis view at the papillary muscle level and 2-D guided M-mode images were recorded. Left ventricular ejection fraction, internal diameter and anterior wall thickness during systole and diastole were measured, and the measurements were performed by an individual who was blinded to the treatment assignments.

Histology. Both formalin-fixed paraffin-embedded (FFPE) and fixed frozen (FF) sections were used in the present study. For FFPE sections, cardiac tissues were fixed in 10% formalin overnight, embedded in paraffin, and cut at 5 μ m in thickness as previously described (40). For FF sections, tissues were fixed in 4% paraformaldehyde for 4 h, dehydrated in 30% sucrose and embedded in OCT compound followed by cryosectioning into 8 μ m in thickness. A total of 3 sections per heart and 6 fields per section were examined. The H&E staining (Thermo Fisher Scientific, Inc.) was performed by following standard procedures. The Sirius Red/Fast Green (Chondrex) staining were performed by following manufacturer's instructions. For TUNEL staining, FF sections or cells growing on slides were fixed in 4% paraformaldehyde for 30 min, then permeabilized with proteinase K solution (20 μ g/ml; for tissue) or 0.2% Triton X-100 (for cells) for 5 min at room temperature. After wash with PBS, the sections were stained with TUNEL reagent (Sigma-Aldrich; Merck KGaA) for 60 min at 37°C. After wash, the nuclei were counter stained with DAPI for 7 min at room temperature. For each section, 4-6 fields were examined under fluorescent microscope. To analyze tissue CXCR4 expression, anti-CXCR4 antibody (Abcam, cat. no. ab124824; 1:200) was used in the histochemical staining, as previously reported by the authors (9). The histological assessments and analyses were performed by an individual blinded to treatment assignments.

Cells. H9C2, a well characterized and widely used cardiomyocyte cell line that was originally derived from rat embryonic heart tissue, was used as the *in vitro* cardiomyocyte model (41). The cells were purchased from the American Type Culture

Collection (cat. no. CRL-1446), maintained in DMEM with 10% FBS, and used until passage 8.

3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) assay. The MTT assay was performed to evaluate the viability of H9C2 cells after treatment with test reagents in 96-well plates by using CellTiter 96 Nonradioactive Cell Proliferation Assay kit (Promega Corporation). First, 15 μ l of dye solution was added to each well, and the plate was returned to incubator for 4 h at 37°C. Then, 100 μ l of Solubilization/Stop Solution was added to each well. The colored formazan product is stable at 4°C, and absorbance was recorded at 570 nm using a 96-well plate reader.

Lenti-CXCR4 shRNA. The lentiviral vector that carries CXCR4 shRNA or non-targeting (NT) shRNA was obtained from Sigma-Aldrich; Merck KGaA with the knockdown specificity and efficiency authenticated by the company. The CXCR4-shRNA base sequence was 5'-GGATCAGCA TCGATTCTTCA-3' and the NT-shRNA base sequence was 5'-TTCTCCGAACGTGTCACGT-3'. The vectors were packaged following the manufacturers' instructions. The multiplicity of infection was 2 for application on H9C2 cells, and the transduced cells were selected in puromycin for 10 days before evaluation for CXCR4 expression and myocyte functions.

Reverse transcription-quantitative (RT-q) PCR. RT-qPCR was performed via standard techniques as previously described by the authors (9). Briefly, total RNA was extracted with RNA Stat-60 (Tel-Test, Inc.), and RNA was reverse transcribed with the Taqman Multiscribe RT kit (Applied Biosystems; Thermo Fisher Scientific, Inc.) according to the manufacturer's instructions. qPCR was performed in duplicate with cDNA from 10 ng of RNA by using the Lightcycler hybridization Probes Master Mix [Roche Diagnostics (Shanghai) Co., Ltd.]; a negative control (lacking a template) was included for each probe set. Relative gene expression was calculated using the $2^{-\Delta\Delta C_q}$ method (42) and normalized to GAPDH. The primer sequences were as follows: mouse CXCR4 forward, 5'-CCT CGCCTTCTTCCACTGTT-3' and reverse, 5'-CTGGG AGAGCTTTTGAAGTTG-3'; and mouse GAPDH forward, 5'-GGGTCCCAGCTTAGGTTTCATC-3' and reverse, 5'-TAC GGCCAAATCCGTTTCA-3'.

Western blot analysis. Western blotting was performed via standard techniques as previously described by the authors (43), using primary antibodies against phosphorylated (p)-Akt, total Akt, β -actin and relevant horseradish peroxidase (HRP)-linked secondary antibodies (all from Cell Signaling Technology, Inc.). Briefly, the membrane was first blotted with anti-p-Akt (cat. no. 9271; 1:1,000) at 4°C for overnight, then with anti-rabbit IgG (cat. no. 7074; 1:5,000) before application of chemiluminescence detection reagents and image taking. Next, the membrane was placed in the stripping solution [25 mM glycine-HCl, pH 2, 1% (w/v) SDS] with agitation for 30 min at room temperature, followed by the second round of immunoblotting with the use of anti-total-Akt (cat. no. 2920; 1:1,000) and anti-Mouse IgG (cat. no. 7076; 1:5,000). β -actin was immunoblotted in some experiments as loading control by

using anti- β -actin (cat. no. 4967; 1:1,000) and then anti-rabbit IgG (cat. no. 7074; 1:5,000).

Statistical analysis. All values are presented as the mean \pm SEM. Unpaired two-tailed Student's t-test was used for comparison between two means. One-way or two-way analysis of variance (ANOVA), followed by Bonferroni post-hoc tests, were used in multiple (>2) group comparisons with one or two independent variables, respectively. GraphPad Prism 8 software (GraphPad Software, Inc.) was used for statistical analysis. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

CXCR4 is expressed in the adult cardiomyocytes. The present investigation was initiated by assessing the expression of CXCR4 in wild-type mouse hearts. Immunohistological staining revealed that CXCR4 protein is expressed in the cardiomyocytes (Fig. S1) in the pattern consistent with plasmalemma localization as previously reported (16). This result is consistent with previous studies and confirmed that CXCR4 is expressed in cardiomyocytes.

Loss of CXCR4 in cardiomyocyte worsens Iso-induced heart failure. To investigate the role of CXCR4 in cardiomyocyte, a cardiomyocyte-specific CXCR4 knockout (^{CM}KO) mouse line was generated, $\alpha MHC-Cre; CXCR4^{f/f}$, by crossing $CXCR4^{f/f}$ mice (44) with $\alpha MHC-Cre$ mice. The resultant ^{CM}KO mice were born at the expected Mendelian ratio and displayed no overt defect in gross appearance or cardiac structure for at least up to 4 months old, suggesting that CXCR4 expression specifically in the cardiomyocyte lineage is not essential for heart development. To induce heart failure, ^{CM}KO mice and control $\alpha MHC-Cre$ (Cre) littermates (8-10 weeks old) were subjected to a 14-day mini-pump delivery of Iso, a well characterized adrenergic agonist that induces heart failure primarily as the result of activated β_1 -adrenergic signaling and Gs stress. Serial echocardiography revealed that Iso treatment induced heart failure in both groups of mice; however, in ^{CM}KO mice, the left ventricular contractility was worse, chamber was more dilated, and end-systolic left-ventricular anterior wall thickness was thinner than in Cre control littermates (Fig. 1). In addition, it was observed that ^{CM}KO mice exhibit a higher heart rate than control Cre littermates under the same anesthesia regimen (Fig. S2). These worsened parameters in ^{CM}KO mice suggested that CXCR4 expression in cardiomyocytes plays a protective role in Gs stress-induced heart failure.

Loss of CXCR4 in cardiomyocytes worsens Iso-induced cardiac adverse remodeling and apoptosis. In another set of experiments, mice were administered Iso or saline via subcutaneous mini-pumps and euthanized at day 14, and the hearts were excised. The heart size and heart-weight (H.W.) to body-weight (B.W.) ratio appeared normal and comparable between ^{CM}KO and Cre mice with Saline treatment (Fig. 2A and B). However, with Iso treatment, these measures were significantly elevated in both groups and significantly deteriorated in ^{CM}KO mice compared with Cre mice (Fig. 2A and B). Histological assessments of cardiac tissue




Figure 1. Loss of CXCR4 in cardiomyocytes worsens Iso-induced cardiac dysfunction. $CMKO$ (αMHC -Cre; $CXCR4^{fl/fl}$) mice and control αMHC -Cre (Cre) littermates were treated with Iso (45 mg/kg BW/day) or saline via a subcutaneously-implanted mini-osmotic pump for 14 days, then cardiac functions were serially evaluated via echocardiography at day 0 (baseline), 3, 7, and 14. (A) Representative image of echocardiography. (B) LVEF. (C) LVID at end systole (S, left panel) and end diastole (D, right panel). (D) Thickness of LVAW at end systole (S, left panel) and end diastole (D, right panel). $n=15$ per group. * $P<0.05$ (B-D: One-way ANOVA). Iso, isoproterenol; $CMKO$, CXCR4 knockout; LVEF, left ventricular ejection fraction; LVID, left ventricular internal diameter; LVAW, left ventricular anterior wall; Iso, isoproterenol.

revealed that Iso induced a significantly increased interstitial fibrosis (Sirius Red/Fast Green staining; Fig. 2C and D) and apoptotic cardiomyocytes (TUNEL staining; Fig. 2E and F) in $CMKO$ mice. Thus, it was identified that loss of CXCR4 in cardiomyocytes increases Iso-induced cardiac cell death and adverse remodeling.

SDF-1 treatment abrogates Iso-induced apoptosis in cultured cardiomyocytes. Since CXCR4 is a Gi protein-coupled receptor, and Gi protein-coupled receptor signaling is inhibitory to Gs effects, it was hypothesized that activation of CXCR4 by SDF-1 counters β -AR/Gas signaling to improve cardiomyocyte survival. In cultured H9C2 cardiomyocytes, treatment with Iso for 24 h significantly reduced cell number and viability (Fig. 3A-C) and increased cell apoptosis (Fig. 3D and E). While treatment with SDF-1 or AMD3100 (CXCR4 antagonist) alone did not show a significant effect, co-treatment with SDF-1 almost completely prevented Iso-induced reduction in cell number and viability (Fig. 3A-C) and significantly attenuated Iso-induced apoptosis (Fig. 3D and E). Consistently, the level of p-Akt, a major downstream effector of Gi signaling-activated PI-3 kinase- β and cell survival, was diminished by Iso treatment (Fig. 3F) however elevated by SDF-1 treatment (Fig. 3G); importantly, co-treatment with SDF-1 attenuated Iso-induced reduction in p-Akt (Fig. 3H). Collectively, these results suggested that CXCR4 protection of Iso-induced cell death is mediated by SDF-1, and that SDF-1/CXCR4 signaling promotes cell survival.

SDF-1/CXCR4 axis promotes cardiomyocyte survival through Gai protein signaling. To understand the role of CXCR4 more definitively, H9C2 cardiomyocytes were transduced with Lenti-CXCR4-shRNA, which knocked down CXCR4 mRNA expression by $\sim 90\%$ (Fig. 4A). While knockdown of CXCR4 alone did not affect cell number and viability, it worsened Iso-induced reduction of cell number and viability and more importantly, abolished SDF-1 mediated protection from Iso-induced reduction in cell number and viability (Fig. 4B and C). Consistently, knockdown of CXCR4 also diminished the ability of SDF-1 to ameliorate Iso-induced p-Akt reduction (Fig. 4D). These results confirmed that SDF-1 mediated cell survival is dependent on CXCR4 expression. To further determine the role of Gi signaling in SDF-1-mediated protection, H9C2 cells were treated with Iso/SDF-1 or Saline (-) in the presence or absence of a Gi-specific inhibitor. Clearly, SDF-1 attenuated the Iso-induced cell number and viability (Fig. 4E and F), and the protection was lost in the presence of Gi inhibitor but not vehicle control, confirming that SDF-1 protection from Iso-induced cell death is mediated by Gi signaling.

Discussion

In the present study, it was identified that loss of CXCR4 in cardiomyocytes renders the mice more vulnerable to Iso-induced heart failure-characterized by worsened LV contractility, greater LV dilation and thinner ventricular wall- and that SDF-1




Figure 2. Loss of CXCR4 in cardiomyocytes worsens Iso-induced cardiac adverse remodeling and apoptosis. Mice were administered Iso or Saline via mini-pump implantation and at day 14, euthanized. (A) Representative gross appearance of the mouse hearts. (B) H.W. to B.W. ratio. (C and D) Representative images of Sirius Red/Fast Green staining [C; Original magnifications, x40 for upper panels, x100 for lower panels) and (D) quantification of the interstitial fibrosis areas]. (E and F) Representative images of (E) TUNEL (red) and DAPI (blue) double staining and (F) quantification of the TUNEL⁺ apoptotic cardiomyocytes. Original magnifications, x100. HPF, high-powered field. n=6 per group. *P<0.05 and **P<0.01 (B: Two-way ANOVA; D and F: unpaired two-tailed t-test). Iso, isoproterenol; H.W., heart weight; B.W., body weight; CM^{KO}, CXCR4 knockout; WT, wild-type; N.S., not significant.

treatment potentially protects cardiomyocytes from Iso-induced cell death in a CXCR4 and Gi-dependent manner. These data suggested that SDF-1/CXCR4 signaling plays a protective role in β -adrenergic stress-induced cell death, cardiac adverse remodeling and progression to dilation, and heart failure.

The results of cardiac functional analysis in CXCR4^{CMKO} mice are consistent with a previous study (45); however, in the present study direct evidence was provided of Iso-induced cardiomyocyte apoptosis *in vivo* and *in vitro* in the absence or downregulation of CXCR4, respectively. Furthermore, it




Figure 3. SDF-1/CXCR4 signaling attenuates Iso-induced apoptosis in cardiomyocytes. H9C2 cardiomyocytes were treated for 24 h with Saline, Iso (100 μ M), SDF-1 (500 ng/ml), AMD3100 (100 ng/ml), Iso + SDF-1, or Iso + AMD3100, then analyzed. (A) Microphotograph of the treated cells. Original magnification, $\times 400$. (B) Manual counts of live cells. (C) MTT assays of viable cells. (D) TUNEL (red) and DAPI (blue) double staining (original magnification, $\times 400$). (E) quantification of the TUNEL⁺ apoptotic cells. (F-H) Western blot analyses of p-Akt and total Akt in cells treated with (F) Iso, (G) SDF-1 and (H) Iso + SDF-1. $n=6$ per group. *** $P<0.001$ (B, C and E: One-way ANOVA). Iso, isoproterenol; p-, phosphorylated.




Figure 4. SDF-1/CXCR4 mediates cardiomyocyte survival via Gi signaling. H9C2 cardiomyocytes were transduced with Lenti-CXCR4-shRNA or Lenti-NT-shRNA and subjected to the following experiments: (A) Reverse transcription-quantitative PCR analysis of CXCR4 mRNA levels. (B and C) Treatment with Saline, Iso (100 μ M), or Iso + SDF-1 (500 ng/ml) for 24 h, followed by (B) manual counting of cell numbers and (C) MTT viability assays. n=6 per group. (D) Treatment with Iso (+) plus SDF-1 (+) or vehicle (-), followed by western blotting of p-Akt and total Akt. (E and F) Treatment with Saline, Iso, or Iso + SDF-1 for 24 h in the presence of Gi inhibitor Pertussis toxin (200 ng/ml) (+) or vehicle (-), followed by (E) manual cell counts and (F) MTT viability assays. n=6 per group. **P<0.01 and ***P<0.001. A, unpaired two-tailed t-test; B and C, E and F, Two-way ANOVA. shRNA, short hairpin RNA; NT, non-targeting; p-, phosphorylated; N.S., not significant.

was demonstrated that this effect is mediated by SDF-1 action and Gi protein signaling. The extent of SDF-1 protection of Iso-induced cardiomyocyte death is particularly striking. Previous studies suggested that SDF-1/CXCR4-mediated cell survival from ischemic stress is linked to G β γ mediated activation of phosphoinositide 3-kinase (likely PI3K β) and

Akt (11,13,46). Indeed, it was found that SDF-1 induces a significantly increased p-Akt in cardiomyocytes and also attenuates Iso-induced downregulation of p-Akt. Nevertheless, since sustained activation of G α s-coupled β -ARs results in apoptotic loss of cardiomyocytes via a cAMP-dependent mechanism (30), SDF-1/CXCR4 signaling, through G α i activation,

may also enhance cell survival by directly downregulating adenylyl cyclase activity. Further experiments would be needed to confirm this assertion. Furthermore, it was observed that during progression of heart failure, Gai protein expression is increased (20). This increase, previously considered as a compensatory effect from altered β -AR signaling, may in fact play an important protective role by substantiating the effect of CXCR4 and potentially other Gai coupled receptors.

Iso is the prototypical β -AR agonist with well documented profile in positive inotropic/chronotropic effects and induction of apoptosis (47), and continuous infusion is commonly used in rodents to induce phenotypic features of myocardial infarction and heart failure (37). Our *in vivo* analyses were particularly focused on the initial 14 days following Iso treatment, which is considered the best time window for detecting direct effects of adrenergic stress on cardiomyocytes in this model (37). Nevertheless, the detrimental effects are primarily mediated via β_1 -AR and G_{α_s} signaling, thus investigation of CXCR4, a G_{α_i} -coupled receptor, with Iso stress has a limitation, because heart failure is a complex disease and involves mechanism beyond β -AR and G_{α_s} stress. Thus, examining the role of SDF-1/CXCR4 with additional heart failure model, such as pressure overload, is warranted in future studies.

Notably, it was observed that CXCR4^{CM}KO mice display an elevated heart rate associated with progression to heart failure. This is likely due to the dysregulated adrenergic signaling, although detailed electrophysiologic features and underlying mechanisms are unclear at present and remain a subject of our ongoing investigations. Notably, upregulation of CXCR4 expression was observed in patients with atrial fibrillation (AF) a decade ago (48). Recently, a flurry of bioinformatics analyses of clinical high-throughput data sets revealed that CXCR4 is a potent hub gene strongly associated with the pathogenesis of AF (49-52). Previous studies with a mouse model of AF suggested that CXCR4 antagonist AMD3100 can treat AF by ameliorating calcium mishandling and tissue inflammation (50,52). Given that CXCR4 signaling can mediate both cardiomyocyte survival and tissue inflammation, which appear to have contrasting effects on AF, strategies that target CXCR4 for treatment of cardiac arrhythmia would need to be carefully evaluated before clinical application.

One of the limitations to the present study is the limited scope of mechanistic study. Gain-of-function experiments were performed only in cultured cardiomyocytes, but not *in vivo* in animals. While the conclusion that ablation of CXCR4 expression in cardiomyocytes exacerbates isoproterenol-induced cell death and heart failure is fully supported by the data, the translational value of the present finding or if CXCR4 signaling can be used for treatment of cardiac adverse remodeling and heart failure remains to be rigorously investigated by using relevant CXCR4 activator/agonists particularly in large animal models with longer term follow-ups.

Lastly, CXCR4-null embryos exhibit dysplasia of the ventricular septum (3). However, a structural or functional defect in our CXCR4^{CM}KO mouse line was not observed for at least 4 months after birth, consistent with a previous study from Agarwal *et al* (53). The findings of the present study suggested that CXCR4 is required for septum formation probably through a non-cardiomyocyte lineage or cardiomyocyte

progenitor cells prior to expressing α -MHC, which remains a subject of future research by the authors.

In conclusion, the present results suggested that SDF-1/CXCR4 signaling promotes cardiomyocyte survival and attenuates heart failure induced by β -AR overactivation.

Acknowledgements

Not applicable.

Funding

This study was supported by NSFC (grant nos. 82070318 and 81570257) and Wuhan Science and Technology program (grant no. 2019020701011455).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

MC conceptualized the study, interpreted the data, and wrote the manuscript. CC, KY, and XL performed the experiments and analyzed the data. ND, QZ and FZ made intellectual contributions and assisted in data interpretation. FZ edited the manuscript. MC and CC confirm the authenticity of all the raw data. All authors read and approved the final version of the manuscript.

Ethics approval and consent to participate

All animal experiments in the present study were approved (IACUC approval no. 3005) by the Animal Care and Use Committee of Huazhong University of Science and Technology (Wuhan, China) and performed in compliance with the 'Guide for the Care and Use of Laboratory Animals' (NIH publication) and all relevant ethical regulations for animal testing and research.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Tachibana K, Hirota S, Iizasa H, Yoshida H, Kawabata K, Kataoka Y, Kitamura Y, Matsushima K, Yoshida N, Nishikawa S, *et al*: The chemokine receptor CXCR4 is essential for vascularization of the gastrointestinal tract. *Nature* 393: 591-594, 1998.
2. Nagasawa T, Hirota S, Tachibana K, Takakura N, Nishikawa S, Kitamura Y, Yoshida N, Kikutani H and Kishimoto T: Defects of B-cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking the CXC chemokine PBSF/SDF-1. *Nature* 382: 635-638, 1996.
3. Zou YR, Kottmann AH, Kuroda M, Taniuchi I and Littman DR: Function of the chemokine receptor CXCR4 in haematopoiesis and in cerebellar development. *Nature* 393: 595-599, 1998.

4. Askari AT, Unzek S, Popovic ZB, Goldman CK, Forudi F, Kiedrowski M, Rovner A, Ellis SG, Thomas JD, DiCorleto PE, *et al*: Effect of stromal-cell-derived factor 1 on stem-cell homing and tissue regeneration in ischaemic cardiomyopathy. *Lancet* 362: 697-703, 2003.
5. Jujo K, Ii M, Sekiguchi H, Hamada H, Thorne T, Klyachko E, Clarke T, Ito A, Misener S, Renault MA and Losordo DW: Bone marrow MMP-9 expression mediates therapeutic effect of selective CXCR4 antagonist AMD3100 on myocardial infarction. *Circulation* 118: S536-S537, 2008.
6. Marin-Juez R, El-Sammak H, Helker CSM, Kamezaki A, Mullapuli ST, Bibli SI, Foglia MJ, Fleming I, Poss KD and Stainier DYR: Coronary revascularization during heart regeneration is regulated by epicardial and endocardial cues and forms a scaffold for cardiomyocyte repopulation. *Dev Cell* 51: 503-515 e504, 2019.
7. Huang FY, Xia TL, Li JL, Li CM, Zhao ZG, Lei WH, Chen L, Liao YB, Xiao D, Peng Y, *et al*: The bifunctional SDF-1-AnxA5 fusion protein protects cardiac function after myocardial infarction. *J Cell Mol Med* 23: 7673-7684, 2019.
8. Abbott JD, Huang Y, Liu D, Hickey R, Krause DS and Giordano FJ: Stromal cell-derived factor-1alpha plays a critical role in stem cell recruitment to the heart after myocardial infarction but is not sufficient to induce homing in the absence of injury. *Circulation* 110: 3300-3305, 2004.
9. Cheng M, Zhou J, Wu M, Boriboun C, Thorne T, Liu T, Xiang Z, Zeng Q, Tanaka T, Tang YL, *et al*: CXCR4-mediated bone marrow progenitor cell maintenance and mobilization are modulated by c-kit activity. *Circ Res* 107: 1083-1093, 2010.
10. Tang YL, Zhu W, Cheng M, Chen L, Zhang J, Sun T, Kishore R, Phillips MI, Losordo DW and Qin G: Hypoxic preconditioning enhances the benefit of cardiac progenitor cell therapy for treatment of myocardial infarction by inducing CXCR4 expression. *Circ Res* 104: 1209-1216, 2009.
11. Saxena A, Fish JE, White MD, Yu S, Smyth JWP, Shaw RM, DiMaio JM and Srivastava D: Stromal cell-derived factor-1alpha is cardioprotective after myocardial infarction. *Circulation* 117: 2224-2231, 2008.
12. Segers VF, Tokunou T, Higgins LJ, MacGillivray C, Gannon J and Lee RT: Local delivery of protease-resistant stromal cell derived factor-1 for stem cell recruitment after myocardial infarction. *Circulation* 116: 1683-1692, 2007.
13. Hu X, Dai S, Wu WJ, Tan W, Zhu X, Mu J, Guo Y, Bolli R and Rokosh G: Stromal cell derived factor-1 alpha confers protection against myocardial ischemia/reperfusion injury: Role of the cardiac stromal cell derived factor-1 alpha CXCR4 axis. *Circulation* 116: 654-663, 2007.
14. Bromage DI, Taferner S, He Z, Ziff OJ, Yellon DM and Davidson SM: Stromal cell-derived factor-1alpha signals via the endothelium to protect the heart against ischaemia-reperfusion injury. *J Mol Cell Cardiol* 128: 187-197, 2019.
15. Xiong H, Luo Y, Yue Y, Zhang J, Ai S, Li X, Wang X, Zhang YL, Wei Y, Li HH, *et al*: Single-cell transcriptomics reveals chemotaxis-mediated intraorgan crosstalk during cardiogenesis. *Circ Res* 125: 398-410, 2019.
16. Segret A, Rucker-Martin C, Pavoine C, Flavigny J, Deroubaix E, Châtel MA, Lombet A and Renaud JF: Structural localization and expression of CXCL12 and CXCR4 in rat heart and isolated cardiac myocytes. *J Histochem Cytochem* 55: 141-150, 2007.
17. Pyo RT, Sui J, Dhume A, Palomeque J, Blaxall BC, Diaz G, Tunstead J, Logothetis DE, Hajjar RJ and Schecter AD: CXCR4 modulates contractility in adult cardiac myocytes. *J Mol Cell Cardiol* 41: 834-844, 2006.
18. LaRocca TJ, Altman P, Jarrah AA, Gordon R, Wang E, Hadri L, Burke MW, Haddad GE, Hajjar RJ and Tarzami ST: CXCR4 cardiac specific knockout mice develop a progressive cardiomyopathy. *Int J Mol Sci* 20: 2267, 2019.
19. Damas JK, Eiken HG, Oie E, Bjerkeli V, Yndestad A, Ueland T, Tonnessen T, Geiran OR, Aass H, Simonsen S, *et al*: Myocardial expression of CC- and CX-chemokines and their receptors in human end-stage heart failure. *Cardiovasc Res* 47: 778-787, 2000.
20. Wang J, Gareri C and Rockman HA: G-protein-coupled receptors in heart disease. *Circ Res* 123: 716-735, 2018.
21. de Lucia C, Eguchi A and Koch WJ: New insights in cardiac β -adrenergic signaling during heart failure and aging. *Front Pharmacol* 9: 904, 2018.
22. Packer M, Carver JR, Rodeheffer RJ, Ivanhoe RJ, DiBianco R, Zeldis SM, Hendrix GH, Bommer WJ, Elkayam U and Kukin ML: Effect of oral milrinone on mortality in severe chronic heart failure. The PROMISE study research group. *N Engl J Med* 325: 1468-1475, 1991.
23. Shizukuda Y, Buttrick PM, Geenen DL, Borczuk AC, Kitsis RN and Sonnenblick EH: Beta-adrenergic stimulation causes cardiocyte apoptosis: Influence of tachycardia and hypertrophy. *Am J Physiol* 275: H961-H968, 1998.
24. Asai K, Yang GP, Geng YJ, Takagi G, Bishop S, Ishikawa Y, Shannon RP, Wagner TE, Vatner DE, Homcy CJ and Vatner SF: Beta-adrenergic receptor blockade arrests myocyte damage and preserves cardiac function in the transgenic G(salpa) mouse. *J Clin Invest* 104: 551-558, 1999.
25. Zhu WZ, Wang SQ, Chakir K, Yang D, Zhang T, Brown JH, Devic E, Kobilka BK, Cheng H and Xiao RP: Linkage of beta1-adrenergic stimulation to apoptotic heart cell death through protein kinase A-independent activation of Ca2+/calmodulin kinase II. *J Clin Invest* 111: 617-625, 2003.
26. Whelan RS, Konstantinidis K, Xiao RP and Kitsis RN: Cardiomyocyte life-death decisions in response to chronic beta-adrenergic signaling. *Circ Res* 112: 408-410, 2013.
27. Packer M, Bristow MR, Cohn JN, Colucci WS, Fowler MB, Gilbert EM and Shusterman NH: The effect of carvedilol on morbidity and mortality in patients with chronic heart failure. U.S. Carvedilol heart failure study group. *N Engl J Med* 334: 1349-1355, 1996.
28. Zhang X, Szeto C, Gao E, Tang M, Jin J, Fu Q, Makarewicz C, Ai X, Li Y, Tang A, *et al*: Cardiotoxic and cardioprotective features of chronic β -adrenergic signaling. *Circ Res* 112: 498-509, 2013.
29. Singh K, Xiao L, Remondino A, Sawyer DB and Colucci WS: Adrenergic regulation of cardiac myocyte apoptosis. *J Cell Physiol* 189: 257-265, 2001.
30. Adams JW and Brown JH: G-proteins in growth and apoptosis: Lessons from the heart. *Oncogene* 20: 1626-1634, 2001.
31. Steinberg SF: Beta1-adrenergic receptor regulation revisited. *Circ Res* 123: 1199-1201, 2018.
32. Rohrer DK, Desai KH, Jasper JR, Stevens ME, Regula DP Jr, Barsh GS, Bernstein D and Kobilka BK: Targeted disruption of the mouse beta1-adrenergic receptor gene: Developmental and cardiovascular effects. *Proc Natl Acad Sci USA* 93: 7375-7380, 1996.
33. Milano CA, Allen LF, Rockman HA, Dolber PC, McMinn TR, Chien KR, Johnson TD, Bond RA and Lefkowitz RJ: Enhanced myocardial function in transgenic mice overexpressing the beta2-adrenergic receptor. *Science* 264: 582-586, 1994.
34. Engelhardt S, Hein L, Wiesmann F and Lohse MJ: Progressive hypertrophy and heart failure in beta1-adrenergic receptor transgenic mice. *Proc Natl Acad Sci USA* 96: 7059-7064, 1999.
35. Iwase M, Bishop SP, Uechi M, Vatner DE, Shannon RP, Kudej RK, Wight DC, Wagner TE, Ishikawa Y, Homcy CK and Vatner SF: Adverse effects of chronic endogenous sympathetic drive induced by cardiac GS alpha overexpression. *Circ Res* 78: 517-524, 1996.
36. Antos CL, Frey N, Marx SO, Reiken S, Gaburjakova M, Richardson JA, Marks AR and Olson EN: Dilated cardiomyopathy and sudden death resulting from constitutive activation of protein kinase a. *Circ Res* 89: 997-1004, 2001.
37. Nichtova Z, Novotova M, Kralova E and Stankovicova T: Morphological and functional characteristics of models of experimental myocardial injury induced by isoproterenol. *Gen Physiol Biophys* 31: 141-151, 2012.
38. Chang SC, Ren S, Rau CD and Wang JJ: Isoproterenol-induced heart failure mouse model using osmotic pump implantation. *Methods Mol Biol* 1816: 207-220, 2018.
39. Tan Y, Li Y, Xiao J, Shao H, Ding C, Arteel GE, Webster KA, Yan J, Yu H, Cai L and Li X: A novel CXCR4 antagonist derived from human SDF-1beta enhances angiogenesis in ischaemic mice. *Cardiovasc Res* 82: 513-521, 2009.
40. Wu M, Zhou J, Cheng M, Boriboun C, Biyashev D, Wang H, Mackie A, Thorne T, Chou J, Wu Y, *et al*: E2F1 suppresses cardiac neovascularization by down-regulating VEGF and PIGF expression. *Cardiovasc Res* 104: 412-422, 2014.
41. Kuznetsov AV, Javadov S, Sickinger S, Frotschnig S and Grimm M: H9c2 and HL-1 cells demonstrate distinct features of energy metabolism, mitochondrial function and sensitivity to hypoxia-reoxygenation. *Biochim Biophys Acta* 1853: 276-284, 2015.
42. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
43. Cheng M, Yang J, Zhao X, Zhang E, Zeng Q, Yu Y, Yang L, Wu B, Yi G, Mao X, *et al*: Circulating myocardial microRNAs from infarcted hearts are carried in exosomes and mobilise bone marrow progenitor cells. *Nat Commun* 10: 959, 2019.

44. Nie Y, Han YC and Zou YR: CXCR4 is required for the quiescence of primitive hematopoietic cells. *J Exp Med* 205: 777-783, 2008.
45. Wang ER, Jarrah AA, Benard L, Chen J, Schwarzkopf M, Hadri L and Tarzami ST: Deletion of CXCR4 in cardiomyocytes exacerbates cardiac dysfunction following isoproterenol administration. *Gene Ther* 21: 496-506, 2014.
46. Bresnick AR and Backer JM: PI3K β -A versatile transducer for GPCR, RTK, and small GTPase signaling. *Endocrinology* 160: 536-555, 2019.
47. Zhu WZ, Zheng M, Koch WJ, Lefkowitz RJ, Kobilka BK and Xiao RP: Dual modulation of cell survival and cell death by beta(2)-adrenergic signaling in adult mouse cardiac myocytes. *Proc Natl Acad Sci USA* 98: 1607-1612, 2001.
48. Wang XX, Zhang FR, Zhu JH, Xie XD and Chen JZ: Up-regulation of CXC chemokine receptor 4 expression in chronic atrial fibrillation patients with mitral valve disease may be attenuated by renin-angiotensin system blockers. *J Int Med Res* 37: 1145-1151, 2009.
49. Liu X, Zeng Y, Liu Z, Li W, Wang L and Wu M: Bioinformatics analysis of the circRNA-miRNA-mRNA network for atrial fibrillation. *Medicine (Baltimore)* 101: e30221, 2022.
50. Zhang YF, Meng LB, Hao ML, Li XY and Zou T: CXCR4 and TYROBP mediate the development of atrial fibrillation via inflammation. *J Cell Mol Med* 26: 3557-3567, 2022.
51. Zhang J, Huang X, Wang X, Gao Y, Liu L, Li Z, Chen X, Zeng J, Ye Z and Li G: Identification of potential crucial genes in atrial fibrillation: A bioinformatic analysis. *BMC Med Genomics* 13: 104, 2020.
52. Liu P, Sun H, Zhou X, Wang Q, Gao F, Fu Y, Li T, Wang Y, Li Y, Fan B, *et al*: CXCL12/CXCR4 axis as a key mediator in atrial fibrillation via bioinformatics analysis and functional identification. *Cell Death Dis* 12: 813, 2021.
53. Agarwal U, Ghalayini W, Dong F, Weber K, Zou YR, Rabbany SY, Rafii S and Penn MS: Role of cardiac myocyte CXCR4 expression in development and left ventricular remodeling after acute myocardial infarction. *Circ Res* 107: 667-676, 2010.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.