

***Radix clematidis* extract inhibits UVB-induced MMP expression by suppressing the NF- κ B pathway in human dermal fibroblasts**

YOUNG-RAE LEE^{1*}, EUN-MI NOH^{1*}, KANG-BEOM KWON², YUN-SIK LEE³, JONG PHIL CHU³,
EUN JIB KIM⁴, YOUNG-SEOK PARK⁵, BYEONG-SOO KIM⁵ and JONG-SUK KIM¹

¹Department of Biochemistry, Institute for Medical Sciences, Chonbuk National University Medical School, Jeonju, Jeonbuk 560-182; ²Department of Physiology, School of Oriental Medicine, Wonkwang University, Iksan, Jeonbuk, 570-749; ³Department of Medical Zoology, College of Medicine, Kyunghee University, Dongdaemoon-gu, Seoul 130-701; ⁴Cheonan Yonam College, Cheonan, Chungnam, 330-709; ⁵Department of Companion and Laboratory Animal Science, Kongju National University, Yesan 340-702, South Korea

Received December 22, 2008; Accepted February 10, 2009

DOI: 10.3892/ijmm_00000180

Abstract. Ultraviolet (UV)B irradiation induces the production of matrix metalloproteinases (MMPs), which are responsible for the degradation of collagenous extracellular matrix in connective tissues, causing skin photoaging. Although *Radix clematidis* is commonly used in Chinese medicine for the treatment of arthralgia, the anti-skin photoaging effects of *Radix clematidis* have not yet been reported. In the present study, we investigated the inhibitory effects of *Radix clematidis* extract (RCE) on MMP-1 and -3 expression of human dermal fibroblast cells via various *in vitro* experiments and elucidated the pathways of inhibition. Western blot analysis and real-time PCR revealed RCE inhibited UVB-induced MMP-1 and -3 expressions in a dose-dependent manner. UVB strongly activated nuclear factor- κ B (NF- κ B) activity, which was determined by I κ B α degradation, nuclear localization of p50 and p65 subunit, and NF- κ B binding activity. However, UVB-induced NF- κ B activation was completely blocked by RCE

pretreatment. These findings suggest that RCE prevents UVB-induced MMP expression through inhibition of NF- κ B activation. In conclusion, RCE is a potential agent for the prevention and treatment of skin photoaging.

Introduction

Skin aging is divided into intrinsic or chronologic aging, which is the process of senescence that affects all body organs, and extrinsic aging (photoaging), which occurs as a consequence of exposure to environmental factors. One of the most important extrinsic aging factors is sunlight, particularly exposure to ultraviolet (UV)B irradiation, which causes skin photoaging. The skin photoaging process increases skin fragility, laxity, roughness, dryness, pigmentation, blister formation, leathery appearance and formation of wrinkles (1). Chronic exposure of human skin to UVB radiation results in photoaging and induces the production of matrix metalloproteinases (MMPs) (2).

MMPs are responsible for the degradation or synthesis inhibition of collagenous extracellular matrix in connective tissues (3). Collagen represents the main component of the extracellular matrix of dermal connective tissue, and its concentration decreases in chrono- and photoaging. MMP-1 preferentially degrades fibrillar collagens, which maintain the tensile strength of fetal membranes, whereas MMP-3 degrades an extremely broad array of extracellular matrix substrates and activates the secreted, zymogenic form of other MMPs (4-6). UVB is known to induce the expression of interstitial collagenase (MMP-1) and stromelysin-1 (MMP-3) in human dermal fibroblasts (HDF) and normal human epidermis *in vivo* (7-9). Among those MMPs, MMP-1 is the most important MMP in the degradation of the extracellular matrix by skin photoaging (10).

Radix clematidis is the name given to the root of *Clematis florida* T_{HUNB}, a member of the *Ranunculaceae* family. *Radix clematidis* is commonly used in traditional Chinese medicine to alleviate joint pain and relieve swelling. In recent studies,

Correspondence to: Dr Byeong-Soo Kim, Department of Companion and Laboratory Animal Science, Kongju National University, Yesan 340-702, South Korea
E-mail: bskim@kongju.ac.kr

Dr Jong-Suk Kim, Department of Biochemistry, Institute for Medical Sciences, Chonbuk National University Medical School, Jeonju, Jeonbuk 560-182, South Korea
E-mail: jsukim@chonbuk.ac.kr

*Contributed equally

Key words: *Radix clematidis*, matrix metalloproteinases, ultraviolet B, NF- κ B, photoaging

Radix clematidis was reported to exert inhibitory effects against β -cell damage by suppressing the NF- κ B pathway and depigmentation of melanocytes (11,12).

In the present study, we evaluated the preventive effects of *Radix clematidis* extract (RCE) on UVB-induced MMPs expression in HDF. RCE was found to block the UVB-induced NF- κ B pathway, thereby inhibiting the MMPs expression. These results indicate that *Radix clematidis* is useful as an anti-skin photoaging agent.

Materials and methods

Materials. Primary antibodies for MMP-1 and -3 were obtained from R&D Systems (Minneapolis, MN, USA). High glucose-containing Dulbecco's modified Eagle's medium (DMEM), fetal bovine serum (FBS) and phosphate-buffered saline (PBS) were obtained from Gibco-BRL (Gaithersburg, MD, USA). 3-(4,5-dimethyl-thiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), dimethyl sulfoxide (DMSO) and anti- β -actin were obtained from Sigma. Antibodies of p50, p65, and I κ B α , PCNA, and horseradish peroxidase (HRP)-conjugated IgG were from Santa Cruz Biochemicals (Santa Cruz, CA, USA).

Preparation of *Radix clematidis* extract. *Radix clematidis* was obtained from Wonkwang Oriental Medical Hospital in Iksan, Jeonbuk, Korea and its identity was confirmed by Ho-Joon Song, keeper of the Herbarium. Voucher samples were preserved for reference in the Herbarium of the Department of Physiology, School of Oriental Medicine, Wonkwang University (Omphy, 2005-58). *Radix clematidis* (200 g) were then ground and extracted in boiling water for 4 h. The extract was centrifuged at 3,000 x g for 20 min, after which the supernatant was concentrated to 200 ml under reduced pressure and then freeze dried to 20.4 g. The sterile extract was then stored at -70°C until use.

Isolation and culture of HDF. HDF was aseptically isolated from a circumcised neonatal foreskin. The epidermis and dermis were separated by incubation in 0.9 units/ml dispase in medium for 16 h at 4°C. After the epidermis and dermis were mechanically separated, the dermis was minced and attached on the surface of a tissue culture flask and fed with DMEM containing 10% FBS for 1-2 weeks. The dermal fibroblasts spread as radial outgrowth from attached pieces of dermis, cultured in DMEM with 10% FBS.

UV irradiation. For UVB irradiation, we used UVB cross-linker (6x8 W, 312 nm, Model CL-508M, Vilber Lourmat, Paris, France). In brief, serum-starved confluent cells were rinsed twice with PBS, and all irradiations were performed under a thin layer of PBS. Immediately after irradiation, fresh serum-free medium was added to the cells. Responses were measured after an incubation period of 24 h. Mock-irradiated controls followed the same schedule of medium changes without UVB irradiation.

Determination of cell viability. The effect of RCE on cell viability of HDF was determined using MTT assay. Briefly, 2x10⁴ cells/well were treated with various concentrations of RCE. After incubation for 24 h, cells were washed twice with

PBS, MTT (0.5 mg/ml PBS) was added to each well and incubated at 37°C for 30 min. The formazan crystals formed were then dissolved by adding DMSO (100 μ l/well) at 570 nm using a microplate reader (Model 3550, Bio-Rad, Richmond, CA, USA).

Western blot analysis. HDF (2x10⁶ cells) was irradiated with UVB (25 mJ/cm²) and then treated with RCE for 24 h. Cells were lysed with 40 μ l of ice-cold M-PER[®] Mammalian Protein Extraction Reagent (Pierce Biotechnology, Rockford, IL). In lysates, the protein concentration was determined using the Bradford method (13). Samples were separated by SDS-PAGE with 10% acrylamide running and 3% acrylamide stacking gels, and then transferred to Hybond[™]-PVDF membranes using a Western blot apparatus. The PVDF membranes were blotted with 1 μ g/ml of primary antibodies for MMP-1, -3, p50, p65, PCNA and β -actin. HRP-conjugated IgG was used as a secondary antibody. The protein expression levels were then determined by analyzing the signals captured on the PVDF membranes using an image analyzer (Las-1000, Fujifilm, Japan).

Quantitative real-time PCR assay. Total RNA was extracted from the cells using FastPure[™] RNA Kit (Takara, Shiga, Japan). The concentration and purity of RNA were determined by absorbance at 260/280 nm. cDNA was synthesized from 1 μ g total RNA using PrimeScript[™] RT reagent Kit (Takara). The expression of MMP-1 and -3 mRNA was determined by real-time PCR using the ABI PRISM 7900 sequence detection system and the SYBR[®] Green (Applied Biosystems, Foster City, CA, USA). The primers were, MMP-1 (NM 002424.2) sense, AGTGACTGGGAAACCAGATGCTGA; antisense, GCTCTTGGCAAATCTGGCCTGTAA and MMP-3 (NM 002422) sense, ATTCCATGGAGCCAGGCTTTC; antisense, CATTTGGGTCAAACCTCCAACCTGTG and GAPDH (NM 002046) sense, ATGGAAATCCCATCACCA TCTT; antisense, CGCCCCACTTGATTTTGG. To control variation in mRNA concentration, all results were normalized to the housekeeping gene, GAPDH. Relative quantitation was performed using the comparative $\Delta\Delta C_t$ method according to the manufacturer's instructions.

Determination of MMP-1 and -3 secretions by ELISA. HDF was seeded in 100 mm culture dishes at a density of 2x10⁶ cells per dish, and then irradiated with UVB (25 mJ/cm²) and then treated with RCE. Following 24 h of incubation, the culture supernatant was collected and centrifuged at 10,000 x g for 5 min to remove the particulate matter, and stored at -80°C in fresh tubes. The active MMP-1 in culture supernatants was quantified by fluorescent assay, using the Fluorokine E Human Active MMP-1 Fluorescent Assay Kit (R&D Systems) and MMP-3 in the cell culture supernatants were then determined using Quantikine ELISA kits (R&D Systems), according to the manufacturer's protocol.

Preparation of nuclear extract. HDF (2x10⁶ cells) was irradiated with UVB (25 mJ/cm²) and then treated with RCE for 24 h. Cells were immediately washed twice, scraped into 1.5 ml of ice-cold PBS (pH 7.4), and then pelleted at 12,000 x g for 30 sec. Cytoplasmic and nuclear extracts were




Figure 1. Effect of RCE on cell viability in HDF. Cells were cultured in 96-well plates until 70% confluence, then incubated with the indicated concentration of RCE for 24 h. MTT assay was used to detect the viability of cells as detailed in Materials and methods. The optical density value of control was regarded as 100%. Data points are the mean \pm SE of three independent experiments.

prepared from cells using the NE-PER[®] Nuclear and Cytoplasmic Extraction Reagents (Pierce Biotechnology).

Electrophoretic mobility shift assay (EMSA). The activation of NF- κ B was assayed by a gel mobility shift assay using nuclear extracts. An oligonucleotide containing the κ -chain binding site (κ B, 5'-CCGGTTAACAGAGGGGGCTTTCCGAG-3') was synthesized and used as a probe for the gel retardation assay. The two complementary strands were then annealed and labeled with [α -³²P]dCTP. Labeled oligonucleotides (10,000 cpm), 10 μ g of nuclear extracts, and binding buffer (10 mM Tris-HCl, pH 7.6, 500 mM KCl, 10 mM EDTA, 50% glycerol, 100 ng poly (dI:dC), 1 mM DTT) were then incubated for 30 min at room temperature in a final volume of 20 μ l. The reaction mixtures were analyzed by electrophoresis on 4% polyacrylamide gels in 0.5 X Tris-borate buffer, and the gels were then dried and examined by autoradiography.




Figure 2. Effect of RCE on UVB-induced MMP-1 and -3 expressions in HDF. HDF in monolayer was incubated with stimulated UVB (25 mJ/cm²) in a time-dependent manner (A). To investigate the effect of RCE, cells were stimulated with UVB (25 mJ/cm²) and the indicated concentrations of RCE for 24 h (B). The cell lysates were analyzed by Western blotting with anti-MMP-1 and -3. The blot was reprobed with anti- β -actin to confirm equal loading. HDF was pretreated with cordycepin for 1 h and stimulated by IL-1 β for 24 h. Total cellular RNA was analyzed by real-time PCR for MMP-1 and -3 (C). Each value represents the mean \pm SEM of three independent experiments. * p <0.01 vs. untreated control; # p <0.01 vs. UVB.




Figure 3. Inhibition of UVB-induced MMP-1 and -3 secretions from HDF by RCE. Cells were stimulated with UVB (25 mJ/cm²) and the indicated concentrations of RCE for 24 h. Level of MMP-1 and -3 in culture media was measured using a commercially available ELISA kit as described in Materials and methods. Each value represents the mean \pm SEM of three independent experiments. **p*<0.01 vs. untreated control; #*p*<0.01 vs. UVB.

Specific binding was controlled by competition with a 50-fold excess of cold κ B oligonucleotide.

Statistical analysis. Statistical analysis of the data was performed using ANOVA and Duncan's test. Differences of *p*<0.05 were considered statistically significant.

Results

Effect of RCE on cell viability of HDF. Cytotoxicity of RCE on HDF has not yet been reported. Therefore, we designed to avoid interference of cytotoxicity from reagent. In this study, the effect of RCE on cellular toxicity of HDF was analyzed using the MTT assay. MTT assays were performed to determine optimal non-toxic concentrations at 24 h after incubation with RCE and no change in morphology. As shown in Fig. 1, treatment of HDF with indicated concentrations of RCE for 24 h resulted in no significant effect in cell viability.

Effect of RCE on UVB-induced MMP-1 and -3 expressions in HDF. To investigate UVB-induced MMP-1 and -3 expression, we performed Western blot analysis and real-time PCR in HDF. Western blot analysis revealed that the irradiation of HDF with UVB (25 mJ/cm²) increased the levels of MMP-1 and -3 in a time-dependent manner (Fig. 2A). Treatment with RCE (50 and 100 μ g/ml) completely blocked the up-regulation of MMP-1 and -3 induced by UVB irradiation (Fig. 2B). Real-time PCR revealed that UVB irradiation increased the level of MMP-1 and -3 in HDF and RCE blocked UVB-induced up-regulation of MMP-1 and -3 in a dose-dependent manner (Fig. 2C). As shown, RCE itself had no effect on either MMP-1 or -3 in HDF. These results indicate




Figure 4. Suppression of IL-1 β -induced DNA binding of NF- κ B, translocation of p65 and p50 to the nucleus, and I κ B α degradation by RCE. Cells were stimulated with UVB (25 mJ/cm²) and the indicated concentrations of RCE. Following 3 h of incubation, DNA binding of NF- κ B was analyzed by electrophoretic mobility shift analysis (A), and the translocation of p65 and p50 to the nucleus and I κ B α degradation in the cytoplasm (B) were determined by Western blotting.

that RCE is a potent inhibitor of UVB-induced MMP-1 and -3 expression in HDF.

Effect of RCE on UVB-induced secretion of MMP-1 and -3. MMP secretion, which is the hallmark of skin aging, is activated by UVB, and 23 members of the MMP family have been identified in humans to date. Of these, MMP-1 and -3 are particularly important to skin aging because they are responsible for the degradation of collagenous extracellular matrix, and because levels of MMP-1 and -3 are significantly higher in aged skin. We used ELISA to investigate the effect of RCE on UVB-induced MMP secretion. Irradiation of HDF with UVB (25 mJ/cm²) resulted in an increase in the secretion of MMP-1 and -3, respectively (Fig. 3). However, RCE (50 or 100 μ g/ml) significantly diminished the UVB-induced MMP-1 and -3 secretions, respectively (Fig. 3).

Suppressive effect of RCE on UVB-induced NF- κ B activation. NF- κ B was implicated in the transcriptional regulation of UVB-induced MMP expression (14). Therefore, we studied the effect of RCE on UVB-stimulated translocation of NF- κ B from the cytoplasmic compartment to the nucleus and on DNA binding in HDF. UVB-irradiated HDF showed increased binding activity of an NF- κ B consensus sequence (Fig. 4A), as well as increased p65 and p50 subunit levels in their nuclei (Fig. 4B) when compared to unstimulated cells. Additionally, UVB-induced NF- κ B activation was significantly suppressed by pretreatment with RCE, which suggests that RCE inhibits MMP-1 and -3 expression through the inhibition of NF- κ B activation. The specificity of the DNA-protein interactions for NF- κ B was demonstrated by performing competition assays using a 50-fold excess of unlabeled oligonucleotide (Fig. 4A, lane 7).

From above, we found that RCE blocked UVB-induced NF- κ B activation. Under basal condition, the cytoplasmic protein I κ B directly binds to p65 and p50 subunits and represses their nuclear translocation (15-17). Therefore, we determined the alteration of I κ B α levels of the cytoplasmic fraction in this study. UVB-irradiated HDF showed a decreased level of I κ B α protein in the cytoplasm when compared to a similar fraction in unstimulated cells, however, the increased I κ B α degradation as a result of UVB stimulation was significantly suppressed by treatment with RCE (Fig. 4B).

Discussion

In this study, we demonstrated that RCE inhibits UVB-induced expression of MMP-1 and -3 in cultured HDF. Our results also showed that RCE blocked UVB-induced activation of NF- κ B, which has an important role in MMP-1 and -3 expressions.

Skin aging is attributed to extrinsic (photoaging) and intrinsic (chronological) aging. Photoaging concerns premature skin aging caused by repeated sun exposure (18-20). UV irradiation of cultured HDF *in vitro* or human skin *in vivo* induces the expression of MMPs which play important roles in the degradation of extracellular matrix components during skin aging (7,21,22). Varani *et al* reported that with increasing age, MMP levels rise and collagen synthesis declines for sun-protected human skin *in vivo* (23). Recently, it was suggested that excessive matrix degradation by UV-induced MMPs secreted by various cells (e.g., keratinocytes, fibroblasts, and inflammatory cells) contributes substantially to the connective tissue damage that occurs during photoaging (20,24,25). The following mechanism of photoaging was proposed. Initially, AP-1 and NF- κ B are activated by UV light, and AP-1 and NF- κ B driven MMPs such as MMP-1 and -3 are induced (24-26). These MMPs then degrade collagen, which results in a collagen deficiency in photodamaged skin and eventually causes skin wrinkling (27).

In this study, we observed an increase in the activation of NF- κ B in the UVB-irradiated HDF, and the ability of RCE to protect against the UVB-induced MMPs expressions, suggesting that RCE inhibits UVB-induced expression of MMPs by suppressing the NF- κ B pathway in HDF. It is well established that a nuclear transcription factor, NF- κ B, is activated upon UV irradiation (10,28,29). In previous studies, it was also reported that UVB-mediated skin photoaging is

prevented by suppression of NF- κ B activation (15,30,31). Therefore, inhibition of the NF- κ B activation pathway is important to avoid UVB-mediated skin damage. In fact, NF- κ B is known to increase MMP-1 in the dermis (14,32,33). NF- κ B is a crucial factor for the immunoinflammatory responses and is also implicated in various skin diseases including allergic dermatitis, psoriasis vulgaris, and skin cancer (34). Hence, although NF- κ B is involved in maintaining the skin homeostasis (35,36), excessive activation is pathogenic. NF- κ B is an inducible dimeric transcription factor that belongs to the Rel/NF- κ B family of transcription factors. NF- κ B consists of two major polypeptides, p65 and p50 (37). NF- κ B is initially located in the cytoplasm in an inactive form complexed with I κ B, an inhibitory factor of NF- κ B. Various inducers such as IL-1, TNF- α and UV cause dissociation of this complex, presumably by phosphorylation of I κ B, resulting in NF- κ B being released from the complex. NF- κ B then translocates to the nucleus, where it interacts with specific DNA recognition sites to mediate gene transcription

In conclusion, the development of MMP inhibitors is considered a promising strategy for skin cancer therapy and photoaging. In recent years, the development of compounds with MMP inhibition activities from natural plants received a great deal of attention. This study demonstrates the inhibitory effect of RCE on the MMP expression via an mRNA assay. Also, our results demonstrate that RCE is a potent inhibitor of UVB-induced MMP expression and strongly blocks the ability of the NF- κ B signaling pathway in HDF. The dose of RCE that inhibits UVB-induced MMP expression are below the toxicity limit. Therefore, we suggest that RCE should be viewed as a potential therapeutic candidate for preventing and treating skin photoaging.

Acknowledgements

This Study was supported by the Technology Development Program for Agriculture and Forestry, Ministry for Agriculture, Forestry and Fisheries, Republic of Korea.

References

- Jenkins G: Molecular mechanisms of skin ageing. *Mech Ageing Dev* 123: 801-810, 2002.
- Ho JN, Lee YH, Park JS, Jun WJ, Kim HK, Hong BS, Shin DH and Cho HY: Protective effects of aucubin isolated from *eucommia ulmoides* against UVB-induced oxidative stress in human skin fibroblasts. *Biol Pharm Bull* 28: 1244-1248, 2005.
- Scharffetter-Kochanek K, Brenneisen P, Wenk J, Herrmann G, Ma W, Kuhr L, Meewes C and Wlaschek M: Photoaging of the skin from phenotype to mechanisms. *Exp Gerontol* 35: 307-316, 2000.
- Ogata Y, Enghild JJ and Nagase H: Matrix metalloproteinase 3 (stromelysin) activates the precursor for the human matrix metalloproteinase 9. *J Biol Chem* 267: 3581-3584, 1992.
- Knauper V, Lopez-Otin C, Smith B, Knight G and Murphy G: Biochemical characterization of human collagenase-3. *J Biol Chem* 271: 1544-1550, 1996.
- Jackson C, Nguyen M, Arkell J and Sambrook P: Selective matrix metalloproteinase (MMP) inhibition in rheumatoid arthritis-targeting gelatinase A activation. *Inflamm Res* 50: 183-186, 2001.
- Brenneisen P, Sies H and Scharffetter-Kochanek K: Ultraviolet-B irradiation and matrix metalloproteinases: From induction via signaling to initial events. *Ann N Y Acad Sci* 973: 31-43, 2002.
- Brenneisen P, Oh J, Wlaschek M, Wenk J, Briviba K, Hommel C, Herrmann G, Sies H and Scharffetter-Kochanek K: Ultraviolet B wavelength dependence for the regulation of two major matrix-metalloproteinases and their inhibitor timp-1 in human dermal fibroblasts. *Photochem Photobiol* 64: 877-885, 1996.

9. Brenneisen P, Wenk J, Wlaschek M, Krieg T and Scharffetter-Kochanek K: Activation of p70 ribosomal protein s6 kinase is an essential step in the DNA damage-dependent signaling pathway responsible for the ultraviolet B-mediated increase in interstitial collagenase (mmp-1) and stromelysin-1 (MMP-3) protein levels in human dermal fibroblasts. *J Biol Chem* 275: 4336-4344, 2000.
10. Wenk J, Schuller J, Hinrichs C, Syrovets T, Azoitei N, Podda M, Wlaschek M, Brenneisen P, Schneider LA, Sabiwalsky A, Peters T, Sulyok S, Dissemond J, Schauen M, Krieg T, Wirth T, Simmet T and Scharffetter-Kochanek K: Overexpression of phospholipid-hydroperoxide glutathione peroxidase in human dermal fibroblasts abrogates UVA irradiation-induced expression of interstitial collagenase/matrix metalloproteinase-1 by suppression of phosphatidylcholine hydroperoxide-mediated NF-kappaB activation and interleukin-6 release. *J Biol Chem* 279: 45634-45642, 2004.
11. Kim EK, Song MY, Hwang TO, Kim HJ, Moon WS, Ryu DG, So HS, Park R, Park JW, Kwon KB and Park BH: *Radix clematidis* extract protects against cytokine- and streptozotocin-induced β -cell damage by suppressing the NF- κ B pathway. *Int J Mol Med* 22: 349-356, 2008.
12. Zhong S, Wu Y, Soo-Mi A, Zhao J, Wang K, Yang S, Jae-Ho Y and Zhu X: Depigmentation of melanocytes by the treatment of extracts from traditional chinese herbs: A cell culture assay. *Biol Pharm Bull* 29: 1947-1951, 2006.
13. Bradford MM: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248-254, 1976.
14. Bond M, Baker AH and Newby AC: Nuclear factor kappaB activity is essential for matrix metalloproteinase-1 and -3 upregulation in rabbit dermal fibroblasts. *Biochem Biophys Res Commun* 264: 561-567, 1999.
15. Adhami VM, Afaq F and Ahmad N: Suppression of ultraviolet B exposure-mediated activation of NF-kappaB in normal human keratinocytes by resveratrol. *Neoplasia* 5: 74-82, 2003.
16. Kim EK, Kwon KB, Koo BS, Han MJ, Song MY, Song EK, Han MK, Park JW, Ryu DG and Park BH: Activation of peroxisome proliferator-activated receptor-gamma protects pancreatic beta-cells from cytokine-induced cytotoxicity via NF-kappaB pathway. *Int J Biochem Cell Biol* 39: 1260-1275, 2007.
17. Afaq F, Ahmad N and Mukhtar H: Suppression of UVB-induced phosphorylation of mitogen-activated protein kinases and nuclear factor kappa B by green tea polyphenol in skh-1 hairless mice. *Oncogene* 22: 9254-9264, 2003.
18. Uitto J: Connective tissue biochemistry of the aging dermis. Age-related alterations in collagen and elastin. *Dermatol Clin* 4: 433-446, 1986.
19. Gilchrist BA: A review of skin ageing and its medical therapy. *Br J Dermatol* 135: 867-875, 1996.
20. Chung JH, Seo JY, Lee MK, Eun HC, Lee JH, Kang S, Fisher GJ and Voorhees JJ: Ultraviolet modulation of human macrophage metalloelastase in human skin in vivo. *J Invest Dermatol* 119: 507-512, 2002.
21. Rittie L and Fisher GJ: UV-light-induced signal cascades and skin aging. *Ageing Res Rev* 1: 705-720, 2002.
22. Chung JH, Hanft VN and Kang S: Aging and photoaging. *J Am Acad Dermatol* 49: 690-697, 2003.
23. Varani J, Perone P, Fligiel SE, Fisher GJ and Voorhees JJ: Inhibition of type I procollagen production in photodamage: Correlation between presence of high molecular weight collagen fragments and reduced procollagen synthesis. *J Invest Dermatol* 119: 122-129, 2002.
24. Fisher GJ, Datta SC, Talwar HS, Wang ZQ, Varani J, Kang S and Voorhees JJ: Molecular basis of sun-induced premature skin ageing and retinoid antagonism. *Nature* 379: 335-339, 1996.
25. Chung JH, Seo JY, Choi HR, Lee MK, Youn CS, Rhie G, Cho KH, Kim KH, Park KC and Eun HC: Modulation of skin collagen metabolism in aged and photoaged human skin in vivo. *J Invest Dermatol* 117: 1218-1224, 2001.
26. Fisher GJ, Wang ZQ, Datta SC, Varani J, Kang S and Voorhees JJ: Pathophysiology of premature skin aging induced by ultraviolet light. *N Engl J Med* 337: 1419-1428, 1997.
27. Gilchrist BA and Yaar M: Ageing and photoageing of the skin: Observations at the cellular and molecular level. *Br J Dermatol* 127 (suppl 41): 25-30, 1992.
28. Saliou C, Kitazawa M, McLaughlin L, Yang JP, Lodge JK, Tetsuka T, Iwasaki K, Cillard J, Okamoto T and Packer L: Antioxidants modulate acute solar ultraviolet radiation-induced NF-kappaB activation in a human keratinocyte cell line. *Free Radic Biol Med* 26: 174-183, 1999.
29. Kim J, Hwang JS, Cho YK, Han Y, Jeon YJ and Yang KH: Protective effects of (-)-epigallocatechin-3-gallate on UVA- and UVB-induced skin damage. *Skin Pharmacol Appl Skin Physiol* 14: 11-19, 2001.
30. Tanaka K, Hasegawa J, Asamitsu K and Okamoto T: Prevention of the ultraviolet B-mediated skin photoaging by a nuclear factor kappaB inhibitor, parthenolide. *J Pharmacol Exp Ther* 315: 624-30, 2005.
31. Tanaka K, Hasegawa J, Asamitsu K and Okamoto T: *Magnolia obovata* extract and its active component magnolol prevent skin photoaging via inhibition of nuclear factor kappaB. *Eur J Pharmacol* 565: 212-219, 2007.
32. Sun HB, Malacinski GM and Yokota H: Promoter competition assay for analyzing gene regulation in joint tissue engineering. *Front Biosci* 7: A169-A174, 2002.
33. Chung JH: Photoaging in Asians. *Photodermatol Photoimmunol Photomed* 19: 109-121, 2003.
34. Bell S, Degitz K, Quirling M, Jilg N, Page S and Brand K: Involvement of NF-kappaB signalling in skin physiology and disease. *Cell Signal* 15: 1-7, 2003.
35. Pasparakis M, Courtois G, Hafner M, Schmidt-Supprian M, Nenci A, Toksoy A, Krampert M, Goebeler M, Gillitzer R, Israel A, Krieg T, Rajewsky K and Haase I: TNF-mediated inflammatory skin disease in mice with epidermis-specific deletion of ikk2. *Nature* 417: 861-866, 2002.
36. Takao J, Yudate T, Das A, Shikano S, Bonkobara M, Ariizumi K and Cruz PD: Expression of NF-kappaB in epidermis and the relationship between NF-kappaB activation and inhibition of keratinocyte growth. *Br J Dermatol* 148: 680-688, 2003.
37. Thanos D, Maniatis T: NF-kappaB: A lesson in family values. *Cell* 80: 529-532, 1995.