

Identification of sex hormone binding globulin-interacting proteins in the brain using phage display screening

M. GNANASEKAR¹, F.G. SULEMAN², K. RAMASWAMY¹ and J.D. CALDWELL³

¹Department of Biomedical Sciences, University of Illinois College of Medicine, 1601 Parkview Avenue, Rockford, IL 61107; ²Thermo Fisher Scientific, 3747 N. Meridian Road, Rockford, IL 61105; ³Department of Pharmacology, Lake Erie College of Osteopathic Medicine, 1858 West Grandview Boulevard, Erie, PA 16509, USA

Received April 16, 2009; Accepted June 2, 2009

DOI: 10.3892/ijmm_00000248

Abstract. The present study reports the identification of human sex hormone binding globulin (SHBG)-interacting proteins in the brain using a phage display-based screening technology. Phage display is a system in which a foreign protein is displayed on the surface of a bacteriophage as a fusion protein with one of the coat proteins of the bacteriophage. T7 phage clones expressing normal human brain proteins (human normal brain phage-display cDNA expression library) were screened using SHBG as bait. The bound phage clones were then identified by DNA sequencing and by BLAST search analysis. Of the twenty binding proteins analyzed, three were found to be membrane-associated proteins: synaptosomal associated protein 25 (SNAP25), Thy-1 cell surface antigen and zonadhesin. Further studies will determine if the interactions of SHBG with these proteins have any role in the internalization and cell signaling events or whether they contribute to steroid delivery to specific cells.

Introduction

Examining the roles of steroid binding globulins as active participants in steroid actions has received attention recently. One reason is that there are membrane-associated receptors for binding globulins. Hryb *et al* (1) showed that sex hormone binding globulin (SHBG) binds with high affinity to a specific membrane receptor in prostate stromal and epithelial cells.

Nakhla *et al* (2,3) demonstrated that SHBG binds to a receptor or receptor complex which then results in increases of intracellular cAMP. Studies by Rosner and colleagues (3,4) suggest that the putative SHBG receptor is a G protein-coupled receptor or functionally linked to one. In these studies (1) a 167-kDa protein was solubilized from membranes of human prostate glands and its binding activity to SHBG was demonstrated using ¹²⁵I-SHBG. Other studies (5,6) also suggested that there are membrane-associated receptors for SHBG in the uterus. These membrane-associated receptors may be important for internalization of SHBG, which was observed in many tissues (7-14) including the brain (15). The Nykjaer and Willnow laboratory demonstrated that a membrane-associated protein called megalin, a 600-kDa member of the LDL receptor gene family, is capable of binding to and facilitating the internalization of both SHBG and its bound steroid (16). It is the protein that the same laboratory previously suggested was involved in the uptake of other steroids and demonstrated that it belonged to a larger class of proteins responsible for the uptake of a broad range of lipophilic molecules (17). Megalin is important in steroid effects. This group demonstrated that without megalin, mice showed a delayed onset of puberty (16). The same laboratory demonstrated the importance of binding globulins in mediating the effects of steroids. Mice lacking a corticosteroid-binding globulin, another steroid-binding globulin found in the brain (18,19), had more free steroids in the blood but showed fewer physiological responses to them, suggesting that the presence of this binding globulin was necessary in order to see the effects of corticosteroids (20).

The first indication that there were SHBG receptors in the brain was when SHBG facilitated female sexual receptivity (21) except when coupled to dihydrotestosterone (22). Our laboratory and others also found that SHBG is produced in the brain, where it is often co-localized with the neuropeptides oxytocin and vasopressin (23-25). Since both SHBG and oxytocin are often found together in synaptic vesicles in axonal varicosities, we concluded that SHBG is likely released into the brain. This suggests the existence of receptors for this released SHBG. We also presented indirect evidence of such SHBG receptors in the hypothalamus (26). After binding to membrane-associated receptors, it is possible that SHBG, its associated steroid, and the SHBG receptor are all internalized into brain cells. Caldwell *et al* (15) demonstrated uptake of SHBG *in vivo*

Correspondence to: Dr Munirathinam Gnanasekar, Department of Biomedical Sciences, University of Illinois College of Medicine, 1601 Parkview Avenue, Rockford, IL 61107, USA
E-mail: mgnanas@uic.edu

Dr Jack D. Caldwell, Department of Pharmacology, Lake Erie College of Osteopathic Medicine, 1858 West Grandview Boulevard, Erie, PA 16509, USA
E-mail: jcaldwell@lecom.edu

Key words: sex hormone binding globulin, brain, phage display, zonadhesin, synaptosomal associated protein 25




Figure 1. Schematic strategy to identify targets of SHBG.

in the choroids plexus, paraventricular and supraoptic nuclei, and in specific periventricular cells. Despite observations that suggest the presence of both central and peripheral SHBG membrane receptors, the identity of brain SHBG receptors has not been elucidated. Another study utilized a yeast two-hybrid system to identify candidate proteins for SHBG binding in prostate (27). In this study, phage display of a human brain cDNA library is used to identify SHBG-interacting proteins.

Materials and methods

Materials. The normal human brain T7Select™ phage library, host strain BLT5403, as well as primers were obtained from Novagen, USA. Qiaquick PCR purification kits were from Qiagen (Valencia, CA). Jumpstart TAQ polymerase was from Sigma (St. Louis, MO).

Biopanning. The strategy to identify the SHBG interacting proteins is presented in Fig. 1. Briefly, biopanning was performed as follows. SHBG was diluted in deionized water to 10 µg/ml; 100 µl aliquots were coated in the wells of a 96-well microtiter plate (Costar, Corning, NY) and incubated at room temperature (RT) for 3 h. The SHBG-coated wells were then washed with Tris-buffered saline (TBS), blocked by adding 200 µl 3% BSA to each well and incubated at RT for 1 h. Excess blocking reagent was removed by washing with deionized water. The T7Select phage of the human normal brain cDNA library (Novagen, USA) was added to wells and incubated at RT for 1 h. Unbound phages were washed off with TBS containing Tween while bound phages were then eluted by incubating each well with 200 µl of 1% SDS in TBS. The eluted phages were then used to infect *E. coli* BLT5403 which amplifies the phages for the next biopanning cycle. Three more biopanning rounds were done to increase the phage binding affinity to the SHBG. Finally, the phages obtained after this fourth amplification were plaque titered with a series of dilutions (10^{-2} to 10^{-14}) on LB agar plates containing 1% agarose and incubated for 2–4 h at 37°C. Single plaques of phages were randomly selected with a pipette tip, resuspended in 100 µl of 0.1X TE buffer, boiled for 10 min and placed on ice for 5 min. To amplify the phage DNA, 1 µl




Figure 2. Size distribution of phage clones obtained from first round of biopanning. Phage bound to SHBG from the first cycle of biopanning were eluted and amplified on BLT5403. The amplified phages were plated with BLT5403 on LB agar plates for plaque formation. Ten plaques were randomly selected for PCR amplification. Amplified PCR products (5 µl) were loaded on 1% agarose gel containing Etbr and electrophoresed for 30 min. The resolved gel was observed on a UV transilluminator and the image was captured using Kodak digital software. Lane 1, control PCR that lacks a template; lanes 2–11 random phage clones.

of the boiled samples were used as DNA templates for PCR amplification.

PCR and DNA sequencing. DNAs of the selected phage clones were amplified by PCR according to the manufacturer's (Novagen) instructions on a Perkin Elmer Thermocycler using the T7Select up 5'-GGAGCTGTCGTATTCCAGTC-3' and T7Select down 5'-GGCTGATACCACCTTCAAG-3' as forward and reverse primers respectively. PCR conditions were 30 cycles of denaturation at 94°C for 30 sec, 55°C annealing for 30 sec, and 72°C extension for 30 sec. A final extension at 72°C was for 5 min. The reaction products were resolved on agarose gel to confirm the presence of amplification products. The PCR products were then purified using the Qiaquick PCR purification kit (Qiagen, Valencia, CA) and the inserts were sequenced on both strands at DNA Services Facility (Research Resources Center, University of Illinois at Chicago, IL) to confirm the authenticity of the genes. The determined DNA sequences were then analyzed by BLAST analyses.

Results

Over twenty phage clones bound to SHBG. Of these, initially we isolated ten phage plaques from the fourth round of biopanning and processed them for sequencing. Agarose gel resolution of the PCR products showed that they were all the same size (data not shown), so we sequenced two clones out of the ten. Sequence analysis demonstrated that SHBG-interacting phage clones encoded for FAU gene products [ubiquitin-like protein (FUBI) and ribosomal protein S30] (28).

Because of the low molecular weight of these two candidate proteins, we decided to screen more plaques from the first biopanning. Through this process we isolated twenty-four more phage clones. All these clones were then processed for sequencing. Fig. 2 shows the size distribution of the phage clones that were obtained after PCR amplification. The size of these clones varied between 570 and 1,265 bases. Each of the sequences was then subjected to a BLAST search to determine their identity. Table I shows the list of candidate proteins encoded by a phage that bound to SHBG. These

Phage clone no.	Encoding proteins
Membrane	
1	Synaptosomal-associated protein (SNAP25), 25 kDa
2	Thy-1 glycoprotein, Thy-1 cell surface antigen
3	Zonadhesin (ZAN)
Dual localization: Membrane and intracellular	
4	GTP-binding protein (RAB3A)
5	Myelin basic protein
6	Vacuolar protein sorting factor 4A
Intracellular	
7	Adenylate kinase 5
8	ADP-ribosylation factor 6 interacting protein 4 (ARL61P4)
9	Cystatin C
10	Cytoplasmic linker 2 (CYLN2), variant 2
11	Fatty acid binding protein 5
12	Fubi (also coded for by the FAU gene)
13	FUS glycin rich protein
14	Lymphoid-enhancer binding factor 1
15	Mitochondrial ribosomal protein S6
16	Phospholipase A2, group IVC (cytosolic calcium-independent) (PLA2G4C) gene
17	PR-domain zinc finger protein 2 variant
18	Ribosomal protein L15
19	Ribosomal protein L10
20	Ribosomal protein L13
21	Ribosomal protein L17, transcript 1
22	Ribosomal protein S30 (FAU),
23	Transcription elongation factor A (SII)-like 3 (TCEAL3)
24	Zinc finger protein 260

include cell membrane associated proteins (SNAP25, Thy-1 cell surface antigen, and zonadhesin), GTP-binding protein (RAB3A), myelin basic protein and vacuolar protein sorting factor 4A.

Discussion

SHBG is a 94-kDa protein found in the blood (10,29,30), brain (23-25), and the male reproductive tract (24,31). One model of SHBG action suggests that it is a passive carrier of steroids in blood, while another model suggests that there are receptors for SHBG on cellular membranes that mediate intracellular actions (1,5,21,22,32-34). In this study, we tested whether we could utilize the phage display technique to identify proteins

that interact with SHBG. This technique identified over twenty binding proteins. Of these identified candidate proteins three are localized to the cell membrane [SNAP25 (35), Thy-1 cell surface antigen (36), and zonadhesin (37)]. Some of the other proteins include GTP-binding protein (RAB3A) (38), myelin basic protein (39) and vacuolar protein sorting factor 4A (40). These findings demonstrate that both cytoplasmic and membrane proteins interact with SHBG. The majority of the identified proteins are intracellular with a good number of them involved in protein synthesis.

Pope and Lee sequenced at least sixty proteins from over 230 interacting proteins they obtained using the yeast two-hybrid system to identify prostatic SHBG binding proteins (27). Two of the metabolic cytosolic proteins we identified, Adenylate kinase 5 and Phospholipase A2, group IVC, were also identified by them (27). Adenylate kinase is involved in the homeostasis of adenine and guanine nucleotides (41), while phospholipase A2 catalyzes the hydrolysis of the 2-acyl bond of 3-n-phosphoglycerides (42). However, despite the high number of membrane proteins they obtained, none of them are high molecular weight proteins. Interestingly, adenylate kinase and phospholipase A2 were shown to be over-expressed in prostatic cancer (43,44) and the putative SHBG receptor was also first described in the prostate (1). Given the association of these proteins with SHBG in the brain it is not known at this time if these proteins are also involved in brain cancer. If so, elucidation of the interaction of SHBG with these proteins may have implications for steroid-initiated tumorigenesis.

Because others have postulated that the SHBG receptor is associated with the cellular membrane, we will first discuss candidate proteins associated with the cellular membrane. Of the three membrane-associated candidate proteins, only zonadhesin has a molecular weight that corresponds to the molecular weight suggested by Hryb *et al* (1) for the putative SHBG receptor, i.e. 167 kDa. It is known that zonadhesin forms two disulfide-bonded monomeric constituents of 105 and 45 kDa, for a total molecular weight of 150 kDa. Zonadhesin is a mosaic protein in sperm membrane fractions that binds directly and in a species-specific manner to the extracellular matrix (zona pellucida; ZP) of the oocytes. Post-translational processing leads to multiple forms of zonadhesin with differing apparent avidities for the ZP (45). In the pig, the p105/45 form binds preferentially to conspecific ZP (45). It is interesting that with the discovery of membrane receptors for progesterone in oocytes (46-48), that rapid steroid effects in oocytes are receiving renewed interest. We found SHBG immunoreactivity in prostatic secretions (24), which suggests that SHBG in the ejaculate serves as a stimulatory agent at oocyte membrane receptors, as with zonadhesin. However, it is not known whether zonadhesin exists in the brain. Using Northern blot analysis, zonadhesin expression was found only within the testes and it was not detected in other organs including the brains of mice (49). However, due to the presence of MAM, mucin, D-, and EGF domains in zonadhesin, Gao and Garbers (49) suggested that apart from binding to zona pellucida, mouse zonadhesin functions in other cell adhesion processes. Our suggestion that zonadhesin exists in the human brain might indicate its involvement in cell adhesion and possibly cell migration during development.

Another indicator that a candidate protein is associated with the membrane is coupling to G proteins. We identified RAB3A as a candidate protein, which is a GTP-binding protein of 24 kDa which belongs to the GTP-binding protein superfamily. RAB3A and its interacting proteins are involved in Ca²⁺-dependent exocytosis (50). RAB3A has been identified in membrane and cytosolic fractions of mouse with the predominant form being membrane-bound. The localization to acrosomal membrane suggests a role for RAB3A in the regulation of zona pellucida-induced acrosomal exocytosis (51) that, along with identification of zonadhesin, may indicate a parallel receptor complex in brain and acrosome.

Soluble N-ethylmaleimide-sensitive factor attachment protein (SNAP-25) is an integral membrane protein localized to the plasma membrane of pre-synaptic axon terminals. As a component of the SNARE (receptor for SNAPs), SNAP-25 appears to play an important role in the docking and fusion of pre-synaptic vesicles during exocytosis of neurotransmitters (52). Pope and Lee (27) identified another protein called SNARE associated protein which is also a component of the SNARE complex (53). SHBG internalization was observed in a number of tissues (7,8,10,11,13,54) including the brain (15), so it is possible that some of the proteins we identified here, in particular SNAP-25, are involved in that internalization process.

We also discovered that one protein binding to SHBG is myelin basic protein. Myelin basic protein (MBP) makes up ~30% of the components of a myelin sheath which electrically insulates axons and aids conduction of nerve impulses. There is significant interest in MBP because of its role in demyelinating diseases, particularly multiple sclerosis (MS). Several studies have shown a role for antibodies against MBP in the pathogenesis of MS (55). It is possible that one function of SHBG in the brain is due to actions at the level of myelin in brain fiber tracts. SHBG internalization was often associated with such tracts in the brain (15). It is also interesting to note that MBP is present in various different cell types including lymphoid cells and all types of myeloid lineage cells (i.e., macrophages, dendritic cells, granulocytes, megakaryocytes, and erythroblasts) (56). At least five isoforms of MBP were identified with molecular weights ranging from 14 to 21 kDa. (57). Again, this is not likely the SHBG receptor, but it could be a component of it. It was demonstrated that supplemental estrogen enhances the efficacy of TCR-based immunotherapy for autoimmune diseases that are predominant in females (58). Estrogen potentiates treatment with T-cell-receptor protein of female mice with experimental encephalomyelitis (EAE). This is a significant finding because the involvement of estrogen in the therapy of EAE (a mouse model for MS) also suggests a role for SHBG in its transportation and hence potential therapy for MS.

Thy-1 glycoproteins were identified as candidate proteins as well. Thy-1 glycoproteins are constituents of thymocytes and neurons. Thy-1 cell surface antigen is a small protein (18 kDa) that may play a role in cell-cell or -ligand interactions during synaptogenesis and other events in the brain (UniProtKB/Swiss-Prot, Institute of Bioinformatics and the EMBL outstation-the European Bioinformatics Institute). It was also shown that despite the fact that Thy-1 glycoprotein is anchored (glycosylphosphatidyl-inositol anchor) to the outer hemi-leaflet of the cell membrane, it is capable of transducing signals across the

plasma membrane of murine T-lymphoma cells (59). To our knowledge, no such activity has been demonstrated in brain cells. In any case, the low molecular weight of Thy-1 suggests that it could only be part of a large receptor complex for SHBG.

Another candidate protein was vacuolar protein sorting factor 4A (VPS4), which is a member of the ATPases Associated with diverse cellular Activities (AAA) ATPase family and a central regulator for early endosome trafficking (60). This protein is well characterized in yeast where it is known to bind to oxysterol binding protein and regulate intracellular lipid and vesicular transport (25). Since VPS4 has a molecular weight of 48 kDa (61), it is not likely to be the anticipated SHBG receptor, however it could be one of the components of the receptor, which after being internalized, functions in the transport of SHBG-bound steroid.

Although FUBI and ribosomal protein S30 are cytosolic low molecular weight proteins, they were obtained at the fourth panning suggesting that they had the highest affinity for SHBG. FUBI is synthesized with ribosomal S30 as its C-terminal extension. It was proposed that the fusion protein is post-translationally processed to generate free FUBI and free ribosomal protein S30. FUBI is a member of the ubiquitin family, and ribosomal protein S30 belongs to the S30E family of ribosomal proteins. Whereas the function of FUBI is currently unknown, ribosomal protein S30 is a component of the 40S subunit of the cytoplasmic ribosome (62). It is also interesting to note the high number of other ribosomal proteins that are involved in binding to SHBG. It is not known what the consequences of the interaction of SHBG with FUBI or ribosomal proteins are or any possible involvement in steroid metabolism. However, it will be interesting to investigate, using cell culture systems, how these proteins are modified in the presence of SHBG and how their deficiencies affect steroid metabolism.

This phage display system did not identify megalin which is a candidate for the SHBG receptor as mentioned above. The failure of SHBG to identify megalin may be due to the inability of phages to fuse large proteins or simply that megalin is absent in the brain.

Overall, it seems likely that the SHBG receptor is made up of several protein subunits and that any one of the proteins we identified could be a component of the large molecule. For example, there are four proteins in the presynaptic terminal involved in neurotransmitter release, RAB3A, SNAP-25, VAMP/synaptobrevin-2 and syntaxin, all form a complex of ~119 kDa (63). Another study that lends credibility to this argument is by Oike *et al* (64) who showed that Group IIA phospholipase A2 is coexpressed with SNAP-25 in rat mature taste receptor cells that possess exocytotic function. If several elements of the SHBG receptor complex are only transiently found in the membrane, it may explain why Pope and Lee (27) were unable to identify a single protein of high molecular weight using their yeast two-hybrid method.

Because the phages of the Phage Display technique display all proteins on their surface, it may be that we here demonstrated SHBG binding to proteins that are normally only able to interact with SHBG within the cell. SHBG internalization has been demonstrated in multiple tissues (7-9,11-13,54) including the brain (15) suggesting that SHBG along with associated



are selectively brought into cells, where cytosolic may bind the SHBG. This work may have identified proteins that are candidates for an intracellular SHBG-binding complex.

Acknowledgements

This work was completed with funding from R21-MH069810 (JDC) and Excellence in Academic Medicine grant of the Swedish American Health System (MG).

References

- Hryb DJ, Khan MS, Romas NA and Rosner W: Solubilization and partial characterization of the sex hormone-binding globulin receptor from human prostate. *J Biol Chem* 264: 5378-5383, 1989.
- Nakhla AM, Khan MS, Romas NP and Rosner W: Estradiol causes the rapid accumulation of cAMP in human prostate. *Proc Natl Acad Sci USA* 91: 5402-5405, 1994.
- Nakhla AM, Leonard J, Hryb DJ and Rosner W: Sex hormone-binding globulin receptor signal transduction proceeds via a G protein. *Steroids* 64: 213-216, 1999.
- Rosner W, Hryb DJ, Khan MS, Nakhla AM and Romas NA: Androgen and estrogen signaling at the cell membrane via G-proteins and cyclic adenosine monophosphate. *Steroids* 64: 100-106, 1999.
- Avvakumov GV, Zhuk NI and Strel'chyonok OA: Subcellular distribution and selectivity of the protein-binding component of the recognition system for sex-hormone-binding protein-estradiol complex in human decidua endometrium. *Biochim Biophys Acta* 881: 489-498, 1986.
- Strel'chyonok OA, Avvakumov GV and Survilo LI: A recognition system for sex-hormone-binding protein-estradiol complex in human decidua endometrium plasma membranes. *Biochim Biophys Acta* 802: 459-466, 1984.
- Gerard A: Endocytosis of androgen-binding protein (ABP) by spermatogenic cells. *J Steroid Biochem Mol Biol* 53: 533-542, 1995.
- Gueant JL, Fremont S, Felden F, Nicolas JP, Gerard A, Leheup B, Gerard H and Grignon G: Evidence that androgen-binding protein endocytosis in vitro is receptor mediated in principal cells of the rat epididymis. *J Mol Endocrinol* 7: 113-122, 1991.
- Meyer S, Brumm C, Stegner HE and Sinnecker GH: Intracellular sex hormone-binding globulin (SHBG) in normal and neoplastic breast tissue--an additional marker for hormone dependency? *Exp Clin Endocrinol* 102: 334-340, 1994.
- Pardridge WM: Transport of protein-bound hormones into tissues in vivo. *Endocr Rev* 2: 103-123, 1981.
- Porto CS, Gunsalus GL, Bardin CW, Phillips DM and Musto NA: Receptor-mediated endocytosis of an extracellular steroid-binding protein (TeBG) in MCF-7 human breast cancer cells. *Endocrinology* 129: 436-445, 1991.
- Porto CS, Lazari MF, Abreu LC, Bardin CW and Gunsalus GL: Receptors for androgen-binding proteins: internalization and intracellular signalling. *J Steroid Biochem Mol Biol* 53: 561-565, 1995.
- Sakiyama R, Pardridge WM and Musto NA: Influx of testosterone-binding globulin (TeBG) and TeBG-bound sex steroid hormones into rat testis and prostate. *J Clin Endocrinol Metab* 67: 98-103, 1988.
- Sinnecker G, Hiort O, Kwan PW and DeLellis RA: Immunohistochemical localization of sex hormone-binding globulin in normal and neoplastic breast tissue. *Horm Metab Res* 22: 47-50, 1990.
- Caldwell JD, Shapiro RA, Jirikowski GF and Suleman F: Internalization of sex hormone-binding globulin into neurons and brain cells in vitro and in vivo. *Neuroendocrinology* 86: 84-93, 2007.
- Hammes A, Andreassen TK, Spoelgen R, Raila J, Hubner N, Schulz H, Metzger J, Schweigert FJ, Luppa PB, Nykjaer A and Willnow TE: Role of endocytosis in cellular uptake of sex steroids. *Cell* 122: 751-762, 2005.
- Willnow TE, Nykjaer A and Herz J: Lipoprotein receptors: new roles for ancient proteins. *Nat Cell Biol* 1: E157-E162, 1999.
- Mopert B, Herbert Z, Caldwell JD and Jirikowski GF: Expression of corticosteroid-binding globulin in the rat hypothalamus. *Horm Metab Res* 38: 246-252, 2006.
- Sivukhina EV, Jirikowski GF, Bernstein HG, Lewis JG and Herbert Z: Expression of corticosteroid-binding protein in the human hypothalamus, co-localization with oxytocin and vasopressin. *Horm Metab Res* 38: 253-259, 2006.
- Petersen HH, Andreassen TK, Breiderhoff T, Brasen JH, Schulz H, Gross V, Grone HJ, Nykjaer A and Willnow TE: Hyporesponsiveness to glucocorticoids in mice genetically deficient for the corticosteroid binding globulin. *Mol Cell Biol* 26: 7236-7245, 2006.
- Caldwell JD, Moe BD, Hoang J and Nguyen T: Sex hormone binding globulin stimulates female sexual receptivity. *Brain Res* 874: 24-29, 2000.
- Caldwell JD, Hofle S and Englof I: Sex hormone binding globulin facilitates female sexual receptivity except when coupled to dihydrotestosterone. *Brain Res* 948: 102-107, 2002.
- Herbert Z, Jirikowski GF, Petrusz P, Englof I and Caldwell JD: Distribution of androgen-binding protein in the rat hypothalamo-neurohypophyseal system, co-localization with oxytocin. *Brain Res* 992: 151-158, 2003.
- Herbert Z, Weigel S, Sendemir E, Marshall A, Caldwell JD, Petrusz P, Peuckert C and Jirikowski GF: Androgen-binding protein is co-expressed with oxytocin in the male reproductive tract. *Anat Histol Embryol* 34: 286-293, 2005.
- Wang YM, Bayliss DA, Millhorn DE, Petrusz P and Joseph DR: The androgen-binding protein gene is expressed in male and female rat brain. *Endocrinology* 127: 3124-3130, 1990.
- Caldwell JD: Evidence of sex hormone binding globulin binding sites in the medial preoptic area and hypothalamus. *Horm Metab Res* 33: 7-9, 2001.
- Pope SN and Lee IR: Yeast two-hybrid identification of prostatic proteins interacting with human sex hormone-binding globulin. *J Steroid Biochem Mol Biol* 94: 203-208, 2005.
- Rossmann TG, Visalli MA and Komissarova EV: *fau* and its ubiquitin-like domain (FUBI) transforms human osteogenic sarcoma (HOS) cells to anchorage-independence. *Oncogene* 22: 1817-1821, 2003.
- Siiteri PK, Murai JT, Hammond GL, Nisker JA, Raymoure WJ and Kuhn RW: The serum transport of steroid hormones. *Recent Prog Horm Res* 38: 457-510, 1982.
- Walsh KA, Titani K, Takio K, Kumar S, Hayes R and Petra PH: Amino acid sequence of the sex steroid binding protein of human blood plasma. *Biochemistry* 25: 7584-7590, 1986.
- Hryb DJ, Nakhla AM, Kahn SM, St George J, Levy NC, Romas NA and Rosner W: Sex hormone-binding globulin in the human prostate is locally synthesized and may act as an autocrine/paracrine effector. *J Biol Chem* 277: 26618-26622, 2002.
- Hryb DJ, Khan MS, Romas NA and Rosner W: The control of the interaction of sex hormone-binding globulin with its receptor by steroid hormones. *J Biol Chem* 265: 6048-6054, 1990.
- Khan MS, Hryb DJ, Hashim GA, Romas NA and Rosner W: Delineation and synthesis of the membrane receptor-binding domain of sex hormone-binding globulin. *J Biol Chem* 265: 18362-18365, 1990.
- Krupenko NI, Avvakumov GV and Strel'chyonok OA: Binding of human sex hormone-binding globulin-androgen complexes to the placental syncytiotrophoblast membrane. *Biochem Biophys Res Commun* 171: 1279-1283, 1990.
- Oyler GA, Polli JW, Wilson MC and Billingsley ML: Developmental expression of the 25-kDa synaptosomal-associated protein (SNAP-25) in rat brain. *Proc Natl Acad Sci USA* 88: 5247-5251, 1991.
- De Hoop MJ and Dotti CG: Membrane traffic in polarized neurons in culture. *J Cell Sci (Suppl)* 17: 85-92, 1993.
- Hardy DM and Garbers DL: A sperm membrane protein that binds in a species-specific manner to the egg extracellular matrix is homologous to von Willebrand factor. *J Biol Chem* 270: 26025-26028, 1995.
- Fischer von Mollard G, Sudhof TC and Jahn R: A small GTP-binding protein dissociates from synaptic vesicles during exocytosis. *Nature* 349: 79-81, 1991.
- Gillespie CS, Wilson R, Davidson A and Brophy PJ: Characterization of a cytoskeletal matrix associated with myelin from rat brain. *Biochem J* 260: 689-696, 1989.
- Beyer A, Scheuring S, Muller S, Mincheva A, Lichter P and Kohrer K: Comparative sequence and expression analyses of four mammalian VPS4 genes. *Gene* 305: 47-59, 2003.
- Yoneda T, Sato M, Maeda M and Takagi H: Identification of a novel adenylate kinase system in the brain: cloning of the fourth adenylate kinase. *Brain Res Mol Brain Res* 62: 187-195, 1998.
- Dennis EA: Diversity of group types, regulation, and function of phospholipase A2. *J Biol Chem* 269: 13057-13060, 1994.

43. Hall M, Mickey DD, Wenger AS and Silverman LM: Adenylate kinase: an onco-developmental marker in an animal model for human prostatic cancer. *Clin Chem* 31: 1689-1691, 1985.
44. Sved P, Scott KF, McLeod D, King NJ, Singh J, Tsatralis T, Nikolov B, Boulas J, Nallan L, Gelb MH, Sajinovic M, Graham GG, Russell PJ and Dong Q: Oncogenic action of secreted phospholipase A2 in prostate cancer. *Cancer Res* 64: 6934-6940, 2004.
45. Hickox JR, Bi M and Hardy DM: Heterogeneous processing and zona pellucida binding activity of pig zonadhesin. *J Biol Chem* 276: 41502-41509, 2001.
46. King Wt, Ghosh S, Thomas P and Sullivan CV: A receptor for the oocyte maturation-inducing hormone 17 α ,20 β ,21-trihydroxy-4-pregnen-3-one on ovarian membranes of striped bass. *Biol Reprod* 56: 266-271, 1997.
47. Thomas P, Zhu Y and Pace M: Progesterone membrane receptors involved in the meiotic maturation of teleost oocytes: a review with some new findings. *Steroids* 67: 511-517, 2002.
48. Zhu Y, Rice CD, Pang Y, Pace M and Thomas P: Cloning, expression, and characterization of a membrane progesterone receptor and evidence it is an intermediary in meiotic maturation of fish oocytes. *Proc Natl Acad Sci USA* 100: 2231-2236, 2003.
49. Gao Z and Garbers DL: Species diversity in the structure of zonadhesin, a sperm-specific membrane protein containing multiple cell adhesion molecule-like domains. *J Biol Chem* 273: 3415-3421, 1998.
50. Takai Y, Sasaki T, Shirataki H and Nakanishi H: Rab3A small GTP-binding protein in Ca(2+)-dependent exocytosis. *Genes Cells* 1: 615-632, 1996.
51. Ward CR, Faundes D and Foster JA: The monomeric GTP binding protein, rab3a, is associated with the acrosome in mouse sperm. *Mol Reprod Dev* 53: 413-421, 1999.
52. Goda Y: SNAREs and regulated vesicle exocytosis. *Proc Natl Acad Sci USA* 94: 769-772, 1997.
53. Ilardi JM, Mochida S and Sheng ZH: Snapin: a SNARE-associated protein implicated in synaptic transmission. *Nat Neurosci* 2: 119-124, 1999.
54. Bordin S and Petra PH: Immunocytochemical localization of the sex steroid-binding protein of plasma in tissues of the adult monkey *Macaca nemestrina*. *Proc Natl Acad Sci USA* 77: 5678-5682, 1980.
55. Berger T, Rubner P, Schautzer F, Egg R, Ulmer H, Mayringer I, Dilitz E, Deisenhammer F and Reindl M: Antimyelin antibodies as a predictor of clinically definite multiple sclerosis after a first demyelinating event. *N Engl J Med* 349: 139-145, 2003.
56. Kim JG, Nonneman D, Vallet JL, Rohrer GA and Christenson RK: Linkage mapping of the porcine myelin basic protein gene to chromosome 1. *Anim Genet* 36: 163-164, 2005.
57. Mathisen PM, Pease S, Garvey J, Hood L and Readhead C: Identification of an embryonic isoform of myelin basic protein that is expressed widely in the mouse embryo. *Proc Natl Acad Sci USA* 90: 10125-10129, 1993.
58. Offner H, Adlard K, Zamora A and Vandenbark AA: Estrogen potentiates treatment with T-cell receptor protein of female mice with experimental encephalomyelitis. *J Clin Invest* 105: 1465-1472, 2000.
59. Lemansky P, Fatemi SH, Gorican B, Meyale S, Rossero R and Tartakoff AM: Dynamics and longevity of the glycolipid-anchored membrane protein, Thy-1. *J Cell Biol* 110: 1525-1531, 1990.
60. Scheuring S, Rohricht RA, Schoning-Burkhardt B, Beyer A, Muller S, Abts HF and Kohrer K: Mammalian cells express two VPS4 proteins both of which are involved in intracellular protein trafficking. *J Mol Biol* 312: 469-480, 2001.
61. Babst M, Sato TK, Banta LM and Emr SD: Endosomal transport function in yeast requires a novel AAA-type ATPase, Vps4p. *EMBO J* 16: 1820-1831, 1997.
62. Kas K, Michiels L and Merregaert J: Genomic structure and expression of the human fau gene: encoding the ribosomal protein S30 fused to a ubiquitin-like protein. *Biochem Biophys Res Commun* 187: 927-933, 1992.
63. Horikawa HP, Saisu H, Ishizuka T, Sekine Y, Tsugita A, Odani S and Abe T: A complex of rab3A, SNAP-25, VAMP/synaptobrevin-2 and syntaxins in brain presynaptic terminals. *FEBS Lett* 330: 236-240, 1993.
64. Oike H, Matsumoto I and Abe K: Group IIA phospholipase A(2) is coexpressed with SNAP-25 in mature taste receptor cells of rat circumvallate papillae. *J Comp Neurol* 494: 876-886, 2006.