

# Deoxyschisandrin inhibits H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death in intestinal epithelial cells through nuclear factor- $\kappa$ B

BON-HEE GU<sup>1</sup>, NGUYEN VAN MINH<sup>1</sup>, SUN-HEE LEE<sup>1</sup>, SUNG-WON LIM<sup>1</sup>,  
YOUNG-MI LEE<sup>2</sup>, KANG-SOO LEE<sup>3</sup> and DAE-KI KIM<sup>1</sup>

<sup>1</sup>Department of Immunology, Institute of Medical Sciences, Chonbuk National University Medical School, Jeonju, Jeonbuk 561-180; <sup>2</sup>Department of Oriental Pharmacy, College of Pharmacy, Wonkwang University, Iksan, Jeonbuk 570-749;

<sup>3</sup>Faculty of Biological Resources Science, Chonbuk National University, Jeonju, Jeonbuk 561-756, Korea

Received April 16, 2010; Accepted June 10, 2010

DOI: 10.3892/ijmm\_00000479

**Abstract.** Oxidative stress is a pathogenesis for a typical inflammatory intestinal disease known as ulcerative colitis (UC) characterized by erosion and mucosal ulceration. For the treatment of UC, many kinds of traditional Asian medical plants have been used. *Schisandra chinensis* fruits (SC) are known to possess anti-ulcer, anti-hepatotoxic and anti-neurotoxic activity. However, its mechanism is still unknown. In the present study, we investigated the cytoprotective effect of deoxyschisandrin, a lignan compound comprised of SC fruits, on H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death in human intestinal epithelial cells (HCT116). In flow cytometry assay using Annexin V and propidium iodide, deoxyschisandrin inhibited H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death. To further evaluate the apoptotic signaling by H<sub>2</sub>O<sub>2</sub>, we detected caspase-3 activation using cleavage of pro-caspase-3. Deoxyschisandrin inhibited H<sub>2</sub>O<sub>2</sub>-induced caspase-3 activation by blocking cleavage of pro-caspase-3. Furthermore, it has been reported that oxidative stress by H<sub>2</sub>O<sub>2</sub> induces an activation of nuclear factor- $\kappa$ B (NF- $\kappa$ B). In our results, H<sub>2</sub>O<sub>2</sub> stimulated the degradation of I $\kappa$ B $\alpha$ , inhibitor of NF- $\kappa$ B, in a concentration-dependent manner. On the contrary, deoxyschisandrin inhibited H<sub>2</sub>O<sub>2</sub>-stimulated degradation of I $\kappa$ B $\alpha$  and activation of NF- $\kappa$ B by blocking translocation of NF- $\kappa$ B to the nucleus. Therefore, we suggest that deoxyschisandrin inhibits H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death.

## Introduction

Intestinal epithelial cells (IECs) constitute a single layer of the intestine and act as a critical physical barrier against pathogen invasion. It is considered to be an important component associated with initiation and regulation of adaptive and innate defense mechanisms in the mucosal immune system (1). In IECs, the activation of the pro-inflammatory gene is related with chronic and acute intestinal inflammation (2-4).

Ulcerative colitis (UC) is a type of inflammatory bowel disease (IBD) characterized by mucosal ulceration, up-regulation of pro-inflammatory cytokines, and infiltration of inflammatory cells into the colonic mucosa (5,6). Although the etiology of UC is not clearly understood, it has been reported that exaggerated immune response by immune cells such as white blood cells (mainly neutrophils) is a considerable pathogenesis factor (7). Responsibility of neutrophils for tissue damage in the colonic mucosal surface has been well described but an explanation of the mechanism has not been proposed (8). However, the possible mechanism has been reported that reactive oxygen species (ROS)-mediated oxidative stress is a major pathogenic mechanism in IBD. ROS, reactive molecules containing the oxygen atom, can initiate inflammatory cascades and cause subsequent tissue damage (9). ROS-mediated oxidative stress provokes oxidative damage, results in neutrophil infiltration into the colonic mucosa. H<sub>2</sub>O<sub>2</sub> is a major contributor to ROS-mediated oxidative stress (10-12).

H<sub>2</sub>O<sub>2</sub> is produced as a toxic by-product in living organisms through the mitochondrial electron transport chain (ETC) respiratory process (13). When H<sub>2</sub>O<sub>2</sub> is produced excessively by aberrant cellular metabolism in intestinal epithelial cells, H<sub>2</sub>O<sub>2</sub> can have an effect on damage of colonic barrier function structures to induce UC (10). Furthermore, H<sub>2</sub>O<sub>2</sub>-induced oxidative stress regulates the intracellular signaling pathway. Representatively, nuclear factor- $\kappa$ B (NF- $\kappa$ B), a heteromeric transcription factor, is activated by H<sub>2</sub>O<sub>2</sub> (14). NF- $\kappa$ B mediates cell survival and inflammation via regulation of apoptotic genes such as caspase-3 and anti-apoptotic genes such as Bcl-2 (15).

*Schisandra Chinensis* fruits (SC) is widely used as a traditional Asian medical plant for diarrhea. It has been reported that SC has anti-ulcer, anti-hepatotoxic and anti-neurotoxic activity (16-18). We also previously demonstrated

---

**Correspondence to:** Dr Dae-Ki Kim, Department of Immunology and Institute of Medical Sciences, Chonbuk National University Medical School, 2-20 Geumam 2-dong, Deokjin-gu, Jeonju, Jeonbuk 561-180, Korea  
E-mail: daekim@jbnu.ac.kr

**Abbreviations:** IBD, inflammatory bowel disease; IECs, intestinal epithelial cells; NF- $\kappa$ B, nuclear factor- $\kappa$ B; ROS, reactive oxygen species; SC, *Schisandra chinensis* fruits; UC, ulcerative colitis

**Key words:** apoptosis, deoxyschisandrin, intestinal epithelial cells, nuclear factor- $\kappa$ B, oxidative stress

inhibitory effects of aqueous extract from SC in dextran sulfate sodium (DSS)-induced colitis mouse model, but the molecular mechanism has still not been identified (19). SC contains a variety of pharmacologically active lignan components (20).

Deoxyschisandrin is one of the lignan components of SC. It consists of dibenzocyclooctadiene which possesses hepatoprotective, neuroprotective, and antioxidant activities (21-24). It was reported that deoxyschisandrin has neuroprotective activity in glutamate-induced neurotoxicity. Deoxyschisandrin attenuated the content of cellular peroxide and enhanced the glutathione antioxidative defense system in rat cortical cells (24).

In the present study, we investigated the effect of deoxyschisandrin on apoptotic cell death of intestinal epithelial cells (HCT116) induced by H<sub>2</sub>O<sub>2</sub>. We attempted to identify the molecular mechanisms. The study demonstrated that deoxyschisandrin suppresses H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death through inhibition of the activities of NF- $\kappa$ B.

## Materials and methods

**Materials and cell culture.** Deoxyschisandrin was purchased from Chromadex (CA, USA). HCT116 cell line was incubated in Dulbecco's modified Eagle's medium (DMEM) (Thermo Scientific Hyclone, UT, USA) with penicillin (100 units/ml) - streptomycin (100  $\mu$ g/ml) (Welgene, Korea), and 10% fetal bovine serum (Thermo Scientific Hyclone, UT, USA) at 37°C under 5% CO<sub>2</sub> humidified air.

**CCK-8 assay.** To evaluate cell viability, we used CCK-8 cell counting kit (Dojindo Molecular Technologies, MD, USA). Cells were prepared in 96-well plates (5 $\times$ 10<sup>3</sup> cells/plate) to treat H<sub>2</sub>O<sub>2</sub> or deoxyschisandrin. Absorbance was measured at 450 nm.

**Determining early apoptosis and necrosis by flow cytometry.** For apoptosis and necrosis analysis of HCT116 cells, flow cytometry assay was performed using FITC-labeled Annexin V (BD Biosciences Pharmingen, CA, USA) and propidium iodide (PI) (Sigma Aldrich, Germany). Cells were seeded into 12-well plates (4 $\times$ 10<sup>4</sup> cells/plate), and 24 h later, were treated with H<sub>2</sub>O<sub>2</sub> and deoxyschisandrin in each concentration. After treatment, cells were gently detached with Detachin (Genlantis, CA, USA) and washed with PBS containing 0.1% sodium azide. Cells were incubated with Annexin V (final concentration, 1  $\mu$ g/ml) for 15 min in the dark, washed one more time, and suspended in PI (final concentration 5  $\mu$ g/ml). Stained cells were analyzed with a FACSCalibur flow cytometer (BD Biosciences, CA, USA). A total 5 $\times$ 10<sup>3</sup> cells were analyzed per sample.

**Western blot analysis.** To detect protein expression levels, cells were detached using cell scrapers and washed with PBS. Lysis buffer (Intron Biotech, Korea) was used to prepare total cell lysates. Lysates were electrophoresed using 10-15% SDS-PAGE and transferred to PVDF membranes (Amersham Pharmacia Biotech, NJ, USA). Membranes were incubated with blocking solution, containing a 1:1,000 dilution of anti-I $\kappa$ B $\alpha$  (Santa Cruz Biotechnology, CA, USA), or anti-caspase-3 (Santa Cruz Biotechnology, CA, USA) antibodies. Antibody of anti-




Figure 1. Effects of deoxyschisandrin on the viability of HCT116 cells. Cells were treated with deoxyschisandrin in various concentrations from 0.01 to 5  $\mu$ g/ml for 24 h. Cell viability was determined using CCK-8 assay. Values are presented as percent of control and results are the mean  $\pm$  SEM of three independent experiments.




Figure 2. Suppression of H<sub>2</sub>O<sub>2</sub> on the viability in HCT116 cells. Cells were treated with H<sub>2</sub>O<sub>2</sub> in different concentrations (0-800  $\mu$ M) for 24 h. Cell viability was determined using CCK-8 assay. Values are represented as percent of control and the results are the mean  $\pm$  SEM of three independent experiments.

GAPDH (Epitomics, CA, USA) diluted 1:2,000 with blocking solution was used as a loading control. Then, the membranes were incubated with blocking solution containing a 1:10,000 dilution of horseradish peroxidase (HRP)-conjugated anti-rabbit secondary antibodies (Amersham Pharmacia Biotech, NJ, USA). An enhanced chemiluminescence detection kit (Amersham Pharmacia Biotech, NJ, USA) was used to detect the epitope on the proteins recognized by the specific antibody.

**Immunofluorescent staining and nuclear staining.** HCT116 cells were grown in culture slides for 24 h, and then treated either with or without H<sub>2</sub>O<sub>2</sub> and deoxyschisandrin. After treatment, cells were treated with 4% paraformaldehyde (PFA) for 15 min. After washing 3 times, cells were blocked with 1% bovine serum albumin (BSA)-PBS for 3 h and incubated with rabbit monoclonal NF- $\kappa$ B p65 antibody (Epitomics, CA, USA) diluted 1:100 with 1% BSA-PBS overnight in 4°C. Incubated cells were washed 3 times, and treated with FITC-conjugated anti-rabbit secondary antibody (Santa Cruz Biotech, Santa Cruz, CA, USA) diluted 1:200 with 1% BSA-PBS for 20 min. Then, cells were stained with DAPI solution for the nuclei staining. The fluorescence and translocation of NF- $\kappa$ B were evaluated by confocal laser scanning microscope (LSM 510 Meta, Carl Zeiss Co., Germany).




Figure 3. Inhibitory effect of deoxyschisandrin on H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death in HCT116 cells. Cells were pretreated with deoxyschisandrin (1 µg/ml) for 30 min and stimulated with H<sub>2</sub>O<sub>2</sub> (400 µM) for 4 h. Flow cytometry assay was performed with Annexin V and PI staining.

## Results

*Deoxyschisandrin has a cytotoxic effect in HCT116 cells.* We performed CCK-8 assay in human intestinal epithelial cell line (HCT116) with deoxyschisandrin (Fig. 1). Deoxyschisandrin was treated with various concentrations for 24 h. As a result, the cell viability was not affected up to 1 µg/ml of deoxyschisandrin, whereas 5 µg/ml of deoxyschisandrin decreased cell viability.

*H<sub>2</sub>O<sub>2</sub> induces apoptotic cell death.* We examined the cytotoxic effect of H<sub>2</sub>O<sub>2</sub> to investigate whether H<sub>2</sub>O<sub>2</sub> induces apoptosis in HCT116 cells using CCK-8 assay (Fig. 2). Cell viability was measured after treatment with various concentrations of H<sub>2</sub>O<sub>2</sub> for 24 h. In an H<sub>2</sub>O<sub>2</sub> concentration-dependent condition, cell viability decreased in over 400 µM of H<sub>2</sub>O<sub>2</sub>. When cells were treated with 400 µM of H<sub>2</sub>O<sub>2</sub> for 24 h, the viability of cells was reduced to ~57% of control, whereas the viability of cells treated with 800 µM of H<sub>2</sub>O<sub>2</sub> was significantly reduced to 13% of control. Next, we confirmed H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death in flow cytometry assay using co-staining with Annexin V and PI (Fig. 3). Annexin V-FITC and PI fluorescent intensity were increased in H<sub>2</sub>O<sub>2</sub>-treated cells more than H<sub>2</sub>O<sub>2</sub>-non-treated cells. The percentage of apoptotic/necrotic cells increased from 1.68% in H<sub>2</sub>O<sub>2</sub> non-treated control to 23.34% in H<sub>2</sub>O<sub>2</sub> treated control.

Activation of caspase-3 has been shown to play a role in oxidative stress-induced apoptotic-cell death (15). Therefore, the examination of the effect of H<sub>2</sub>O<sub>2</sub>-induced apoptosis focuses on activation of caspase-3, we measured the cleavage level of pro-caspase-3. We exposed cells to 400 µM of H<sub>2</sub>O<sub>2</sub> at

various times (Fig. 4A). Expression of pro-caspase-3 was time-dependently decreased. In the treatment of cells for 4 h, expression level of pro-caspase-3 was highly decreased. It reveals that caspase-3 was activated by stimulation of H<sub>2</sub>O<sub>2</sub>. These findings support that H<sub>2</sub>O<sub>2</sub> induces apoptotic cell death in the intestinal epithelial cell line.

### *Deoxyschisandrin inhibits H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death.*

To determine whether deoxyschisandrin has an inhibitory effect on H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death, we first analyzed cell apoptosis using flow cytometry assay. FACS was used to quantitatively determine and distinguish the percentage of viable, apoptotic, and necrotic cells after pretreatment of deoxyschisandrin (1 µg/ml) for 30 min and H<sub>2</sub>O<sub>2</sub> (400 µM) treatment for 4 h (Fig. 3). The percentage of apoptotic/necrotic cells pre-treated with deoxyschisandrin was significantly attenuated up to 4.67% compared to H<sub>2</sub>O<sub>2</sub>-treated control. It was similar with H<sub>2</sub>O<sub>2</sub> non-treated cells. This shows that deoxyschisandrin can protect cells from H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death. Furthermore, we detected activation of caspase-3 in Western blot analysis (Fig. 4B). In cells treated with deoxyschisandrin and H<sub>2</sub>O<sub>2</sub>, expression of pro-caspase-3 was higher than in just H<sub>2</sub>O<sub>2</sub>-treated cells. Therefore, activation of caspase-3 was inhibited by deoxyschisandrin.

### *Deoxyschisandrin inhibits activation of NF-κB in stimulation by H<sub>2</sub>O<sub>2</sub>.*

Oxidative stress by H<sub>2</sub>O<sub>2</sub> induces apoptotic cell death. H<sub>2</sub>O<sub>2</sub> is known as a stimulant of NF-κB-translocation from the cytoplasm to the nucleus. We examined whether stimulation by H<sub>2</sub>O<sub>2</sub> activates NF-κB through detection of the presence level for IκBα in cells (Fig. 5A). H<sub>2</sub>O<sub>2</sub> (400 µM) was




Figure 4. Effects of H<sub>2</sub>O<sub>2</sub> (A) and deoxyschisandrin (B) on caspase-3 activation. Cells were treated with H<sub>2</sub>O<sub>2</sub> (400 µM) in a time-dependent manner (A). Cells were pretreated with deoxyschisandrin (1 µg/ml) for 30 min and stimulated with H<sub>2</sub>O<sub>2</sub> (400 µM) for 4 h (B). Blots were examined with anti-caspase-3 antibody.




Figure 5. Effects of deoxyschisandrin on IκB degradation by H<sub>2</sub>O<sub>2</sub> exposure. Cells were treated with H<sub>2</sub>O<sub>2</sub> (400 µM) in a time-dependent manner (A). Cells were pretreated with deoxyschisandrin (1 µg/ml) for 30 min and stimulated with H<sub>2</sub>O<sub>2</sub> (400 µM) for 4 h (B). Blots were examined with anti-IκBα antibody.




Figure 6. Effects of deoxyschisandrin on NF-κB translocation by H<sub>2</sub>O<sub>2</sub> exposure. Cells were treated with H<sub>2</sub>O<sub>2</sub> (400 µM) with or without deoxyschisandrin (1 µg/ml) for 4 h. In order to perform NF-κB translocation analysis, cells were stained with FITC-conjugated anti-NF-κB antibody (green) and DAPI solution (blue) for nuclei staining. Fluorescence was visualized using confocal laser scanning microscope.

treated in a time-dependent manner, and degradation level of IκBα was detected using Western blot analysis. Degradation of IκBα by phosphorylation was decreased from 60 min, and significantly decreased after 4 h treatment with H<sub>2</sub>O<sub>2</sub>. This result supports that H<sub>2</sub>O<sub>2</sub> induced the activation of NF-κB through degradation of phosphorylated IκBα, inhibitor of NF-κB. To determine the effect of deoxyschisandrin on H<sub>2</sub>O<sub>2</sub>-induced NF-κB, cells were treated with deoxyschisandrin (1 µg/ml) prior to H<sub>2</sub>O<sub>2</sub> (400 µM) stimulation (Fig. 5B). Interestingly, cells treated with deoxyschisandrin before H<sub>2</sub>O<sub>2</sub> treatment showed a similar expression level of IκBα with

non-stimulated cells. It showed that deoxyschisandrin inhibited degradation of IκBα to prevent activation of NF-κB. To confirm the nuclear translocation of the p65 subunit of NF-κB, we performed an immunofluorescence assay (Fig. 6). Subsequent to H<sub>2</sub>O<sub>2</sub> (400 µM) exposure for 4 h, nuclear translocation of the NF-κB p65 subunit was confirmed. DAPI staining was used to inspect the nucleus. In contrast, treatment with deoxyschisandrin (1 µg/ml), which was identified to have an inhibitory effect for degradation of IκBα, inhibited translocation of the NF-κB p65 subunit from the cytoplasm to the nucleus in HCT116 cells. These data suggest



## Discussion

IECs are considered an important physical barrier in response to many inflammatory stimuli in intestinal inflammation, because most pro-inflammatory genes are expressed in the activated IECs (3,4). In IECs, it has been demonstrated that ROS-induced oxidative stress is involved in the development of IBD (8). Excess oxidative stress by H<sub>2</sub>O<sub>2</sub>, a major contributor, induces infiltration of immune cells, and then, immune cells activate NF-κB to produce pro-inflammatory mediators such as tumor necrosis factor α (TNF-α) and pro-apoptosis factors in IECs. Subsequently, the colonic tissue is destroyed leading to UC with diarrhea and rectal bleeding.

In the present study, we identified whether deoxyschisandrin has an inhibitory effect on H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death by stimulation in IECs. We exposed exogenous H<sub>2</sub>O<sub>2</sub> (400 μM) to HCT116. Apoptotic cell death was induced time-dependently by H<sub>2</sub>O<sub>2</sub>. Caspase-3 is known as a central component of the apoptosis machinery (25). We verified that H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death was related to activation of caspase-3. Interestingly, the percentage of apoptotic/necrotic cells was significantly decreased with deoxyschisandrin treatment in flow cytometry assay using Annexin V and PI. Deoxyschisandrin attenuated activation of caspase-3 by H<sub>2</sub>O<sub>2</sub> in IECs. In addition, NF-κB plays an essential role as a transcriptional factor involved in inflammatory processes and cell apoptosis. In the resting state, NF-κB is sequestered in the cytoplasm in an inactive form by inhibitory protein IκB. In response to oxidative stress by H<sub>2</sub>O<sub>2</sub>, IκB is phosphorylated by IκB kinases (IKKs) and degraded by proteasomes (26-28). NF-κB translocates subsequently to the nucleus in an active form (26). We tried to clarify whether H<sub>2</sub>O<sub>2</sub>-induced apoptotic cell death is related to NF-κB in IECs. In stimulation by H<sub>2</sub>O<sub>2</sub>, NF-κB was activated and translocated into the nucleus. On the contrary, pretreatment with deoxyschisandrin inhibited the activation and translocation of NF-κB by H<sub>2</sub>O<sub>2</sub> through blocking degradation of IκBα. It is likely that deoxyschisandrin prevents degradation of IκBα by the ubiquitin-proteasome pathway to inhibit activation of NF-κB. Furthermore, these results support our previous *in vivo* data that SC significantly inhibited the development of UC in the mouse model (19).

When the balance between pro-inflammatory and anti-inflammatory mediators is impaired, IBD breaks out. NF-κB is a key factor in IBD including UC, because it regulates transcriptional activity for various pro-inflammatory mediators like TNF-α, interleukin-6 (IL-6), and interleukin-1 (IL-1) as well as pro-apoptotic factors. Although exact mechanisms were not identified, it has been reported that factors such as TNF act to stimulate NF-κB for regulation of cell survival and pro-apoptotic signaling in UC (29). In fact, the expression level of NF-κB subunit p65 was increased in epithelial cells of UC (30). Activation of NF-κB was significantly augmented in severe intestinal inflammation (31). Furthermore, some therapeutic drugs for IBD like sulfasalazine, methotrexate, corticosteroids and sucalfate were established to improve the anti-inflammatory effect by inhibition of NF-κB activation (32-36).

In conclusion, we focused on oxidative stress as a pathogenesis of UC. This study suggests that deoxyschisandrin may be a useful therapeutic agent for oxidative stress-induced UC through inhibiting activation of NF-κB. NF-κB is involved in transcription of diverse genes. Therefore, further studies are necessary to identify what kind of factors are mediated by inhibitory effect of deoxyschisandrin for NF-κB activation in ROS-induced oxidative stress manner in *in vivo* model.

## Acknowledgements

This work was supported by grants from Jangsu Omija Cluster in 2010.

## References

1. Werner T and Haller D: Intestinal epithelial cell signalling and chronic inflammation: From the proteome to specific molecular mechanisms. *Mutat Res* 622: 42-57, 2007.
2. Dotan I and Mayer L: Immunopathogenesis of inflammatory bowel disease. *Curr Opin Gastroenterol* 18: 421-427, 2002.
3. McCormack G, Moriarty D, O'Donoghue DP, McCormick PA, Sheahan K and Baird AW: Tissue cytokine and chemokine expression in inflammatory bowel disease. *Inflamm Res* 50: 491-495, 2001.
4. Dijkstra G, Moshage H, van Dullemen HM, *et al*: Expression of nitric oxide synthases and formation of nitrotyrosine and reactive oxygen species in inflammatory bowel disease. *J Pathol* 186: 416-421, 1998.
5. Ukil A, Maity S, Karmakar S, Datta N, Vedasiromoni JR and Das PK: Curcumin, the major component of food flavour turmeric, reduces mucosal injury in trinitrobenzene sulphonic acid-induced colitis. *Br J Pharmacol* 319: 209-218, 2003.
6. Hendrickson BA, Gokhale R and Cho JH: Clinical aspects and pathophysiology of inflammatory bowel disease. *Clin Microbiol Rev* 15: 79-94, 2002.
7. Owczarek D, Cibor D, Szczepanek M and Mach T: Biological therapy of inflammatory bowel disease. *Pol Arch Med Wewn* 119: 84-88, 2009.
8. Grisham MB: Oxidants and free radicals in inflammatory bowel disease. *Lancet* 344: 859-861, 1994.
9. Kaplan M, Mutlu EA, Benson M, Fields JZ, Banan A and Keshavarzian A: Use of herbal preparations in the treatment of oxidant-mediated inflammatory disorders. *Complement Ther Med* 15: 207-216, 2007.
10. Pravda J: Radical induction theory of ulcerative colitis. *World J Gastroenterol* 11: 2371-2384, 2005.
11. Verspaget HW, Elmgreen J, Weterman IT, Peña AS, Riis P and Lamers CB: Impaired activation of the neutrophil oxidative metabolism in chronic inflammatory bowel disease. *Scand J Gastroenterol* 21: 1124-1130, 1986.
12. Sedghi S, Fields JZ, Klamut M, *et al*: Increased production of luminol enhanced chemiluminescence by the inflamed colonic mucosa in patients with ulcerative colitis. *Gut* 34: 1191-1197, 1993.
13. Thannickal VJ and Fanburg BL: Reactive oxygen species in cell signaling. *Am J Physiol Lung Cell Mol Physiol* 279: L1005-L1028, 2000.
14. Storz P: Reactive oxygen species in tumor progression. *Front Biosci* 10: 1881-1896, 2005.
15. Kefaloyianni E, Gaitanaki C and Beis I: ERK1/2 and p38-MAPK signaling pathways, through MSK1, are involved in NF-κB transactivation during oxidative stress in skeletal myoblasts. *Cell Signal* 18: 2238-2251, 2006.
16. Daniel EH, Juan LH and Georg W: Evaluation of the antiulcer and antisecretory activity of extracts of *Aralia elata* root and *Schizandra chinensis* fruit in the rat. *J Ethnopharmacol* 23: 109-114, 1998.
17. Long ZZ and Xie SS: Experimental study on the enhancement of the immunosuppressive effect of cortisone by wuorchun, an extract of *Schizandra chinensis* Baill. *Zhong Xi Yi Jie He Za Zhi* 5: 361-362, 1985.
18. Kang CS, Tae J, Ham SH, Kim DK, Lee YM, Lee KS and Yun YG: Administration of aqueous extract of *Schizandra chinensis* fruit inhibits the experimental colitis in mice. *Natural Product Sci* 13: 78-84, 2007.

19. Zhu M, Lin KF, Yeung RY and Li RC: Evaluation of the protective effects of *Schisandra chinensis* on Phase I drug metabolism using a CCl<sub>4</sub> intoxication model. *J Ethnopharmacol* 67: 61-68, 1999.
20. Choi YW, Takamatsu S, Khan SI, Srinivas PV, Ferreira D, Zhao J and Khan IA: Schisandrene, a dibenzocyclooctadiene lignan from *Schisandra chinensis*: structure-antioxidant activity relationships of dibenzocyclooctadiene lignans. *J Nat Prod* 69: 356-359, 2006.
21. Lee IS, Jung KY, Oh SR, Park SH, Ahn KS and Lee HK: Structure-activity relationships of lignans from *Schisandra chinensis* as platelet activating factor antagonists. *Biol Pharm Bull* 22: 265-267, 1999.
22. Kim SR, Lee MK, Koo KA, *et al*: Dibenzocyclooctadiene lignans from *Schisandra chinensis* protect primary cultures of rat cortical cells from glutamate-induced toxicity. *J Neurosci Res* 76: 397-405, 2004.
23. Panossian A and Wikman G: Pharmacology of *Schisandra chinensis* Bail.: an overview of Russian research and uses in medicine. *J Ethnopharmacol* 118: 183-212, 2008.
24. Fu M, Sun ZH, Zong M, He XP, Zuo HC and Xie ZP: Deoxyschisandrins modulates synchronized Ca<sup>2+</sup> oscillations and spontaneous synaptic transmission of cultured hippocampal neurons. *Acta Pharmacol Sin* 29: 891-898, 2008.
25. Li PF, Dietz R and von Harsdorf R: Differential effect of hydrogen peroxide and superoxide anion on apoptosis and proliferation of vascular smooth muscle cells. *Circulation* 96: 3602-3609, 1997.
26. Karin M and Ben-Neriah Y: Phosphorylation meets ubiquitination: the control of NF-[kappa]B activity. *Annu Rev Immunol* 18: 621-663, 2000.
27. Oliveira-Marques V, Marinho HS, Cyrne L and Antunes F: Role of hydrogen peroxide in NF-kappaB activation: from inducer to modulator. *Antioxid Redox Signal* 11: 2223-2243, 2009.
28. Del Rio MJ and Velez-Pardo C: Monoamine neurotoxins-induced apoptosis in lymphocytes by a common oxidative stress mechanism: involvement of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), caspase-3, and nuclear factor kappa-B (NF-kappaB), p53, c-Jun transcription factors. *Biochem Pharmacol* 63: 677-688, 2002.
29. Earnshaw WC, Martins LM and Kaufmann SH: Mammalian caspases: structure, activation, substrates, and functions during apoptosis. *Annu Rev Biochem* 68: 383-424, 1999.
30. Frey MR, Edelblum KL, Mullane MT, Liang D and Polk DB: The ErbB4 growth factor receptor is required for colon epithelial cell survival in the presence of TNF. *Gastroenterology* 136: 217-226, 2009.
31. Neurath MF, Pettersson S, Meyer zum Buschenfelde KH and Strober W: Local administration of antisense phosphorothioate oligo-nucleotides to the p65 subunit of NF-kappa B abrogates established experimental colitis in mice. *Nat Med* 2: 998-1004, 1996.
32. Rogler G, Brand K, Vogl D, *et al*: Nuclear factor kappaB is activated in macrophages and epithelial cells of inflamed intestinal mucosa. *Gastroenterology* 115: 357-369, 1998.
33. Majumdar S and Aggarwal BB: Methotrexate suppressed NF-kappaB activation through inhibition of IkappaBalpha phosphorylation and degradation. *J Immunol* 167: 2911-2920, 2001.
34. Weber CK, Liptay S, Wirth T, Adler G and Schmid RM: Suppression of NF-kappaB activity by sulfasalazine is mediated by direct inhibition of IkappaB kinases alpha and beta. *Gastroenterology* 119: 1209-1218, 2000.
35. Shindo K, Iizuka M, Sasaki K, Konno S, Ito H, Horie Y and Watanabe S: Sucralfate prevents the delay of wound repair in intestinal epithelial cells by hydrogen peroxide through NF-kappaB pathway. *J Gastroenterol* 41: 450-461, 2006.
36. Kim JM, Kang HW, Cha MY, *et al*: Novel guggulsterone derivative GG-52 inhibits NF-kappaB signaling in intestinal epithelial cells and attenuates acute murine colitis. *Lab Invest* Mar 2010. (Epub ahead of print).