

Expression and localization of prominin-1 in isoproterenol-treated rat parotid gland

FABIO D'AMICO, EVANGELIA SKARMOUTSOU, KATIA MANGANO, GRAZIA MALAPONTE,
FRANCA STIVALA and MARIA CLORINDA MAZZARINO

Department of Biomedical Sciences, University of Catania, Catania, Italy

Received May 25, 2010; Accepted June 28, 2010

DOI: 10.3892/ijmm_00000492

Abstract. Prominin-1 (CD133) is a pentaspan cholesterol-binding membrane glycoprotein. Chronic treatment with isoproterenol, a β -receptor agonist, induces several dramatic effects on salivary glands, such as enhanced DNA synthesis and proliferation of salivary acinar cells. In addition, the biosynthetic pathways of membrane lipids may be altered by the isoproterenol stimulation. The purpose of this study was to investigate the effect of isoproterenol administration on prominin-1 expression profiles in rat parotid gland by means of PCR and immunohistochemistry. Rats were chronically treated with the β -adrenergic agonist for 1, 3, and 7 days. Our results showed that isoproterenol-treatment caused a down-regulation of prominin-1 on day 3 and 7 of treatment, as well as a differential immunostaining distribution pattern. This study suggests that isoproterenol-treatment may represent a useful tool to explore the molecular mechanisms involved in the synthesis and release of prominin-1. Such efforts could contribute to the development of diagnostic tools based on the detection of prominin-1 in biological fluids, such as saliva.

Introduction

Prominin-1 (CD133) is a pentaspan cholesterol-binding membrane glycoprotein belonging to the prominin family (1). Mammalian prominin-1 is broadly expressed in many embryonic and adult tissues (2-4), and it is specifically associated with plasma membrane evaginations, such as microvilli and stereocilia. Moreover, biological fluids, such as saliva and tears, are enriched in prominin-1, which is associated with small membrane vesicles (2,5-8). The functions and ligands of the prominins are still unclear and elusive, and little is known about mechanisms involved in delivery of prominin-1 to cell membrane (9). Now it is emerging that prominin-1 may have an important role in the regulation of

plasma membrane structure, maintaining an appropriate lipid composition (1). Finally, prominin-1 has been shown to be a potential marker for cancer stem cells in several tumours (10).

In mammalian salivary glands, the stimulation of β -adrenergic receptors promotes protein secretion, postnatal development and growth (11). Activation of β -adrenergic receptor causes the production of cAMP and the activation of cAMP-dependent protein kinases, stimulating several cascades, such as MAPKs (12).

It is well known that the treatment with the β -adrenergic agonist isoproterenol causes enhanced DNA synthesis and proliferation of salivary acinar cells, which leads to hyperplastic and hypertrophic gland enlargement (13,14). Furthermore, isoproterenol administration may alter the biosynthetic pathways of membrane lipids (15).

The purpose of this study was to investigate the effect of isoproterenol on prominin-1 expression profiles in rat parotid gland by means of PCR and immunohistochemistry. Rats were chronically treated with the β -adrenergic agonist for 1, 3, and 7 days.

Materials and methods

Materials. d,l Isoproterenol was obtained from Sigma Chemicals Co. (St. Louis, MO). Female Wistar rats, weighing 180-200 g each, were purchased from Harlan (Italy). Polyclonal rabbit antibody to prominin-1 (ab19898) was purchased from Abcam (UK).

Animal treatment and tissue preparation. Sets of female Wistar rats (n=5) received single daily (between 8 and 9 a.m.) intraperitoneal injections of isoproterenol at a dose of 25 mg/kg body weight, at days 0, 1, 3, and 7. Control animals received injections of 0.5 ml saline under the same regimen of treatment. Animals were kept under standard laboratory conditions (non-specific pathogen-free) with food and water *ad libitum*. Twenty-four hours after the last dose, rats were euthanized by carbon dioxide inhalation and parotid glands were excised, trimmed of fat and lymph nodes washed with PBS (pH 7.4), weighed and immediately fixed in buffered 3.7% (para) formaldehyde, pH 7.4, for 24 h. After washing and dehydration, specimens were embedded in paraffin. The handling of animals and the study protocol were in accordance with international guidelines and approved by the local Institutional Animal Care and Use Committee.

Correspondence to: Dr Fabio D'Amico, Department of Biomedical Sciences, University of Catania, via Androne 83, 95124 Catania, Italy
E-mail: f.damico@unict.it

Key words: prominin-1, CD133, isoproterenol, parotid gland, immunohistochemistry

RNA preparation and RT-PCR analysis of mRNA expression. Total RNA was isolated from paraffin sections of parotid gland tissue (10 μ m thickness) by using the PureLink FFPE Total RNA Isolation kit (Invitrogen, Carlsbad, CA, USA), according to the manufacturer's instructions. After treatment with DNase I (Invitrogen), reverse transcription reaction and PCR amplification were performed with Eppendorf AG22331 (Hamburg, Germany) thermal cycler reactor. Total RNA (1.4 μ g) was heated at 65°C for 5 min and then cooled on ice. It was combined with 0.15 μ g of random primers (Invitrogen), transcription buffer, RNase OUT (40 U/ μ l; Invitrogen), dNTPs (10 mM each) and M-MLV reverse transcriptase (200 U; Invitrogen) in a final volume of 20 μ l. Incubation was performed at 37°C for 50 min.

Amplification were performed with 3 μ l cDNA in a total volume of 50 μ l containing 200 mM Tris-HCl, pH 8.4, 500 mM KCl, 50 mM MgCl₂, 10 mM dNTP mix, 10 μ M of specific primers and 5 U/ μ l of Taq DNA polymerase (Invitrogen). Primer pairs of rat prominin-1 were: FW: 5-ATCTGGCTC CGAGTGCTTGCC-3; RV: 5-ATCATAGCCAGAGATGG ATACC-3. For semi-quantitative amplification each cycle were carried out at 93°C for 60 sec, 64°C for 60 sec, and 72°C for 60 sec. The reaction was stopped after 35 cycles. Normalisation was performed by RT-PCR amplification of glucose-6-phosphate dehydrogenase (G6PD). The sequences of these primers were: FW: 5-ACGTGATGCAGAACCACCTA CTG-3; RV: 5-ACGACGGCTGCAAAAGTGGCG-3. The amplified products were obtained as described for prominin-1. PCR product were separated on a 2% agarose gel, stained with ethidium bromide, and photographed under ultraviolet illumination.

For quantitation, gels were scanned, and the pixel intensity for each band was determined using the Quantity-One software (Bio-Rad, Richmond, CA, USA) and normalized to the amount of glyceraldehyde-3-phosphate dehydrogenase (GAPDH). All values were determined in triplicate. Negative controls consisting of RNase-free water substituted for template cDNA, were included to ensure no crossover DNA contamination.

Immunohistochemistry. Five- μ m-thick sections from paraffin-embedded tissue blocks were placed on silanized slides. After section deparaffinization and rehydration through a graded ethanol series at room temperature, antigen retrieval was performed in citrate buffer (pH 6.0) for 30 min at 95°C using a water bath. Then, sections were treated with 0.3% hydrogen peroxide for 15 min at room temperature to quench the endogenous peroxidase activity, and incubated with diluted normal blocking serum for 20 min at room temperature. Subsequently, the sections were incubated with rabbit polyclonal primary antibodies anti-prominin-1, diluted 1:200, for 1 h at room temperature. After washing with phosphate buffered saline, the detection was carried out using the streptavidin-biotin-peroxidase complex system (Immuno Cruz™ Staining System; SantaCruz Biotechnology Inc., Santa Cruz, CA, USA), according to the manufacturer's instructions. Finally, sections were counterstained with Mayer's hematoxylin. Slides were mounted and observed by conventional light microscopy. Negative control experiments were performed substituting the primary antibody with non immune serum and by omitting the primary antibody, thus




Figure 1. Changes in weight of parotid glands during the isoproterenol treatment. The most striking increase occurs between 0 and 3 days of treatment. Each value was statistically significant ($p < 0.05$). Values are expressed as mean \pm SD.

incubating slides only with buffer. For standard histological evaluation, sections were also stained with hematoxylin-eosin.

Image analysis. Immunolabelled and random systematically sampled sections were observed using a Leica DMRB microscope (Leica, Wetzlar, Germany) and photographed with a Canon G-9 camera (Canon, Japan). All images were obtained using a x40 objective. A total of 30 fields were analyzed using ImageJ software, version 1.42g (16). Immunolabelling intensity was analyzed with the colour deconvolution plug-in (17). Subsequently, the immunostained fraction area (fa) was used (percent area) according the following formula:

$$Fa = \frac{\sum Sa}{\sum A}$$

where Sa is the stained area, A is the total area of the image field.

Statistical analysis. For mass gland measurements, semi-quantitative RT-PCR and immunohistochemical analysis, statistics was performed by using the unpaired Student's t-test. Differences were considered statistically significant at a level of $p < 0.05$.

Results

Parotid mass increase and histological findings of the parotid gland during isoproterenol treatment. As shown in Fig. 1, parotid gland wet weight markedly increased from 0 to 7 days of treatment. The overall increase was of 196% ($p < 0.01$). Mass increase between 0 and 1 day was approximately 97%, between 1 and 3 days was about 35%, and between 3 and 7 days was about 11% ($p < 0.05$). The most striking increase in weight was observed between 0 and 3 days of treatment.




Figure 2. Representative micrographs of histological changes in normal (a; bar: 100 μ m) and isoproterenol parotid glands (b and c; bars: 60 μ m) stained with hematoxylin and eosin. The size of acinar cells is increased by isoproterenol treatment at day 1 and 7 (b and c, respectively).




Figure 3. Expression of prominin-1 mRNA in normal and isoproterenol-treated rat. Representative blots (a) and densitometric quantification of mRNA expression normalized by GAPDH levels (b). Isoproterenol treatment induces a decrease of prominin-1 mRNA expression from day 3 of treatment. Histogram bars correspond to treatment days 0, 1, 3, and 7, respectively. Values are expressed as mean \pm SEM. * P <0.05 vs. control.

Fig. 2 shows a representative morphological comparison between H&E stained sections of normal parotid gland tissue and isoproterenol-treated glands. The main effect of isoproterenol was the enlargement of the acinar cells, because of the increased number of secretory granules.

Semiquantitative analysis of mRNA expression of prominin-1 in normal and isoproterenol treated rat parotid gland. The results of PCR assay are summarized in Fig. 3. The levels of mRNA were given as relative amounts with normalization against GAPDH. Expression of prominin-1 was revealed in normal parotid glands as well as in isoproterenol-treated tissues. However, there was a time-dependent and inhibitory

effect of isoproterenol treatment on prominin expression. In fact, a significant decrease of prominin-1 expression to approximately 84% and 70% of control values (p <0.05) was observed in treated rats after 3 and 7 days of treatment, respectively (p <0.05).

Immunohistochemical localization of prominin-1 in normal and isoproterenol treated rat parotid gland. Immunohistochemical localization of prominin-1 in normal and isoproterenol-treated rat parotid glands is shown in Fig. 4. Both untreated and treated glands express prominin-1. In normal parotid, prominin-1 was mainly localized on the apical and lateral surfaces of acinar cells (Fig. 4a), as well as in their cytoplasm. In addition, ductal epithelial cells showed a discrete immunostaining at the apical membrane facing the lumen (Fig. 4b). On day 1 of treatment with isoproterenol, acinar cells showed a similar immunostaining pattern in comparison to the untreated tissue: prominin-1 was observed at cytoplasmic level and, although less frequently, over apical and lateral cell membrane domains (Fig. 4c). Ducts epithelial cells maintained their characteristic staining pattern as in untreated glands (inset of Fig. 4c). On days 3 and 7 of treatment (Fig. 4d and e, respectively), parotid acinar cells exhibited a discrete immunostaining in their secretory granules, localized on their membranes (Fig. 4d). However, at 7 days of treatment the duct luminal surface seemed to be unstained for the protein (Fig. 4e). Appropriate negative controls showed no immunoreaction (Fig. 4f).

As showed in Fig. 5, semi-quantitative image analysis of immunohistochemical results showed that isoproterenol treatment induced a decreased expression of prominin-1. In particular, significant decreases (p <0.05) of prominin-1 expression were observed after 3 and 7 days of treatment (42% and 52% of control value, respectively).

Discussion

We described here the expression profiles of prominin-1 in normal and isoproterenol-treated rat parotid gland. The main localization of prominin-1 in the apical and basal membrane domains of acinar cells, as well as, in membrane of ductal epithelial cells facing the lumen, is in accordance with the known topological occurrence of prominin-1 in various plasmalemmal protrusions (2). It is known, in fact, that apical and lateral free surfaces of acinar cells of salivary gland




Figure 4. Immunolocalization of prominin-1 in normal and isoproterenol-treated rat parotid gland. In normal rat parotid gland, prominin-1 is mainly localized on the apical and lateral zones of acinar cells. In addition, the cytoplasm of acinar cells is immunolabelled (a). Ductal epithelial cells of untreated gland show discrete immunostaining at the apical membrane facing the lumen (b). After 1 day of treatment with isoproterenol, acinar cells show a similar staining pattern in comparison to the untreated tissue: a cytoplasmic staining for the prominin-1, and, although less frequently, a staining over the apical and lateral zones of acinar cells (c). The ducts maintain their characteristic staining pattern as in untreated glands (c, inset). At days 3 and 7 of treatment (d and e, respectively), prominin-1 immunostaining is mainly localized over membranes of secretory granules in acinar cells. However, at 7 days of treatment the duct luminal surface seems to be unstained for the protein under study (e). A representative micrograph of a negative control in untreated tissue is shown (f). [bars: 50 μ m, except for inset (25 μ m)].




Figure 5. Semi-quantitative image analysis of prominin-1 immunolabelling (area density) in normal and isoproterenol treated rat parotid gland. Isoproterenol treatment induces a decreased expression of prominin-1. Significant decreases are observed after 3 and 7 days of treatment (approximately 42% and 52% of control value, respectively). Histograms bars correspond to treatment days 0, 1, 3, and 7, respectively. Values are expressed as mean and standard errors. * $p < 0.05$ vs. control.

contain such protrusions (18). In particular, prominin-1 has been recently immunolocalized on human and mouse parotid glands (19,20). Furthermore, many other duct epithelia, such as those of pancreas, mammary gland lactiferous and epididymal ducts, contain this molecule (21,22). The major finding of this study was that isoproterenol-treatment caused a time-dependent downregulation of prominin-1 mRNA and protein, as well as, a different immunostaining distribution pattern.

It is known that, among the dramatic effects of isoproterenol administration, there are an alteration of the biosynthetic pathways of membrane lipids (15), as well as, a cell damage in rodent salivary glands, with an appearance of large lipid droplets within cells (23). Since it is known that sorting of prominin-1 involves a cholesterol-based lipid raft (24), and that β -adrenergic receptors are dynamically associated with membrane rafts (25), the disturbing effect of isoproterenol on lipid biochemistry could result in the decreased transcription and expression of prominin-1. On days 3 and 7 of treatment with the β -agonist, prominin-1 is mainly localized over secretory granule membranes of acinar



SPANDIDOS PUBLICATIONS

h induced change in protein distribution could be the isoproterenol treatment, which would alter the mechanisms of the regulated secretion in rat parotid glands, with a consequent routing of apical secretion pathway to the basolateral one (26). Moreover, changes in cell polarity have been shown to occur in salivary acinar cell treated with isoproterenol (27), and prominin-1 has been suggested to influence cell polarity (9). Finally, experimentally induced changes in lipid composition, such as in cholesterol, was found alter the release of membrane vesicle containing prominin-1 in Caco-2 cells (7,28).

In addition to differentiated cells, the turnover of salivary glands may be ensured by a stem cell population, and prominin-1 has been shown to be a marker for stem cell in several tissues (5,29-31). However, our study shows that prominin-1 is not restricted to a single cell population and its expression is broadly localized among the acinar and ductal cells. Thus, we can conclude that prominin-1 is not a suitable marker for stem cell in parotid gland. For this reason other molecules, such as keratin 18 (K18), α -smooth muscle actin (α -SMA), achaete scute homolog like 3 (Ascl3), DNp63, PSCA (prostate stem cell antigen), stem cell antigen-1 (Sca-1), c-Kit, Musashi-1, are more suitable markers or promising candidates for stem cell population in salivary glands (32-35). However, the differences of staining patterns of prominin-1 between normal and isoproterenol-treated glands may reflect a different stage of differentiation induced by the β -agonist.

In conclusion, our study suggests the isoproterenol treatment may represent a useful tool to explore the molecular mechanisms involved in the synthesis and release of the membrane vesicles containing prominin-1. Such efforts could contribute to the development of diagnostic tools based on the detection of prominin-1 in biological fluids, such as saliva. In fact, recent investigations suggest the presence of an association between the release of prominin-1 from cells and several diseases (1).

References

- Corbeil D, Marzesco AM, Fargeas CA and Huttner WB: Prominin-1: a distinct cholesterol-binding membrane protein and the organisation of the apical plasma membrane of epithelial cells. *Subcell Biochem* 51: 399-423, 2010.
- Weigmann A, Corbeil D, Hellwig A and Huttner WB: Prominin, a novel microvilli-specific polytopic membrane protein of the apical surface of epithelial cells, is targeted to plasmalemmal protrusions of non-epithelial cells. *Proc Natl Acad Sci USA* 94: 12425-12430, 1997.
- Corbeil D, Röper K, Hellwig A, Tavian M, Miraglia S, Watt SM, Simmons PJ, Peault B, Buck DW and Huttner WB: The human AC133 hematopoietic stem cell antigen is also expressed in epithelial cells and targeted to plasma membrane protrusions. *J Biol Chem* 275: 5512-5520, 2000.
- Corbeil D, Joester A, Fargeas CA, Jászai J, Garwood J, Hellwig A, Werner HB and Huttner WB: Expression of distinct splice variants of the stem cell marker prominin-1 (CD133) in glial cells. *Glia* 57: 860-874, 2009.
- Miraglia S, Godfrey W, Yin AH, Atkins K, Warnke R, Holden JT, Bray RA, Waller EK and Buck DW: A novel five-transmembrane hematopoietic stem cell antigen: isolation, characterization, and molecular cloning. *Blood* 90: 5013-5021, 1997.
- Fargeas CA, Fonseca AV, Huttner WB and Corbeil D: Prominin-1 (CD133) - from progenitor cells to human diseases. *Future Lipidol* 1: 213-225, 2006.
- Marzesco AM, Wilsch-Bräuninger M, Dubreuil V, Janich P, Langenfeld K, Thiele C, Huttner WB and Corbeil D: Release of extracellular membrane vesicles from microvilli of epithelial cells is enhanced by depleting membrane cholesterol. *FEBS Lett* 583: 897-902, 2009.
- Dubreuil V, Marzesco AM, Corbeil D, Huttner WB and Wilsch-Bräuninger M: Midbody and primary cilium of neural progenitors release extracellular membrane particles enriched in the stem cell marker prominin-1. *J Cell Biol* 176: 483-495, 2007.
- Shmelkov SV, St Clair R, Lyden D and Rafii S: AC133/CD133/Prominin-1. *Int J Biochem Cell Biol* 37: 715-719, 2005.
- Mizrak D, Brittan M and Alison MR: CD133: molecule of the moment. *J Pathol* 214: 3-9, 2008.
- Barka T: β -Adrenergic effects on salivary glands: growth and gene regulation. *Acta Histochem Cytochem* 23: 245-255, 1990.
- Quissell DO, Watson E and Dowd F: Signal transduction mechanisms involved in salivary gland regulated exocytosis. *Crit Rev Oral Biol Med* 3: 83-107, 1992.
- Selye HR, Veilleux R and Cantin M: Excessive stimulation of salivary gland growth by isoproterenol. *Science* 133: 44-45, 1961.
- Lillie JH and Han SS: Secretory protein synthesis in the stimulated rat parotid gland. Temporal dissociation of the maximal response from secretion. *J Cell Biol* 59: 708-721, 1973.
- Yashiro K, Kameyama Y, Mizuno M and Yokota Y: Alteration of membrane phospholipids in hypertrophied rat salivary glands induced by chronic administration of isoproterenol. *Arch Oral Biol* 32: 799-805, 1987.
- Rasband WS: ImageJ, US National Institutes of Health, Bethesda, MD, USA, <http://rsb.info.nih.gov/ij/>, 1997-2009.
- Ruifrok AC and Johnston DA: Quantification of histochemical staining by color deconvolution. *Anal Quant Cytol Histol* 23: 291-299, 2001.
- D'Amico F and Skarmoutsou E: Immunolocalization of E-cadherin and α E-catenin in rat parotid acinar cell under chronic stimulation of isoproterenol. *Arch Oral Biol* 52: 161-167, 2007.
- Jászai J, Janich P, Farkas LM, Fargeas CA, Huttner WB and Corbeil D: Differential expression of Prominin-1 (CD133) and Prominin-2 in major cephalic exocrine glands of adult mice. *Histochem Cell Biol* 128: 409-419, 2007.
- Karbanová J, Missol-Kolka E and Fonseca AV: The stem cell marker CD133 (Prominin-1) is expressed in various human glandular epithelia. *J Histochem Cytochem* 56: 977-993, 2008.
- Fargeas CA, Joester A, Missol-Kolka E, Hellwig A, Huttner WB and Corbeil D: Identification of novel prominin-1/CD133 splice variants with alternative C-termini and their expression in epididymis and testis. *J Cell Sci* 117: 4301-4311, 2004.
- Lardon J, Corbeil D, Huttner WB, Ling Z and Bouwens L: Stem cell marker prominin-1/AC133 is expressed in duct cells of the adult human pancreas. *Pancreas* 36: E1-6, 2008.
- Simson JA: Evidence of cell damage in rat salivary glands after isoproterenol. *Anat Rec* 173: 437-451, 1972.
- Roper K, Corbeil D and Huttner WB: Retention of prominin in microvilli reveals distinct cholesterol-based lipid microdomains in the apical plasma membrane. *Nat Cell Biol* 2: 582-592, 2000.
- Willoughby D and Cooper DM: Organization and Ca^{2+} regulation of adenylyl cyclases in cAMP microdomains. *Physiol Rev* 87: 965-1010, 2007.
- Venkatesh SG, Tan J, Gorr SU and Darling DS: Isoproterenol increases sorting of parotid gland cargo proteins to the basolateral pathway. *Am J Physiol Cell Physiol* 293: C558-C565, 2007.
- Tamaki H and Yamashina S: Changes in cell polarity during mitosis in rat parotid acinar cells. *J Histochem Cytochem* 39: 1077-1087, 1991.
- Hansen GH, Niels-Christiansen LL, Thorsen E, Immerdal L and Danielsen EM: Cholesterol depletion of enterocytes. Effect on the Golgi complex and apical membrane trafficking. *J Biol Chem* 275: 5136-5142, 2000.
- Yin AH, Miraglia S, Zanjani ED, Almeida-Porada G, Ogawa M, Leary AG, Olweus J, Kearney J and Buck DW: AC133, a novel marker for human hematopoietic stem and progenitor cells. *Blood* 90: 5002-5012, 1997.

30. Rizzo S, Attard G and Hudson DL: Prostate epithelial stem cells. *Cell Prolif* 38: 363-374, 2005.
31. Corti S, Nizzardo M, Nardini M, Donadoni C, Locatelli F, Papadimitriou D, Salani S, Del Bo R, Ghezzi S, Strazzer S, Bresolin N and Comi GP: Isolation and characterization of murine neural stem/progenitor cells based on Prominin-1 expression. *Exp Neurol* 205: 547-562, 2007.
32. Kawanami T, Matsuzaki Y, Sawaki T, Sakai T, Jin ZX, Masaki Y, Fukushima T, Tanaka M and Umehara H: Identification of human salivary stem cells from cultured labial minor salivary cells. *Nihon Rinsho Meneki Gakkai Kaishi* 30: 455-460, 2007.
33. Bullard T, Koek L, Roztocil E, Kingsley PD, Mirels L and Ovitt CE: Ascl3 expression marks a progenitor population of both acinar and ductal cells in mouse salivary glands. *Dev Biol* 320: 72-78, 2008.
34. Kimoto M, Yura Y, Kishino M, Toyosawa S and Ogawa Y: Label-retaining cells in the rat submandibular gland. *J Histochem Cytochem* 56: 15-24, 2008.
35. Lombaert IM, Brunsting JF, Wierenga PK, Faber H, Stokman MA, Kok T, Visser WH, Kampinga HH, de Haan G and Coppes RP: Rescue of salivary gland function after stem cell transplantation in irradiated glands. *PLoS One* 30: E2063, 2008.