

Enhanced enteric invasion of scrapie agents into the villous columnar epithelium via maternal immunoglobulin

YASUHISA ANO^{1,2}, AKIKAZU SAKUDO³, RYUTA URAKI¹, YUKITA SATO⁴, JYURI KONO⁴,
KATSUAKI SUGIURA⁵, TAKASHI YOKOYAMA⁶, SHIGEYOSHI ITOHARA⁷,
HIROYUKI NAKAYAMA¹, MASAYOSHI YUKAWA⁴ and TAKASHI ONODERA¹

¹Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657;

²Central Laboratories for Frontier Technology, Kirin Holdings Co., Ltd., 1-13-5 Fukuura Kanazawa-ku, Yokohama,

Kanagawa 236-0004; ³Research Institute for Microbial Diseases, Osaka University, Yamadaoka, Suita,

Osaka 565-0871; ⁴Department of Veterinary Medicine, College of Biosource Sciences, Nihon University,

Kameino 1866, Fujisawa, Kanagawa 252-8510; ⁵Food and Agricultural Materials Inspection Centre,

Saitama 330-9731; ⁶Prion Disease Research Center, National Institute of Animal Health, Ibaraki 305-0856;

⁷Laboratory for Behavioral Genetics, Brain Science Institute, RIKEN, Wako, Saitama 351-0198, Japan

Received June 22, 2010; Accepted July 31, 2010

DOI: 10.3892/ijmm_00000533

Abstract. Transmissible spongiform encephalopathies (TSE) are caused by dietary oral exposure to infectious prion proteins (PrP^{Sc}); however, the mechanism behind the uptake of PrP^{Sc} in the intestines is poorly understood. In addition, epidemiological studies of BSE showed that most cattle are exposed to the agents in the first 6 months of life, during the suckling and weaning periods. In the present study, to elucidate the enteric invasion mechanism of prions and to investigate the age-dependent transmission mechanism suggested by epidemiological studies, wild-type and SCID mice were orally administered brain homogenate from scrapie (Tsukuba 1)-infected mice during the suckling and weaning stages, before being analyzed histopathologically. PrP^{Sc} was found to be incorporated into the villous columnar epithelial cells and was also detected in the villous lacteal of 15-day-old suckling mice. However, no such uptake of PrP^{Sc} was observed in the weaned mice at 25-days-old. Four different strains of mice were tested. There was no mouse strain difference in the frequency of PrP^{Sc} positive columnar epithelial cells. In addition, the uptake of PrP^{Sc} in suckling SCID mice lacking maternal antibodies was significantly lower than that in the wild-type suckling mice, and the uptake of PrP^{Sc} was enhanced

by dilution with purified IgG. In the present study, it was suggested that the weaning period and maternal immunoglobulin are important risk factors for the oral transmission of PrP^{Sc}.

Introduction

Transmissible spongiform encephalopathies (TSE) are a group of fatal neurodegenerative diseases characterized by the abundant accumulation of abnormal prion proteins (PrP^{Sc}) with β -sheet structures (PrP^{Sc}) (1). The most probable entry site of PrP^{Sc} is the intestinal tract. After entry, the transmissible agents pass through one or several biological barriers and finally reach the brain (2). The peroral route of entry is widely assumed to be the most important in the natural pathogenesis of bovine spongiform encephalopathy (BSE), scrapie (3), variant Creutzfeldt-Jakob disease (vCJD) (4) and other TSE (5). Such transmission has also been shown to be caused by dietary exposure to PrP^{Sc}-contaminated food; however, the uptake process and the movement of infectious agents from the intestines to the central nervous system (CNS) are poorly understood.

Experimental models of the oral transmission of BSE agent in mink (6), mice (7), sheep (8,9) and non-human primates (10,11) and experimental models of the oral transmission of scrapie agent in hamsters (12,13) and mice (14) have been established. These models focused on the accumulation of PrP^{Sc} in gut-associated lymphoid tissue (GALT) and the neuro-invasion of PrP^{Sc} into the peripheral nervous system. Invasion through the intestinal epithelial barrier is the first critical step in oral transmission, but the mechanism of intestinal epithelial invasion by PrP^{Sc} is poorly understood.

PrP^{Sc} was shown to accumulate around the follicular dendritic cells (FDC) of GALT and in tangible-body macrophages of lymphoid nodules (15). PrP^{Sc}-positive cells with the morphology of dendritic cells and macrophages were also found to be scattered throughout the dome region of intestinal

Correspondence to: Professor Takashi Onodera, Department of Molecular Immunology, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Bunkyo-ku, Tokyo 113-8657, Japan

E-mail: aonoder@mail.ecc.u-tokyo.ac.jp

Key words: columnar epithelial cells, maternal immunoglobulin, prion protein, SCID mice, scrapie, suckling period

Peyer's patches (16). This suggests that M cells in the follicle-associated epithelium are the entry site of transmissible agents (17); however, no reports on the entry sites from the gut lumen or the distribution of PrP^{Sc} in the early stage of infection have been published.

There is a report dealing with PrP^{Sc} transport across the intestinal mucosa in sheep that had been operated on to form an intestinal loop (18) and inoculated with PrP^{Sc} into the lumen of the loop. The report shows that trans-epithelial passage through the intact villous epithelium was more probable than that through M cells in Peyer's patches.

In addition, epidemiological studies and simulation models of BSE showed that most cattle are exposed to the agents in the first 6 months of life and that younger cattle are more likely to be infected than older cattle (19). The first 6 months of life in cattle represent the suckling and weaning periods, when the gastrointestinal tracts are still immature. During the suckling period, the intestinal epithelium easily takes up proteins such as immunoglobulins and growth factors from milk. Therefore, the intestinal epithelium may play a role in the incorporation of PrP^{Sc} during the suckling period.

We have already demonstrated that amyloid- β protein is incorporated into the villous epithelium during the suckling and weaning periods in mice (20) and cattle (21). In the present study, to elucidate the enteric invasion mechanism of prions and examine the age-dependent transmission mechanism suggested by previous epidemiological studies, scrapie agents were orally administered to mice during the suckling or weaning period, and then the mice were subjected to histopathological analysis.

Materials and methods

Experimental animals. Fifteen-, 20- and 25-day-old CD-1, BALB/c and C57BL/6 mice and 15-day-old CD-1.SCID mice (Japan CLEA, Tokyo, Japan) were housed in SPF conditions under an alternating 14 h/10 h light/dark cycle. The animals were given free access to standard laboratory food (Oriental Yeast Co., Ltd., Tokyo, Japan) and tap water and were treated in accordance with the procedures authorized by the Animal Experiment Committee of Nihon University College of Biosource Sciences.

Administration of PrP^{Sc} and preparation of tissue specimens.

To observe the differences in uptake between the suckling and weaning periods, mice (age, 15, 20 or 25 days; $n=3$ for each age and each administered substance) were administered with mouse brain homogenates from mice in the terminal stage of infection with mouse-adapted scrapie (Tsukuba 1 strain) (22). Ten percent brain homogenates (w/v) of scrapie infected or normal mouse brains in sterile PBS were prepared by a previously described method (23), and then 15-, 20- and 25-day-old mice were orally administered the brain homogenates. One gram of brain material was positive by Western blotting until 1:1000000 dilution. The administration was repeated 3 h later, and at 1 h post-administration (p.a.) the mice were euthanized with ether. The administered amount for the 15-day-old mice was 100 μ l and those of the 20- and 25-day-old mice were determined in proportion to the weight of each mouse. Next, to elucidate the involvement of maternal

immunoglobulin in the intestinal invasion of scrapie agents, 15-day-old suckling SCID mice were administered 100 μ l of 10% homogenate (w/v) composed of scrapie-infected brains diluted with PBS ($n=5$) or diluted with PBS containing 5 mg/ml purified mouse IgG (Beckman, Fullerton, CA, USA, $n=3$) or normal mouse brains diluted with PBS ($n=3$). The administration was repeated 3 h later, and at 1 h post-administration (p.a.) the mice were euthanized with ether. Their intestines were then removed and fixed by immersion in PBS containing 4% paraformaldehyde for 2 h before being washed in PBS containing 6.8% sucrose. After dehydration in 100% acetone for 1 h, tissue samples were embedded in resin (Technovit 8100; Heraeus Kulzer, Wehrheim, Germany) in accordance with the manufacturer's instructions and sectioned at a thickness of 4 μ m.

Identification of PrP^{Sc}. PrP^{Sc}-deposition in the brains of mice at the terminal stage of Tsukuba 1 infection was confirmed using rabbit anti-PrP polyclonal antibody (P8, 1:100), (24) and mouse anti-PrP monoclonal antibody (T2, 10 μ g/ml), (25) using a conventional immunohistochemical procedure (26). The PrP^{Sc} in the intestines was identified by immunohistochemistry. Resin sections obtained from mice treated with PrP^{Sc} were pretreated with 0.1% CaCl₂ at pH 7.8 containing 0.01% trypsin for 10 min at 37°C and were then quenched in 0.3% hydrogen peroxide in methanol for 30 min. After incubation with rabbit anti-PrP polyclonal antibody (P8, 1:100) or mouse anti-PrP monoclonal antibody (T2, 10 μ g/ml) at 37°C for 2 h and secondary horseradish peroxidase-coupled goat anti-rabbit IgG antibody or goat anti-mouse IgG antibody (4 μ g/ml; Nichirei, Tokyo, Japan), respectively, at room temperature for 30 min, diaminobenzidine (DAB; Wako, Osaka, Japan) was applied for 10 min. The sections were then counterstained with hematoxylin for 1 min. The number of PrP^{Sc} positive cells in each microscope-visual field in the villous epithelium was counted at five random points. Cell counts were expressed as the mean \pm SD of the microscopic fields viewed at x400 magnification. Intestinal epithelial cells were selected for determining the intensity of infection, which was not known to the observer measuring the respective intestinal sections. Statistical analysis ($P<0.01$) was performed to assess the differences between groups using the Student's *t*-test.

Lectin staining to identify the targeted cells. To analyze the PrP^{Sc} positive cells, the resin-fixed sections were treated with PBS containing 1% BSA for 30 min. Then, they were incubated with rabbit anti-PrP polyclonal antibody (P8, 1:100) for 2 h at 37°C and secondary Alexa Fluor® 546-coupled goat anti-rabbit IgG antibody (5 μ g/ml Molecular Probes, Eugene, OR, USA). Ulex europaeus agglutinin (UEA-1) conjugated to rhodamine (Vector Labs, Burlingame, CA, USA) and wheat germ agglutinin (WGA) conjugated to Alexa Fluor® 350 (Molecular Probes) were used. The sections were incubated at room temperature for 1 h with the lectins (10 μ g/ml).

Results

Detection of PrP^{Sc} using two anti-PrP antibodies. Scrapie agents of the Tsukuba 1 strain were successfully detected




Figure 1. Detection of PrP^{Sc} by two anti-PrP antibodies. Mouse brains from CD-1 mice at the terminal stage of infection with scrapie prion [Tsukuba 1 strain, (a-c)] and normal CD-1 mouse brains (d-f). PrP^{Sc} was detected by immunohistochemistry using rabbit anti-PrP polyclonal antibody [P8, (a, d)] and mouse anti-PrP monoclonal antibody [T2, (b, e)], as well by an immunofluorescent assay using P8 (red) and T2 (green) antibodies. PrP^{Sc} was detected (a, b), and the same PrP^{Sc} was identified by both antibodies (c).




Figure 2. Uptake of PrP^{Sc} through the villi. The duodenum (a), jejunum (b-e), and Peyer's patch (f) in 15-day-old CD-1 mice. The mice were orally administered the brain homogenates of scrapie (Tsukuba 1) infected mice (a-d, f) and normal mice (e). PrP^{Sc} was incorporated into the villous epithelial cells in the jejunum [arrows, (b, d)] but not into those of the duodenum (a). PrP^{Sc} was also detected in the dome epithelium of Peyer's patches (f) and in the villous lacteal (f, arrows). PrP^{Sc} was detected by P8 (b) and T2 (d) anti-prion protein antibodies, whereas no PrP was detected in the CD-1 mice administered a homogenate composed of normal CD-1 mouse brains. The lectins used were UEA-1 conjugated with rhodamine and WGA conjugated with Alexa Fluor® 350. The luminal surface of the columnar epithelial cells in the villus is shown in blue (g, h). The columnar cells (arrows) took up PrP^{Sc} (green). The scale bars represent 50 μ m (a-f, g and h) and 20 μ m (magnification of f).

using rabbit anti-mouse PrP polyclonal antibody (P8, Fig. 1a) and mouse anti-PrP monoclonal antibody (T2, Fig. 1b) in the brains of CD-1 mice at the terminal stage of scrapie-infection. These two antibodies identified the same PrP^{Sc} (Fig. 1c). In the normal CD-1 mouse brains, no PrP^{Sc} was detected (Fig. 1d-f).

Incorporation through the villi. PrP^{Sc} was detected by immunohistochemistry using both rabbit anti-PrP polyclonal antibody (P8, Fig. 2a-c) and mouse anti-PrP monoclonal antibody (T2, Fig. 2d). PrP^{Sc} was not incorporated into the villous epithelium in the duodenum (Fig. 2a) but was present




Figure 3. Age-dependent uptake of PrP^{Sc}. Intestinal villi of 15- (a), 20- (b) and 25-day-old mice (c) that had been orally administered PrP^{Sc}. PrP^{Sc} was incorporated readily into the villi of the 15-day-old CD-1 mice, incorporated slightly in the 20-day-old CD-1 mice, but not incorporated at all in the 25-day-old CD-1 mice. The number of ileal epithelial cells that had incorporated PrP^{Sc} was significantly higher in the 15-day-old CD-1 mice than in the 20- or 25-day-old CD-1 mice (d). The results are expressed as the mean \pm SD. The statistical significance of differences was determined by the Student's t-test. **P<0.01. Scale bar, 50 μ m.

in the jejunum (Fig. 2b and d, arrows) of 15-day-old CD-1 mice and was also detected in the villous lacteal (Fig. 2c, arrow). On the other hand, PrP was not detected in the CD-1 mice orally administered the homogenate of normal mice brains (Fig. 2e). In addition, the sections that were only reacted with the secondary antibodies did not show PrP^{Sc} on immunohistochemistry. In the Peyer's patches, some PrP^{Sc} was detected in the dome epithelium (Fig. 2f, arrows) but none was found in the subepithelial dome or germinal center regions.

Incorporation across absorptive epithelial cells. Four kinds of cells were identified in the intestinal villous epithelium of the 15-day-old CD-1 mice: UEA-1-positive and WGA-negative (UEA-1⁺/WGA⁻), UEA-1-positive and WGA-positive (UEA-1⁺/WGA⁺), UEA-1-negative and WGA-positive (UEA-1⁻/WGA⁺), and UEA-1-negative and WGA-negative (UEA-1⁻/WGA⁻) cells (data not shown). PrP^{Sc} was detected in the UEA-1⁺/WGA⁺ cells (Fig. 2g and h).

Age-dependent incorporation. To determine whether aged mice were less reactive to scrapie agent ingestion, intestinal sections were observed using immunohistochemistry. PrP^{Sc} was incorporated into the villous epithelium and the lamina propria in the 15-day-old CD-1 mice (Fig. 3a). On the other hand, PrP^{Sc} was attached to the luminal surface of villous epithelial cells but was scarcely detected in the villous columnar epithelial cells in the 20-day-old CD-1 mice (Fig. 3b)

and was not found at all in the 25-day-old CD-1 mice (Fig. 3c). Significantly more PrP^{Sc} was incorporated into the villous epithelium in the 15-day-old CD-1 mice than in the 20-day-old CD-1 mice and in the 20-day-old CD-1 mice compared with the 25-day-old CD-1 mice (Fig. 3d).

The same experiments were repeated in BALB/c and C57BL/c mice to see if there were any mouse strain differences in the intestinal incorporation of PrP^{Sc}. As shown in Fig. 4, more PrP^{Sc} was incorporated into the villous epithelium in the 15-day-old BALB/c and C57BL/6 mice than in the 20-day-old BALB/c and C57BL/6 mice (Fig. 4b and c, respectively). More PrP^{Sc} was observed in the villous epithelium in the 20-day-old BALB/c and C57BL/6 mice than in the 25-day-old BALB/c and C57BL/c mice. There were no distinct mouse strain differences in the ratio of PrP^{Sc} incorporation among the CD-1 (Fig. 4a), BALB/c (Fig. 4b) and C57BL/6 (Fig. 4c) mice.

Differences in PrP^{Sc} uptake between suckling wild-type and SCID mice. To determine whether immunodeficient mice were less reactive to scrapie agent ingestion, intestinal sections from CD-1.SCID mice were observed using immunohistochemistry. PrP^{Sc} uptake into the villous epithelial cells was observed in the CD-1.SCID mice (Fig. 5b). Compared with the wild-type CD-1 mice (Fig. 5a), the number of cells incorporating PrP^{Sc} in the CD-1.SCID mice was significantly lower (Fig. 5e). In the Peyer's patches, PrP^{Sc} was incorporated into the dome epithelial cells (Fig. 5d).




Figure 4. Age-dependent uptake of PrP^{Sc} in CD-1, BALB/c and C57BL/6 mice. PrP^{Sc} was incorporated readily into the villi of the 15-day-old mice, incorporated slightly in the 20-day-old mice, but not incorporated at all in the 25-day-old mice [CD-1, (a); BALB/c, (b); C57BL/6, (c)]. The number of ileal epithelial cells that incorporated PrP^{Sc} was significantly higher in the 15-day-old mice than in the 20- or 25-day-old mice. The results are expressed as the mean \pm SD. The statistical significance of differences was determined by the Student's t-test. **P<0.01. Scale bar, 50 μ m.

On the contrary, the PrP^{Sc} uptake into the villi in the CD-1.SCID mice administered PrP^{Sc} containing abundant mouse IgG (Fig. 5c) was enhanced compared with that in CD-1.SCID mice administered only PrP^{Sc} (Fig. 5b). As a control, mouse IgG was given orally to mice without PrP^{Sc}. Mouse intestinal sections were stained for PrP^{Sc} using anti-PrP polyclonal rabbit antibody (P8) and secondary stained

with anti-rabbit IgG by conventional immunohistochemistry (26). In these sections, PrP^{Sc} was not detected using anti-PrP polyclonal antibody and no cross-reactive staining was observed for the ingested mouse IgG using anti-rabbit IgG polyclonal secondary antibody (Fig. 5f).

Discussion

PrP^{Sc} was detected after being incorporated by villous epithelial cells. PrP^{Sc} is resistant to degradation by gastric juices and intestinal enzymes because of its abundant stable β -sheet structure. It was revealed previously that murine and bovine β -amyloid proteins, which also contain many β -sheet structures, were incorporated into villous columnar epithelial cells in mice (20) and cows (21). These proteins are able to resist digestion by gastric juices and intestinal enzymes in a similar manner to amyloid proteins such as PrP^{Sc} and are incorporated into the villous epithelial cells by a common mechanism.

Villous epithelial cells are composed of absorptive, goblet, endocrine, basal granular and Paneth cells (27). Among the murine intestinal epithelial cells, the UEA-1⁺/WGA⁻, UEA-1⁺/WGA⁺, UEA-1⁻/WGA⁺ and UEA-1⁻/WGA⁻ cells correspond to the M, goblet, columnar, and other cells (e.g., endocrine cells), respectively (28). The villous epithelium of 15-day-old suckling mice is composed of columnar, goblet and other types of cells. The dome epithelium in Peyer's patches mainly consists of M cells. In the present experiment, PrP^{Sc} was incorporated mainly through villous columnar cells, which play a primary role in nutritional absorption, rather than through M cells, which are responsible for the uptake of foreign substances (29). PrP^{Sc} was thought to be selectively taken up by the M cells of Peyer's patches (30), although the possibility of the incorporation of PrP^{Sc} across columnar epithelial cells was reported recently (8). In addition, it was shown that amyloid proteins were incorporated via columnar epithelial cells in mice (20) and cows (21). This possibility was strongly supported by the present finding of the incorporation of PrP^{Sc} via villous columnar epithelial cells.

In cases of oral or intragastric challenge with scrapie agents in hamsters, it is thought that the infectious agent first accumulates in Peyer's patches (31). These observations suggest that uptake is mediated through M cells in the dome epithelium of Peyer's patches. In the Peyer's patches in the present study, some PrP^{Sc} was detected in the dome epithelium, but most was incorporated in the villous epithelium. In addition, during the suckling and weaning periods, the Peyer's patches have not fully developed. So, it was suggested that uptake through the villi is important for the intestinal epithelial invasion of PrP^{Sc}. PrP^{Sc} was also detected in the villous lacteal, but no PrP^{Sc} was found in the dome regions of Peyer's patches at 4 h p.a.; however, PrP^{Sc} was detected in the dome regions at 7 days p.a. (31). PrP^{Sc} might be transmitted to lymphatic tissues through lacteal circulation after being taken up into the villous epithelium and then accumulate in the dome regions of Peyer's patches during the period after suckling and weaning.

The present experiment also showed a difference in uptake between before and after weaning. The uptake of PrP^{Sc} through the villi observed in suckling mice was not recognized in 25-day-old CD-1, BALB/c or C57BL/6 mice.




Figure 5. Uptake of PrP^{Sc} in normal and SCID mice. Intestinal villi of 15-day-old wild-type CD-1 (a) and CD-1.SCID (b, c) mice that had been orally administered PrP^{Sc} diluted with PBS (a, b) or PBS containing purified mouse IgG (c). PrP^{Sc} was incorporated into the villi of the CD-1 and CD-1.SCID mice lacking maternal antibodies. The number of ileal epithelial cells incorporating PrP^{Sc} was significantly higher in the CD-1 mice than in the CD-1.SCID mice, and the uptake was significantly enhanced by dilution with IgG (e). The statistical significance of differences was determined by the Student's t-test. ***P*<0.01. Scale bar, 50 μ m. PrP^{Sc} was also detected in the dome epithelium of Peyer's patches in the CD-1.SCID mice (d). As a control, the intestinal villi of 15-day-old wild-type CD-1 mice that had been orally administered with PBS containing purified mouse IgG and immune-stained for PrP^{Sc} (f). No immune-positive cells were detected in the mice not administered PrP^{Sc}.

Proteins such as the immunoglobulins and growth factors in breast milk (32) are incorporated into the body through the villous epithelium (33,34) during the suckling period without losing their original biological activity (35). Other macromolecules may also be absorbed non-specifically during the suckling period (36). In addition, during the suckling period the possibility of environmental antigenic challenge at the intestinal epithelium is lower, whereas the intestinal epithelium needs to take up more trophic and immunomodulatory factors (37). It was reported that the uptake of inert macroparticles in suckling SCID mice was significantly lower than that in normal mice (38). The neonatal Fc receptor for IgG (FcRn) binds to maternal IgG and is transcytosed in acidic conditions (39). In mice, FcRn is exposed on the cell surface brush border and is released at neutral pH on the neonatal side (40). A recent study showed that in the adult gut enterocytes transcytose IgG into the gut lumen where it binds to antigen. The IgG-antigen complex is then delivered to the lamina propria dendritic cells either directly or by reverse transcytosis across the epithelial T-cell barrier (40). In the present study, the levels of PrP^{Sc} incorporated by suckling SCID mice lacking maternal immunoglobulins (41) were significantly lower than those incorporated by wild-type suckling mice, whereas the uptake of PrP^{Sc} was enhanced by immunoglobulin. So, maternal immunoglobulin plays an important role in the enteric invasion of PrP^{Sc} into epithelial cells. It was also recently reported that the oral transmissibility of PrP^{Sc} was enhanced by its binding to soil particles (42). In that report, it was suggested that the association of PrP^{Sc} with soil minerals and organic carbon enhanced the oral transmissibility of prion

disease relative to the unbound agent. So, it is suggested in the present study that the binding of PrP^{Sc} to maternal immunoglobulin might enhance the enteric invasion of PrP^{Sc} relative to unbound PrP^{Sc}. Ongoing experiments have shown that an Fc receptor blocking agent (Z- ϵ -aminocaproic acid) significantly decreased PrP^{Sc} incorporation into the intestinal epithelia of CD-1 suckling mice. In this experiment, the incorporation of IgG into the intestinal epithelium was also blocked by Z- ϵ -aminocaproic acid. Therefore, the addition of such blocking agents to animal feed might be useful for preventing prion infection in younger animals (data not shown).

Cows younger than 6 months old are thought to be at the highest risk of infection from dietary PrP^{Sc} during the suckling and weaning period (19). Therefore, the villous epithelium may possess a specialized mechanism for the incorporation of foreign proteins like PrP^{Sc} during the suckling and weaning period, thereby increasing the risk of transmission across it.

The mechanism revealed in the present study using scrapie agents is a novel finding that reveals the oral transmission mechanism of prion diseases, and in particular, the mechanism by which they invade the intestinal epithelium.

Acknowledgements

This study was supported by Grant-in-Aids from the Food Safety Commission of the Japanese Government Cabinet Office and the Research Committee of Prion disease and Slow Virus Infection, the Ministry of Health, Labour and Welfare of Japan.

References

- Prusiner SB: Prion diseases and the BSE crisis. *Science* 278: 245-251, 1997.
- Ano Y, Sakudo A, Nakayama H and Onodera T: Uptake and distribution of infectious prion protein after oral exposure. *Protein Peptide Lett* 16: 247-255, 2009.
- Hoinville LJ: A review of the epidemiology of scrapie in sheep. *Rev Sci Tech* 15: 827-852, 1996.
- Will RG: Acquired prion disease: iatrogenic CJD, variant CJD, kuru. *Br Med Bull* 66: 255-265, 2003.
- Press CM, Heggebo R and Espenes A: Involvement of gut-associated lymphoid tissue of ruminants in the spread of transmissible spongiform encephalopathies. *Adv Drug Deliv* 56: 885-899, 2004.
- Robinson MM, Hadlow WJ, Huff TP, Wells GA, Dawson M, Marsh RF and Gorham JR: Experimental infection of mink with bovine spongiform encephalopathy. *J Gen Virol* 75: 2151-2155, 1994.
- Gonzalez L, Terry L and Jeffrey M: Expression of prion protein in the gut of mice infected orally with the 301V murine strain of the bovine spongiform encephalopathy agent. *J Comp Pathol* 132: 273-282, 2005.
- Foster JD, Paruham DW, Hunter N and Bruce M: Distribution of the prion protein in sheep terminally affected with BSE following experimental oral transmission. *J Gen Virol* 82: 2319-2326, 2001.
- Jeffrey M, Ryder S, Martin S, Hawkins SA, Terry L, Berthelin-Baker C and Bellworthy SJ: Oral inoculation of sheep with the agent of bovine spongiform encephalopathy (BSE). 1. Onset and distribution of disease-specific PrP accumulation in brain and viscera. *J Comp Pathol* 124: 280-289, 2001.
- Bons N, Mestre-Frances N, Belli P, Cathala F, Gajdusek DC and Brown P: Natural and experimental oral infection of non-human primates by bovine spongiform encephalopathy agents. *Proc Natl Acad Sci USA* 96: 4046-4051, 1999.
- Herzog C, Sales N, Etchegary N, Charbonnier A, Freire S, Dormont D, Deslys JP and Lasmezas CI: Tissue distribution of bovine spongiform encephalopathy agent in primates after intravenous or oral infection. *Lancet* 363: 422-428, 2004.
- Prusiner SB, Cochran SP and Alpers MP: Transmission of scrapie in hamsters. *J Infect Dis* 152: 971-978, 1985.
- McBride PA and Beekes M: Pathological PrP is abundant in sympathetic and sensory ganglia of hamsters fed with scrapie. *Neurosci Lett* 265: 135-138, 1999.
- Maigne T, Lasmezas CI, Beringue V, Dormont D and Deslys JP: Pathogenesis of the oral route of infection of mice with scrapie and bovine spongiform encephalopathy agents. *J Gen Virol* 80: 3035-3042, 1999.
- Jeffrey M, McGovern G, Goodsir CM, Brown KL and Bruce ME: Sites of prion protein accumulation in scrapie-infected mouse spleen revealed by immuno-electron microscopy. *J Pathol* 191: 323-332, 2000.
- Heggebo R, Press CM, Gunnes G, Lie KI, Tranulis MA, Ulvund M, Groschup MH and Landsverk T: Distribution of prion protein in the ileal Peyer's patch of scrapie-free lambs and lambs naturally and experimentally exposed to the scrapie agent. *J Gen Virol* 81: 2327-2337, 2000.
- Mabbott NA and Bruce ME: Follicular dendritic cells as targets for intervention in transmissible spongiform encephalopathies. *Semin Immunol* 14: 285-293, 2002.
- Jeffrey M, Gonzales L, Espenes A, *et al*: Transportation of prion protein across the intestinal mucosa of scrapie-susceptible and scrapie-resistant sheep. *J Pathol* 209: 4-14, 2006.
- Arnold ME and Wilesmith JW: Estimation of the age-dependent risk of infection to BSE of dairy cattle in Great Britain. *Prev Vet Med* 66: 35-47, 2004.
- Ano Y, Nakayama H, Sakudo A, Sawano Y, Tanokura M, Itohara S and Onodera T: Intestinal uptake of amyloid β protein through columnar epithelial cells in suckling mice. *Histol Histopathol* 24: 283-292, 2009.
- Ano Y, Nakayama H, Sakai Y, *et al*: Incorporation of β -amyloid protein through the bovine ileal epithelium before and after weaning: model for orally transmitted amyloidosis. *Microbiol Immunol* 52: 429-434, 2008.
- Hirogari Y, Kubo M, Kimura KM, Haritani M and Yokoyama T: Two different scrapie prions isolated in Japanese sheep flocks. *Microbiol Immunol* 47: 871-876, 2003.
- Yokoyama T, Masujin K, Iwamura Y, Imamura M and Mohri S: Alteration of the biological and biochemical characteristics of bovine spongiform encephalopathy prions during interspecies transmission in transgenic mice models. *J Gen Virol* 90: 261-268, 2009.
- Inoue Y, Yamakawa Y, Sakudo A, *et al*: Infection route-independent accumulation of splenic abnormal prion protein. *Jpn J Infect Dis* 58: 78-82, 2005.
- Yokoyama T, Kimura KM, Ushiki Y, *et al*: In vivo conversion of cellular prion protein to pathogenic isoforms, as monitored by conformation-specific antibodies. *J Biol Chem* 276: 11265-11271, 2001.
- Hosokawa T, Tsuchiya K, Sato I, *et al*: A monoclonal antibody (1D12) defines novel distribution patterns of prion protein (PrP) as granules in nucleus. *Biochem Biophys Res Commun* 366: 657-663, 2008.
- Hunyady B, Mezey E and Palkovits M: Gastrointestinal immunology: cell types in the lamina propria a morphological review. *Acta Physiol Hung* 87: 305-328, 2002.
- Clark MA, Jepson MA, Simmons NL, Booth TA and Hirst BH: Differential expression of lectin-binding sites defines mouse intestinal M-cells. *J Histochem Cytochem* 41: 1679-1687, 1993.
- Neutra MR, Frey A and Kraehenbuhl JP: Epithelial M cells: gateways for mucosal infection and immunization. *Cell* 86: 345-348, 1996.
- Heppner FL, Christ AD, Klein MA, Prinz M, Fried M, Kraehenbuhl JP and Aguzzi A: Transepithelial prion transport by M cells. *Nat Med* 7: 976-977, 2002.
- Beekes M and McBride PA: Early accumulation of pathological PrP in the enteric nervous system and gut-associated lymphoid tissue of hamster orally infected with scrapie. *Neurosci Lett* 278: 181-184, 2000.
- Guyer RL, Koshland ME and Knopf PM: Immunoglobulin binding by mouse intestinal epithelial cell receptors. *J Immunol* 117: 587-593, 1976.
- Udall JN, Colony P, Fritze L, Pang K, Trier JS and Walker WA: Development of gastrointestinal mucosal barrier: the effect of natural versus artificial feeding on intestinal permeability to macromolecules. *Pediatr Res* 15: 245-249, 1981.
- Axelsson L, Jakobsson L, Lindberg T, Polberger S, Benediktsson B and R  ih   N: Macromolecular absorption in preterm and term infant. *Acta Paediatr Scand* 78: 532-537, 1989.
- Chu SH and Walker WA: Growth factor signal transduction in human intestinal cells. *Adv Exp Med Biol* 310: 107-112, 1991.
- Teichberg S, Wapnir RA, Moyse J and Lifshitz F: Development of the neonatal rat small intestinal barrier to non-specific macromolecular absorption. Role of dietary corticosterone. *Pediatr Res* 32: 50-57, 1992.
- Rumbo M and Schiffrin EJ: Ontogeny of intestinal epithelium immune functions: developmental and environmental regulation. *Cell Mol Life Sci* 62: 1288-1296, 2005.
- Smyth SH, Feldhaus S, Schumacher U and Carr KE: Uptake of inert microparticles in normal and immune deficient mice. *Int J Pharm* 346: 109-118, 2008.
- Israel EJ, Patel VK, Taylor SF, Marshak-Rothstein A and Simister NE: Requirement for a beta 2-microglobulin-associated Fc receptor for acquisition of maternal IgG by fetal and neonatal mice. *J Immunol* 154: 6246-6251, 1995.
- Roopenian DC and Akilesh S: FcRn: the neonatal Fc receptor comes of age. *Nat Rev Immunol* 7: 715-725, 2007.
- Kramer DR and Cebra JJ: Early appearance of 'natural' mucosal IgA responses and germinal centers in suckling mice developing in the absence of maternal antibodies. *J Immunol* 154: 2051-2062, 1995.
- Johnson CJ, Pedersen JA, Chappell RJ, McKenzie D and Aiken JM: Oral transmissibility of prion disease is enhanced by binding to soil particles. *PLoS Pathog* 3: e93, 2007.