

Substrate stiffness affects epithelial-mesenchymal transition of cervical cancer cells through miR-106b and its target protein DAB2

JINLAN PIAO¹, KE YOU¹, YANLI GUO¹, YOUYI ZHANG¹, ZIJIAN LI² and LI GENG¹

¹Department of Gynecology and Obstetrics, Peking University Third Hospital; ²Institute of Vascular Medicine, Peking University Third Hospital, Beijing Key Laboratory of Cardiovascular Receptors Research, Key Laboratory of Cardiovascular Molecular Biology and Regulatory Peptides and Ministry of Health, Key Laboratory of Molecular Cardiovascular Sciences, Ministry of Education, Haidian, Beijing 100191, P.R. China

Received December 13, 2016; Accepted March 9, 2017

DOI: 10.3892/ijo.2017.3978

Abstract. The effects of different substrate stiffness were investigated on epithelial-mesenchymal transition (EMT) of cervical cancer cell lines and the role of miR-106b and its target protein DAB2 therein. Cervical cancer cell lines HeLa and SiHa were cultured on artificial substrates with different stiffness prepared using different ratios of acrylamide and bis-acrylamide. Changes of microRNA profiles were detected using microRNA chip analysis, and the expression levels of EMT-related markers E-cadherin and vimentin were detected using western blotting and real-time PCR. In addition, the effects of miR-106b overexpression as well as miR-106b and DAB2 knockdown on expression of E-cadherin and vimentin were also examined using western blotting and real-time PCR. The results showed that i) cervical cancer cell lines SiHa and HeLa cultured on substrate with stiffness of 20 kPa had the strongest EMT ability, showed the highest levels of vimentin and lowest levels of E-cadherin, compared with cells cultured on substrate with stiffness of 1 kPa; ii) miR-106b knockdown reversed the effects of substrate stiffness on EMT of cervical cancer cells, while miR-106 overexpression and DAB2 knockdown induced EMT of cervical cancer cells cultured on

substrate with stiffness of 20 kPa. Overall, the results indicated that substrate stiffness could regulate EMT of cervical cancer cell lines HeLa and SiHa at least partially through miR-106b and its downstream target DAB2.

Introduction

Cervical cancer is the world's fourth most common female malignancy and ranks the second in the developing countries and the third in the developed countries, seriously affecting the health of women worldwide. Each year there are around 500,000 new cervical cancer cases and 266,000 deaths, of which more than 80% are in developing countries. It is estimated that there are approximately 62,000 new cases and 30,000 dead cases each year in China, and its annual incidence increases by 2-3%. In addition, the onset age of cervical cancer patients in recent years has decreased, and the incidence of cervical cancer increases sharply in the age group of 25 to 34 years, causing great concern world-wide (1-3).

According to recent epidemiological investigation, clinical stage is one of the risk factors affecting prognosis of patients with cervical cancer: the later the stage, the worse the prognosis of the patients (4). These findings indicate that invasion and metastasis are the main causes for poor prognosis and high mortality of patients with cervical cancer and these difficult problems need to be solved. Studies have found that epithelial-mesenchymal transition (EMT) is the primary step causing cervical cancer invasion and metastasis and its relationship to tumor progression is gradually drawing more attention (5-7). EMT is a phenomenon where epithelial cells lose their polarity and gradually shift to mesenchymal cell morphology under certain pathological stimulations. The process is accompanied with cell morphological changes and increased ability to alter cell motility, thus further affecting cell migration (8,9).

Recent studies have found that changes in the mechanical properties of the extracellular matrix, i.e. the substrate stiffness, could regulate EMT and other biological functions of tumor cells. In other words, mechanical properties of extracellular matrix play a major role in tumor progression. Studies

Correspondence to: Dr Zijian Li, Institute of Vascular Medicine, Peking University Third Hospital, Beijing Key Laboratory of Cardiovascular Receptors Research, Key Laboratory of Cardiovascular Molecular Biology and Regulatory Peptides and Ministry of Health, Key Laboratory of Molecular Cardiovascular Sciences, Ministry of Education, 49 North Garden Road, Haidian, Beijing 100191, P.R. China
E-mail: lzi1995@126.com

Dr Li Geng, Department of Gynecology and Obstetrics, Peking University Third Hospital, 49 North Garden Road, Haidian, Beijing 100191, P.R. China
E-mail: gengli1957@bjmu.edu.cn

Key words: cervical cancer, substrate stiffness, epithelial-mesenchymal transition, microRNA

have found that changes in substrate stiffness can affect EMT of breast cancer cell lines, lung cancer cell lines and hepatoma cell lines (10,11). In addition, human tissue stiffness changes significantly with disease progressing. For example, the stiffness of normal breast tissues is around 4 kPa, while that of cancerous breast tissues is approximately 12 kPa (7,12-16). Moreover, studies have shown that tumor-associated fibroblasts and interstitial fibrosis components which is produced by fibroblasts, such as α -SMA, laminin and fibronectin increase significantly with cervical cancer progression, suggesting that the stiffness of cervical cancer tissues is higher than that of normal cervical tissues (16). However, whether the mechanical properties of extracellular matrix can influence EMT of cervical cancer cells has not been reported.

MicroRNA is a small RNA with post-transcriptional regulation functions. MicroRNA regulate expression levels of important oncogenes and tumor suppressor genes, and further regulate cancer cell invasion and metastasis. Previous studies have found that the mechanical signals of extracellular matrix can alter microRNA expression in cells (17-19). For example, miR-29 expression was significantly increased by bladder obstruction-induced increase in bladder tissue stiffness (19); microRNA expression changes in alveolar epithelium and mesothelial cells under different mechanical shear stress. In other words, different substrate stiffness can affect some microRNA expression in cells, therefore regulating cell functions by targeting degradation of important proteins (20). Mouw *et al* found that mechanical stiffness of extracellular matrix could regulate PTEN expression and its downstream PI3K/AKT signaling pathway and promote breast cancer occurrence through microRNA, indicating that microRNA is an important key node in transducing mechanical signals to cells (21).

Our previous study established a related miRNA-mRNA regulatory network diagram through mRNA and microRNA chips analysis of normal cervical tissues and cervical cancer tissues and found that miR-106b is a key node in the network and closely related to cervical cancer cell migration. In addition, our previous study also found that miR-106b was significantly upregulated in cervical cancer tissues and could promote EMT and migration of cancer cells by degrading its target protein DAB2 (22).

In this study, we explored whether substrate stiffness can affect EMT of cervical cancer cells, and the role of miR-106b and its target protein DAB2 in this process.

Materials and methods

Artificial substrate preparation. Artificial substrates with different stiffness were prepared using acrylamide and bisacrylamide at different ratios and laid on 24x24 mm slides. After irradiated with UV light for 2 h, they were used to grow cervical cancer cell lines.

Cell culture. The cervical cancer cell lines SiHa and HeLa were cultured in DMEM medium supplemented with 10% FBS, 100 U/ml penicillin and 100 mg/ml streptomycin at 37°C in a humidified incubator supplemented with 5% CO₂. When cells reached 70-80% confluency, they were transferred on artificial substrates.

Table I. Sequences of primers used in real-time PCR.

Gene	Primer (5'-3')
DAB2	F: AGCAAGGTTCAAAGGCGATG R: GTTGTCCCTGAGACCGACC
Vimentin	F: TGCCGTTGAAGCTGCTAACTA R: CCAGAGGGAGTGAATCCAGATTA
N-cadherin	F: AGCCAACCTTAACTGAGGAGT R: GGCAAGTTGATTGGAGGGATG
GAPDH	F: CCTCCGGGAAACTGTGGCGTGATGG R: AGACGGCAGGTCAGGTCCACCACTG

F, forward; R, reverse.

Western blotting. Proteins were extracted from cervical cancer cell lines HeLa and SiHa and semi-quantitated with Coomassie brilliant blue. A total of 10 μ g protein of each sample was separated on 10% SDS-PAGE and transferred onto PVDF membrane. The membrane was then blocked with 5% skim milk at 37°C for 1 h. Target proteins on the membrane were recognized by incubating with appropriate primary antibodies (rabbit anti-human E-cadherin, mouse anti-human vimentin, rabbit anti-human Twist1 and rabbit anti-human Eif-5 antibodies, respectively) at room temperature for 4 h. After rinsed with TBST three times for 5 min each time, the protein-antibody complexes were recognized by alkaline phosphatase-labeled anti-IgG (1:1000) antibody by incubating at room temperature for 1 h. After rinsed with TBST three times for 10 min each, the complexes were visualized by incubation with NBT/BCIP. The gray scale images were scanned and protein expression was analyzed using Eif-5 as the internal control and expressed as relative RGS. Each experiment was repeated 3 times.

Real-time PCR. Total RNA were extracted using TRIzol from SiHa cells cultured on substrate with different stiffness. mRNA were reverse transcribed into cDNA and subjected to fluorescence quantitative PCR at the condition of denature at 95°C for 2 min followed by 40 cycles of 95°C for 30 sec, 55°C for 30 sec and 72°C for 30 sec using primers shown in Table I on Bio-Rad PCR real-time PCR instrument. Expression levels of DAB2 and vimentin were normalized to internal control Eif-5. Each experiment was repeated three times.

MicroRNA chip analysis. Total RNA was extracted using TRIzol from SiHa cells cultured on substrates with stiffness of 1 kPa and 20 kPa as well as on a glass plate were reverse transcribed with the Qiagen RNeasy Mini kit and subjected to microRNA chip analysis using Genisphere FlashTag Labeling kit from Beijing CapitalBio Corp.

Statistical analysis. The results are expressed as mean \pm SEM. Differences among multiple groups were compared using one way ANOVA. Samples with different variance between two groups were compared using non-parametric tests.




Figure 1. Morphological changes of SiHa and HeLa cells cultured on glass slide and substrates with stiffness of 1 and 20 kPa, respectively. Both types of cells are in round shape when cultured on substrate with stiffness of 1 kPa, but form obvious pseudopodia when cultured on substrate with stiffness of 20 kPa.




Figure 2. Real-time PCR results showing the mRNA levels of EMT epithelial marker E-cadherin and EMT mesenchymal marker vimentin (B) in cervical cancer cell lines cultured on substrates with different stiffness (1, 10, 20, 30 kPa) and on glass. (A) mRNA level of E-cadherin SiHa cells; (B) mRNA level of vimentin in SiHa cells; (C) mRNA level of E-cadherin HeLa cells; (D) mRNA level of vimentin in HeLa cells. * $P<0.05$; ** $P<0.01$.

$P<0.05$ was considered to indicate a statistically significant difference. Graphs were plotted using GraphPad Prism 5.0 software (GraphPad Software Inc., La Jolla, CA, USA).

Results

Selection of substrate stiffness range. It was reported that the average stiffness of human solid tumors is approximately 20 kPa. The stiffness of most solid tumor tissues before carcinogenesis is higher than that of normal tissues (7). Therefore, in this study, the artificial substrates were prepared to have stiffness of 1, 10, 20 and 30 kPa, respectively, and glass plate was used as control.

Effects of substrate stiffness on the morphology of cervical cancer cell lines SiHa and HeLa. SiHa and HeLa cells were cultured on artificial substrate with stiffness of 1 kPa to represent their normal condition and on artificial substrate with stiffness of 20 kPa to represent cervical tumor tissues. Morphological observation showed that cells cultured on artificial substrate with stiffness of 1 kPa were spherical and hardly had pseudopodia, while cells cultured on artificial substrate with stiffness of 20 kPa were rich in pseudopodia (Fig. 1), indicating that

SiHa and HeLa cells have stronger migration ability when cultured on artificial substrate with stiffness of 20 kPa than cultured on artificial substrate with stiffness of 1 kPa.

Effects of substrate stiffness on EMT of cervical cancer cell lines SiHa and HeLa. First, we detected the expression level of EMT markers in SiHa and HeLa cultured at artificial substrate with different stiffness using real-time PCR. The results showed that mRNA levels of vimentin were significantly lower in SiHa and HeLa cells cultured on artificial substrate with stiffness of 20 kPa than cultured on substrate with stiffness of 1 kPa, and the expression level of E-cadherin higher in stiffness of 20 kPa than 1 kPa (Fig. 2).

Expression levels of EMT markers E-cadherin and vimentin in SiHa and HeLa cultured at artificial substrate with different stiffness was also detected using western blotting. The results showed that protein levels of E-cadherin is significantly lower in SiHa and HeLa cells cultured on artificial substrate with stiffness of 20 kPa than cultured on substrate with stiffness of 1 kPa, and expression levels vimentin is significantly higher in stiffness of 20 kPa than 1 kPa (Fig. 3).

The above results indicate that cervical cancer cell line SiHa and HeLa had stronger EMT ability when cultured on




Figure 3. Western blot results showing protein levels of EMT epithelial marker E-cadherin and EMT mesenchymal marker vimentin (B) in cervical cancer cell lines cultured on substrates with different stiffness (1, 10, 20, 30 kPa) and on glass. (A) E-cadherin and vimentin in SiHa cells; (B) E-cadherin and vimentin in HeLa cells; (C) statistical results of E-cadherin protein levels in SiHa cells; (D) statistical results of vimentin in SiHa cells; (E) statistical results of E-cadherin protein levels in HeLa cells; (F) statistical results of vimentin in HeLa cells. ** $P < 0.01$.




Figure 4. Effects of Twist1 expression of cervical cancer cell line SiHa cultured on glass or substrate with stiffness of 1, 10, 20, 30 kPa. (A) Expression levels of Twist1 detected by western blotting in cervical cancer cell line SiHa; (B) statistical results of Twist protein levels in SiHa cells; (C) expression levels of Twist1 detected by PCR in cervical cancer cell line SiHa. * $P < 0.05$.

artificial substrate with stiffness of 20 kPa than cultured on substrate with stiffness of 1 kPa.

Effects of substrate stiffness on Twist1 expression of cervical cancer cell line SiHa. We detected the expression level of Twist1 in SiHa which were cultured at artificial substrate with different stiffness using western blotting. The results assessed by western blotting showed that Twist1 expression was the

highest in 30 kPa on glass, lowest in 10 kPa situation. However, when tested by Real-time PCR the Twist1 expression was the lowest in 20 kPa situation excluding on glass. Expression of Twist1 of the cervical cancer cell line SiHa is not consistent with EMT tendency (Fig. 4).

Effects of substrate stiffness on microRNA expression in cervical cancer cell lines SiHa and HeLa. To explore whether




Figure 5. MicroRNA expression profile in cervical cancer cell line SiHa cultured on substrates with different stiffness.

microRNA is involved in regulating EMT of cervical cancer cell lines cultured on substrates with different stiffness, we tested differential expression of microRNA using microarray test. SiHa cells cultured on substrates with stiffness of 1 kPa and 20 kPa as well as on glass plates for 36 h. Their total RNA was extracted and used for microRNA chip analysis to detect differential expression of microRNA. Cluster analysis clearly showed that 173 microRNA were differentially expressed in cervical cancer cell line SiHa cultured on substrates with different stiffness (Fig. 5). Among them, 70 microRNA including miR-106b were upregulated and 103 microRNA were downregulated in SiHa cells cultured on substrates with stiffness of 20 kPa ($p < 0.05$, $FRD < 0.05$). Tables II and III list the microRNA that were mostly upregulated and down-regulated in SiHa cells cultured on substrate with stiffness of 20 kPa compared with cells cultured on substrate with stiffness of 1 kPa, respectively. Obviously, substrate stiffness can affect microRNA expression in cervical cancer SiHa cells, wherein miR-106b may play an important regulatory role.

Effects of substrate stiffness on miR-106b expression in cervical cancer cell line SiHa. To understand whether




Figure 6. Expression of miR-106b in cervical carcinoma cell line SiHa cultured on substrate with different stiffness. * $P < 0.05$.




Figure 7. Expression of DAB2 in cervical cancer cell line SiHa cultured on substrates with different stiffness. (A) A representative image of western blot detecting DAB2 expression in cervical cancer cell line SiHa cultured on substrates with different stiffness; (B) statistical analysis of expression of DAB2 normalized to control Eif-5 in SiHa cells cultured on substrate with different stiffness. ** $P < 0.01$.

substrate stiffness affects EMT of cervical cancer cell lines through miR-106b, miR-106b expression in cervical cancer cells was further measured and verified using real-time PCR. The results (Fig. 6) confirmed that miR-106b expression was indeed upregulated in cervical cancer cell line SiHa cultured on substrate with stiffness of 20 kPa. Considering that SiHa cultured on substrate with stiffness of 20 kPa had stronger EMT ability than SiHa cultured on substrate with stiffness of 1 kPa, the result suggests that substrate stiffness might regulate EMT of cervical carcinoma cell line SiHa by regulating miR-106 expression.

Effects of substrate stiffness on expression of tumor suppressor protein DAB2 in cervical cancer cell line SiHa. To further investigate whether substrate stiffness affects EMT of cervical




Figure 8. Effects of miR-106b overexpression on cervical cancer cell line SiHa cultured on glass or substrates with stiffness of 1 and 20 kPa, respectively. (A) miR-106b levels detected by real-time PCR in cervical cancer cell line SiHa cultured on glass and substrates with stiffness of 1 and 20 kPa, respectively, before and after miR-106b overexpression. (B) The expression levels of miR-106b target protein DAB2 and EMT markers E-cadherin and vimentin detected by western blotting in cervical cancer cell line SiHa cultured on glass and substrates with stiffness of 1 and 20 kPa, respectively, before and after miR-106b overexpression; (C) changes of DAB2 in SiHa cells cultured on glass and substrates with different stiffness before and after miR-106b overexpression; (D) changes of E-cadherin in SiHa cells cultured on glass and substrates with different stiffness before and after miR-106b overexpression; (E) changes of vimentin levels in SiHa cells cultured on glass and substrates with different stiffness before and after miR-106b overexpression. * $P < 0.05$; ** $P < 0.01$.

Table II. Top differentially upregulated microRNAs in cervical cancer cells cultured on substrates with stiffness of 1 and 20 kPa, respectively.

MicroRNA	Fold change	MicroRNA	Fold change
hsa-miR-125a	17.554	hsa-miR75b	3.8857
hsa-miR-21	3.8289	hsa-miR-595	1.6495
hsa-miR-150	1.067	hsa-miR-6785-3p	8.487
hsa-miR-6507-5p	3.5284	hsa-miR-548u	2.8511
hsa-miR-211	2.6768	hsa-miR-106b	2.1564

Table III. Top differentially downregulated microRNAs in cervical cancer cells cultured on substrates with stiffness of 1 and 20 kPa, respectively.

MicroRNA	Fold change	MicroRNA	Fold change
hsa-miR-506b	-30.9045	hsa-miR-218	-1.2745
hsa-miR-200b	-0.5418	hsa-miR-107	-0.0175
hsa-miR-183	-0.0517	hsa-miR-7-1-3p	-0.0197
hsa-miR-1279	-0.0933	hsa-miR-522-3p	-0.011
hsa-miR-101-5p	-0.0791	hsa-miR-5001-3p	-0.0127

carcinoma cell line SiHa through miR-106b and its target protein DAB2, the level of DAB2 in cervical cancer cell line

SiHa was examined by western blotting. The results (Fig. 7) showed that the expression level of DAB2 was significantly




Figure 9. Effects of miR-106b knockdown on cervical cancer cell line SiHa cultured on glass or substrates with stiffness of 1 and 20 kPa, respectively. (A) miR-106b levels detected by real-time PCR in cervical cancer cell line SiHa cultured on glass and substrates with stiffness of 1 and 20 kPa, respectively, before and after miR-106b knockdown. (B) The expression levels of miR-106b target protein DAB2 and EMT markers E-cadherin and vimentin detected by western blotting in cervical cancer cell line SiHa cultured on glass and substrates with stiffness of 1 and 20 kPa, respectively, before and after miR-106b knockdown; (C) changes of DAB2 in SiHa cells cultured on glass and substrates with different stiffness before and after miR-106b knockdown; (D) changes of E-cadherin in SiHa cells cultured on glass and substrates with different stiffness before and after miR-106b knockdown; (E) changes of vimentin levels in SiHa cells cultured on glass and substrates with different stiffness before and after miR-106b knockdown. *P<0.05; **P<0.01.

different in SiHa cells cultured on substrate with different stiffness and lower in cells cultured on substrate with stiffness of 20 kPa, suggesting that substrate stiffness may regulate EMT through miR-106b targeted DAB2 degradation.

miR-106b is involved in substrate stiffness regulated EMT of cervical cancer cell line SiHa. To investigate the specific mechanism, substrate stiffness of 1 and 20 kPa was selected to represent normal cervical tissue stiffness and cervical cancer tissue stiffness, respectively, and the changes in expression levels of miR-106b target protein DAB2 and EMT-related markers vimentin and E-cadherin after miR-106b overexpression and silencing were detected using western blotting. Fig. 8 shows that miR-106b overexpression significantly decreased the expression of DAB2 and vimentin, but increased the expression of E-cadherin in cervical cancer cell line SiHa cultured on substrate with stiffness of 20 kPa.

Fig. 9 shows that miR-106b silencing significantly increased the expression level of DAB2 and vimentin and decreased the expression level of E-cadherin in cervical cancer cell line SiHa cultured on substrate with stiffness of 20 kPa. Together, these data suggest that substrate stiffness regulates EMT of cervical carcinoma cell line SiHa through miR-106b and its target protein DAB2.

DAB2 is involved in regulating EMT of cervical cancer cell SiHa. To further investigate whether DAB2 is involved in regulating EMT of cervical cancer cells in different substrate stiffness, we explored the effect of DAB2 knockdown on the expression of E-cadherin and vimentin in cervical cancer cell SiHa cultured on substrate with the stiffness of 1 kPa using western blotting. The results showed that DAB2 knockdown significantly decreased EMT-associated E-cadherin level and increased vimentin level, i.e. enhanced EMT, indirectly




Figure 10. Effects of DAB2 knockdown on cervical cancer cell line SiHa cultured on glass or substrate with stiffness of 1 kPa. (A) Expression levels of DAB2 and EMT markers E-cadherin and vimentin detected by western blotting in cervical cancer cell line SiHa cultured on glass and substrate with stiffness of 1 kPa before and after DAB2 knockdown; (B) Changes of DAB2 in SiHa cells cultured on glass and substrate with different stiffness before and after DAB2 knockdown; (C) Changes of vimentin in SiHa cells cultured on glass and substrate with stiffness of 1 kPa before and after DAB2 knockdown; (D) Changes of E-cadherin levels in SiHa cells cultured on glass and substrate with stiffness of 1 kPa before and after DAB2 knockdown. * $P < 0.05$; ** $P < 0.01$.

suggesting that DAB2 is involved in substrate stiffness regulation of EMT of cervical cancer cell line SiHa (Fig. 10).

Discussion

Substrate stiffness affects EMT of cervical cancer cells. In this study, we explored the effects of substrate stiffness on EMT of cervical cancer cells. Changes in substrate stiffness are important signs of many epithelial tumors and can regulate EMT of many important cells such as liver epithelial cells and different types of cells. EMT of hepatocytes is stronger on substrate with stiffness of 60 kPa (hard substrate) than on substrate with stiffness of 11 kPa (soft substrate) (23); EMT of mouse mammary epithelial cells is stronger on substrate with stiffness of 11 kPa (hard substrate) than on substrate with stiffness of 4 kPa (soft substrates) (24); EMT of alveolar epithelial cells is stronger on substrate with stiffness of 17 kPa than on substrate with stiffness of 4 kPa. All these results indicated that EMT of different types of tumor cells requires different extracellular conditions. In our study, we found that cervical cancer cell lines are more prone to EMT on substrate with stiffness of 20 kPa than 1 kPa, and less prone to EMT on substrate with stiffness of 30 kPa than 20 kPa. Substrate stiffness of 20 kPa is currently considered as average stiffness of human solid tumor tissues and very close to the stiffness of cervical cancer tissue, suggesting that cervical cancer cells are prone to EMT at certain substrate stiffness while are not prone to EMT at other conditions.

Substrate stiffness affects EMT of cervical cancer cells through miR-106b and its target DAB2. In recent years, many transcription factors and signal pathways have been shown closely related with EMT of cancer cells. Research on transcription factors have not made substantial progress (5). In our investigation we also tested the transcription factor Twist1 which has a close relation with EMT of cervical cancer. We found the expression of Twist1 1 kPa condition the lowest and the expression of Twist1 the highest in 30 kPa condition. Thus, the expression of Twist1 does not match the EMT tendency (Fig. 10). The impact of microRNA on tumorigenesis has gradually become a hot topic of research. It is well acknowledged that microRNA is closely related to pathological process of several kinds of cancers such as cervical cancer (17,18).

Importantly some studies have found that specific microRNA can regulate EMT of tumor cells (25-32). In the present study, we found that miR-106b and its target protein DAB2 play significantly important regulatory roles in EMT process of cervical cancer cell lines SiHa and HeLa. In our study, we found that miR-106b and DAB2 play important roles in regulating EMT of cervical cancer SiHa and HeLa cells which were caused by changes in substrate stiffness. In other words, with substrate stiffness changing, miR-106b expression in cervical cancer cell lines changes significantly. Suppression or overexpression of miR-106b correspondingly alters the expression of EMT markers, indicating that substrate stiffness may affect EMT of cervical cancer cells through microRNA. This view is also supported by other studies. Mouw *et al* found

that changes in substrate stiffness increased miR-18a expression and promoted breast cancer development by regulating expression of its target protein PTEN (21). It is clear that miR-106b is an important key node in mechanical signaling transduction in cervical cancer HeLa and SiHa cells and microRNA can be the key component in mechanical signal transduction process.

It is currently accepted that integrin is the cell membrane receptor of mechanic signals. It sensors extracellular mechanical signals and transfers the signals into intracellular signals to further regulate expression of important genes and cellular functions such as growth, proliferation, migration and EMT. It has been reported that some microRNA may further affect integrin receptor expression or activity through their target proteins and associated signaling pathways to influence cell EMT, migration, invasion and other cell functions (30,31). It is possible that changes in miR-106b expression due to substrate stiffness may also affect expression or activity of certain specific cell membrane integrin receptors and further the transduction of extracellular mechanical signals into intracellular signals. This hypothesis need to be further explored.

Moreover, microRNA usually regulate the expression of its target proteins by post-translational degradation. Our results found that miR-106b target protein DAB2 could regulate EMT of cervical cancer cell lines cultured on substrates with different stiffness. Many studies have found that loss of expression of tumor suppressor protein DAB2 can significantly promote EMT of many tumor cells, including cervical cancer cells. To our knowledge, whether different substrate stiffness can regulate DAB2 expression has not been reported previously. Our results showed that substrate stiffness could affect DAB2 expression. DAB2 expression is the lowest in cervical cancer cell lines cultured on substrate with stiffness of 20 kPa and the highest in cervical cancer cell lines cultured on substrate with stiffness of 1 kPa.

The characteristics that tumor tissues have different stiffness in ultrasound and elastography examinations have been used in clinical medicine for preliminary tumor diagnosis (32-34). Our results showed that substrate stiffness can regulate EMT of cancer cells. We have previously shown that miR-106b expression is higher in cervical tumor tissues than in normal cervical tissues, and high miR-106b expression promotes cervical carcinogenesis. In this study, we found that at cytological level, miR-106b plays an important regulatory role in EMT of cervical cancer cell lines cultured on substrates with different stiffness, suggesting that miR-106b and substrate stiffness play important roles in cervical cancer progression. Our results also suggest that changes in substrate stiffness can regulate specific molecular signaling pathways, miR-106b and its downstream target proteins DAB2, and changes in substrate stiffness and miR-106b can be targets for treatment of cervical cancer. The study indicates a new way for treatment of cervical cancer.

Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (no. 81472429, 81471893, 81270157) and National Basic Research Program of China (no. 973 Program, 2013CB933702, 2014CBA02003).

References

- Poljak M, Kocjan BJ, Oštrbenk A and Seme K: Commercially available molecular tests for human papillomaviruses (HPV): 2015 update. *J Clin Virol* 76 (Suppl 1): S3-S13, 2016.
- Oh J-K and Weiderpass E: Infection and cancer: Global distribution and burden of diseases. *Ann Glob Health* 80: 384-392, 2014.
- Parkin DM, Bray F, Ferlay J and Pisani P: Estimating the world cancer burden: Globocan 2000. *Int J Cancer* 94: 153-156, 2001.
- Wang J, Xu K, Shi B, Zhang K, Zhou P, Liao X, Liu G and Zhang Y: Prognosis of 4374 cases of cervical cancer and its affecting factors. *China Cancer* 23: 181-188, 2014.
- Lamouille S, Xu J and Derynck R: Molecular mechanisms of epithelial-mesenchymal transition. *Nat Rev Mol Cell Biol* 15: 178-196, 2014.
- Wei SC and Yang J: Forcing through tumor metastasis: The interplay between tissue rigidity and epithelial-mesenchymal transition. *Trends Cell Biol* 26: 111-120, 2016.
- Nagelkerke A, Bussink J, Rowan AE and Span PN: The mechanical microenvironment in cancer: How physics affects tumours. *Semin Cancer Biol* 35: 62-70, 2015.
- Mani SA, Guo W, Liao MJ, Eaton EN, Ayyanan A, Zhou AY, Brooks M, Reinhard F, Zhang CC, Shipitsin M, *et al*: The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell* 133: 704-715, 2008.
- Zhang L, Huang G, Li X, Zhang Y, Jiang Y, Shen J, Liu J, Wang Q, Zhu J, Feng X, *et al*: Hypoxia induces epithelial-mesenchymal transition via activation of SNAIL by hypoxia-inducible factor-1 α in hepatocellular carcinoma. *BMC Cancer* 13: 108, 2013.
- Brown AC, Fiore VF, Sulchek TA and Barker TH: Physical and chemical microenvironmental cues orthogonally control the degree and duration of fibrosis-associated epithelial-to-mesenchymal transitions. *J Pathol* 229: 25-35, 2013.
- Young JL, Holle AW and Spatz JP: Nanoscale and mechanical properties of the physiological cell-ECM microenvironment. *Exp Cell Res* 343: 3-6, 2016.
- Tilghman RW, Cowan CR, Mih JD, Koryakina Y, Gioeli D, Slack-Davis JK, Blackman BR, Tschumperlin DJ and Parsons JT: Matrix rigidity regulates cancer cell growth and cellular phenotype. *PLoS One* 5: e12905, 2010.
- Das T, Safferling K, Rausch S, Grabe N, Boehm H and Spatz JP: A molecular mechanotransduction pathway regulates collective migration of epithelial cells. *Nat Cell Biol* 17: 276-287, 2015.
- Mierke CT: The fundamental role of mechanical properties in the progression of cancer disease and inflammation. *Rep Prog Phys* 77: 076602, 2014.
- Parekh A and Weaver AM: Regulation of invadopodia by mechanical signaling. *Exp Cell Res* 343: 89-95, 2016.
- Fullár A, Dudás J, Oláh L, Hollósi P, Papp Z, Sobel G, Karácsi K, Paku S, Baghy K and Kovalszky I: Remodeling of extracellular matrix by normal and tumor-associated fibroblasts promotes cervical cancer progression. *BMC Cancer* 15: 256, 2015.
- Yehya N, Yerrapureddy A, Tobias J and Margulies SS: MicroRNA modulate alveolar epithelial response to cyclic stretch. *BMC Genomics* 13: 154, 2012.
- Valastyan S and Weinberg RA: Roles for microRNAs in the regulation of cell adhesion molecules. *J Cell Sci* 124: 999-1006, 2011.
- Ekman M, Bhattachariya A, Dahan D, Uvelius B, Albinsson S and Swärd K: Mir-29 repression in bladder outlet obstruction contributes to matrix remodeling and altered stiffness. *PLoS One* 8: e82308, 2013.
- Neth P, Nazari-Jahantigh M, Schober A and Weber C: MicroRNAs in flow-dependent vascular remodelling. *Cardiovasc Res* 99: 294-303, 2013.
- Mouw JK, Yui Y, Damiano L, Bainer RO, Lakins JN, Acerbi I, Ou G, Wijekoon AC, Levental KR, Gilbert PM, *et al*: Tissue mechanics modulate microRNA-dependent PTEN expression to regulate malignant progression. *Nat Med* 20: 360-367, 2014.
- Cheng Y, Guo Y, Zhang Y, You K, Li Z and Geng L: MicroRNA-106b is involved in transforming growth factor β 1-induced cell migration by targeting disabled homolog 2 in cervical carcinoma. *J Exp Clin Cancer Res* 35: 11, 2016.
- Bonnans C, Chou J and Werb Z: Remodelling the extracellular matrix in development and disease. *Nat Rev Mol Cell Biol* 15: 786-801, 2014.
- Lee K, Chen QK, Lui C, Cichon MA, Radisky DC and Nelson CM: Matrix compliance regulates Rac1b localization, NADPH oxidase assembly, and epithelial-mesenchymal transition. *Mol Biol Cell* 23: 4097-4108, 2012.

25. Cheng YX, Chen GT, Chen C, Zhang QF, Pan F, Hu M and Li BS: MicroRNA-200b inhibits epithelial-mesenchymal transition and migration of cervical cancer cells by directly targeting RhoE. *Mol Med Rep* 13: 3139-3146, 2016.
26. Yao J, Deng B, Zheng L, Dou L, Guo Y and Guo K: miR-27b is upregulated in cervical carcinogenesis and promotes cell growth and invasion by regulating CDH11 and epithelial-mesenchymal transition. *Oncol Rep* 35: 1645-1651, 2016.
27. Peralta-Zaragoza O, Deas J, Meneses-Acosta A, De la O-Gómez F, Fernández-Tilapa G, Gómez-Cerón C, Benítez-Boijseauneau O, Burguete-García A, Torres-Poveda K, Bermúdez-Morales VH, *et al*: Relevance of miR-21 in regulation of tumor suppressor gene PTEN in human cervical cancer cells. *BMC Cancer* 16: 215, 2016.
28. Qin X, Wan Y, Wang S and Xue M: MicroRNA-125a-5p modulates human cervical carcinoma proliferation and migration by targeting ABL2. *Drug Des Devel Ther* 10: 71-79, 2015.
29. Cheng Y, Ma D, Zhang Y, Li Z and Geng L: Cervical squamous cancer mRNA profiles reveal the key genes of metastasis and invasion. *Eur J Gynaecol Oncol* 36: 309-317, 2015.
30. Zhang J, Na S, Liu C, Pan S, Cai J and Qiu J: MicroRNA-125b suppresses the epithelial-mesenchymal transition and cell invasion by targeting ITGA9 in melanoma. *Tumour Biol* 37: 5941-5949, 2016.
31. Liu X, Liang Z, Gao K, Li H, Zhao G, Wang S and Fang J: MicroRNA-128 inhibits EMT of human osteosarcoma cells by directly targeting integrin $\alpha 2$. *Tumour Biol* 37: 7951-7957, 2016.
32. Fasching PA, Heusinger K, Loehberg CR, Wenkel E, Lux MP, Schrauder M, Koscheck T, Bautz W, Schulz-Wendtland R, Beckmann MW, *et al*: Influence of mammographic density on the diagnostic accuracy of tumor size assessment and association with breast cancer tumor characteristics. *Eur J Radiol* 60: 398-404, 2006.
33. Chang JM, Park IA, Lee SH, Kim WH, Bae MS, Koo HR, Yi A, Kim SJ, Cho N and Moon WK: Stiffness of tumours measured by shear-wave elastography correlated with subtypes of breast cancer. *Eur Radiol* 23: 2450-2458, 2013.
34. Evans A, Whelehan P, Thomson K, McLean D, Brauer K, Purdie C, Jordan L, Baker L and Thompson A: Quantitative shear wave ultrasound elastography: Initial experience in solid breast masses. *Breast Cancer Res* 12: R104, 2010.