Secreted amphiregulin promotes vincristine resistance in oral squamous cell carcinoma

MING-JU HSIEH^{1-4*}, YIN-HONG CHEN^{5*}, I-NENG LEE⁶, CHENG HUANG^{7,8}, YU-JU KU⁹ and JUI-CHIEH CHEN¹⁰

¹Oral Cancer Research Center, Changhua Christian Hospital, Changhua 500; ²Institute of Medicine,

Chung Shan Medical University, Taichung 402; ³Graduate Institute of Biomedical Sciences,

China Medical University, Taichung 404; ⁴Department of Holistic Wellness, Mingdao University, Changhua 52345;

⁵Department of Otorhinolaryngology-Head and Neck Surgery, Changhua Christian Hospital, Changhua 500;

⁶Department of Medical Research, Chang Gung Memorial Hospital, Chiayi 61363; ⁷Department of Biotechnology and

Laboratory Science in Medicine, National Yang-Ming University, Taipei 112; ⁸Department of Earth and Life Sciences,

University of Taipei, Taipei 100; ⁹The Center for General Education of China Medical University, China Medical University,

Taichung 404; ¹⁰Department of Biochemical Science and Technology, National Chiayi University,

Chiayi 60004, Taiwan, R.O.C.

Received February 27, 2019; Accepted August 1, 2019

DOI: 10.3892/ijo.2019.4866

Abstract. Oral squamous cell carcinoma (OSCC) is the most common type of oral cancer. Despite advances in surgery, radiotherapy and chemotherapy, the overall 5-year survival rate of patients with OSCC has not significantly improved. In addition, the prognosis of patients with advanced-stage OSCC remains poor. Therefore, it is necessary to develop novel therapeutic modalities. Vincristine (VCR), a naturally occurring vinca alkaloid, is a classical microtubule-destabilizing agent and is widely used in the treatment of a number of cancers. Despite the proven antitumor benefits of VCR treatment, one of the major reasons for the failure of treatment is drug resistance. Changes in the tumor microenvironment are responsible for cross-talk between cells, which may facilitate drug resistance in cancers; secreted proteins may promote communication between cancer cells to induce the development of resistance. To identify the secreted proteins involved in VCR resistance, conditioned media was obtained, and an antibody array was conducted to screen a comprehensive secretion profile between VCR-resistant (SAS-VCR) and parental (SAS) OSCC cell lines. The results showed that amphiregulin (AREG) was highly expressed and secreted in SAS-VCR cells. Pretreatment with exogenous recombinant AREG markedly increased drug

*Contributed equally

Key words: oral cancer, oral squamous cell carcinoma, vincristine, resistance, secreted proteins, amphiregulin

resistance against VCR in OSCC cells, as assessed by an MTT assay. Colony formation, MTT and western blot assays were performed to investigate the effects of AREG knockdown on VCR sensitivity. The results indicated that AREG expression can regulate VCR resistance in OSCC cells; overexpression of AREG increased VCR resistance in parental cells, whereas AREG knockdown decreased the VCR resistance of resistant cells. In addition, it was also demonstrated that the glycogen synthase kinase- 3β pathway may be involved in AREG-induced VCR resistance. These findings may provide rationale to combine VCR with blockade of AREG-related pathways for the effective treatment of OSCC.

Introduction

Oral cancer refers any cancerous cells that are located in the oral cavity. It is a type of head and neck cancer, accounting for most head and neck cancers and leading to >145,400 cases/year of mortality globally (1). Oral squamous cell carcinoma (OSCC) is the most common type of malignancy in the oral cavity (2). Conventional treatment of OSCC includes surgery, radiotherapy and chemotherapy (3). Although the clinical outcome of patients with OSCC has gradually improved in the last few years, the prognosis of patients with advanced-stage disease remains poor, reflecting limited advances in present understanding of the pathogenesis of this disorder (4). Therefore, it is necessary to develop novel therapeutic approaches for patients with advanced and unresectable OSCC.

Vincristine (VCR), a naturally occurring *Vinca* alkaloid, is a classical microtubule-destabilizing agent (5). It is widely used for hematologic malignancies and certain solid tumors (6-8). However, despite the proven antitumor activity of VCR treatment, the efficacy of chemotherapy is often limited by the rapid emergence of acquired resistance and patient relapse following initial response.

Correspondence to: Professor Jui-Chieh Chen, Department of Biochemical Science and Technology, National Chiayi University, 300 Syuefu Road, Chiayi 60004, Taiwan, R.O.C. E-mail: jcc@mail.ncyu.edu.tw

Tumors consist of a complex microenvironment composed of cancer cells, stroma and immune cells (9). Drug resistance is a complex process involving reciprocal interplay between different types of cells. Secreted proteins are responsible for the cross-talk among cells, which may facilitate drug resistance in tumors (10-12).

Several studies have indicated that soluble mediators from the microenvironment can promote cancer growth and therapy resistance (13-15). Amphiregulin (AREG) (16), a ligand of the epidermal growth factor (EGF) receptor (EGFR), is synthesized as a transmembrane precursor that undergoes a series of proteolytic steps to produce mature forms for secretion (16). AREG has been reported to induce oncogenic effects in numerous cancer cell types, including breast, liver, pancreatic and colorectal cancer cells (17-20) and to be implicated in drug resistance (10,18,21,22); however, the effects of AREG and its mechanisms of action in OSCC cells remain unknown. Understanding the complex mechanisms underlying its effects may reveal potential therapeutic opportunities.

Antibody arrays possess valuable applications in cancer research to identify biomarkers or molecules that are potentially relevant for diagnosis, prognosis, treatment and drug development (23). To elucidate the association between secreted proteins and VCR resistance in OSCC cells, a VCR-resistant SAS subline (SAS-VCR) was established by exposure to an increasing drug concentration gradient. Conditioned medium (CM) was collected from parental and VCR-resistant cells, and the secreted proteins were assessed using an antibody array. In the present study, comprehensive secretion profiling was performed to provide novel insight in the mechanisms of VCR resistance. Understanding the relationship between secreted proteins and drug resistance may contribute to the development of novel therapeutic strategies and biomarkers in OSCC.

Materials and methods

Chemicals, reagents and antibodies. VCR, 5-fluorouracil (5-FU), cisplatin, MTT and dimethyl sulfoxide (DMSO) were purchased from Sigma-Aldrich (Merck KGaA). VCR was dissolved in sterile PBS and diluted in cell culture medium to the required concentration prior to use. 5-FU was dissolved in DMSO. Cisplatin was dissolved in dimethylformamide. A potent glycogen synthase kinase-3 (GSK-3) inhibitor, LY2090314 (Selleck Chemicals), was dissolved in DMSO. The antibodies used in this study were as follows: Cleaved poly (ADP-ribose) polymerase (1:1,000; PARP; cat. no. 9541) was purchased from Cell Signaling Technology, Inc.; AREG (1:200; cat. no. sc-74501), Bcl-2 (1:200; cat. no. sc-7382), phosphorylated (p)-GSK-3ß (1:200; cat. no. sc-135653) and GSK-3ß (1:200; cat. no. sc-9166) were purchased from Santa Cruz Biotechnology, Inc.; α-tubulin (1:10,000; cat. no. 05-829) was purchased from EMD Millipore. Human recombinant AREG (rAREG) was purchased from R&D Systems, Inc.

Cell culture. The human OSCC cell lines, SAS and SCC9, were kindly provided by Dr Ming-Chang Hsieh (Chung Shan Medical University Hospital, Taichung, Taiwan). All cells were cultured in DMEM/F12 medium (Gibco; Thermo Fisher Scientific, Inc.) supplemented with 10% fetal bovine serum (FBS; Gibco; Thermo Fisher Scientific, Inc.), 1%

penicillin-streptomycin (10,000 U/ml penicillin and 10 mg/ml streptomycin) and 2 mM glutamine, and maintained at 37° C in a humidified atmosphere of 5% CO₂. To investigate the mechanism of VCR resistance in OSCC, SAS-VCR and SCC9-VCR cells were established over ~6 months by gradually increasing the concentration of VCR in the culture medium by 0.5 to 16 nM and 0.125 to 4 nM, respectively.

Cell viability assay and treatments. An MTT assay was conducted to evaluate the effects of VCR on the viability of OSCC cells. Briefly, cells were seeded into the wells of 96-well plates at a density of 5,000 cells in 100 μ l of culture medium. Following overnight incubation to allow the attachment of cells, cells were incubated with 0-64 nM VCR in serum-free medium at 37°C. At 0-48 h time intervals following VCR treatment, 30 μ l of MTT (5 mg/ml) was added to each well and incubated for a further 4 h at 37°C. The supernatant was then discarded, and 100 μ l DMSO was added to each well to dissolve the formazan crystals. The optical density was evaluated by measuring the absorbance, with a test wavelength of 490 nm and a reference wavelength of 630 nm.

To investigate whether CM can enhance VCR resistance, CM was collected as described in the 'Collection of CM' section. SAS cells were activated by CM from SAS and SAS-VCR cells for 8 h, and then treated with 8 μ M VCR for 24 h.

To further determine the relationship between AREG and VCR resistance, SAS cells were pretreated with SAS-VCR-CM in combination with or without AREG-neutralizing antibody (1:500) for 8 h, and then treated with 8 μ M VCR for 24 h.

For the effect of rAREG on VCR sensitivity, SAS cells were pretreated with rAREG (50 ng/ml) for 4 h and then stimulated with 16 μ M VCR for 48 h. SCC9 cells were pretreated with rAREG (100 ng/ml) for 4 h and then stimulated with 4 μ M VCR for 24.

To analyze whether serum starvation can induce expression of AREG, 5x10⁵ cells/well were seeded in 6-well plates and cultured overnight. Following attachment, cells were washed twice with PBS, and then serum-free DMEM/F12 was added. Cells were analyzed by MTT and western blot assays after 12, 24 and 48 h.

To investigate whether inhibition of GSK3- β activation can block AREG-induced VCR resistance, cells were pretreated with LY2090314 (20 nM) for 30 min, followed by treatment with rAREG (50 ng/ml) for 4 h and then treatment with 4 μ M VCR for 24 h. Cell viability was assessed by an MTT assay.

To investigate the sensitivity of SAS and SAS-VCR cells to 5-FU and cisplatin, cells were treated with 0-160 μ M 5-FU for 48 h or 0-20 μ M cisplatin for 24 h, and then analyzed using an MTT assay.

Colony formation assay. Cells were cultured a 6-well plate at a density of $5x10^4$ cells/well with regular medium. Cells were treated with the 0-64 nM VCR concentrations for 24 h. Then, the cells were seeded at $1x10^5$ cells/well into a separate 6-well plate. Cells were allowed to grow until colonies were visible (5-6 days) and then fixed with methanol for 30 min and stained with 0.5% crystal violet for 30 min at room temperature (RT). Images of the colonies were acquired using a digital camera. Colonies were counted using ImageJ 1.52a software (National Institutes of Health). Western blot analysis. Cells were lysed in RIPA buffer containing protease inhibitor cocktail (Thermo Fisher Scientific, Inc.). The protein concentration was determined using a PierceTM BCA Protein Assay kit (Thermo Fisher Scientific, Inc.). Proteins from the total cell lysates or CM (40 μ g/lane) were separated by via 8-15% SDS-PAGE and electrotransferred onto PVDF membranes. The membranes were blocked with 5% skim milk for 1 h at RT and then probed with the indicated primary antibodies for 1 h at RT. Following three washes with TBS-0.1% Tween 20, the membranes were incubated with the HRP-conjugated anti-mouse (cat. no. 20102) or anti-rabbit IgG antibody (cat. no. 20202; both 1:5,000; Leadgene Biomedical, Inc.) for 1 h at RT. The blots were visualized using ECL reagent (PerkinElmer, Inc.) and autoradiography.

Collection of CM. Cells seeded in 10-cm dishes were grown to 80% confluence and then washed with PBS twice. Cells were subsequently incubated in serum-free media for 48 h. CM was collected via gentle aspiration and then centrifuged at 875 x g at RT for 10 min to remove cell debris. The CM was further concentrated using Amicon[®] Ultra 15 ml centrifugal filters with a 3-kDa cut-off (EMD Millipore) at 4,000 x g and 4°C to a total volume of ~150-200 ul. The CM was aliquoted and stored at -20°C prior to use.

Growth factor human antibody array. A human growth factor antibody array (cat. no. ab134002; Abcam) was used according to the manufacturer's protocols; all reagents listed below were included in this array unless otherwise specified. Briefly, the concentrated CM (200 μ g total protein) was mixed with blocking buffer and incubated with membranes at 4°C overnight. Membranes were then washed and incubated with biotin-conjugated anti-cytokines for 2 h at RT, followed by washing and incubation with HRP-conjugated streptavidin for 2 h at RT. Membranes were then washed again, and bound antibodies were visualized using ECL reagents and autoradiography. The relative expression was determined using UN-SCAN-IT gel 6.1 software (Silk Scientific, Inc.). The average signal of a pair of duplicate spots was normalized using negative control spots as a background value. The relative intensities in SAS-VCR cells were determined by comparing the corresponding signals to SAS cells.

Oncomine database. The Oncomine Cancer Microarray database (http://www.oncomine.org/) was used to study the expression levels of AREG in human oral tumor and normal tissues obtained from separate individuals. The gene expression data were log transformed, median centered per array, and the standard deviation was normalized to 1 per array. A gene was considered as overexpressed when its mean value in tumor samples was significantly increased compared with its mean value in normal tissue as determined using a t-test (P \leq 0.05) and the fold change was \geq 1.5.

Lentivirus infection and short hairpin (sh)RNA knockdown. The pLKO.1-puro-based lentiviral plasmids containing TRCN0000117995-shAREG (sequence: 5'-CCGGGAACG AAAGAAACTTCGACAACTCGAGTTGTCGAAGTTTCT TTCGTTCTTTTG-3')andpLKO.1-shScramble(sequence:5'-CCG GCCTAAGGTTAAGTCGCCCTCGCTCGAGCGAGGGCG ACTTAACCTTAGGTTTTT-3') were obtained from National RNAi Core Facility (Academia Sinica, Taipei, Taiwan). All plasmids (4 μ g lentiviral plasmid; 4 μ g pCMV Δ R8.91; 0.4 μ g pMD; all Academia Sinica) were cotransfected into 293T cells (5x10⁵; cat. no. 632180; Clontech Laboratories, Inc.) in 6-cm dishes using TurboFect (Thermo Fisher Scientific, Inc.), according to the manufacturer's protocols. The lentivirus-containing supernatants were harvested at 48 h post-transfection. SAS-VCR cells (1x10⁵) were infected using the lentivirus-containing supernatant (12,762 RIU/ μ l). For stable cell lines, the infected cells were selected by puromycin (5 μ g/ml) within 1 week.

AREG overexpression. An AREG overexpression plasmid (pCMV3-AREG) and negative control (pCMV3) were purchased from Sino Biological Inc. Plasmids were transfected into the two OSCC cell lines using TurboFect according to the manufacturer's protocol. Cells ($1x10^5$) were seeded in each well of a 24-well plate. After culturing for 24 h, cells were transiently transfected under optimized transfection conditions. Briefly, 1 μ g of DNA plasmid DNA was diluted in 100 μ l of serum-free DMEM/F12, and mixed with 2 μ l of transfection reagent followed by incubation at RT for 20 min. The mixture was then added dropwise to the cells and incubated for an additional 72 h at 37°C in a humidified atmosphere and 5% CO₂.

Statistical analysis. All values represent the mean \pm SEM from at least 3 independent experiments. Student's t-test was used when comparing two independent groups. Statistical comparisons of >2 groups were performed using one-way analysis of variance (ANOVA) with Bonferroni's *post hoc* test. In all cases, P<0.05 was considered to indicate a statistically significant difference.

Results

Establishment and characterization of VCR-resistant SAS cells. As presented in Fig. 1A, SAS-VCR cells (IC₅₀>1,024 nM) were more resistant to VCR than their respective parental SAS cells (IC₅₀= 63.96 ± 0.25 nM). In addition, a colony formation assay was also conducted. As presented in Fig. 1B, the colony numbers of SAS cells were significantly decreased compared with control treatment in dose-dependent manner; however, the SAS-VCR colony number was significantly decreased compared with the control only in the high dose group (64 nM). Next, the effects of VCR on the expression of cleaved PARP, a marker of apoptosis, and Bcl-2 (an antiapoptotic protein) were evaluated via western blotting. As shown in Fig. 1C, the expression of cleaved PARP was notably induced in VCR-treated SAS cells. In contrast, the expression of cleaved PARP was not observed in SAS-VCR cells following exposure to 8 or 16 nM VCR for 24 h, indicating that SAS-VCR cells were highly resistant to VCR compared with SAS cells. The expression of Bcl-2 was increased in a dose-dependent manner in VCR-treated SAS-VCR cells, whereas the expression of Bcl-2 was decreased in a dose-dependent manner in VCR-treated SAS cells. These results indicated that SAS-VCR cells were resistant to VCR compared with SAS cells. Furthermore, the



Figure 1. Cytotoxic effects of VCR on SAS and SAS-VCR cells. (A) Cells were treated with various concentrations of VCR for 48 h and cell viability was measured by an MTT assay. Data are presented as the mean \pm SEM of four independent experiments. *P<0.05 vs. SAS. (B) Long-term effects of VCR were assessed using a colony formation assay. Above, representative images showing that SAS cells formed fewer colonies compared with SAS-VCR cells. Below, densitometric analysis of the clonogenic growth of SAS and SAS-VCR cells. Data are presented as the mean \pm SEM of three independent experiments. *P<0.05 vs. 0 μ M of the respective cell type. (C) Cells were treated with VCR (0, 8 or 16 μ M) for 24 h, and the expression levels of apoptosis-associated proteins (cleaved PARP and Bcl-2) were examined via western blotting. α -tubulin was used as a loading control. PARP, poly (ADP-ribose) polymerase; SAS-VCR, VCR-resistant SAS cells; VCR, vincristine.

sensitivity of the SAS-VCR cell line to 5-FU and cisplatin was also explored, and it was revealed that only resistance to 5-FU was observed in these cells, suggesting a potential link between resistance to VCR and 5-FU resistance (Fig. S1).

Comparison of growth factor profiles in the secretomes of SAS and SAS-VCR cells. To evaluate whether the secreted proteins from resistant cells were associated with the induction of drug resistance, CM was obtained from SAS and SAS-VCR cells. Notably, the CM from SAS-VCR cells significantly increased VCR resistance when applied to parental SAS cells (Fig. 2A). Subsequently, antibody arrays were used to analyze the differences in the secretomes of CM from SAS and SAS-VCR cell lines. As presented in Fig. 2B, a total of 22 secreted proteins were identified whose expression was changed >1.5-fold between the two media. The levels of 17 secreted proteins were increased, and those of 5 secreted proteins was decreased in SAS-VCR. Of these, the levels of AREG, basic fibroblast growth factor, heparin-binding EGF-like growth factor, platelet-derived growth factor-AB, placental growth factor and vascular endothelial growth factor D secretion were most notably upregulated (>4-fold) in SAS-VCR CM, indicating that the secreted proteins may be important mediators of VCR resistance in OSCC cells.

AREG is highly expressed and secreted to promote VCR resistance in OSCC cells. To determine the clinical relevance of these secreted proteins, the Oncomine database was employed to select the appropriate target for further study. The results revealed that there were only clinical data concerning

the expression of AREG; its expression was significantly upregulated in carcinoma tissue compared normal oral cavity tissue (Fig. 2C). In addition, among the six proteins, numerous studies indicated that AREG serves a critical role in OSCC (24-26); however, its role in VCR sensitivity is yet to be described in the literature. To further confirm the antibody array results, western blotting was performed to determine if AREG was highly expressed and secreted in SAS-VCR cells. As shown in Fig. 2D, the expression of AREG in the cell lysates was analyzed, and the results indicated that expression of AREG was increased in SAS-VCR cells compared with in SAS cells. In addition, the levels of AREG in CM were also significantly elevated for SAS-VCR cells compared with SAS cells (Fig. 2E). To provide further evidence that AREG mediates VCR sensitivity, AREG activity was blocked using a neutralizing antibody. The results showed that pretreatment of CM with neutralizing antibodies against AREG restored VCR sensitivity in SAS cells (Fig. 2F).

AREG also modulates VCR sensitivity in SCC9 cells. To further validate the role of AREG in VCR sensitivity, SAS cells were pretreated with rAREG and then stimulated with VCR. The results showed that AREG markedly increased resistance to VCR in SAS cells (Fig. 3A). Similarly, rAREG also enhanced VCR resistance in SCC9 cells (Fig. 3B). Furthermore, the expression levels of AREG and VCR resistance in the two cell lines were measured. The results showed that the expression of AREG was higher in the more VCR-resistant cell line, SAS (Fig. 3C). To clarify whether AREG expression is associated with VCR



Figure 2. Growth factors secreted by SAS-VCR cells may promote resistance to VCR. (A) Secreted substances of SAS-VCR cells can promote VCR resistance in SAS cells. Data are presented as the mean \pm SEM of three independent experiments. *P<0.05 vs. Non; #P<0.05 vs. VCR. (B) Comparison of growth factors in CM from SAS and SAS-VCR cells was performed using a human growth factor antibody array. The expression of the marked molecules was altered. Below, densitometric analysis of the pair of duplicate spots representing each marked protein. (C) Box plots derived from gene expression data from the Oncomine cancer database comparing expression of AREG gene in normal and carcinoma tissue. The fold change is 2.668 and P-value is 6.15x10⁻⁹. (D) Analysis of AREG levels in total cell lysate of SAS and SAS-VCR as determined via western blotting. α -Tubulin was used as a loading control. (E) Equal volumes of CM from SAS and SAS-VCR cells was analyzed for secreted AREG via western blotting. Loading quantities were shown on left side by Ponceau S staining. (F) Cells were pretreated with SAS-VCR-CM and/or AREG-neutralizing antibody, and then treated with 8 μ M VCR for 24 h. Cell viability was determined by an MTT assay. Data are presented as the mean \pm SEM of at least four independent experiments. *P<0.05 vs. Non; #P<0.05 vs. VCR; &P<0.05 vs. SAS-VCR -CM + VCR. Ab, antibody; AREG, amphiregulin; CM, condition medium; Non, untreated cells; SAS-VCR, VCR-resistant SAS cells; VCR, vincristine.

sensitivity in SCC9 cells, a VCR-resistant SCC9 subline termed SCC9-VCR was established. According to the results of the MTT assay, SCC9-VCR cells were significantly more viable following VCR treatment compared with SCC9 cells, as assessed at 24 and 48 h (Fig. 3D and E). Furthermore, SCC9 cells exhibited increased expression of cleaved PARP compared with SCC9-VCR cells following VCR treatment (Fig. 3F). To further evaluate the association between AREG levels and VCR resistance, SCC9 and SCC9-VCR cells were treated with increasing doses of VCR, and then the expression of AREG was analyzed via western blotting. The results revealed that SCC9-VCR cells exhibited upregulated AREG expression compared with SCC9 cells. Furthermore, an increase in the expression levels of AREG was observed in SCC9-VCR cells, but not SCC9 cells, after treatment with increasing doses of VCR (Fig. 3G).

Knockdown of AREG restores VCR sensitivity, and overexpression of AREG confers resistance to VCR. To determine the impact of AREG expression on VCR sensitivity, AREG expression was suppressed in SAS-VCR cells using a lentivirus-mediated RNA interference system. The knockdown efficiency of shAREG was evaluated via western blotting (Fig. 4A). Silencing AREG expression resulted in elevated sensitivity to VCR in a dose-dependent manner compared with shControl cells, as determined by MTT (Fig. 4B) and colony formation assays (Fig. 4C). Furthermore, AREG knockdown notably promoted cell apoptosis in SAS-VCR cells, as



Figure 3. AREG is involved in VCR resistance in oral squamous cell carcinoma cells. Pretreatment of rAREG confers VCR resistance in (A) SAS and (B) SCC9 cells. SAS cells were pretreated with rAREG (50 ng/ml) for 4 h and then stimulated with 16 μ M VCR for 48 h (n=6). SCC9 cells were pretreated with rAREG (100 ng/ml) for 4 h and then stimulated with 4 μ M VCR for 24 h (n=3). Cell viability was examined by an MTT assay. Data are presented as the mean ± SEM. *P<0.05 vs. Non; *P<0.05 vs. VCR. (C) Left, AREG levels in cell lysates from SCC9 and SAS cells were analyzed via western blotting. Right, SCC9 and SAS cells were exposed to indicated doses of VCR for 24 h and assayed for survival by an MTT assay. Data are presented as the mean ± sEM independent experiments. *P<0.05 vs. SCC9. Comparison of VCR sensitivity between parental SCC9 and SCC9-VCR cells at (D) 24 and (E) 48 h as determined by an MTT assay (n=4). *P<0.05 vs. SCC9. (F) SCC9 and SCC9-VCR cells were treated with various concentrations of VCR (0-16 μ M), for 24 h and the expression levels of cleaved PARP were examined via western blotting. (G) Dose-dependent effects of VCR on the expression of AREG in SCC9 and SCC9-VCR cells were analyzed via western blotting. α -tubulin was used as a loading control. AREG, amphiregulin; Non, untreated cells; PARP, poly (ADP-ribose) polymerase; r, recombinant; SAS-VCR/SCC9-VAR, VCR-resistant SAS/SCC9 cells; VCR, vincristine.

evidenced by an increase in the cleavage of PARP at a concentration of 64 nM VCR for 24 h (Fig. 4D). The results indicated that the knockdown of AREG may increase VCR-induced apoptosis in OSCC cells. To verify the relevance of AREG in mediating resistance to VCR, cells were then transiently transfected with AREG expression vector or control vector. AREG was overexpressed in SAS and SCC9 cells following transfection, as determined via western blotting (Fig. 4E). As presented in Fig. 4F, overexpression of AREG in SAS and SCC9 cells significantly increased resistance to VCR, as assessed using an MTT assay. Furthermore, whether AREG overexpression protected against starvation-induced death in OSCC cells was analyzed. The result revealed that overexpression of AREG significantly increased the viability of serum-starved SAS cells at 48 h compared with cells transfected with a control vector (Fig. 4G). Furthermore, whether serum starvation can affect the expression of AREG, which may prevent cell death and promote resistance to harsh environments such as drug treatment, was evaluated. As presented in Fig. 4H, serum deprivation notably induced AREG expression in SAS and SCC9 cells in a time-dependent manner. These results indicated that AREG was involved in protecting OSCC cells against various stresses, including VCR treatment.

AREG regulates VCR sensitivity in OSCC cells via activation of GSK- 3β . A previously study reported that AREG can modulate the GSK-3 β pathway to regulate cell functions; GSK-3 β is known to be a potential therapeutic target for cancer treatment (27). In addition, it has been demonstrated that targeting the GSK-3 β pathway may be beneficial for the treatment of oral cancer (28). Therefore, the activation of GSK-3^β was analyzed after treatment of SAS and SAS-VCR cells with increasing concentrations of VCR. The results showed that VCR induced a dose-dependent decrease in GSK-3^β phosphorylation in SAS cells, whereas GSK-3^β maintained sustained activation in SAS-VCR cells (Fig. 5A). In addition, a marked downregulation of p-GSK-3β was also observed in VCR-treated SAS-VCR/shAREG cells compared with SAS-VCR/shControl cells (Fig. 5B). To further confirm that AREG can indeed activate the GSK-3ß pathway, GSK-3ß phosphorylation was directly analyzed in response to rAREG. The results revealed that treatment of SAS cells with rAREG can induce an increase in the phosphorylation of GSK-3 β in a time-dependent manner (Fig. 5C). Then, a GSK-3ß inhibitor, LY2090314, was used to interfere with AREG-induced GSK-3ß activation and observe whether blocking GSK-3ß activation would affect the role of AREG in VCR resistance. It was demonstrated that rAREG induced a significant increase in VCR resistance; however, this effect was significantly attenuated by LY2090314 (Fig. 5D). These findings suggested that AREG can regulate the activation of GSK-3 to promote VCR resistance in OSCC cells.



Figure 4. Effects of knockdown or overexpression of AREG on VCR sensitivity in oral squamous cell carcinoma cells. (A) Efficiency of AREG knockdown was verified via western blotting. (B) Cells were exposed to increasing concentrations of VCR for 48 h and subsequently evaluated by an MTT assay (n=4). *P<0.05 vs. SAS-VCR/shAREG. (C) SAS-VCR/shAREG and SAS-VCR/shControl were treated with the indicated concentrations of VCR and then subjected to colony formation assays. (D) SAS-VCR/shControl and SAS-VCR/shAREG cells were treated with increasing concentrations of VCR for 24 h, and cleaved PARP expression was analyzed via western blotting. (E) Expression of AREG was analyzed in whole cell lysates using western bloting. α-Tubulin was used as a loading control. (F) Overexpression of AREG induces VCR resistance in SAS and SCC9 cells. SAS cells were treated with the indicated concentrations of VCR for 48 h. SCC9 cells were treated with the indicated concentrations of VCR for 24 h. Data are presented as the mean ± SEM of four independent experiments. *P<0.05 vs. pCMV. (G) MTT assay to determine the effects of AREG overexpression on the viability of serum-starved SAS cells (n=4). *P<0.05 vs. pCMV at 48 h. (H) Effects of serum starvation on AREG expression in OSCC cells. Cells were cultured in serum-free medium for various durations and then analyzed via western blotting. AREG, amphiregulin; PARP, poly (ADP-ribose) polymerase; SAS-VCR, VCR-resistant SAS cells; sh, short hairpin (RNA); VCR, vincristine.

Discussion

A substantial body of evidence has revealed that upregulated expression of AREG is associated with cancer progression in a wide variety of cancers, including lung (29,30), breast (17,31), ovarian (32,33), liver (18,34), pancreatic (19,35) and colorectal cancers (20,36). At present, however, there has been no study into the function of ARGE in relation to drug resistance in OSCC. In the present study, an antibody array was performed to explore the secretion profile of OSCC cells, and it was revealed that AREG was highly expressed and secreted in VCR-resistant cells. In addition, it was also suggested that the GSK-3 β pathway may be involved in AREG-induced VCR resistance. These findings may aid the development of novel therapeutic targets for OSCC treatment and improved prognosis.

Chemotherapy is widely used in the treatment of OSCC, and VCR is a classic microtubule-destabilizing agent that is effective and widely used in hematological malignancies and certain solid tumors (37). As VCR exhibits substantial anticancer activity, it may be a potential treatment for OSCC. However, several studies have indicated that high-dose VCR is associated with a significant risk of severe gastrointestinal toxicity; mortality from treatment-related toxicity has been previously reported in patients (38,39). Therefore, combining VCR with other agents or molecules is may improve the clinical management of oral cancers. Certain drugs with antitumor activity have been reported to increase the VCR sensitization of VCR-resistant oral epidermoid carcinoma cells (40-42). Though great efforts have been made in developing novel anticancer drugs with increased curative potential or the ability to reverse drug resistance, the results remain satisfactory, due to either a lack of potency or unacceptable side effects.

Proteins secreted, shed or leaking from cancer cells, collectively termed the cancer secretome, are considered promising biomarkers, as they may be detectable in blood or other body fluids (43-45). Previous studies indicated that cancer cells can secrete soluble mediators in response to drug therapy, which contribute to the promotion of drug resistance and tumor progression (46,47). In addition, secreted proteins



Figure 5. GSK3- β signaling pathways may be involved in the effects of AREG on VCR sensitivity. (A) SAS and SAS-VCR cells were treated with different concentrations of VCR (0-64 nM) for 24 h. (B) SAS-VCR/shControl and SAS-VCR/shAREG cells were exposed to various concentrations of VCR for 24 h. (C) SAS cells were incubated with rAREG (50 ng/ml) for the indicated time intervals. p-GSK3- β and GSK3- β were analyzed via western blotting. (D) Cells were pretreated with Ly (20 nM) for 30 min, followed by treatment with rAREG (50 ng/ml) for 4 h, and then treatment with 4 μ M VCR for 24 h. Cell viability was evaluated by an MTT assay. Data are presented as the mean ± SEM (n=5). *P<0.05 vs. Non; *P<0.05 vs. VCR; &P<0.05 vs. rAREG + VCR. AREG, amphiregulin; GSK-3 β , glycogen synthase kinase-3 β ; Ly, LY2090314; Non, untreated cells; p, phosphorylated; r, recombinant; SAS-VCR, VCR-resistant SAS cells; VCR, vincristine.

are also considered good candidate serological tumor markers, as they are released by the cells and thus exhibit the greatest possibility of entering the circulation (48). Therefore, secreted proteins are regarded as a rich source of potential markers and drug targets for cancer treatment (49).

AREG is an EGF-like ligand that has been identified as a ligand responsible for EGFR-ERK signaling activation, which can lead to cancer progression (19,50-52). Regarding drug sensitivity, AREG is upregulated in non-responding patients compared with patients who do respond to gefitinib (21). In animal models, a previous study has shown that AREG silencing can reduce the size of tumor growth and increase drug sensitivity in glioma cells (53). As AREG is a secreted protein, it can enter the circulatory system. A separate study showed that circulating AREG could be clinically relevant as an indicator of unfavorable response to gefitinib in NSCLC (54). In Taiwan, oral cancer accounts for the fourth highest incidence of malignancy in males (55). Numerous studies indicate that EGFR is frequently overexpressed in human OSCC (56-59).

High EGFR expression has been associated with resistance to chemotherapeutic agents used in the treatment of OSCC, including cisplatin, 5-FU, cyclophosphamide and doxorubicin, suggesting that EGFR signaling may be a promising target for OSCC therapy (58-60). Cetuximab is a chimeric IgG1-human antibody targeted against the extracellular domain of EGFR (61). It was approved by the FDA in 2006 as a component of combination therapy along with radiation and/or chemotherapy to treat OSCC; however, clinical use of cetuximab is limited, as EGFR expression levels have not been associated with response levels to cetuximab (62,63). At present, in OSCC, clinically relevant mechanisms of cetuximab resistance have not been clearly elucidated.

GSK- 3β , a multifunctional serine/threonine kinase, was originally discovered as a key regulator of glycogen metabolism (64). Accumulating evidence suggests that it involved in tumorigenesis, migration, invasion, chemotherapy and drug resistance (65,66). Therefore, GSK-3β has emerged as a potential therapeutic target for different types of cancers (67,68). Recent research in OSCC has shown that GSK-3ß can regulate matrix metalloproteinase-9 activity to promote cancer progression and invasion (69). In clinical specimens, another study also showed that the levels of p-GSK-3β and GSK-3β in OSCC tissues are upregulated compared with in controls, and are positively associated with tumor metastasis and poor survival in patients (70). Although a number of studies have indicated that multidrug resistance can be reversed by inhibiting GSK-3 β (71-73), whether this occurs in OSCC remains unclear. The relationship of GSK-3 β with the drug treatment of OSCC remains to be further explored in future studies.

In conclusion, to improve understanding of the mechanisms underlying drug resistance in OSCC, a VCR-resistant OSCC cell line was established, and the secreted proteins were analyzed using an antibody microarray. This study is the first, to our knowledge, to characterize changes in the secretome of VCR-resistant OSCC cells. The results indicated that AREG was highly expressed and secreted in VCR-resistant cells compared with VCR-sensitive cells. Pretreatment with exogenous rAREG markedly increased drug resistance against VCR in OSCC cells. Furthermore, knockdown of AREG increased VCR sensitivity, whereas overexpression of AREG further promoted VCR resistance. The results indicated that AREG contributes to VCR resistance in OSCC cells. Additionally, it was also demonstrated that the GSK- 3β pathway may be involved in AREG-induced VCR resistance. These findings may provide valuable insight for the development of effective treatments against OSCC.

Acknowledgements

The author would like to thank Ms Yu-Ping Wu and Ms Wen-Ling Li at the National Chiayi University for their assistance.

Funding

This work was supported by the Ministry of Science and Technology (grant no. 107-2314-B-415-002-) and grant nos. 107-CCH-HCR-046 and 107-CCH-HCR-047 from Changhua Christian Hospital, Taiwan, R.O.C.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

JCC was involved in the study design, data analysis and drafting the manuscript. MJH and YHC performed the experiments and data acquisition. INL, CH and YJK contributed to data analysis and interpretation.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- 1. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J and Jemal A: Global cancer statistics, 2012. CA Cancer J Clin 65: 87-108, 2015.
- 2. Curado MP and Hashibe M: Recent changes in the epidemiology of head and neck cancer. Curr Opin Oncol 21: 194-200, 2009.
- 3. Scully C and Bagan J: Oral squamous cell carcinoma overview. Oral Oncol 45: 301-308, 2009.
- 4. Schwam ZG and Judson BL: Improved prognosis for patients with oral cavity squamous cell carcinoma: Analysis of the National Cancer Database 1998-2006. Oral Oncol 52: 45-51, 2016.
- Dumontet C and Jordan MA: Microtubule-binding agents: A dynamic field of cancer therapeutics. Nat Rev Drug Discov 9: 790-803, 2010.
- Silverman JA, Reynolds L and Deitcher SR: Pharmacokinetics and pharmacodynamics of vincristine sulfate liposome injection (VSLI) in adults with acute lymphoblastic leukemia. J Clin Pharmacol 53: 1139-1145, 2013.
 Yan Z, Zhu ZL, Qian ZZ, Hu G, Wang HQ, Liu WH and Cheng G:
- Yan Z, Zhu ZL, Qian ZZ, Hu G, Wang HQ, Liu WH and Cheng G: Pharmacokinetic characteristics of vincristine sulfate liposomes in patients with advanced solid tumors. Acta Pharmacol Sin 33: 852-858, 2012.
- Olasz L, Orsi E, Marko T and Szalma J: Induction chemotherapy response and recurrence rates in correlation with N0 or N+ stage in oral squamous cell cancer (OSCC). Cancer Metastasis Rev 29: 607-611, 2010.

- 9. Villanueva J and Herlyn M: Melanoma and the tumor microenvironment. Curr Oncol Rep 10: 439-446, 2008.
- Chen JC, Lee IN, Huang C, Wu YP, Chung CY, Lee MH, Lin MH and Yang JT: Valproic acid-induced amphiregulin secretion confers resistance to temozolomide treatment in human glioma cells. BMC Cancer 19: 756, 2019.
- 11. Milman N, Ginini L and Gil Z: Exosomes and their role in tumorigenesis and anticancer drug resistance. Drug Resist Updat 45: 1-12, 2019.
- 12. Siveen KS, Raza A, Ahmed EI, Khan AQ, Prabhu KS, Kuttikrishnan S, Mateo JM, Zayed H, Rasul K, Azizi F, et al: The role of extracellular vesicles as modulators of the tumor microenvironment, metastasis and drug resistance in colorectal cancer. Cancers (Basel) 11: pii: E746, 2019.
- Wilson TR, Fridlyand J, Yan Y, Penuel E, Burton L, Chan E, Peng J, Lin E, Wang Y, Sosman J, *et al*: Widespread potential for growth-factor-driven resistance to anticancer kinase inhibitors. Nature 487: 505-509, 2012.
- 14. Straussman R, Morikawa T, Shee K, Barzily-Rokni M, Qian ZR, Du J, Davis A, Mongare MM, Gould J, Frederick DT, *et al*: Tumour micro-environment elicits innate resistance to RAF inhibitors through HGF secretion. Nature 487: 500-504, 2012.
- Acharyya S, Oskarsson T, Vanharanta S, Malladi S, Kim J, Morris PG, Manova-Todorova K, Leversha M, Hogg N, Seshan VE, *et al*: A CXCL1 paracrine network links cancer chemoresistance and metastasis. Cell 150: 165-178, 2012.
- Berasain C and Avila MA: Amphiregulin. Semin Cell Dev Biol 28: 31-41, 2014.
- 17. Willmarth NE and Ethier SP: Autocrine and juxtacrine effects of amphiregulin on the proliferative, invasive, and migratory properties of normal and neoplastic human mammary epithelial cells. J Biol Chem 281: 37728-37737, 2006.
- Berasain C, Castillo J, Perugorria MJ, Prieto J and Avila MA: Amphiregulin: A new growth factor in hepatocarcinogenesis. Cancer Lett 254: 30-41, 2007.
 Yotsumoto F, Fukami T, Yagi H, Funakoshi A, Yoshizato T,
- Yotsumoto F, Fukami T, Yagi H, Funakoshi A, Yoshizato T, Kuroki M and Miyamoto S: Amphiregulin regulates the activation of ERK and Akt through epidermal growth factor receptor and HER3 signals involved in the progression of pancreatic cancer. Cancer Sci 101: 2351-2360, 2010.
- 20. Kuramochi H, Nakajima G, Kaneko Y, Nakamura A, Inoue Y, Yamamoto M and Hayashi K: Amphiregulin and Epiregulin mRNA expression in primary colorectal cancer and corresponding liver metastases. BMC Cancer 12: 88, 2012.
- Chen JC, Huang C, Lee IN, Wu YP and Tang CH: Amphiregulin enhances cell migration and resistance to doxorubicin in chondrosarcoma cells through the MAPK pathway. Mol Carcinog 57: 1816-1824, 2018.
- 22. Tokunaga S, Nagano T, Kobayashi K, Katsurada M, Nakata K, Yamamoto M, Tachihara M, Kamiryo H, Yokozaki H and Nishimura Y: Amphiregulin as a novel resistance factor for amrubicin in lung cancer cells. Anticancer Res 37: 2225-2231, 2017.
- 23. Haab BB: Antibody arrays in cancer research. Mol Cell Proteomics 4: 377-383, 2005.
- 24. Tsai ST, Yang KY, Jin YT, Lin YC, Chang MT and Wu LW: Amphiregulin as a tumor promoter for oral squamous cell carcinoma: Involvement of cyclooxygenase 2. Oral Oncol 42: 381-390, 2006.
- 25. Zhang J, Wang Y, Chen X, Zhou Y, Jiang F, Chen J, Wang L and Zhang WF: MiR-34a suppresses amphiregulin and tumor metastatic potential of head and neck squamous cell carcinoma (HNSCC). Oncotarget 6: 7454-7469, 2015.
- 26. Gao J, Ulekleiv CH and Halstensen TS: Epidermal growth factor (EGF) receptor-ligand based molecular staging predicts prognosis in head and neck squamous cell carcinoma partly due to deregulated EGF-induced amphiregulin expression. J Exp Clin Cancer Res 35: 151, 2016.
- 27. Wang S, Zhang Y, Wang Y, Ye P, Li J, Li H, Ding Q and Xia J: Amphiregulin confers regulatory T cell suppressive function and tumor invasion via the EGFR/GSK-3β/Foxp3 Axis. J Biol Chem 291: 21085-21095, 2016.
- Mishra R: Glycogen synthase kinase 3 beta: Can it be a target for oral cancer. Mol Cancer 9: 144, 2010.
- Addison CL, Ding K, Zhao H, Le Maître A, Goss GD, Seymour L, Tsao MS, Shepherd FA and Bradbury PA: Plasma transforming growth factor alpha and amphiregulin protein levels in NCIC Clinical Trials Group BR.21. J Clin Oncol 28: 5247-5256, 2010.
- Hsu YL, Huang MS, Cheng DE, Hung JY, Yang CJ, Chou SH and Kuo PL: Lung tumor-associated dendritic cell-derived amphiregulin increased cancer progression. J Immunol 187: 1733-1744, 2011.

- McBryan J, Howlin J, Napoletano S and Martin F: Amphiregulin: Role in mammary gland development and breast cancer. J Mammary Gland Biol Neoplasia 13: 159-169, 2008.
- 32. D'Antonio A, Losito S, Pignata S, Grassi M, Perrone F, De Luca A, Tambaro R, Bianco C, Gullick WJ, Johnson GR, *et al*: Transforming growth factor alpha, amphiregulin and cripto-1 are frequently expressed in advanced human ovarian carcinomas. Int J Oncol 21: 941-948, 2002.
- 33. Freimann S, Ben-Ami I, Hirsh L, Dantes A, Halperin R and Amsterdam A: Drug development for ovarian hyper-stimulation and anti-cancer treatment: Blocking of gonadotropin signaling for epiregulin and amphiregulin biosynthesis. Biochem Pharmacol 68: 989-996, 2004.
- 34. Castillo J, Erroba E, Perugorria MJ, Santamaría M, Lee DC, Prieto J, Avila MA and Berasain C: Amphiregulin contributes to the transformed phenotype of human hepatocellular carcinoma cells. Cancer Res 66: 6129-6138, 2006.
- Ebert M, Yokoyama M, Kobrin MS, Friess H, Lopez ME, Büchler MW, Johnson GR and Korc M: Induction and expression of amphiregulin in human pancreatic cancer. Cancer Res 54: 3959-3962, 1994.
- 36. Yamada M, Ichikawa Y, Yamagishi S, Momiyama N, Ota M, Fujii S, Tanaka K, Togo S, Ohki S and Shimada H: Amphiregulin is a promising prognostic marker for liver metastases of colorectal cancer. Clin Cancer Res 14: 2351-2356, 2008.
- Gidding CE, Kellie SJ, Kamps WA and de Graaf SS: Vincristine revisited. Crit Rev Oncol Hematol 29: 267-287, 1999.
- 38. Corbett R, Pinkerton R, Pritchard J, Meller S, Lewis I, Kingston J and McElwain T: Pilot study of high-dose vincristine, etoposide, carboplatin and melphalan with autologous bone marrow rescue in advanced neuroblastoma. Eur J Cancer 28A: 1324-1328, 1992.
- 39. Gordon SJ, Pearson AD, Reid MM and Craft AW: Toxicity of single-day high-dose vincristine, melphalan, etoposide and carboplatin consolidation with autologous bone marrow rescue in advanced neuroblastoma. Eur J Cancer 28A: 1319-1323, 1992.
- 40. Xi GM, Sun B, Jiang HH, Kong F, Yuan HQ and Lou HX: Bisbibenzyl derivatives sensitize vincristine-resistant KB/VCR cells to chemotherapeutic agents by retarding P-gp activity. Bioorg Med Chem 18: 6725-6733, 2010.
- Yan YX, Li WZ, Huang YQ and Liao WX: The COX-2 inhibitor Celecoxib enhances the sensitivity of KB/VCR oral cancer cell lines to Vincristine by down-regulating P-glycoprotein expression and function. Prostaglandins Other Lipid Mediat 97: 29-35, 2012.
- 42. Yuan Z, Wang H, Hu Z, Huang Y, Yao F, Sun S and Wu B: Quercetin inhibits proliferation and drug resistance in KB/VCR oral cancer cells and enhances its sensitivity to vincristine. Nutr Cancer 67: 126-136, 2015.
- 43. Canto LMD, Cury SS, Barros-Filho MC, Kupper BEC, Begnami MDFS, Scapulatempo-Neto C, Carvalho RF, Marchi FA, Olsen DA, Madsen JS, *et al*: Locally advanced rectal cancer transcriptomic-based secretome analysis reveals novel biomarkers useful to identify patients according to neoadjuvant chemoradiotherapy response. Sci Rep 9: 8702, 2019.
- 44. Lin M, Zhou C, He S, Yu H, Guo T, Ye J, Feng X and Bian X: The research advances of exosomes in esophageal cancer. Biomark Med 13: 685-695, 2019.
- 45. Osborne DG, Domenico J, Luo Y, Reid AL, Amato C, Zhai Z, Gao D, Ziman M, Dinarello CA, Robinson WA and Fujita M: Interleukin-37 is highly expressed in regulatory T cells of melanoma patients and enhanced by melanoma cell secretome. Mol Carcinog 58: 1670-1679, 2019.
- 46. Obenauf AC, Zou Y, Ji AL, Vanharanta S, Shu W, Shi H, Kong X, Bosenberg MC, Wiesner T, Rosen N, *et al*: Therapy-induced tumour secretomes promote resistance and tumour progression. Nature 520: 368-372, 2015.
- 47. Amrutkar M, Aasrum M, Verbeke CS and Gladhaug IP: Secretion of fibronectin by human pancreatic stellate cells promotes chemoresistance to gemcitabine in pancreatic cancer cells. BMC Cancer 19: 596, 2019.
- Kulasingam V and Diamandis EP: Strategies for discovering novel cancer biomarkers through utilization of emerging technologies. Nat Clin Pract Oncol 5: 588-599, 2008.
- 49. May M: From cells, secrets of the secretome leak out. Nat Med 15: 828, 2009.
- 50. Wang X, Masri S, Phung S and Chen S: The role of amphiregulin in exemestane-resistant breast cancer cells: Evidence of an autocrine loop. Cancer Res 68: 2259-2265, 2008.

- Chen JC, Chen YJ, Lin CY, Fong YC, Hsu CJ, Tsai CH, Su JL and Tang CH: Amphiregulin enhances alpha6beta1 integrin expression and cell motility in human chondrosarcoma cells through Ras/Raf/MEK/ERK/AP-1 pathway. Oncotarget 6: 11434-11446, 2015.
- 52. Kakiuchi S, Daigo Y, Ishikawa N, Furukawa C, Tsunoda T, Yano S, Nakagawa K, Tsuruo T, Kohno N, Fukuoka M, *et al*: Prediction of sensitivity of advanced non-small cell lung cancers to gefitinib (Iressa, ZD1839). Hum Mol Genet 13: 3029-3043, 2004.
- 53. Lorente M, Carracedo A, Torres S, Natali F, Egia A, Hernández-Tiedra S, Salazar M, Blázquez C, Guzmán M and Velasco G: Amphiregulin is a factor for resistance of glioma cells to cannabinoid-induced apoptosis. Glia 57: 1374-1385, 2009.
- 54. Ishikawa N, Daigo Y, Takano A, Taniwaki M, Kato T, Hayama S, Murakami H, Takeshima Y, Inai K, Nishimura H, *et al*: Increases of amphiregulin and transforming growth factor-alpha in serum as predictors of poor response to gefitinib among patients with advanced non-small cell lung cancers. Cancer Res 65: 9176-9184, 2005.
- 55. Kao SY and Lim E: An overview of detection and screening of oral cancer in Taiwan. Chin J Dent Res 18: 7-12, 2015.56. Chiang WF, Liu SY, Yen CY, Lin CN, Chen YC, Lin SC and
- 56. Chiang WF, Liu SY, Yen CY, Lin CN, Chen YC, Lin SC and Chang KW: Association of epidermal growth factor receptor (EGFR) gene copy number amplification with neck lymph node metastasis in areca-associated oral carcinomas. Oral Oncol 44: 270-276, 2008.
- 57. Huang SF, Chuang WY, Chen IH, Liao CT, Wang HM and Hsieh LL: EGFR protein overexpression and mutation in areca quid-associated oral cavity squamous cell carcinoma in Taiwan. Head Neck 31: 1068-1077, 2009.
- 58. Huang SF, Chien HT, Cheng SD, Chuang WY, Liao CT and Wang HM: EGFR copy number alterations in primary tumors, metastatic lymph nodes, and recurrent and multiple primary tumors in oral cavity squamous cell carcinoma. BMC Cancer 17: 592, 2017.
- 59. Huang SF, Chien HT, Chuang WY, Lai CH, Cheng SD, Liao CT and Wang HM: Epidermal growth factor receptor intron-1 CA repeat polymorphism on protein expression and clinical outcome in Taiwanese oral squamous cell carcinoma. Sci Rep 7: 4963, 2017.
- 60. Rubin Grandis J, Melhem MF, Gooding WE, Day R, Holst VA, Wagener MM, Drenning SD and Tweardy DJ: Levels of TGF-alpha and EGFR protein in head and neck squamous cell carcinoma and patient survival. J Natl Cancer Inst 90: 824-832, 1998.
- 61. Harding J and Burtness B: Cetuximab: An epidermal growth factor receptor chemeric human-murine monoclonal antibody. Drugs Today (Barc) 41: 107-127, 2005.
- 62. Bonner JA, Harari PM, Giralt J, Azarnia N, Shin DM, Cohen RB, Jones CU, Sur R, Raben D, Jassem J, *et al*: Radiotherapy plus cetuximab for squamous-cell carcinoma of the head and neck. N Engl J Med 354: 567-578, 2006.
- 63. Vermorken JB, Mesia R, Rivera F, Remenar E, Kawecki A, Rottey S, Erfan J, Zabolotnyy D, Kienzer HR, Cupissol D, *et al*: Platinum-based chemotherapy plus cetuximab in head and neck cancer. N Engl J Med 359: 1116-1127, 2008.
- 64. Doble BW and Woodgett JR: GSK-3: Tricks of the trade for a multi-tasking kinase. J Cell Sci 116: 1175-1186, 2003.
- Luo J: Glycogen synthase kinase 3beta (GSK3beta) in tumorigenesis and cancer chemotherapy. Cancer Lett 273: 194-200, 2009.
 Domoto T, Pyko IV, Furuta T, Miyashita K, Uehara M,
- 66. Domoto T, Pyko IV, Furuta T, Miyashita K, Uehara M, Shimasaki T, Nakada M and Minamoto T: Glycogen synthase kinase-3β is a pivotal mediator of cancer invasion and resistance to therapy. Cancer Sci 107: 1363-1372, 2016.
- 67. Maqbool M and Hoda N: GSK3 inhibitors in the therapeutic development of diabetes, cancer and neurodegeneration: Past, present and future. Curr Pharm Des 23: 4332-4350, 2017.
- 68. Walz A, Ugolkov A, Chandra S, Kozikowski A, Carneiro BA, O'Halloran TV, Giles FJ, Billadeau DD and Mazar AP: Molecular pathways: Revisiting glycogen synthase kinase-3β as a target for the treatment of cancer. Clin Cancer Res 23: 1891-1897, 2017.
- 69. Pramanik KK, Nagini S, Singh AK, Mishra P, Kashyap T, Nath N, Alam M, Rana A and Mishra R: Glycogen synthase kinase-3β mediated regulation of matrix metalloproteinase-9 and its involvement in oral squamous cell carcinoma progression and invasion. Cell Oncol (Dordr) 41: 47-60, 2018.
- 70. Matsuo FS, Andrade MF, Loyola AM, da Silva SJ, Silva MJB, Cardoso SV and de Faria PR: Pathologic significance of AKT, mTOR, and GSK3β proteins in oral squamous cell carcinoma-affected patients. Virchows Arch 472: 983-997, 2018.

- 71. Huang GL, Shen DY, Cai CF, Zhang QY, Ren HY and Chen QX: β-escin reverses multidrug resistance through inhibition of the GSK3β/β-catenin pathway in cholangiocarcinoma. World J Gastroenterol 21: 1148-1157, 2015.
- 72. Ugolkov A, Gaisina I, Zhang JS, Billadeau DD, White K, Kozikowski A, Jain S, Cristofanilli M, Giles F, O'Halloran T, *et al*: GSK-3 inhibition overcomes chemoresistance in human breast cancer. Cancer Lett 380: 384-392, 2016.
- 73. Ugolkov A, Qiang W, Bondarenko G, Procissi D, Gaisina I, James CD, Chandler J, Kozikowski A, Gunosewoyo H, O'Halloran T, *et al*: Combination treatment with the GSK-3 inhibitor 9-ING-41 and CCNU cures orthotopic chemoresistant glioblastoma in patient-derived xenograft models. Transl Oncol 10: 669-678, 2017.