

Prolyl hydroxylase 2 inhibits glycolytic activity in colorectal cancer via the NF- κ B signaling pathway

LISHA XIANG^{1*}, HAO WEI^{1*}, WENTAO YE², SHUANG WU³ and GANFENG XIE³

¹Division of Thoracic Tumor Multimodality Treatment and Department of Medical Oncology, Cancer Center, State Key Laboratory of Biotherapy and ²West China Medical School, West China Medical Center, West China Hospital, Sichuan University, Chengdu, Sichuan 610041; ³Department of Oncology and Southwest Cancer Center, Southwest Hospital, Third Military Medical University (Army Medical University), Chongqing 400038, P.R. China

Received September 17, 2023; Accepted October 30, 2023

DOI: 10.3892/ijo.2023.5590

Abstract. A variety of malignancies preferentially meet energy demands through the glycolytic pathway. Hypoxia-induced cancer cell adaptations are essential for tumor development. However, in cancerous glycolysis, the functional importance and underlying molecular mechanism of prolyl hydroxylase domain protein 2 (PHD2) have not been fully elucidated. Gain- and loss-of-function assays were conducted to evaluate PHD2 functions in colon cancer cells. Glucose uptake, lactate production and intracellular adenosine-5'-triphosphate/adenosine diphosphate ratio were measured to determine glycolytic activities. Protein and gene expression levels were measured by western blot analysis and reverse transcription-quantitative PCR, respectively. The human colon cancer xenograft model was used to confirm the role of PHD2 in tumor progression *in vivo*. Functionally, the data demonstrated that PHD2 knockdown leads to increased glycolysis, while PHD2 overexpression resulted in suppressed glycolysis in colorectal cancer cells. In addition, the glycolytic activity was enhanced without PHD2 and normalized after PHD2 reconstitution. PHD2 was shown to inhibit colorectal tumor growth, suppress cancer cell proliferation and improve tumor-bearing mice survival *in vivo*. Mechanically, it was found that PHD2 inhibits the expression of critical glycolytic enzymes (glucose transporter 1, hexokinase 2 and phosphoinositide-dependent protein kinase 1). In addition, PHD2 inhibited Ikk β -mediated NF- κ B activation in a hypoxia-inducible factor-1 α -independent manner. In conclusion, the data demonstrated that PHD2/Ikk β /NF- κ B signaling

has critical roles in regulating glycolysis and suggests that PHD2 potentially suppresses colorectal cancer.

Introduction

Unlike non-malignant cells, cancer cells utilize a different program for glucose metabolism. Due to the altered tumor microenvironment, cancer cells have to elevate glycolytic activity to generate adequate adenosine-5'-triphosphate (ATP) within a hypoxic microenvironment to maintain rapid cell proliferation. This phenomenon of continuously producing high lactate in cancer cells in the presence of oxygen is known as the Warburg effect (1-3).

During tumor expansion, uncontrollable cell proliferation and abnormal tumor angiogenesis impede the nutrient supply for cancer cells (4). Consequently, hypoxic regions, where oxygen concentration is low, are frequently found in solid tumors. To adapt to this hypoxic environment, tumor cells usually adopt glycolysis for their energy supply to facilitate angiogenesis, which, in turn, contributes to the continued growth and survival of a tumor (5). Therefore, oxygen homeostasis and regulating downstream cellular metabolic events are critical during tumor progression. Members from the family of prolyl hydroxylase (PHD), a group of well-established oxygen sensors, are potentially the major participants that sense oxygen and regulate glucose metabolism in tumors (6-8).

PHDs mediate hypoxia-inducible factor-1 α (HIF-1 α) hydroxylation at 2 prolyl residues in the oxygen-dependent degradation domain, enabling it to be captured, ubiquitinated and ultimately degraded in the proteasome (9). There are three known PHD isoforms in humans; PHD1, PHD2 and PHD3. PHD2 is one of the key enzymes mediating HIF-1 α degradation under normoxia (10-12). In addition, PHD2 is downregulated or even barely detected in numerous cancer cells and tumor tissues (13). The decreased expression of PHD2 is significantly correlated with some genetic disorders, which ultimately develop malignancies (12). Therefore, such findings render PHD2 a potential tumor suppressor (14). In addition, PHD2 has been associated with tumor vasculature, which occurs after tumor glycolysis (15-17). Nevertheless, the role of PHD2 in tumor glycolysis is still unknown. The present study focused on investigating glycolysis, including glucose uptake,

Correspondence to: Dr Ganfeng Xie, Department of Oncology and Southwest Cancer Center, Southwest Hospital, Third Military Medical University (Army Medical University), 30 Gaotanyan Street, Chongqing 400038, P.R. China
E-mail: xiegf@aliyun.com

*Contributed equally

Key words: prolyl hydroxylase 2, glycolysis, nuclear factor- κ B, colorectal cancer, hypoxia-inducible factor-1 α

lactate production and energetic molecular quantification and confirming the regulatory effect of PHD2 in glycolysis in colon cancer cells by gain- and loss-of-functional assays.

Materials and methods

Cell lines. HEK293T and human colorectal carcinoma cell lines (HT-29, HCT116, Ls174t, LoVo, SW480, SW620, RKO and Caco-2) were obtained from the American Type Culture Collection. Cells were maintained in Dulbecco's Modified Eagle's Medium (DMEM) containing 10% fetal bovine serum (FBS), supplemented with 100 U/ml penicillin and 0.1 mg/ml streptomycin at 37°C in an atmosphere containing 5% CO₂. All reagents for cell culturing were purchased from Invitrogen (Thermo Fisher Scientific, Inc.).

PHD2 knockdown and overexpression. For PHD2 silencing, the pGCL-GFP expressing shRNA against PHD2 [short hairpin (sh)PHD2; target sequence: GGATGGAATCCATGA GCTA] or non-targeting shRNA (Mock; TTCTCCGAACGT GTCACGT) were designed and synthesized by Shanghai GeneChem Co., Ltd. For PHD2 overexpression, the open reading frame of PHD2 was constructed into the pGC-FU lentiviral vector (pGC-FU-PHD2) (Shanghai GeneChem Co., Ltd.), while the empty vector was used as vehicle control (vehicle).

HEK293T cells were co-transfected with helper plasmids (pHelper 1.0 and pHelper 2.0; Shanghai GeneChem Co., Ltd.) and lentiviral plasmids (pGCL-GFP-shPHD2 or pGC-FU-PHD2) using Lipofectamine[®]2000 (Invitrogen; Thermo Fisher Scientific, Inc.) at 37°C for 24 h for lentivirus production. Lentivirus was harvested at 48 and 72 h following transfection. Cell debris was removed by centrifugation (300 x g; 10 min; room temperature) and the virus was then filtered through a 0.45 μm cellulose acetate filter and collected by ultracentrifugation (100,000 x g; 4 h; 4°C).

For lentivirus infection (37°C for 12 h), Ls174t and SW480 cells were transduced with PHD2 silencing (PHD2-KD), overexpressing (PHD2-OE) lentivirus, or control lentivirus (represented as MOCK for non-targeting shRNA and vehicle for empty vector), separately, at the multiplicity of infection of 10 (Ls174t) or 20 (SW480). At four days following infection, GFP-positive cells were isolated by a FACSCalibur flow cytometer (BD Biosciences). PHD2 overexpression and silencing were confirmed by western blotting.

HIF-1α lentiviruses transduction. For HIF-1α silencing, the pLKO.1-puro lentiviral vectors encoding shRNA targeting HIF-1α (HIF-1α-KD) or non-targeting control shRNA (MOCK; MilliporeSigma) were obtained for lentiviruses packaging. The cells were transduced with viral supernatant containing 8 μg/ml of Polybrene (MilliporeSigma) and then selected with medium containing 0.6 μg/ml of puromycin (MilliporeSigma) after 12 h (18).

Reverse transcription-quantitative (RT-q) PCR. Human cancerous and adjacent normal colorectum were collected from specimens directly after surgery. Total RNA was extracted using the TRIzol[®] (Thermo Fisher Scientific, Inc.) method and the complementary DNA was synthesized using

a commercially available kit (Invitrogen; Thermo Fisher Scientific, Inc.) according to the manufacturer's protocols. RT-qPCR was carried out using an iCycler system (Bio-Rad Laboratories, Inc.) with SYBR[®] Green PCR master mix (Applied Biosystems; Thermo Fisher Scientific, Inc.). PCR amplification was performed with the following cycling conditions: i) Initial activation at 95°C for 10 min, and subsequently 40 cycles of denaturation at 95°C for 15 sec; ii) Annealing at 55°C for 30 sec and extension at 72°C for 30 sec (40 cycles). Primers used were as follows: Phd2: 5'-CGATCCGGTGAC TTTTCCAC-3' and 5'-GTAGGAGGTCTCCGTCCT-3'; hexokinase-2 (Hk2): 5'-AAGATGCTGCCCCACCTACG-3' and 5'-TCGCTTCCCATTCCTCACA-3'; PHD kinase 1 (Pdk1): 5'-AAGATGAGTGACCGAGGAGGT-3' and 5'-CCA TAACCAAACCAGCCAGAG-3'; glucose transporter 1 (Glut1): 5'-GTGCTCCTGGTTCTGTTCTTCA-3' and 5'-GCC AGAAGCAATCTCATCGAA-3'; β-actin: 5'-ACCCCGTGC TGCTGACCGAG-3' and 5'-TCCCGGCCAGCCAGGTCC A-3'. The expression levels of detected genes were normalized to the level of β-actin mRNA. The relative expression of the target genes was calculated using the following formula: $\text{gene expression} = 2^{-(\Delta Cq_{\text{normal}} - \Delta Cq_{\text{tumor}})}$ (19). The experiment was independently repeated three times.

ELISA assay. Cells were washed with ice-cold PBS, detached and pelleted by centrifugation (1,000 x g; 5 min; 4°C). The cells were resuspended in 300 μl of hypotonic buffer (10 mM Hepes; pH 7.6; 60 mM KCl; 1 mM EDTA; 1 mM dithiothreitol) containing phenylmethylsulfonyl fluoride (PMSF; 1 mM) and protease inhibitor cocktail (10 μl/ml) and incubated on ice for 15 min. Then, 20 μl of 10% NP-40 was added and the tube was vortexed for 10 sec. The lysates were cleared by centrifugation (13,000 x g; 1 min; 4°C) and supernatants (cytosolic fractions) were collected and stored at -80°C.

Nuclear contents were obtained from the pellets. First, the pellets were resuspended in extraction buffer (20 mM Tris-HCl; pH 8; 420 mM NaCl; 1.5 mM MgCl₂; 0.2 mM EDTA) supplemented with PMSF (0.5 mM), protease inhibitors cocktail (10 μl/ml) and glycerol (25% v/v). After 30 min at 4°C, the debris was removed by centrifugation (13,000 x g; 15 min; 4°C). The supernatant containing nuclear proteins was aliquoted and stored at -80°C for further analysis. Protein concentrations were determined using a Bradford protein assay (Bio-Rad Laboratories, Inc.).

For the p50 detection, the levels of p50 were measured using a TransAM NF-κB p50 ELISA kit (cat. no. 31101; Active Motif, Inc.) according to the manufacturer's instructions with a microplate reader (Bio-Rad Laboratories, Inc.). The experiment was conducted in triplicates and was independently repeated three times.

Western blotting and co-immunoprecipitation. Western blotting was conducted using lysates extracted from human tissues or cells as previously described (20). In general, tissues or cells were solubilized in RIPA buffer (BioTeke Corporation) containing a proteinase inhibitor cocktail (1 mM EDTA; 1 mM PMSF; 1 mM iodoacetic acid). After the protein concentration was determined with the Bradford method, identical amounts of proteins (50 μg) were separated on a 10% SDS-PAGE gel and transferred to polyvinylidene

fluoride membranes. The membranes were then blocked with 5% skimmed milk for 2 h at room temperature followed by overnight incubation with antibodies against PHD2 (1:2,000; cat. no. NBP1-30328; Novus Biologicals, LLC), glucose transporter 1 (GLUT1; 1:500; cat. no. sc-377228; Santa Cruz Biotechnology, Inc.), PDK1 (1:500; cat. no. sc-515944; Santa Cruz Biotechnology, Inc.), hexokinase (HK) 2 (1:1,000; cat. no. sc-374091; Santa Cruz Biotechnology, Inc.), p65 (1:1,000; cat. no. sc-8008; Santa Cruz Biotechnology, Inc.) and GAPDH (1:1,000; cat. no. sc-47724; Santa Cruz Biotechnology, Inc.) at 4°C. The following day, membranes were rinsed and incubated with HRP-conjugated secondary antibodies (1:2,000, cat. no. BA1082; Wuhan Boster Biological Technology, Ltd.) at room temperature for 1 h. SuperSignal West Femto Maximum Sensitivity Substrate (Pierce; Thermo Fisher Scientific, Inc.) was used for development and images were captured using a Gel Doc XR system (Bio-Rad Laboratories, Inc.). β -actin was used as the loading control. ImageJ software (NIH; version 1.8.0) was used for densitometry.

For immunoprecipitation assay, fresh cell lysates using RIPA buffer (cat. no. PP1901; BioTek Instruments, Inc.) containing a proteinase inhibitor cocktail (1 mM EDTA; 1 mM PMSF; 1 mM iodoacetic acid) (500 μ g) were incubated with primary antibody (1 μ g) at 4°C for 3 h. Then, 20 μ l of protein A/G PLUS agarose beads (cat. no. sc-2003; Santa Cruz Biotechnology, Inc.) was supplemented for overnight incubation. The following day, the beads were pelleted and collected by centrifugation (10,000 x g; 15 sec; 4°C), followed by washing with RIPA buffer as aforementioned. Then, immobilized proteins were eluted in the boiled SDS buffer for electrophoresis and immunoblotting. Primary antibodies were used to detect the HA-tag (1:1,000; cat. no. sc-805; Santa Cruz Biotechnology, Inc.), Flag-tag (1:4,000; cat. no. F3165; MilliporeSigma), IKK α (1:1,000; cat. no. 2682; Cell Signaling Technology, Inc.), anti-IKK β (1:1,000; cat. no. 2684; Cell Signaling Technology, Inc.), HRP-conjugated anti-rabbit (1:5,000; cat. no. 7074; Cell Signaling Technology, Inc.), -mouse (1:5,000; cat. no. 7076; Cell Signaling Technology, Inc.), or -goat (1:5,000; cat. no. 7075; Cell Signaling Technology, Inc.) secondary antibodies were used for incubation for 1 h at room temperature.

Electrophoretic mobility shift assay. An oligonucleotide containing the I κ B-binding site was radiolabeled with [γ -³²P] dATP using T4 polynucleotide kinase at 37°C for 1 h. The labeled oligonucleotide was purified with the QIAquick Nucleotide Removal Kit (cat. no. 28304; Qiagen GmbH) according to the manufacturer's instruction. The binding between probe (1 ng) and nuclear extracts (5 ng) was allowed to form in 20 μ l reaction buffer (1 μ g poly (dI-dC); 0.1 μ g BSA; 1 mM DTT; 60 mM KCl; 10% glycerol and 20 mM HEPES, pH 8.4) at 30°C for 30 min. Then, 1 μ g of p65 antibody (cat. no. 05-361; Upstate Biotechnology, Inc.) was added to immobilize the oligonucleotide/protein complexes, which were then separated by native electrophoresis in a 6% polyacrylamide gel that was exposed to an X-ray film (Amersham; Cytiva) for 96 h at -80°C.

2-deoxy-D-glucose uptake assay. Cells (1x10⁷ cells/ml) were washed twice with PBS and resuspended and incubated in

100 μ l PBS supplemented with 5 mM of 2-deoxy-D-glucose [containing 2 μ Ci/ml 2-deoxy-(1-³H) glucose] at 37°C for 30 min. The glucose uptake was terminated by adding 100 μ l of ice-cold PBS containing 100 μ M phloretin. The cells were subjected to brief centrifugation (20,000 x g; 30 sec; room temperature). Then, the cell pellets were washed once and transferred into a scintillation vial containing 10 ml of Optifluor (Perkin-Elmer Inc.). The incorporated radioactivity was measured by a liquid scintillation counter (Beckman Coulter, Inc.). The uptaken 2-deoxy-[1-³H] glucose concentrations were presented as nmol/10⁴ cells.

L-lactate assay. The day before the assay, cells were seeded (1.2x10⁵ per well) in 12-well plates in complete medium in quadruplicate. The medium was replaced with starve medium (DMEM containing 2% FBS) on the next day for 48 h. Then, the medium was collected and the lactate concentration was determined using a fluorometric lactate assay kit (cat. no. K607; BioVision, Inc.). Finally, the results were normalized to protein content.

Measurement of intracellular adenine nucleotide levels. For ATP and adenosine diphosphate (ADP) measurement, cell lysate samples were deproteinized with 5% trichloroacetic acid at various times before and after the glucose addition. Following centrifugation (14,000 x g; 10 sec; 4°C), the supernatants were neutralized by ether extraction, lyophilized and stored at -80°C until further analysis. Then, the nucleotide extracts were analyzed by high-performance liquid chromatography using an anion exchange hypersil C18 column (4.6x250 mm) at a flow rate of 0.9 ml/min over a 10-min buffer gradient (0.1 mol/l KH₂PO₄, pH 6.0). The eluting flow was continuously monitored by UV absorption at 254 nm and the peaks of nucleoside triphosphates were quantified by electronic integration with external standards used as reference. Intracellular ATP contents were quantified and normalized to cell number. The results are presented as the ATP/ADP ratio, which is unaffected by sample volume loss during ether extraction or cell numbers.

Human colorectal cancer xenografts in immunodeficient mice. A total of 48 combined immunodeficient (SCID) mice (male, 6-8 weeks old, 20-25 g) were purchased from the Army Medical University Institute of Experimental Animal. Mice were fed at an ambient of 45-55% relative humidity with the constant temperature of 21±2°C under a 12-h light/dark cycle. Lentivirus for PHD2-silencing or -overexpression and their corresponding controls were used to transduce SW480 (for PHD2 overexpression) or Ls174t cells (for PHD2 depletion). SCID mice were randomly divided into 3 groups (n=16) for PHD2 overexpression, silencing and control. Stably transduced SW480 and Ls174t cells or the corresponding parental cells (~2x10⁶) in 200 μ l of PBS were admixed with 200 μ l of Matrigel (BD Biosciences). The cells were delivered to the mice through subcutaneous injection. At three days after injection, the mice were subjected to measurements of tumor size, body weight and other side effects twice a week. Tumor volume (mm³) was calculated by the formula: 0.52 x length x width x thickness. After 40 days, mice were sacrificed (n=6) and tumors were excised for further analysis. The rest of the

mice were subjected to monitoring until they reached the humane endpoint for survival analysis. The humane endpoint was defined as weight loss of more than 20% and mice were sacrificed by CO₂ asphyxiation (at the rate of 30-70% of the chamber volume per minute) followed by cervical dislocation. Survival rate was analyzed using the Kaplan-Meier method. Animal studies were performed under a protocol approved by the Army Medical University Institutional Animal Care and Use Committee (approval no. AMUWEC20232707).

Ki-67 proliferation index. Tumor tissues harvested from the xenografts were fixed in 10% formalin for 24 h at room temperature, gradual dehydration in increasing alcohol concentrations and then embedded within paraffin for 8 h at room temperature as previously described (21). Tissue sections were then incubated with antibodies against Ki67 (1:100; cat. no. NB500-170; Novus Biologicals, LLC). Nuclei were counterstained with Mayer's hematoxylin (MilliporeSigma) for 5 min at room temperature. Ki-67-positive cell from 6-8 random views of each section was counted and images captured under a light inverted microscope (XSZ-D2; Olympus Corporation) by a researcher who was blinded to this study. ImageJ software (NIH; v1.8.0) was used to process the results and the data were presented as the average percentage of Ki-67-positive cells per view (22).

Statistical analysis. The results were analyzed using SPSS 13.0 software (v13.0; SPSS, Inc.). One-way analysis of variance with Tukey's post hoc test was performed to compare results from different groups. The quantitative data were expressed as the mean \pm SEM. All experiments were repeated three to five times independently. The survival analysis was performed using the log-rank test. Pearson's correlation coefficient was used to evaluate the relationship between the expression of PHD2 and glycolysis-related genes in cancerous and adjacent normal colon tissues. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

PHD2 is decreased in colorectal tumors. A total of 45 paired human colorectal tumors and adjacent normal tissues were interrogated for PHD2 expression. PHD2 reduction was observed in 29 of 45 (64.4%) tumors compared with the paired normal colorectal tissues (Fig. 1A). In addition, univariate analysis revealed that PHD2 expression is significantly different in tumor and paired normal tissues at the transcriptional level ($P < 0.001$). Notably, our previous study demonstrated that the decreased PHD2 expression in tumor tissue favors higher tumor grades and poorer overall survival of colorectal cancer patients (23). In agreement with the findings at the mRNA level, decreased PHD2 expression at the protein level was also observed in all tested paired samples (Fig. 1B). In addition, results from our cell-based immunoblotting assay also revealed that PHD2 was barely detected in most tested colon cancer cells, except for Ls174t cells (Fig. 1C).

Low expression levels of PHD2 correlate with high expression levels of glycolysis-related genes. Genes closely related to glycolysis, such as HK2, PDK1 and GLUT1, were




Figure 1. Evaluation of PHD2 expression in patient samples and colorectal cancer cell lines. (A) Results of RT-qPCR detecting PHD2 mRNA level. $\Delta Cq(T)$: GAPDH Cq value subtracted from PHD2 Cq value in normal tissue [$\Delta Cq(N)$] or tumor [$\Delta Cq(T)$]. Bar value: $(\Delta Cq(N) - \Delta Cq(T))$ value indicated the PHD2 mRNA difference in normal tissue and paired tumor and the differences are expressed in log₂ scale. (B) Determination of PHD2 protein level in human samples. (C) Determination of PHD2 expression in human colorectal cancer cell lines. T, tumor tissue; N, paired adjacent normal tissue; PHD2, prolyl hydroxylase domain protein 2; RT-qPCR, reverse transcription-quantitative PCR.

also evaluated in tumor and paired normal adjacent tissues. Consistent with previous reports (24-26), the mRNA levels of those genes, including HK2 ($P < 0.01$), PDK1 ($P < 0.05$) and GLUT1 ($P < 0.001$), were significantly higher in tumor tissues than in paired normal tissues (Fig. 2A). To elucidate the potential relationships between the expression levels of PHD2 and glycolysis-related genes, we plotted PHD2 mRNA levels against HK2, PDK1 and GLUT1 mRNA levels in colorectal tissues from both tumor and adjacent normal colorectum. As a result, the expression of HK2 ($P < 0.00001$), PDK1 ($P = 0.00217$) and GLUT1 ($P < 0.0019$) were significantly inversely-correlated with that of PHD2 in colorectal tumors (Fig. 2B). Together, these findings suggested that PHD2 is downregulated in colorectal tumor tissues and may be involved in tumor-associated glycolysis.




Figure 2. PHD2 is negatively correlated with glycolysis-related genes in human colorectal cancers. (A) Results of RT-qPCR detecting the expression of *HK2*, *PDK1*, or *GLUT1*. $\Delta Cq(N)$ and $\Delta Cq(T)$ indicate the Cq value of GAPDH was subtracted from Cq value of *HK2*, *PDK1*, or *GLUT1* in normal tissue or tumor tissue, respectively. Bar value: $(\Delta Cq(N) - \Delta Cq(T))$ depicts the fold-changes of *HK2*, *PDK1*, or *GLUT1* mRNA in tumor tissues vs. normal tissues and the difference is expressed in log2 scale. (B) Significant negative correlation between the relative mRNA expression level of *PHD2* (x-axis) and *HK2*, *PDK1*, or *GLUT1* (y-axis) in the colorectal cancer samples. PHD2, prolyl hydroxylase domain protein 2; RT-qPCR, reverse transcription-quantitative PCR.

Inhibition of PHD2 stimulates glycolytic activity in colorectal cancer cells. To confirm the role of PHD2 in tumor-associated glycolysis, *in vitro* glycolytic activity was examined in different human colorectal cancer cells with altered PHD2 expression levels. Among the tested colon cancer cell lines, SW480 cells presented the lowest, while Ls174t cells expressed the highest PHD2 protein expression (Fig. 1C). Therefore, PHD2 was overexpressed in SW480 cells (PHD2-OE) and PHD2 silenced in Ls174t cells (PHD2-KD) via lentiviral-based modification, respectively. Transduction efficiency and PHD2 expression were evaluated by immunoblotting analysis (Fig. 3A). As the data indicated a negative correlations between PHD2 and GLUT1, HK2 and PDK1, the protein level of these proteins was assessed in PHD2-OE cells and control cells (Fig. 3B). Compared with control counterparts, PHD2 overexpression significantly suppressed the expression of GLUT1, HK2 and PDK1 proteins.

Notably, PHD2 can also be inhibited *in vitro* by the competitive inhibitor dimethyloxalylglycine (DMOG), a cell-permeable 2-oxoglutarate analog (27). It was observed that 24 h exposure to DMOG led to a dose-dependent

PHD2 reduction in colon cancer cells (0, 0.125, 0.25, 0.50, or 1.00 mM DMOG; Fig. 3C). Then, the role of PHD2 in glycolytic activity was further investigated by evaluating glucose uptake, lactate production and intracellular ATP/ADP ratio. It was found the glycolytic activity increased greatly in PHD2-KD cells and DMOG (0.5 mM) treated cells compared with the control group (Fig. 3D-F). By contrast, opposite effects could be observed in PHD2-OE cells (Fig. 3G-I). In addition, not only was a dose-dependent inhibitory effect of DMOG on PHD2 expression discovered but it was also noted that DMOG promoted glucose uptake, lactate production and intracellular ATP/ADP ratio in colon cancer cells in the same manner (Fig. 3J and K). Taken together, the results suggested that PHD2 may be involved in regulating glycolysis in colon cancer cells.

Regulation of glycolysis by PHD2 is independent of HIF-1 α . As HIF-1 α is the functionally characterized target of PHD2 (28), it was next determined whether the effects of PHD2 on glycolysis were dependent on HIF-1 α . For this purpose, the present study generated HIF-1 α deficient and HIF-1 α /PHD2 double




Figure 3. Inhibition of PHD2 promotes glycolysis in colorectal cancer cells. (A) PHD2 overexpression and silencing were confirmed by western blot analysis. (B) Expression of GLUT1, HK2 and PDK1 in SW480 cells with altered PHD2 expression. (C) Expression of PHD2 in Ls174t treated with DMOG at indicated concentrations for 24 h. (D and G) Quantitative analysis of the ATP/ADP ratio, (E and H) glucose uptake and (F and I) extracellular lactate in control, (0.5 mM) DMOG treated and PHD2-KD colon cancer cells. (J) Glucose uptake, (K) lactate production and (L) the ratio of ATP/ADP, were quantitatively analyzed in Ls174t cells preincubated in the medium containing 0-1 mM DMOG for 24 h (n=3; *P<0.05 and **P<0.01 vs. Mock). PHD2, prolyl hydroxylase domain protein 2; GLUT1, glucose transporter 1; HK2, hexokinase-2; PDK1, PHD kinase 1; DMOG, dimethylxalylglycine; ATP, adenosine-5'-triphosphate; ADP, adenosine diphosphate; NT, non-transduced; Mock, non-targeting shRNA control (for PHD2 knockdown) or vehicle control (for PHD2 overexpression); PHD2-OE, PHD2 overexpression; PHD2-KD, PHD2-knockdown.

deficient colon cancer cells and the expression of HIF-1 α and PHD2 was evaluated. HIF-1 α expression was significantly suppressed by lentivirus-mediated silencing and PHD2 depletion elevated the expression of HIF-1 α (Fig. 4A). In addition, it was noticed that cells with HIF-1 α silenced did not promote glycolysis activity as reflected by no significant change in glucose uptake, lactate production and intracellular ATP/ADP ratio compared with parental cells (Fig. 4B-D). It also found that further knockdown of HIF-1 α in colon cancer cells did not affect the PHD2 depletion-mediated promotive effects on

colon cancer glycolysis activity (Fig. 4B-D). These findings prompted the hypothesis that PHD2-promoted glycolysis is independent of HIF-1 α in colon cancer cells.

PHD2 inhibits glycolytic activity through IKK β /NF- κ B. Accumulating evidence has revealed that PHD2 has novel targets other than HIF-1 α (29). NF- κ B signaling components are notable candidate targets (30,31). Therefore, the present study determined how PHD2 affects NF- κ B signaling activity and found that depletion of PHD2 led to upregulated nuclear p65




Figure 4. Regulation of glycolysis by PHD2 is HIF-1 α -independent. (A) Representative results of western blot analysis detecting the expression of PHD2 and HIF-1 α in Ls174t cells under indicated conditions. The ratio of (B) glucose uptake, (C) lactate production and (D) ATP/ADP ratio were quantitatively analyzed in control, HIF-1 α silenced, PHD2 silenced and HIF-1 α and PHD2 dual-silenced Ls174t cells (n=3; *P<0.05 and **P<0.01 vs. Mock or vs. HIF-1 α -KD). PHD2, prolyl hydroxylase domain protein 2; HIF-1 α hypoxia-inducible factor-1 α ; Mock, non-targeting shRNA control; PHD2-KD, PHD2 knockdown; HIF-1 α -KD, HIF-1 α knockdown.

subunit (Fig. 5A). Also, DMOG-mediated PHD2 suppression increased nuclear translocation of the p50 subunit (Fig. 5B). Notably, employing the IKK β inhibitor (PDTC) or preventing the proteasomal degradation of I κ B significantly attenuated the promotive effects of PHD2 knockdown (Fig. 5C-E) or PHD2 inhibition (Fig. 5F-H) on glycolysis in colon cancer cells. These findings indicated that PHD2 inhibits glycolytic activity through the IKK β /NF- κ B pathway.

PHD2 regulates the DNA binding properties of NF- κ B. Prior studies have shown that NF- κ B relies on I κ B degradation to enter the nucleus and initiate transcription (32,33). In addition, NF- κ B-dependent I κ B synthesis is critical for NF- κ B signaling termination (34). Furthermore, IKK β has been reported to be the target of PHD2 (30,31). Thus, PHD2 may modulate NF- κ B signaling in an IKK β -dependent manner. Through immunoprecipitation assay, it was noted that, among the IKK isoforms, PHD2 preferably interacts with IKK β (Fig. 6A). The present study further confirmed the interaction between PHD2 and IKK β in cells with altered PHD2 expression. As shown, silencing or inhibiting PHD2 attenuated the PHD2-IKK β interaction, while enhanced binding between PHD2 and IKK β was observed in PHD2-OE cells (Fig. 6B and C). Meanwhile, the expression of p65 in the cytosol and nucleus was influenced by PHD2 overexpression (Fig. 6D), suggesting that PHD2 exerts its effect dose-dependently on IKK β , presumably after p65 nuclear translocation.

PHD2 inhibits colorectal tumor growth in vivo and prolongs animal survival. To further characterize the functional significance of PHD2 on tumor growth *in vivo*, a subcutaneous xenograft model of colorectal cancer was established in SCID mice. Parental

colon cancer cells (SW480 cells or Ls174t) or parental cells with or without PHD2 expression modulation (PHD2-OE, PHD2-KD, or Mock control) were subcutaneously inoculated into SCID mice. The data showed that colon cancer cells expressing control vectors grow into sizable tumors and were comparable to non-transduced cells (Fig. 7A and B). In addition, mice transplanted with Ls174t cells, which express a higher level of PHD2, developed smaller tumors than those implanted with SW480 cells (Fig. 7A and B). Notably, the data also demonstrated that tumor growth was significantly inhibited by PHD2 overexpression (Fig. 7A) but was dramatically promoted by PHD2 silencing (Fig. 7B). The result suggests a suppressive effect of exogenous PHD2 on human colorectal cancer progression *in vivo*. Additionally, the findings further revealed that Ki-67-positive cell proportion, representing the tumor proliferation index, was significantly reduced in mice transplanted with PHD2 overexpressing cancer cells and significantly higher in mice inoculated with PHD2 silencing cancer cells compared with the corresponding control mice (Fig. 7C-F). Kaplan-Meier curves were used to determine survival in each mouse group. The log-rank test demonstrated that PHD2 overexpression significantly prolonged the survival of tumor-bearing mice compared with the wild-type or mock control groups (P<0.01) (Fig. 7G). Again, as shown in Fig. 7H, PHD2 deficiency reduced the survival of tumor-bearing mice (P<0.01). Taken together, the data indicated that PHD2 markedly delayed tumor growth and increases survival of mice in a model of colorectal cancer.

Discussion

A promising therapeutic target selective for cancer cells may focus on their dependence on aerobic glycolysis for energy




Figure 5. PHD2 inhibits glycolytic activity through IKK β /NF- κ B. (A) Nuclear p65 expression were determined by western blotting in Ls174t cells. (B) Expression of nuclear p50 was measured by ELISA assay. Quantified (C) glucose uptake (D) lactate production and (E) ATP/ADP ratio in PHD2-deficient Ls174t cells under indicated treatments (n=3; *P<0.05 and **P<0.01 vs. un-treated cells). Quantified (F) glucose uptake (G) lactate production and (H) ATP/ADP ratio in PHD2 inhibited Ls174t cells under indicated conditions (n=3; *P<0.05 and **P<0.01 vs. untreated cells). PHD2, prolyl hydroxylase domain protein 2; ATP, adenosine-5'-triphosphate; ADP, adenosine diphosphate; NT, non-transduced; Mock, non-targeting shRNA control; PHD2-KD, PHD2 knockdown.

metabolism (35). An increasing body of evidence has revealed that elevated glycolysis can enhance the O-glycosylation of IKK β that subsequently triggers the activation of NF- κ B signaling (36-38). Other studies have implicated IKK β /NF- κ B signaling in cancer cell metabolic stress adaption and energy metabolism, including colorectal cancer (39-42). In addition, PHD2-mediated activation of NF- κ B signaling has been reported via the I κ B-dependent pathway (31). However, whether PHD2 regulates glycolysis by modulating NF- κ B signaling in colorectal cancer has not been well studied. The present study explored the crucial role of PHD2 in colon cancer cell glycolysis and investigated the potential regulatory effects of PHD2 in NF- κ B signaling during glycolysis both *in vitro* and *in vivo*.

By interrogating PHD2 expression in human tissues and cell lines, it was found that PHD2 expression was significantly downregulated in cancerous tissues and cells, in agreement with previous findings (29,43). In association with PHD2 downregulation, the present study also observed elevated

critical glycolytic enzymes, HK2, PDK1 and GLUT1, in human colorectal cancer tissues, negatively correlated with PHD2 expression level. The present *in vitro* study consistently reported that PHD2 overexpression decreased, while PHD2 depletion increased, the expression of GLUT1, HK2 and PDK1 in colon cancer cells. Similar findings have also been observed by other researchers (44). These findings highlight the importance of PHD2 in cancer glucose metabolism and shed light on the tumor-suppressive role of PHD2.

PHD2 has been reported to regulate HIF-1 α proteasomal degradation via hydroxylation in normoxia (9). Meanwhile, in the presence of DMOG, or during hypoxia, PHD2 is suppressed. In turn, HIF-1 α enters nuclear and mediates glycolysis (45). Similarly, upregulated HIF-1 α in PHD2 depleted cells was observed. However, in contrast to the expectation that silencing HIF-1 α in PHD2 deficient cells would affect the glucose metabolism, it was found that the PHD2-deficiency-enhanced glycolysis did not depend on the absence of HIF-1 α . Notably, a previous study has demonstrated that depleting PHD2 in




Figure 7. Evaluating the role of PHD2 in tumor progression *in vivo*. (A and B) Results of tumor volumes observed in mice transplanted with colon cancer cells with indicated PHD2 modulations. (n=6; ***P<0.001 vs. NT or Mock). (C-F) Representative images of immunohistochemistry analysis and quantitative results demonstrating the Ki67 expression in xenografts harvested from indicated groups. Scale bar, 200 μ m. (n=6; ***P<0.001 vs. control groups). Kaplan-Meier survival curves were calculated for (G) SW480 subclones and (H) Ls174t subclones formed tumors as aforementioned (n=10; P<0.01 vs. control groups). PHD2, prolyl hydroxylase domain protein 2; NT, non-transduced; Mock, non-targeting shRNA control (for PHD2 knockdown) or vehicle control (for PHD2 overexpression).

NF- κ B signaling-induced cancer cell survival and therapy resistance have been reported in numerous malignancies, including colorectal cancer. Elevated NF- κ B activity has been observed in colorectal cancer cells treated with chemotherapeutics (54); therefore, targeting NF- κ B is expected to potentiate conventional therapeutic regimens in comprehensive strategies. Furthermore, an increasing number of studies indicate that metabolic pathways

may be favorable targets (55,56). For example, the present study found that PHD2 inhibited NF- κ B and knockdown of PHD2 led to upregulation of glycolysis in colorectal cancer cells, which essentially increases tumor cell proliferation and reduces animal survival. Taken together, evaluating PHD2 level may facilitate the selection of a targeted therapeutic regime for colorectal cancer with elevated NF- κ B activity.

Acknowledgements

The authors gratefully acknowledge Dr Xin Liu from the Department of Central Laboratory, Third Military Medical University (Army Medical University), for their kind help in providing RKO and LoVo cell lines. The authors would like to express their sincere gratitude to Pro Houjie Liang from the Third Military Medical University (Army Medical University) for generously providing the HT-29 cells used in the present study.

Funding

The present study was supported by the Sichuan Science and Technology Program (grant no. 2023YFS0318) and the Chengdu Science and Technology Program (grant no. 2022-YF05-01438-SN).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

GX and LX designed the study and wrote the manuscript. HW, LX, WY, SW and GX performed the experiments. LX, HW and SW analyzed the data. LX and GX advised on experimental procedures and revision of the paper. GX and LX confirm the authenticity of all the raw data. All authors contributed to this manuscript. All authors read and approved the final manuscript.

Ethics approval and consent to participate

The experimental protocol was established, according to the ethical guidelines of the 1964 Declaration of Helsinki. The protocol of immunohistochemistry for patient tissues was approved by the Ethics Committee of the First Affiliated Hospital (Southwest Hospital), the Third Military Medical University [Army Medical University; approval no. 2012 (12)] and all patients or family members involved provided written informed consent. Animal studies were performed under a protocol approved by the Army Medical University Institutional Animal Care and Use Committee (approval no. AMUWEC20232707).

Patient consent for publication

Written informed consent for publication was obtained from all participants.

Competing interests

The authors declare that they have no competing interests.

Authors' information

Dr Ganfeng Xie: ORCID: 0000-0003-0466-7710.

References

- Dang CV: Cancer metabolism: The known, unknowns. *Biochim Biophys Acta Rev Cancer* 1870: 1, 2018.
- Martinez-Reyes I and Chandel NS: Cancer metabolism: Looking forward. *Nat Rev Cancer* 21: 669-680, 2021.
- Trefts E and Shaw RJ: AMPK: Restoring metabolic homeostasis over space and time. *Mol Cell* 81: 3677-3690, 2021.
- Hanahan D and Weinberg RA: Hallmarks of cancer: The next generation. *Cell* 144: 646-674, 2011.
- Zecchin A, Kalucka J, Dubois C and Carmeliet P: How endothelial cells adapt their metabolism to form vessels in tumors. *Front Immunol* 8: 1750, 2017.
- Losman JA, Koivunen P and Kaelin WG Jr: 2-Oxoglutarate-dependent dioxygenases in cancer. *Nat Rev Cancer* 20: 710-726, 2020.
- Liao C and Zhang Q: Understanding the oxygen-sensing pathway and its therapeutic implications in diseases. *Am J Pathol* 190: 1584-1595, 2020.
- Semenza GL: The Genomics and genetics of oxygen homeostasis. *Annu Rev Genomics Hum Genet* 21: 183-204, 2020.
- Koivunen P and Kietzmann T: Hypoxia-inducible factor prolyl 4-hydroxylases and metabolism. *Trends Mol Med* 24: 1021-1035, 2018.
- Wong BW, Kuchnio A, Bruning U and Carmeliet P: Emerging novel functions of the oxygen-sensing prolyl hydroxylase domain enzymes. *Trends Biochem Sci* 38: 3-11, 2013.
- Maxwell PH and Eckardt KU: HIF prolyl hydroxylase inhibitors for the treatment of renal anaemia and beyond. *Nat Rev Nephrol* 12: 157-168, 2016.
- Singh L, Aldosary S, Saeedan AS, Ansari MN and Kaithwas G: Prolylhydroxylase 2: A promising target to inhibit hypoxia-induced cellular metabolism in cancer cells. *Drug Discov Today* 23: 1873-1882, 2018.
- Kato H, Inoue T, Asanoma K, Nishimura C, Matsuda T and Wake N: Induction of human endometrial cancer cell senescence through modulation of HIF-1 α activity by EGLN1. *Int J Cancer* 118: 1144-1153, 2006.
- Kaelin WG Jr: The VHL tumor suppressor gene: Insights into oxygen sensing and cancer. *Trans Am Clin Climatol Assoc* 128: 298-307, 2017.
- Chen F, Chen J, Yang L, Liu J, Zhang X, Zhang Y, Tu Q, Yin D, Lin D, Wong PP, *et al*: Extracellular vesicle-packaged HIF-1 α -stabilizing lncRNA from tumour-associated macrophages regulates aerobic glycolysis of breast cancer cells. *Nat Cell Biol* 21: 498-510, 2019.
- Sadiku P, Willson JA, Dickinson RS, Murphy F, Harris AJ, Lewis A, Sammut D, Mirchandani AS, Ryan E, Watts ER, *et al*: Prolyl hydroxylase 2 inactivation enhances glycogen storage and promotes excessive neutrophilic responses. *J Clin Invest* 127: 3407-3420, 2017.
- Guentsch A, Beneke A, Swain L, Farhat K, Nagarajan S, Wielockx B, Raithatha K, Dudek J, Rehling P, Ziesenis A, *et al*: PHD2 is a regulator for glycolytic reprogramming in macrophages. *Mol Cell Biol* 37: e00236-16, 2016.
- Xiang L, Gilkes DM, Hu H, Takano N, Luo W, Lu H, Bullen JW, Samanta D, Liang H and Semenza GL: Hypoxia-inducible factor 1 mediates TAZ expression and nuclear localization to induce the breast cancer stem cell phenotype. *Oncotarget* 5: 12509-12527, 2014.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
- Farooq SM, Hou Y, Li H, O'Meara M, Wang Y, Li C and Wang JM: Disruption of GPR35 exacerbates dextran sulfate sodium-induced colitis in mice. *Dig Dis Sci* 63: 2910-2922, 2018.
- Wang S, Wu Y, Hou Y, Guan X, Castelveter MP, Oblak JJ, Banerjee S, Filtz TM, Sarkar FH, Chen X, *et al*: CXCR2 macromolecular complex in pancreatic cancer: A potential therapeutic target in tumor growth. *Transl Oncol* 6: 216-225, 2013.
- Ruifrok AC and Johnston DA: Quantification of histochemical staining by color deconvolution. *Anal Quant Cytol Histol* 23: 291-299, 2001.
- Xie G, Zheng L, Ou J, Huang H, He J, Li J, Pan F and Liang H: Low expression of prolyl hydroxylase 2 is associated with tumor grade and poor prognosis in patients with colorectal cancer. *Exp Biol Med (Maywood)* 237: 860-866, 2012.
- Zhou L, Wang Y, Zhou M, Zhang Y, Wang P, Li X, Yang J, Wang H and Ding Z: HOXA9 inhibits HIF-1 α -mediated glycolysis through interacting with CRIP2 to repress cutaneous squamous cell carcinoma development. *Nat Commun* 9: 1480, 2018.

25. Dupuy F, Tabariès S, Andrzejewski S, Dong Z, Blagih J, Annis MG, Omeroglu A, Gao D, Leung S, Amir E, *et al*: PDK1-dependent metabolic reprogramming dictates metastatic potential in breast cancer. *Cell Metab* 22: 577-589, 2015.
26. Denko NC: Hypoxia, HIF1 and glucose metabolism in the solid tumour. *Nat Rev Cancer* 8: 705-713, 2008.
27. Jaakkola P, Mole DR, Tian YM, Wilson MI, Gielbert J, Gaskell SJ, von Kriegsheim A, Hebestreit HF, Mukherji M, Schofield CJ, *et al*: Targeting of HIF- α to the von Hippel-Lindau ubiquitylation complex by O₂-regulated prolyl hydroxylation. *Science* 292: 468-472, 2001.
28. Vidimar V, Licona C, Cerón-Camacho R, Guerin E, Coliat P, Venkatasamy A, Ali M, Guenot D, Le Lagadec R, Jung AC, *et al*: A redox ruthenium compound directly targets PHD2 and inhibits the HIF1 pathway to reduce tumor angiogenesis independently of p53. *Cancer Lett* 440-441: 145-155, 2019.
29. Meneses AM and Wielockx B: PHD2: From hypoxia regulation to disease progression. *Hypoxia (Auckl)* 4: 53-67, 2016.
30. Cummins EP, Berra E, Comerford KM, Ginouves A, Fitzgerald KT, Seeballuck F, Godson C, Nielsen JE, Moynagh P, Pouyssegur J and Taylor CT: Prolyl hydroxylase-1 negatively regulates IkkappaB kinase-beta, giving insight into hypoxia-induced NFkappaB activity. *Proc Natl Acad Sci USA* 103: 18154-18159, 2006.
31. Chan DA, Kawahara TLA, Sutphin PD, Chang HY, Chi JT and Giaccia AJ: Tumor vasculature is regulated by PHD2-mediated angiogenesis and bone marrow-derived cell recruitment. *Cancer Cell* 15: 527-538, 2009.
32. Hoesel B and Schmid JA: The complexity of NF- κ B signaling in inflammation and cancer. *Mol Cancer* 12: 86, 2013.
33. Birbach A, Gold P, Binder BR, Hofer E, de Martin R and Schmid JA: Signaling molecules of the NF-kappa B pathway shuttle constitutively between cytoplasm and nucleus. *J Biol Chem* 277: 10842-10851, 2002.
34. Rückert F, Grützmann R and Pilarsky C: Feedback within the inter-cellular communication and tumorigenesis in carcinomas. *PLoS One* 7: e36719, 2012.
35. Stine ZE, Walton ZE, Altman BJ, Hsieh AL and Dang CV: MYC, metabolism, and cancer. *Cancer Discov* 5: 1024-1039, 2015.
36. Obacz J, Pastorekova S, Vojtesek B and Hrstka R: Cross-talk between HIF and p53 as mediators of molecular responses to physiological and genotoxic stresses. *Mol Cancer* 12: 93, 2013.
37. Johnson RF and Perkins ND: Nuclear factor- κ B, p53, and mitochondria: Regulation of cellular metabolism and the Warburg effect. *Trends Biochem Sci* 37: 317-324, 2012.
38. Wilde L, Roche M, Domingo-Vidal M, Tanson K, Philp N, Curry J and Martinez-Outschoorn U: Metabolic coupling and the reverse Warburg effect in cancer: Implications for novel biomarker and anticancer agent development. *Semin Oncol* 44: 198-203, 2017.
39. Wang X, Liu R, Qu X, Yu H, Chu H, Zhang Y, Zhu W, Wu X, Gao H, Tao B, *et al*: α -Ketoglutarate-activated NF- κ B signaling promotes compensatory glucose uptake and brain tumor development. *Mol Cell* 76: 148-162.e7, 2019.
40. Ishak Gabra MB, Yang Y, Lowman XH, Reid MA, Tran TQ and Kong M: IKK β activates p53 to promote cancer cell adaptation to glutamine deprivation. *Oncogenesis* 7: 93, 2018.
41. Ruf M, Pitzen T, Meyer C and Drumm J: Atlantoaxial rotatory dislocation: Delayed diagnose will result in more invasive treatment options. *J Neurol Surg A Cent Eur Neurosurg* 82: 1-8, 2021.
42. Taniguchi K and Karin M: NF- κ B, inflammation, immunity and cancer: Coming of age. *Nat Rev Immunol* 18: 309-324, 2018.
43. Gaete D, Rodriguez D, Watts D, Sormendi S, Chavakis T and Wielockx B: HIF-prolyl hydroxylase domain proteins (PHDs) in cancer-potential targets for anti-tumor therapy? *Cancers (Basel)* 13: 988, 2021.
44. Chen T, Zhou Q, Tang H, Bozkanat M, Yuan JX, Raj JU and Zhou G: miR-17/20 controls prolyl hydroxylase 2 (PHD2)/hypoxia-inducible factor 1 (HIF1) to regulate pulmonary artery smooth muscle cell proliferation. *J Am Heart Assoc* 5: e004510, 2016.
45. Lee G, Won HS, Lee YM, Choi JW, Oh TI, Jang JH, Choi DK, Lim BO, Kim YJ, Park JW, *et al*: Oxidative dimerization of PHD2 is responsible for its inactivation and contributes to metabolic reprogramming via HIF- α activation. *Sci Rep* 6: 18928, 2016.
46. Li J, Yuan W, Jiang S, Ye W, Yang H, Shapiro IM and Risbud MV: Prolyl-4-hydroxylase domain protein 2 controls NF- κ B/p65 transactivation and enhances the catabolic effects of inflammatory cytokines on cells of the nucleus pulposus. *J Biol Chem* 290: 7195-7207, 2015.
47. Wang L, Niu Z, Wang X, Li Z, Liu Y, Luo F and Yan X: PHD2 exerts anti-cancer and anti-inflammatory effects in colon cancer xenografts mice via attenuating NF- κ B activity. *Life Sci* 242: 117167, 2020.
48. Lee DC, Sohn HA, Park ZY, Oh S, Kang YK, Lee KM, Kang M, Jang YJ, Yang SJ and Hong YK: A lactate-induced response to hypoxia. *Cell* 161: 595-609, 2015.
49. Romero-Ruiz A, Bautista L, Navarro V, Heras-Garvín A, March-Díaz R, Castellano A, Gómez-Díaz R, Castro MJ, Berra E, López-Barneo J and Pascual A: Prolyl hydroxylase-dependent modulation of eukaryotic elongation factor 2 activity and protein translation under acute hypoxia. *J Biol Chem* 287: 9651-9658, 2012.
50. Wang X, Xie J and Proud CG: Eukaryotic elongation factor 2 kinase (eEF2K) in cancer. *Cancers* 9: 162, 2017.
51. Xue J, Li X, Jiao S, Wei Y, Wu G and Fang J: Prolyl hydroxylase-3 is down-regulated in colorectal cancer cells and inhibits IKKbeta independent of hydroxylase activity. *Gastroenterology* 138: 606-615, 2010.
52. Lind DS, Hochwald SN, Malaty J, Rekkas S, Hebig P, Mishra G, Moldawer LL, Copeland EM III and Mackay S: Nuclear factor-kappa B is upregulated in colorectal cancer. *Surgery* 130: 363-369, 2001.
53. Wang Z, Sheng C, Kan G, Yao C, Geng R and Chen S: RNAi screening identifies that TEX10 promotes the proliferation of colorectal cancer cells by increasing NF- κ B activation. *Adv Sci (Weinh)* 7: 2000593, 2020.
54. Goka ET, Chaturvedi P, Lopez DTM, Garza A and Lippman ME: RAC1b overexpression confers resistance to chemotherapy treatment in colorectal cancer. *Mol Cancer Ther* 18: 957-968, 2019.
55. Gyamfi J, Kim J and Choi J: Cancer as a metabolic disorder. *Int J Mol Sci* 23: 1155, 2022.
56. Ge T, Gu X, Jia R, Ge S, Chai P, Zhuang A and Fan X: Crosstalk between metabolic reprogramming and epigenetics in cancer: Updates on mechanisms and therapeutic opportunities. *Cancer Commun (Lond)* 42: 1049-1082, 2022.



Copyright © 2023 Xiang et al. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.