

CORRIGENDUM

DOI: 10.3892/mco.2017.1315

High-dose chemotherapy with autologous peripheral blood stem cell transplantation for choriocarcinoma: A case report and literature review

EIKO YAMAMOTO, KAORU NIIMI, KAYO FUJIKAKE, TETSUYANISHIDA, MAKOTO MURATA, AYAKO MITSUMA, YUICHI ANDO and FUMITAKA KIKKAWA

Mol Clin Oncol 5: 660-664, 2016; DOI: 10.3892/mco.2016.1011

Following the publication of the article, the authors noticed the following error: In Table I, the dose of Ifosfamide in mini ICE regimen should be 1,000 mg/m² instead of 2,000 mg/m². This had occurred due to a typing error on behalf of the authors upon submission. Shown below is the new version of Table I:

Table I. Mini and high-dose ifosfamide, carboplatin and etoposide (ICE) regimens.

Agents	Dose	Hours	Days
Mini ICE			
Ifosfamide	1,000 mg/m ² d.i.v.	1	1-2
Etoposide	150 mg/m ² d.i.v.	11	1-2
Carboplatin	200 mg/m ² d.i.v.	1	1-2
Etoposide	150 mg/m ² d.i.v.	11	1-2
High-dose ICE			
Ifosfamide	3,000 mg/m ² d.i.v.	1	1-4
Etoposide	200 mg/m ² d.i.v.	11	1-3
Carboplatin	300 mg/m ² d.i.v.	1	1-4
Etoposide	200 mg/m ² d.i.v.	11	1-3

d.i.v., drip intravenous infusion.

In addition, the authors noted that the concentration unit of G-CSF mentioned on page 661 should be µg/kg/day instead of mg/m²/day or g/kg/day.