# Dual matrilineal geographic distribution of Korean type 2 diabetes mellitus-associated -11,377 G adiponectin allele

JEE-HYE CHOI<sup>1,2</sup>, NA YOUNG MIN<sup>2</sup>, SANG KIL PARK<sup>1</sup>, LKHAGVASUREN GAVAACHIMED<sup>3</sup>, YOUNG JONG KO<sup>2</sup>, SUNG HOON HAN<sup>2</sup>, KYUNG YONG KIM<sup>4</sup>, KIJUNG KIM<sup>4</sup>, KWANG HO LEE<sup>2</sup> and AE JA PARK<sup>1</sup>

<sup>1</sup>Department of Laboratory Medicine, College of Medicine; <sup>2</sup>Department of Life Science, College of Natural Science, <sup>3</sup>Department of Science and Culture, Graduate School; <sup>4</sup>Department of Microbiology, College of Medicine, Chung-Ang University, Dongjak-gu, Seoul 156-756, South Korea

Received October 22, 2013; Accepted June 26, 2014

DOI: 10.3892/mmr.2014.2639

Abstract. The present study was performed to identify the susceptible single nucleotide polymorphisms (SNPs) for the prediction of Korean type 2 diabetes mellitus (T2DM) and to clarify the matrilineal origin of Korean T2DM-specific SNPs. Fourteen SNPs from the adiponectin (ADIPOQ), hepatocyte nuclear factor  $4\alpha$ , phosphoenolpyruvate carboxykinase 1 and glucokinase genes in the Korean population were analyzed. Only one SNP, -11,377 C/G on the ADIPOQ gene, was finally determined to be responsible for the incidence of Korean T2DM (P=0.028). The G-T-T-A haplotype at positions -11,377, +45, +276 and +349 on the ADIPOQ gene was also associated with a high incidence of Korean T2DM (P=0.023). In addition, the susceptibility of Korean individuals to T2DM appears to be affected by their matrilineal origin. Of note, the group of Southern origin, consisting of mitochondrial DNA macrohaplogroups F and R, was predisposed to T2DM, whereas the group of Northern origin, consisting of haplogroups A and Y, was resistant to T2DM. This implied that the differential genetics between the two groups, which were formed from the initial peopling of the proto-Korean population via Southern and Northern routes to the present time, may explain their differing susceptibility to T2DM. In conclusion, from Southern Asia Northward, a matrilineal origin of Korean individuals

Professor Kwang-Ho Lee, Department of Life Science, College of Natural Science, Chung-Ang University, 221 Heuksuk-Dong, Dongjak-Ku, Seoul 156-756, South Korea E-mail: leemanse@cau.ac.kr

*Key words:* type 2 diabetes mellitus, single nucleotide polymorphism, haplotype, adiponectin, hepatocyte nuclear factor  $4\alpha$ , phosphoenolpyruvate carboxykinase 1, glucokinase, mitochondrial DNA

appears to be responsible for the prevalence of Korean T2DM caused by the -11,377 G allele.

#### Introduction

Type 2 diabetes mellitus (T2DM), one of the most common types of diabetes, is a multifactorial disease that appears to differ in prevalence based on ethnicity. To date, ~40 susceptibility loci for T2DM have been discovered through genome-wide association studies (1-3); however, the majority of the genetic loci have been investigated in Caucasian European subjects and the results are not always replicable in other ethnic groups, including Asian populations. For example, +45 T/G and +276 G/T single nucleotide polymorphisms (SNPs) of adiponectin (ADIPOO, 3q27) are highly involved SNPs as candidate risk variants for T2DM in Asian populations, whereas these same SNPs are not correlated with a risk of T2DM in Europeans, including Italian, French and Swedish individuals (4-8). Gong et al (9) described in detail the ethnic differences in SNPs (-11,391 A/G and -11,377 C/G of ADIPOQ) associated with T2DM on the basis of published data obtained from MEDLINE, EMBASE and Science citation index expanded databases. According to this study, a G allele at the position -11,377 of ADIPOQ increased the risk of T2DM in European populations, but did not appear to affect the incidence of T2DM in Asian individuals. Therefore, the results from European studies do not appear to coincide with those from other ethnic groups with different genetic backgrounds and environmental factors. Gong et al (9) suggested that ethnic differences in the -11,377 G ADIPOQ allele in correlation with the susceptibility to T2DM may be due to the combined effect of the subjects' discrete genetic components and somatic compliance affected by environmental factors, as well as the lower size and heterogeneity of the samples examined, which had been formed over long period from initial peopling to the ethnic distribution of the present time. However, the effect of human migration on the peopling of ethnic groups and the formation of ethnicity-specific genetic components has been underestimated in the majority of the association studies between SNPs and the susceptibility to certain diseases, including diabetes.

Gong et al (9) performed a meta-analysis for the association between ADIPOQ SNPs and the incidence of T2DM using data from a large population of European and Asian subjects

*Correspondence to:* Professor Ae Ja Park, Department of Laboratory Medicine, College of Medicine, Chung-Ang University, 221 Heuksuk-Dong, Dongjak-gu, Seoul 156-756, South Korea E-mail: ajcp@unitel.co.kr

who had already been screened for SNPs susceptible to T2DM in several studies. The authors proposed that there existed a statistically significant heterogeneity (P=0.043) in seven Asian studies consisting of Chinese and Japanese populations, and a considerable level of homogeneity (P=0.133) was prominent in six European studies, including British, French, German, Swedish and Swiss subjects. The -11,377 G ADIPOQ allele was proven to be closely correlated with T2DM in the majority of the European populations screened, but demonstrated varied association with the incidence of T2DM in the Chinese and Japanese ethnic groups. A series of association studies for the identification of T2DM-associated SNPs in a Han Chinese population, which may have consisted of different ethnic groups each with a unique history of initial peopling and residing in different regions of China, produced contradictory results in linking ethnic-specific SNPs with the occurrence of T2DM, suggesting the importance of determining the ethnic origin of subjects in an association study (10). These findings implied that the ethnic-specific physical and genetic background during ethnogenesis may impact the susceptibility to T2DM, even in Asian populations who are expected to be genetically similar to each other, reflecting that the ethnic origin of T2DM-associated SNPs should be a concern when identifying the ethnic-specific SNPs responsible for T2DM. Therefore, it is apparent from several studies that the ethnogenesis of individuals, as well as the sample size and genome heterogeneity, need to be carefully considered in an association study.

To assess the correlation of the genotype as SNPs and haplotypes with the occurrence of T2DM, the present study aimed to identify Korean-specific SNPs and haplotypes, which are associated with a predisposition to T2DM, and to define the matrilineal origin of Korean-specific, T2DM-associated SNPs.

# Materials and methods

Subjects. Blood samples were collected from 194 normal subjects and 328 T2DM patients who visited Chung-Ang University Hospital (Seoul, South Korea). Unassociated normal subjects were randomly recruited from a population undergoing routine health check-ups and diabetic subjects were randomly recruited from patients at the outpatient clinic of the Department of Metabolic Diseases (Chung-Ang University Hospital). The diabetes status was determined according to the World Health Organization criteria (11). Whole blood samples were collected from all of the subjects, and the fasting levels of glucose, insulin and glycated hemoglobin (HbA1c) were measured. All subjects enrolled in the present study were of full Korean ethnicity and their clinical characteristics are summarized in Table I. Written informed consent was obtained from all non-diabetic and diabetic subjects. The study procedures was approved by the Institutional Review Board of Chung-Ang University Hospital.

*Genotyping of polymorphisms*. Genomic DNA was extracted from blood samples with EDTA using the Qiagen DNA isolation kit (Qiagen, Chatsworth, CA, USA). The extracted DNA was quantified using a NanoDrop ND-1000 spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA), and its quality was assessed by agarose gel (Bioneer, Dajeon, South Korea) electrophoresis and staining with ethidium bromide (Bioneer). A total of nine sites on the nuclear genome (two sites on ADIPOQ, four sites on hepatocyte nuclear factor (HNF)4 $\alpha$ , two sites on PCK1 and one site on GCK) that are already known to be associated with T2DM according to previous studies, were screened to identify mutations or polymorphisms by polymerase chain reaction (PCR)-directed sequencing (12-15). Primer information is presented in Table II. PCR was performed in 50 µl Accupower PCR premix (Bioneer). PCR cycling conditions were as follows: Denaturation at 94°C for 40 sec, followed by 35 cycles of denaturation for 40 sec, annealing at 54°C for 40 sec and extension at 72°C for 40 sec, with a final extension at 72°C. Following PCR amplification, the PCR products were purified with a Gel/PCR DNA Extraction kit (Real Biotech Corp., Taipei, Taiwan) and then sequenced by a Macrogen Service Center (Macrogen Corp., Seoul, Korea). To identify the SNPs, the sequencing results were analyzed using SeqMan<sup>TM</sup>II software (DNASTAR, Madison, WI, USA).

Mitochondrial (mt)DNA haplogroup analysis. To determine the mtDNA haplogroup of Korean subjects with the -11,377 G ADIPOQ allele, hypervariable region 1 (HV1) and HV2 were analyzed for their DNA sequence. PCR was firstly performed for DNA sequencing using the following conditions: 95°C for 10 min, 40 cycles of 30 sec at 95°C, 1 min at the annealing temperature, 1 min at 72°C, followed by a final extension step of 7 min at 72°C. Following PCR amplification, PCR products were purified with a Gel/PCR DNA Extraction kit (Real Biotech Corp.) and subjected to TA cloning. Four colonies obtained from TA cloning products were sequenced by a Macrogen Service Center (Macrogen Corp.). Sequencing results were analyzed using SeqMan<sup>TM</sup>II software (DNASTAR) and compared with the revised Cambridge reference sequence (rCRS) (16). Haplogroups were established with HV1 and HV2 polymorphisms using the well-established web-based programs, Haplogroup Prediction tool and mtDNA Manager (http://mtmanager.yonsei.ac.kr).

Statistical analysis. The frequencies of genotypes or alleles were compared by  $\chi^2$  analysis. The odds ratios (OR) and 95% confidence intervals (CIs) with adjustment for age and sex were calculated by logistic regression analysis. P<0.05 was considered to indicate a statistically significant result. Association analyses for allele types and genotypes were conducted using SPSS version 13.0 (SPSS, Inc., Chicago, IL, USA). Hardy-Weinberg equilibrium (HWE) and linkage disequilibrium (LD) statistics were obtained through the use of Haploview software version 3.2 (Broad Institute, Cambridge, MA, USA) and the Haplo.stats Package for R was used to predict haplotypes and to directly estimate their frequencies for the haplotype association with T2DM (17). Haplo.stats analysis was restricted to haplotypes with an inferred frequency >1%. Principal component analysis (PCA) was performed with the mtDNA haplogroup frequencies of 33 Asian populations using SPSS software, which were obtained from the literature and the data of the present study (18-27).

	T2DM	l subjects	Normal subjects		
Characteristic	Male (n=120)	Female (n=208)	Male (n=85)	Female (n=109)	
Fasting blood glucose (mg/dl)	140±45.8	125.7±47.2	93.4±9.4	90.56±7.7	
Total cholesterol (mg/dl)	176.3±40.7	190.9±38.9	184.9±25.1	185.5±27.3	
HDL-cholesterol (mg/dl)	44.5±11	49.5±12.4	50.3±9.9	55.4±11.6	
LDL-cholesterol (mg/dl)	103.9±31.9	113.5±30.5	114±23.5	110.1±24.7	
HbA1c (%)	7.2±1.6	6.8±1.5	5.1±0.5	4.8±1.3	
Triglycerides (mg/dl)	158.6±137.1	114.9±66.7	90±28.6	83.3±34.6	

Table I. Clinical characteristics of Korean T2DM pa	atients and normal control subjects.
---	--------------------------------------

T2DM, type 2 diabetes mellitus; DHL, high-density lipoprotein; LDL, low-density lipoprotein; HbA1c, glycated hemoglobin.

Table II. Primers for PCR amplification of T2DM-associated genes.

Gene	Primer name Locus		Primer sequence	Product size (bp)	
ADIPOQ A pr		Promoter	5'-CACTTGCCCTGCCTCTGTC-3'		
~	-1		5'-TATGCCTATTCTGTCTCTTGTGGA-3'	296	
	A_ex	Exon 2 and intron 3	5'-CATGGGAAAATTAGAGGAGTGTC-3'		
			5'-GGAATAGGGATGAGGGTGAAGATG-3'	767	
PCK1	P_J1	Promoter	5'-GCCAAGCCCCCTCCACTCCCTAAA-3'	428	
			5'-CGCCCCTGCCCCGATCTAAAC-3'		
	P_J2	Exon	5'-GGGGCAGGGCTATTCTTTTCCACA-3'	596	
			5'-GGGGGCCGAGCTTACCAGTTG-3'		
HNF-4 $\alpha$	H_J1	P2 promoter	5'-CCGCTTTCATTCGTCTCATAACAG-3'	387	
		-	5'-GAGACCGAGGCAGAGATAGAGAAA-3'		
	H_J2	P1 promoter	5'-CCCGCCGGTATTTGGTCTC-3'	358	
			5'-TCTGCCCTGCTTAGCTTTATTCTG-3'		
	H_J4	Intron 1	5'-CTCCCAGGCCCTTCCAGTCC-3'	368	
			5'-TCACCCCACCCATACAACCTACAT-3'		
	H_J5	Intron 2	5'-GCTCATGGCCCCAAGGTCTGTCT-3'	345	
			5'-AACGGCCTGGGGTCAATCTCCTAT-3'		
GCK	G_pr	Promoter	5'-GACTGTGTCTCTCACATCCT-4'	258	
			5'-TAAGGCGATTGAGTGGTC-4'		
HV1	FN15977	15977	5'-CCACCATTAGCACCCAAAGC-4'	423	
	RN16399	16399	5'-TCAAGGGACCCCTATCTGA-4'		
HV2	FN29	29	5'-CTCACGGGAGCTCTCCAT-4'	353	
	R381	381	5'-GCTGGTGTTAGGGTTCTTTG-4'		

T2DM, type 2 diabetes mellitus; *ADIPOQ*, adiponectin; HNF, hepatocyte nuclear factor; PCK1, phosphoenolpyruvate carboxykinase 1; GCK, glucokinase; HV, hypervariable region; PCR, polymerase chain reaction.

# Results

Identification of T2DM-associated SNPs in a Korean population. DNA sequences of nine regions on ADIPOQ, HNF4 $\alpha$ , phosphoenolpyruvate carboxykinase (PCK)1 and glucokinase (GCK) genes that are well established as closely associated with the incidence of T2DM in several ethnic groups, were analyzed to identify the SNPs susceptible to Korean T2DM. As a result, fourteen SNPs of the four genes were identified in 328 Korean T2DM patients and 194 normal individuals, and statistically analyzed to define a correlation with the risk of T2DM (Table III).

The fourteen SNPs were located in variable regions, including the promoter, exon and intron of each gene (Table III). Five SNPs from *ADIPOQ* were separately located on the promoter (-11,426 and -11,377), exon 2 (+45), and intron 2 (+276 and +349) regions, and seven SNPs from *HNF4a* were on the promoter (-4031, -3927 and +43582), intron 1 (+45980

			Allele freque					Genotype frequencies		
dbSNPs	(locations)	Alleles	T2DM	Normal	OR (95%CI)	P-value	Genotypes	OR (95%CI)	P-value	HWE P-value
rs16861194	ADIPOQ	А	0.77	0.75	0.98.	0.882	AA	0.70	0.419	0.992
	-11,426	G	0.23	0.25	(0.71-1.34)		AG and GG	(0.30-1.66)		
rs266729	ADIPOQ	С	0.73	0.80	1.51	0.028ª	CC	1.75	0.017ª	0.158
	-11,377	G	0.27	0.20	(1.05-2.18)		CG and GG	(1.10-2.76)		
rs2241766	ADIPOQ	G	0.72	0.67	1.10	0.567	GG	0.87	0.553	0.662
	+45	Т	0.28	0.33	(0.79-1.55)		GT and TT	(0.56-1.36)		
rs1501299	ADIPOQ	G	0.69	0.71	1.14	0.441	GG	1.08	0.741	0.206
	+276	Т	0.31	0.29	(0.82 - 1.60)		GT and TT	(0.69-1.68)		
rs2241767	ADIPOQ	А	0.75	0.82	1.00	0.993	AA	1.17	0.514	< 0.001
	+349	G	0.25	0.18	(0.69-1.45)		AG and GG	(0.73-1.86)		
rs1884613	$HNF4\alpha$	С	0.49	0.51	1.03	0.852	CC	0.73	0.527	0.480
	-4,031	G	0.51	0.49	(0.75-1.41)		CG and GG	(0.28-1.91)		
rs1884614	HNF4a	С	0.49	0.53	1.33	0.077	CC	1.45	0.166	0.151
	-3,927	Т	0.51	0.47	(0.97-1.81)		CT and TT	(0.86-2.45)		
rs2425640	$HNF4\alpha$	G	0.78	0.77	0.96	0.813	GG	0.84	0.704	0.911
	+43,582	А	0.22	0.23	(0.70-1.32)		GA and AA	(0.34-2.06)		
rs2071197	$HNF4\alpha$	G	0.53	0.52	1.00	0.994	GG	0.97	0.947	0.965
	+45,980	А	0.47	0.48	(0.73-1.36)		GA and AA	(0.37-2.52)		
rs2071198	$HNF4\alpha$	А	0.98	0.98	1.40	0.541	AA	1.66	0.359	0.581
	+46,029	Т	0.02	0.02	(0.48 - 4.08)		AT and TT	(0.52 - 5.34)		
rs3212183	HNF4a	Т	0.93	0.92	1.09	0.625	TT	0.84	0.776	0.169
	+55,462	С	0.07	0.08	(0.78-1.51)		TC and CC	(0.25 - 2.85)		
rs3212184	HNF4a	С	0.97	0.98	1.18	0.342	CC	3.33	0.215	0.612
	+55,484	G	0.03	0.02	(0.84-1.64)		CG and GG	(0.50-22.34)		
rs1799884	GCK	G	0.83	0.81	0.97	0.845	GG	0.82	0.680	0.070
	-30	А	0.17	0.19	(0.70-1.33)		GA and AA	(0.33-2.07)		
rs1042521	PCK1	А	0.62	0.56	0.98	0.890	AA	1.18	0.500	0.327
	-232	G	0.38	0.44	(0.71-1.34)	-	AG and GG	(0.73-1.88)	-	

Table III. Allele frequencies of SNPs in Korean T2DM subjects and normal individuals.

<sup>a</sup>Significant P-values (P<0.05). T2DM, type 2 diabetes mellitus; *ADIPOQ*, adiponectin; HWE, Hardy-Weinberg equilibrium; SNPs, single nucleotide polymorphisms; HNF, hepatocyte nuclear factor; PCK1, phosphoenolpyruvate carboxykinase 1; GCK, clucokinase; OR, odds ratio; 95% CI, 95% confidence interval.

and +46029) and intron 2 (+55462 and +55484) regions. The two remaining SNPs identified on *GCK* and *PCK1* were at the -30 and -232 sites on the promoter region of the each gene, respectively. All SNPs identified were concordant with variants that are well established as potential genetic makers correlated with the prevalence of T2DM in European and Japanese populations (6,7,28,29). The allele frequencies of the fourteen SNPs are provided in Table III. Among them, 13 SNPs, except +349 SNP on *ADIPOQ*, were in HWE. Severe violation of the HWE (P<0.001) on the +349 SNP was observed in the normal Korean group, while it was not observed in the T2DM patients.

To identify the Korean T2DM-specific SNPs, odds ratios (ORs) and 95% confidence intervals (CIs), according to the distribution of allelotypes, were calculated by multivariate statistical analysis, compensating for gender and age among the subjects examined (Table III). The results demonstrated that only one SNP, -11,377 C/G of *ADIPOQ*, was closely associated

with the incidence of T2DM (P=0.028), implying that the -11,377 G allele may be a putative genetic marker to predict the predisposition of Korean individuals to T2DM. The correlation of the -11,377 G allele with the risk of Korean individuals for T2DM was confirmed by genotype analysis (Table III). According to the genotype analysis, Korean subjects of the CG or GG genotypes at the -11,377 site of *ADIPOQ* are 1.75x more vulnerable to developing T2DM than Korean individuals of the CC type (P=0.017; 95% CI, 1.10-2.76). However, all other SNPs (except for the SNP at the -11,377 site of *ADIPOQ*) demonstrated no statistical difference between T2DM and normal subjects (P>0.05).

Identification of the T2DM-associated ADIPOQ haplotype in a Korean population. Although none of the 13 SNPs except for -11,377 C/G of ADIPOQ were associated with T2DM, haplotypes, a set of closely linked SNPs on the same gene, may be

Table IV. ADIPOQ	haplotypes an	d risk of T2DM	in Korean	individuals

Haplotype -11,426/-11,377/+45/+276/+349	Freq.	OR	95% CI	P-value
+45/+276				
TG	0.395	1	Reference haplotype	
GG	0.302	1.02	(0.69-1.50)	0.929
TT	0.303	1.18	(0.81-1.73)	0.389
+45/+276/+349				
TGA	0.396	1	Reference l	haplotype
GGA	0.078	0.67	(0.32-1.41)	0.293
GGG	0.223	1.08	(0.72-1.63)	0.712
TTA	0.304	1.17	(0.80-1.71)	0.423
-11,426/-11,377				
AC	0.529	1	Reference l	haplotype
AG	0.232	0.91	(0.43-1.92)	0.803
GC	0.236	0.75	(0.34-1.68)	0.483
-11,377/+45/+276/+349				
CTGA	0.256	1	Reference haplotype	
CGGA	0.069	0.8	(0.33-1.93)	0.619
CGGG	0.206	1.26	(0.77-2.05)	0.353
CTTA	0.227	1.15	(0.69-1.89)	0.595
GGGG	0.016	1.03	(0.21-5.07)	0.969
GTGA	0.139	1.59	(0.87-2.89)	0.132
GTTA	0.0379	2.55	(1.14-5.73)	0.023ª
-11,426/-11,377/+45/+276/+349				
ACTGA	0.241	1	Reference l	haplotype
ACGGA	0.104	0.82	(0.20-3.37)	0.788
ACGGG	0.104	1.47	(0.26-8.19)	0.660
ACTTA	0.074	0.49	(0.11-2.21)	0.356
AGGGA	0.020	0.18	(0.01-4.25)	0.290
AGGGG	0.013	4.54	(0.02-896.53)	0.575
AGTGA	0.132	0.45	(0.13-1.53)	0.202
AGTTA	0.068	2.78	(0.36-21.27)	0.326
GCGGA	0.037	0.44	(0.04-4.53)	0.494
GCGGG	0.026	0.3	(0.02-3.76)	0.351
GCTGA	0.037	0.41	(0.04-4.15)	0.451
GCTTA	0.141	0.77	(0.23-2.59)	0.676

<sup>a</sup>Significant P-values (P<0.05); OR, odds ratio; 95% CI, 95% confidence interval; T2DM, type 2 diabetes mellitus; *ADIPOQ*, adiponectin; SNPs, single nucleotide polymorphisms.

more informative in studying T2DM in Korean individuals. All 12 SNPs on the *ADIPOQ* and *HNF4a* that were identified in the T2DM and normal groups were thus combined to determine the haploblock through LD analysis. In the case of *HNF4a*, a haploblock that consisted of -4,031 C/G and -3,927 was found to be highly linked to T2DM (D'=1), but haplotypes from various combinations of a total of seven SNPs, including two SNPs involved in the haploblock, were all proven not to be associated with the incidence of T2DM (data not shown). By contrast, the +45 T/G, +276 G/T and +349 A/G SNPs of *ADIPOQ* were found to be highly linked, as determined by the LD test (D'=1). There was low LD between the SNPs on the promoter and the SNPs

on the exon and intron (Fig. 1). Considering the results of the LD analysis, haplotype analysis was performed for the haploblock combined with +45 T/G, +276 G/T and +349 A/G SNPs. The four haplotypes (T-G-A, G-G-A, G-G-G and T-T-A) with a frequency >1% were not associated with T2DM (Table IV). Although other SNPs were not in the high LD block, an additional 25 haplotypes with a frequency of >1% were detected in *ADIPOQ* through the Haplo Stats package for ambiguous haplotypes (Table IV). Of the haplotypes, the most common haplotype was handled as a reference haplotype to analyze the correlation of the haplotypes with Korean T2DM. The results of the haplotype analysis demonstrated that the G-T-T-A haplotype

Characteristic	Haplogroup	Sub-haplogroup	T2DM (n=23)	Normal (n=15)	
Mostly found in Northeast Asia	А	A5a	-	1 (7%)	
	D	D4 (D4, D4a, D4b1, D4b2b, D4c, D4e1, D4i, D4j*, D4j3, D4k) D5a2	8 (35%)	4 (27%)	
	G	G2a1	2 (9%)	1 (7%)	
	Y	Y Y1	-	3 (20%)	
		Sub total	10 (43%)	9 (60%)	
Mostly found in Southeast Asia	М	M7 (M7a1, M7b2 M7c1)	7 (30%)	4 (27%)	
		M51° M37a M10b			
	Ν	N9 (N9a, N9b)	1 (4%)	1 (7%)	
	В	B4 (B4a, B4b1)	3 (13%)	1 (7%)	
	Б	BDD2 F2a	1(407)		
	Г D	ГJа D20e <sup>*</sup>	1(4%) 1(4%)	-	
	К	Sub total	13 (57%)	6 (40%)	

Table V. Distribution of mitochondrial DNA haplogroup frequencies in Korean individuals with the -11,377 GG genotype of *ADIPOQ*.

T2DM, type 2 diabetes mellitus; ADIPOQ, adiponectin.



Figure 1. Principal component analysis of mtDNA haplogroup frequencies in Korean T2DM and normal subjects in 33 Asian populations. The Principal component analysis of mtDNA haplotype frequencies demonstrated that Southern Asian populations, including those from China, Okinawa and Hong Kong, were largely grouped with Korean T2DM subjects, whereas northern Asian populations, including Eskimo, Yakut and Chukchi, tended to be grouped with normal Korean subjects. T2DM, type 2 diabetes mellitus; mtDNA, mitochondrial DNA.



Figure 2. Migratory history and geographical distribution of mtDNA haplogroups attributed to peopling of the Korean population. The black arrows represent migration routes and divergence of the mtDNA haplogroup from Africa to Asia. Dotted lines represent areas that were proven to be correlated with normal and T2DM Korean groups based on the principal component analysis results. T2DM, type 2 diabetes mellitus; mtDNA, mitochondrial DNA.



Figure 3. Linkage disequilibrium for the single nucleotide polymorphisms on (A) ADIPOQ and (B) HNF4a identified in Koreans. ADIPOQ, adiponectin; HNF, hepatocyte nuclear factor.

at the -11,377, +45, +276 and +394 SNP positions in *ADIPOQ* was closely correlated with Korean T2DM (OR, 2.55; 95% CI, 1.14-5.73; P=0.023). These results indicated that Koreans with the G-T-T-A haplotype of *ADIPOQ* are more susceptible to T2DM than Korean individuals with any other haplotype.

Dual matrilineal geographic distribution of the -11,377 GG genotype of ADIPOQ in a Korean population. As demonstrated

in Tables III and IV, the -11,377 GG genotype of *ADIPOQ* was closely linked with Korean T2DM. Of the 38 subjects demonstrating the -11,377 GG genotype of *ADIPOQ*, 24 subjects were T2DM patients and 14 were T2DM negative. Korean-specific susceptibility to T2DM was determined by varied mutations on the genome that were accumulated during the formation of the Korean population. To analyze the cause of differential susceptibility to T2DM in Korean subjects with the identical

-11,377 GG genotype of *ADIPOQ*, the matrilineal origin of the 38 subjects was traced by determining haplogroups of mtDNA, which are diverged with consecutive mutational steps on the genome, and reliably reflect the ancient migration history and recent admixture of populations. The mtDNA haplogroups of 38 subjects were determined through sequencing of the HV1 (16024~16380) and HV2 (47~360) regions of mtDNA. The haplogroup frequencies of Korean T2DM and normal groups are summarized in Table V. Based on these haplogroup assignments, subjects were divided into two large lineages with the Southern (M7, N9, B, R and F) and the Northern (D, G, A and Y) haplogroup complexes of East Asia, reflecting that the Korean population was formed by a mixture of dual matrilineal origins via Northward and Southward routes.

Nine mtDNA haplogroups (A, D, G, Y, M, N, B, F and R) were identified in 38 subjects demonstrating the -11,377 GG genotype of ADIPOQ. The nine mtDNA haplogroups were divided into 15 subclades (A5a, D4, D5a2, G2a1, Y, Y1, M7, M51\*, M37a, M10b, N9, B4, B5b2, F3a and R30a; Table V). Of them, the A, D and Y haplogroups were widespread in Northeast Asia, Northeast Siberia and America, representing the Northern origin of Asian populations, including Eskimo, Na-denes, Chukchis, Nivkhs and Ainus populations (30,31). A total of 19/38 subjects (50%) demonstrated these haplogroups. The M, N, B, F and R haplogroups are found in Southeast Asia, including Bangladesh, Indonesia, Malaysia, Vietnam and Japan (32), representing the Southern origin of the Asian population. Those haplogroups were also found in 19/38 subjects (50%) in the present study. This result is in concordance with the theory that the Korean population was formed by dual matrilineal origins via Northward and Southward routes (33,34). The Northern haplogroups account for 57 and 40% of the T2DM and normal groups, respectively, whereas the Southern haplogroups constituted 43 and 60%, respectively. In particular, haplogroups A5a and Y, which are mostly prevalent in Northeast Asia and Southeast Siberia, were found exclusively in the normal subjects (Table V). Southeast Asian-prevalent mtDNA lineages represented by haplogroups B, F3a and R were found almost exclusively in Korean subjects with T2DM (Table V). These findings suggested that a matrilineal lineage of Southern origin affects susceptibility to T2DM among Korean subjects with the -11,377 GG genotype of ADIPOQ.

In addition, haplogroups of the Korean subjects were investigated by principal component analysis (PCA) based on data retrieved from the literature (18-27), in which haplogroups of 33 Asian populations were investigated. In the PCA plot, Korean subjects with T2DM demonstrated a close correlation with South Asian populations, including China, Okinawa and Hong Kong individuals (Fig. 2), whereas normal Korean subjects were grouped together with Northern Asian populations, including Chukchi, Eskimo, Tuva, Yukaghir and Yakut. This result indicated that Koreans with an identical -11,377 GG *ADIPOQ* genotype would have a differential susceptibility to T2DM depending on matrilineal origin.

### Discussion

ADIPOQ is located on chromosome 3q27 and encodes adiponectin proteins. Adiponectin is specifically and abundantly

secreted by adipocytes and circulates in the blood at high concentrations (35). It is also known as 'adipocyte complement related protein 30' (*ACRP30*) and 'gelatin-binding protein 28' (*GBP28*) (36). It has previously been demonstrated that plasma concentrations of adiponectin are reduced in obese individuals, as compared with lean subjects (37), and increasing concentrations of adiponectin enhance insulin sensitivity by reducing the risk for T2DM (38,39). Therefore, *ADIPOQ* has been suggested as a causative gene for metabolic diseases, including T2DM.

The -11,377 C/G SNP of *ADIPOQ* is located in the promoter region of *ADIPOQ* and may have a functional role in the regulation of adiponectin expression (7,8). The promoter region of *ADIPOQ* has four SP1 binding sites that control *ADIPOQ* expression. The -11,377 C/G SNP is located in one of the SP1 binding sites on the *ADIPOQ* promoter. It alters the sequence for the SP1 binding site and thereby may inhibit promoter activity, resulting in a reduction of *ADIPOQ* expression (40,41).

Previously, numerous molecular epidemiological studies have been performed to identify the T2DM-specific ADIPOQ SNPs and their ethnic specificity. These investigations have identified T2DM-associated ADIPOQ SNPs, which are commonly found in several ethnic groups, including -11,377 C/G (rs266729), +45 T/G (rs2241766), +276 G/T (rs1501299) and +349 C/G (rs2241767). However, these SNPs, either independently or as a haplotype, do not always demonstrate a correlation with the occurrence of T2DM (42). In the present study, only one SNP, -11,377 C/G on the ADIPOQ gene, among the 14 SNPs identified from four genes, was closely associated with T2DM in Korean subjects. This result is consistent with previously reported results on Korean individuals with T2DM (43); however, these data are not concordant with those from studies on other Asian populations, including Japanese and Chinese (9). Rather, +45 T and +276 T alleles of ADIPOQ, which were proven not to be correlated with the incidence of Korean T2DM in the present study, were reported to demonstrate a strong correlation with T2DM in Japanese and Chinese individuals. That is, in several studies, the -11,377 G allele of ADIPOQ was not associated with T2DM in Japanese and Chinese subjects (28,44-47). Of note, in Caucasoid Europeans, including French and Swedish subjects, living geographically distant from East Asia and having genetic lineages that are different from those of mongoloids, including Japanese and Chinese populations who are expected to be physically and genetically similar to Koreans, the -11,377 G allele was found to be associated with T2DM, as is the case of the Korean subjects in the present study (6,7). Furthermore, 38 Korean subjects who were all expected to be T2DM patients since they all had the -11,377 G allele of ADIPOQ that inhibited promoter activity, were classified into two discrete groups: One for T2DM patients and the other for non-diabetic normal subjects. The results implied that the identical SNP on the ADIPOQ promoter may function differently in the occurrence of T2DM according to ethnic-specific genetic background, and ethnic-specific SNPs alone; therefore, it may be insufficient to predict the predisposition to T2DM in various ethnic groups, suggesting additional genetic markers are required.

Haplotype analysis of *ADIPOQ* has also been performed in several ethnic groups to determine susceptible haplotypes associated with T2DM (45,48,49). The T-G haplotype at the +45 and +276 sites of *ADIPOQ* is most frequently reported as a genetic marker associated with T2DM in various ethnic groups, but the correlation between the haplotype and the incidence of T2DM remains controversial (50,51). The present study identified that the T-G haplotype at positions +45 and +276 is not associated with the prevalence of Korean T2DM. By contrast, the present haplotype analysis, using various combinations of SNPs that were supposed to be associated with the incidence of T2DM, demonstrated that the G-T-T-A haplotype at SNP positions -11,377, +45, +276 and +349 is closely associated with T2DM in Korean subjects. These results strongly suggested that population ethnicity should be considered to identify the specific SNP or haplotype as a pre-diagnostic marker for T2DM prevalence.

The present study focused on matrilineal migration in the peopling of the Korean population as a candidate factor involved in the differential susceptibility of SNPs associated with T2DM. Koreans geographically belong to a Northeast Asian group, and anthropological and archeological evidence suggests that the early Korean population was related to Mongolian ethnic groups who inhabited the broad area of the Altai Mountains and Lake Baikal regions of Southeast Siberia (52). However, recent studies through hierarchical genotyping of Y-chromosomal DNA and mtDNA, which are reliable indicators of migration events in the peopling history of populations, revealed that the Korean population contains lineages typical of both Southern and Northern East Asian populations, although genetic components of Koreans are overall more similar to those of the Northeast Asians than Southeast Asians (32,33). In the present study, the mtDNA haplogroup composition of Korean T2DM and normal subjects also demonstrated dual lineages, including Southeast and Northeast Asian lineages. T2DM patients were mainly composed of M, B, R and F haplogroups of the Southeast Asian-prevalent mtDNA lineages, whereas the haplogroup composition of normal individuals was skewed toward Northeast Asian-prevalent mtDNA lineages consisting of A, D, G and Y haplogroups. These findings suggested that a matrilineal lineage with Southern origin increases susceptibility to T2DM in Korean subjects with an identical -11,377 GG ADIPOQ genotype. Furthermore, the PCA results supported the finding that the Southern matrilineal origin of the -11,377 GG genotype affects the incidence of T2DM in Korean individuals. That is, Korean subjects with T2DM lay entirely within the cluster of Southeast Asian populations, including China, Okinawa and Hong Kong, whereas populations clustered with T2DM negative Korean subjects were predominantly distributed over Northeast Asia. This result also indicated that Korean subjects with the identical -11,377 GG genotype would have different susceptibilities to T2DM according to their matrilineal origin.

In conclusion, the -11,377 C/G SNP of *ADIPOQ* appeared to be closely associated with T2DM in Korean subjects and the G-T-T-A haplotype in *ADIPOQ*, a combination of four SNPs at the -11,377, +45, +276 and +349 sites, may be regarded as a predictive factor for Korean T2DM. Additionally, the prevalence of T2DM in Korean subjects demonstrating identical allelotypes (-11,377 G allele of *ADIPOQ*) appeared to be affected by their matrilineal origin.

## Acknowledgements

The present study was supported by the National Research Foundation (NRF) grant funded by the Korea government(MEST) (grant nos. 2012R1A2A2A03047236 and KRF-D00518) and the Chung-Ang University Excellent Student Scholarship.

#### References

- 1. Permutt MA, Wasson JC, Suarez BK, *et al*: A genome scan for type 2 diabetes susceptibility loci in a genetically isolated population. Diabetes 50: 681-685, 2001.
- Mori Y, Otabe S, Dina C, *et al*: Genome-wide search for type 2 diabetes in Japanese affected sib-pairs confirms susceptibility genes on 3q, 15q, and 20q and identifies two new candidate Loci on 7p and 11p. Diabetes 51: 1247-1255, 2002.
- Salonen JT, Uimari P, Aalto JM, et al: Type 2 diabetes whole-genome association study in four populations: the DiaGen consortium. Am J Hum Genet 81: 338-345, 2007.
- 4. Mohammadzadeh G and Zarghami N: Associations between single-nucleotide polymorphisms of the adiponectin gene, serum adiponectin levels and increased risk of type 2 diabetes mellitus in Iranian obese individuals. Scand J Clin Lab Invest 69: 764-771, 2009.
- Li LL, Kang XL, Ran XJ, et al: Associations between 45T/G polymorphism of the adiponectin gene and plasma adiponectin levels with type 2 diabetes. Clin Exp Pharmacol Physiol 34: 1287-1290, 2007.
- 6. Gu HF, Abulaiti A, Ostenson CG, *et al*: Single nucleotide polymorphisms in the proximal promoter region of the adiponectin (APM1) gene are associated with type 2 diabetes in Swedish caucasians. Diabetes 53 (Suppl 1): S31-S35, 2004.
- Vasseur F, Helbecque N, Dina C, et al: Single-nucleotide polymorphism haplotypes in the both proximal promoter and exon 3 of the APM1 gene modulate adipocyte-secreted adiponectin hormone levels and contribute to the genetic risk for type 2 diabetes in French Caucasians. Hum Mol Genet 11: 2607-2614, 2002.
- Vasseur F, Helbecque N, Lobbens S, *et al*: Hypoadiponectinaemia and high risk of type 2 diabetes are associated with adiponectin-encoding (ACDC) gene promoter variants in morbid obesity: evidence for a role of ACDC in diabesity. Diabetologia 48: 892-899, 2005.
- Gong M, Long J, Liu Q and Deng HC: Association of the ADIPOQ rs17360539 and rs266729 polymorphisms with type 2 diabetes: a meta-analysis. Mol Cell Endocrinol 325: 78-83, 2010.
- Li Y, Li X, Shi L, *et al*: Association of adiponectin SNP+45 and SNP+276 with type 2 diabetes in Han Chinese populations: a meta-analysis of 26 case-control studies. PLoS One 6: e19686, 2011.
- World Health Organization: Definition and diagnosis of diabetes mellitus and intermediate hyperglycemia: Report of a WHO/IDF consultation. Geneva, World Health Org., 2006.
- 12. Wang H, Liu L, Zhao J, *et al*: Large scale meta-analyses of fasting plasma glucose raising variants in GCK, GCKR, MTNR1B and G6PC2 and their impacts on type 2 diabetes mellitus risk. PLoS One 8: e67665, 2013.
- 13. Cao H, van der Veer E, Ban MR, *et al*: Promoter polymorphism in PCK1 (phosphoenolpyruvate carboxykinase gene) associated with type 2 diabetes mellitus. J Clin Endocrinol Metab 89: 898-903, 2004.
- 14. Saif-Ali R, Harun R, Al-Jassabi S and Wan Ngah WZ: Hepatocyte nuclear factor 4 alpha P2 promoter variants associate with insulin resistance. Acta Biochim Pol 58: 179-186, 2011.
- 15. Winckler W, Graham RR, de Bakker PI, *et al*: Association testing of variants in the hepatocyte nuclear factor 4alpha gene with risk of type 2 diabetes in 7,883 people. Diabetes 54: 886-892, 2005.
- Andrews RM, Kubacka I, Chinnery PF, et al: Reanalysis and revision of the Cambridge reference sequence for human mitochondrial DNA. Nat Genet 23: 147, 1999.
- Lake SL, Lyon H, Tantisira K, et al: Estimation and tests of haplotype-environment interaction when linkage phase is ambiguous. Hum Hered 55: 56-65, 2003.

- Yao YG, Kong QP, Bandelt HJ, Kivisild T and Zhang YP: Phylogeographic differentiation of mitochondrial DNA in Han Chinese. Am J Hum Genet 70: 635-651, 2002.
- 19. Umetsu K, Tanaka M, Yuasa I, *et al*: Multiplex amplified product-length polymorphism analysis of 36 mitochondrial single-nucleotide polymorphisms for haplogrouping of East Asian populations. Electrophoresis 26: 91-98, 2005.
- 20. Pakendorf B, Novgorodov IN, Osakovskij VL, Danilova AP, Protod'jakonov AP and Stoneking M: Investigating the effects of prehistoric migrations in Siberia: genetic variation and the origins of Yakuts. Hum Genet 120: 334-353, 2006.
- Mooder KP, Schurr TG, Bamforth FJ, Bazaliiski VI and Savel'ev NA: Population affinities of Neolithic Siberians: a snapshot from prehistoric Lake Baikal. Am J Phys Anthropol 129: 349-361, 2006.
- 22. Derbeneva OA, Starikovskaya EB, Wallace DC and Sukernik RI: Traces of early Eurasians in the Mansi of northwest Siberia revealed by mitochondrial DNA analysis. Am J Hum Genet 70: 1009-1014, 2002.
- Kong QP, Yao YG, Liu M, *et al*: Mitochondrial DNA sequence polymorphisms of five ethnic populations from northern China. Hum Genet 113: 391-405, 2003.
- 24. Starikovskaya EB, Sukernik RI, Derbeneva OA, *et al*: Mitochondrial DNA diversity in indigenous populations of the southern extent of Siberia, and the origins of Native American haplogroups. Ann Hum Genet 69: 67-89, 2005.
- 25. Starikovskaya YB, Sukernik RI, Schurr TG, Kogelnik AM and Wallace DC: mtDNA diversity in Chukchi and Siberian Eskimos: implications for the genetic history of Ancient Beringia and the peopling of the New World. Am J Hum Genet 63: 1473-1491, 1998.
- 26. Fedorova SA, Stepanov AD, Adoian M, et al: Phylogenetic analysis of ancient mitochondrial DNA lineages of human remains found in Yakutia. Mol Biol (Mosk) 42: 445-453, 2008 (In Russian).
- 27. Derbeneva OA, Starikovskaia EB, Volod'ko NV, Wallace DC and Sukernik RI: Mitochondrial DNA variation in Kets and Nganasans and the early peoples of Northern Eurasia. Genetika 38: 1554-1560, 2002 (In Russian).
- Hara K, Boutin P, Mori Y, *et al*: Genetic variation in the gene encoding adiponectin is associated with an increased risk of type 2 diabetes in the Japanese population. Diabetes 51: 536-540, 2002.
- 29. Schwarz PE, Govindarajalu S, Towers W, *et al*: Haplotypes in the promoter region of the ADIPOQ gene are associated with increased diabetes risk in a German Caucasian population. Horm Metab Res 38: 447-451, 2006.
- Tanaka M, Cabrera VM, González AM, *et al*: Mitochondrial genome variation in eastern Asia and the peopling of Japan. Genome Res 14: 1832-1850, 2004.
- 31. Adachi N, Shinoda K, Umetsu K and Matsumura H: Mitochondrial DNA analysis of Jomon skeletons from the Funadomari site, Hokkaido, and its implication for the origins of Native American. Am J Phys Anthropol 138: 255-265, 2009.
- Am J Phys Anthropol 138: 255-265, 2009.
  32. Schurr TG and Wallace DC: Mitochondrial DNA diversity in Southeast Asian populations. Hum Biol 74: 431-452, 2002.
- 33. Jin HJ, Kwak KD, Hammer MF, et al: Y-chromosomal DNA haplogroups and their implications for the dual origins of the Koreans. Hum Genet 114: 27-35, 2003.
- 34. Jin HJ, Tyler-Smith C and Kim W: The peopling of Korea revealed by analyses of mitochondrial DNA and Y-chromosomal markers. PLoS One 4: e4210, 2009.
- 35. Gable DR, Hurel SJ and Humphries SE: Adiponectin and its gene variants as risk factors for insulin resistance, the metabolic syndrome and cardiovascular disease. Atherosclerosis 188: 231-244, 2006.

- Nakano Y, Tobe T, *et al*: Isolation and characterization of GBP28, a novel gelatin-binding protein purified from human plasma. J Biochem 120: 803-812, 1996.
- Oh DK, Ciaraldi T and Henry RR: Adiponectin in health and disease. Diabetes Obes Metab 9: 282-289, 2007.
- Ahima RS: Adipose tissue as an endocrine organ. Obesity (Silver Spring) 14 (Suppl 5) 242S-249S, 2006.
- Heidemann C, Sun Q, van Dam RM, *et al*: Total and high-molecular-weight adiponectin and resistin in relation to the risk for type 2 diabetes in women. Ann Intern Med 149: 307-316, 2008.
- 40. Zhang D, Ma J, Brismar K, Efendic S and Gu HF: A single nucleotide polymorphism alters the sequence of SP1 binding site in the adiponectin promoter region and is associated with diabetic nephropathy among type 1 diabetic patients in the Genetics of Kidneys in Diabetes Study. J Diabetes Complications 23: 265-272, 2009.
- 41. Barth N, Langmann T, Schölmerich J, Schmitz G and Schäffler A: Identification of regulatory elements in the human adipose most abundant gene transcript-1 (apM-1) promoter: role of SP1/SP3 and TNF-alpha as regulatory pathways. Diabetologia 45: 1425-1433, 2002.
- Menzaghi C, Trischitta V and Doria A: Genetic influences of adiponectin on insulin resistance, type 2 diabetes, and cardiovascular disease. Diabetes 56: 1198-1209, 2007.
- Lee YY, Lee NS, Cho YM, et al: Genetic association study of adiponectin polymorphisms with risk of Type 2 diabetes mellitus in Korean population. Diabet Med 22: 569-575, 2005.
- 44. Sun H, Gong ZC, Yin JY, *et al*: The association of adiponectin allele 45T/G and -11377C/G polymorphisms with Type 2 diabetes and rosiglitazone response in Chinese patients. Br J Clin Pharmacol 65: 917-926, 2008.
- 45. Wang Y, Zhang D, Liu Y, *et al*: Association study of the single nucleotide polymorphisms in adiponectin-associated genes with type 2 diabetes in Han Chinese. J Genet Genomics 36: 417-423, 2009.
- 46. Wang X, Zhang S, Chen Y, et al: APM1 gene variants -11377C/G and 4545G/C are associated respectively with obesity and with non-obesity in Chinese type 2 diabetes. Diabetes Res Clin Pract 84: 205-210, 2009.
- 47. Yang M, Qiu CC, Chen W, et al: Identification of a regulatory single nucleotide polymorphism in the adiponectin (APM1) gene associated with type 2 diabetes in Han nationality. Biomed Environ Sci 21: 454-459, 2008.
- 48. Heid IM, Wagner SA, Gohlke H, *et al*: Genetic architecture of the APM1 gene and its influence on adiponectin plasma levels and parameters of the metabolic syndrome in 1,727 healthy Caucasians. Diabetes 55: 375-384, 2006.
- 49. Woo JG, Dolan LM, Deka R, *et al*: Interactions between noncontiguous haplotypes in the adiponectin gene ACDC are associated with plasma adiponectin. Diabetes 55: 523-529, 2006.
  50. Jang Y, Lee JH, Kim OY, *et al*: The SNP276G>T polymorphism
- 50. Jang Y, Lee JH, Kim OY, et al: The SNP276G>T polymorphism in the adiponectin (ACDC) gene is more strongly associated with insulin resistance and cardiovascular disease risk than SNP45T>G in nonobese/nondiabetic Korean men independent of abdominal adiposity and circulating plasma adiponectin. Metabolism 55: 59-66, 2006.
- 51. Zacharova J, Chiasson JL and Laakso M; STOP-NIDDM Study Group: The common polymorphisms (single nucleotide polymorphism [SNP] +45 and SNP +276) of the adiponectin gene predict the conversion from impaired glucose tolerance to type 2 diabetes: the STOP-NIDDM trial. Diabetes 54: 893-899, 2005.
- 52. Kim K (ed): People and language. In Korea Its People and Cultures. Hakwon-sa Ltd., Seoul, pp. 10–20, 1970.