

# Identification of suitable reference genes for gene expression studies using quantitative polymerase chain reaction in lung cancer *in vitro*

HASSAN ALI<sup>1,2</sup>, ZHENWU DU<sup>1,3</sup>, XIUYING LI<sup>4</sup>, QIWEI YANG<sup>4</sup>,  
YU CHENG ZHANG<sup>4</sup>, MEI WU<sup>4</sup>, YI LI<sup>2</sup> and GUIZHEN ZHANG<sup>1,3</sup>

<sup>1</sup>Department of Central Laboratory, Second Hospital, Jilin University, Changchun, Jilin 130041;

<sup>2</sup>Department of Immunology, Norman Bethune College of Medicine, Jilin University, Changchun, Jilin 130021;

<sup>3</sup>Institute of Orthopedics, Second Hospital, Jilin University, Changchun, Jilin 130041;

<sup>4</sup>Department of Central Research, China-Japan Hospital, Jilin University, Changchun, Jilin 130033, P.R. China

Received January 15, 2014; Accepted June 5, 2014

DOI: 10.3892/mmr.2015.3159

**Abstract.** The present study aimed to examine 10 housekeeping genes (HKGs), including 18s ribosomal RNA (18S), glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*), ribosomal protein large P0 (*RPLP0*),  $\beta$ -actin (*ACTB*), peptidylprolyl isomerase A (*PPIA*), phosphoglycerate kinase-1 (*PGK1*),  $\beta$ -2-microglobulin (*B2M*), ribosomal protein L13a (*RPL13A*), hypoxanthine phosphoribosyl transferase-1 (*HPRT1*) and TATA box binding protein (*TBP*) in order to identify the most stable and suitable reference genes for use in expression studies in non-small cell lung cancer. The mRNA expression encoding the panel of the 10 HKGs was determined using reverse transcription-quantitative PCR (RT-qPCR) in human lung cancer cell lines. Three software programs, BestKeeper, NormFinder and geNorm, were used to ascertain the most suitable reference genes to normalize the RNA input. The present study examined three lung cancer cell lines (A549, NCI-H446 and NCI-H460). The analysis of the experimental data using BestKeeper software revealed that all 10 HKGs were stable, with *GADPH*, followed by 18S being the most stable genes and *PPIA* and *HPRT1* being the least stable genes. The NormFinder software results demonstrated that *PPIA* followed by *ACTB* were the most stable and *B2M* and *RPLP0* were the least stable. The geNorm software results revealed that *ACTB* and *PGK1*, followed by *PPIA* were the most stable genes and *B2M* and *RPLP0* were identified as the least stable genes. Due to discrepancies in the ranking orders

of the reference genes obtained by different analyzing software programs, it was not possible to determine a single universal reference gene. The suitability of selected reference genes requires unconditional validation prior to each study. Based on the three analyzing programs, *ACTB*, *PPIA* and *PGK1* were the most stable reference genes in lung cancer cell lines.

## Introduction

Lung cancer is the most commonly diagnosed malignancy and is the leading cause of mortality among all types of cancer. Every year, lung cancer contributes to more than one million mortalities worldwide, among which non-small cell lung cancer (NSCLC) accounts for 85% of cases (1,2). NSCLC can be divided into three types, including adenocarcinoma, large cell lung carcinoma and squamous cell carcinoma (3). Each of these share a common set of carcinoma characteristics. Cell lines derived from each of the main lung tumor types are widely used as experimental models in lung cancer biology (4). Reverse transcription-quantitative polymerase chain reaction (RT-qPCR) has revolutionized the field of gene expression analysis in living organisms (5). The main advantages of RT-qPCR are its superior specificity, sensitivity and broad quantification range (6,7). Despite being a useful technique, there are challenges coupled with its use, an important one being the normalization with an accurate and reliable reference gene, referred to as a housekeeping gene (HKG) (8,9). The term housekeeping gene was initially used to describe genes that are essential for cell function. Ideal HKGs are stably expressed in each cell type, do not respond to external stimuli and exhibit little or no run-to-run or sample-to-sample RT-qPCR variation. They are an internal reference to which target gene expression can be associated in order to correct unspecific variation caused by an imprecise amount of input RNA, RNA degradation or the presence of reaction inhibitors (8,10). Reference genes are often selected from the literature and are used across several experimental conditions, some of which may enhance the differences in the expression of a reference

---

*Correspondence to:* Professor Guizhen Zhang, Department of Central Laboratory, Second Hospital, Jilin University, 9 Ziqiang Street, Changchun, Jilin 130041, P.R. China  
E-mail: zhanguizhenjlu@163.com

*Key words:* reverse transcription-quantitative polymerase chain reaction, housekeeping genes, normalization, lung cancer

Table I. Candidate reference genes and their respective symbols and functions used in the present study.

Symbol	Name	Function
18S	18s ribosomal RNA	Component of the 40s ribosomal subunit
<i>GAPDH</i>	Glyceraldehyde-3-phosphate dehydrogenase	Oxidoreductase in glycolysis and gluconeogenesis
<i>RPLP0</i>	Ribosomal protein large P0	Component of the 60s ribosomal subunit
<i>ACTB</i>	$\beta$ -actin	Protein involved in various types of cell motility
<i>PPIA</i>	Peptidylprolyl isomerase A, cydophilin A, romatase A	Accelerates the folding of proteins, catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides
<i>PGK1</i>	Phosphoglycerate kinase-1	Enzyme involved in glycolysis
<i>B2M</i>	$\beta$ -2-microglobulin	Component of major histocompatibility complex class 1 molecules
<i>RPL13A</i>	Ribosomal protein LI3a	Structural component of the large 60S ribosomal subunit
<i>HPRT1</i>	Hypoxanthine phosphoribosyl transferase-1	Enzyme involved in purine synthesis in salvage pathway
<i>TBP</i>	TATA box binding protein	General RNA polymerase II transcription factor

gene under certain conditions. Previous studies have indicated that certain commonly used HKGs, including  $\beta$ -actin (*ACTB*) and glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) are differentially expressed in various tissues (11-13). The reliability of normalized data is reliant on the robustness of reference genes. If unrecognized, unexpected changes in the expression of reference genes could result in flawed conclusions of real biological effects. Therefore, identification of stable and reliable reference genes is a prerequisite to any reliable analysis of RT-qPCR data. Numerous reference genes, including *GADPH*, *ACTB*,  $\beta$ -2-microglobulin (*B2M*) and ribosomal protein large P0 (*RPLP0*) have been identified, and their suitability for gene expression studies in diverse human tissue and cell types has been validated (14-17). RT-qPCR has been used in lung cancer studies to enumerate the expression of predictive and or prognostic targets (18). In the present study, three types of lung cancer cell lines (NCI-H A549, NCI-H446 and NCI-H460) were assembled and 10 common HKGs, including 18S, *GAPDH*, *RPLP0*, *ACTB*, peptidylprolyl isomerase A (*PPIA*), phosphoglycerate kinase-1 (*PGK1*), *B2M*, ribosomal protein LI3a (*RPL13A*), hypoxanthine phosphoribosyl transferase-1 (*HPRT1*) and TATA box binding protein (*TBP*) (Table I) were selected in order to examine their stability and suitability for RT-qPCR normalization in NSCLC using three common statistical algorithms, NormFinder, geNorm and BestKeeper. Candidate HKGs were selected on the basis of two criteria: i) their previous use as a reference gene and ii) their ability to cover a wide expression spectrum.

## Materials and methods

**Cell lines.** Human lung cancer cell lines A549 and NCI-H446 were purchased from the American Type Culture Collection (Manassas, VA, USA) and NCI-H460 was provided by the Central Gene Therapy Department of China-Japan Union Hospital, Jilin University (Changchun, Jilin, China). Cells were cultured in RPMI-1640 medium (Gibco-BRL, Carlsbad, CA, USA), supplemented with 10% fetal bovine serum (Gibco-BRL) and 100 units of penicillin (Sigma-Aldrich,

St. Louis, MO, USA), and maintained at 37°C in a 5% CO<sub>2</sub> humidified atmosphere.

**RNA extraction.** The cell lines A549, NCI-H446 and NCI-H460 were cultured for 72 h, and total RNA was extracted from each cell using TRIzol reagent (Takara Bio, Inc., Shiga, Japan) according to the manufacturer's instructions. Briefly, 1 ml of TRIzol reagent was used to homogenize the cells (~2x10<sup>6</sup>). Samples were thoroughly mixed and incubated at room temperature for 5 min. The samples were then treated with 0.2 ml chloroform (Haodeng Industrial Co., Ltd, Shanghai, China) by reverse mixing. Phase separation was performed by placing the samples at room temperature for 5 min followed by centrifugation at 12,000 x g and 4°C for 15 min. The aqueous layer was mixed with 0.5 ml isopropanol (Haodeng Industrial Co., Ltd) to precipitate the RNA. Samples were placed at room temperature for 10 min and centrifuged at 12,000 x g and 4°C for 10 min. The RNA pellet was washed with 1 ml 75% alcohol and centrifuged at 10,000 x g and 4°C for 5 min. The pellet was air dried and resuspended with DNA/RNase free water. The purity and concentration of RNA was determined using NanoDrop 1000 (Thermo Scientific, Waltham, MA, USA) spectrophotometry.

**Complementary DNA (cDNA) synthesis.** Total RNA (1  $\mu$ g) from each cell group was reverse-transcribed to cDNA using a First Strand cDNA Synthesis kit (GeneCopoeia, Guangzhou, China) according to the manufacturer's instructions. The cDNA was stored at -20°C.

**Quantitative PCR.** For RT-qPCR analysis, SYBR Green Premix EX Taq (Takara Bio, Inc.) was used in a reaction mixture that comprised 5 pmol of each gene-specific primer and 40 ng of cDNA sample, in a final volume of 20  $\mu$ l. The primer sequences used (Table II) were synthesized by Sangon Biotech Co., Ltd. (Shanghai, China). RT-qPCR was performed using an AB Prism 7500 PCR detection system (Applied Biosystems, Foster City, CA, USA), under the following conditions: 30 sec of polymerase activation at 95°C followed

Table II. Primer sequences, product sizes and PCR efficiencies of candidate reference genes.

Symbol	Forward primer	Reverse primer	Product size (bp)	PCR efficiency
18S	GTGGAGCGATTTGTCTGGTT	AACGCCACTTGTCCCTCTAA	115	1.90
<i>GAPDH</i>	ATGGGGAAGGTGAAGGTCG	GGGGTCATTGATGGCAACAATA	108	1.99
<i>RPLP0</i>	CTGGAAGTCCAACACTTTCCT	CATCATGGTGTTCCTTGCCCAT	160	2.74
<i>ACTB</i>	GAAGATCAAGATCATTGCTCCT	TACTCCTGCTTGCTGATCCA	111	1.89
<i>PPIA</i>	TCCTGGCATCTTGTCCAT	TGCTGGTCTTGCCATTCT	179	2.10
<i>PGK1</i>	GCCACTTGCTGTGCCAAATG	CCCAGGAAGGACTTTACCTT	102	2.62
<i>B2M</i>	CTATCCAGCGTACTCCAAAG	GAAAGACCAGTCCTTGCTGA	188	2.08
<i>RPL13A</i>	CGAGGTTGGCTGGAAGTACC	CTTCTCGGCCTGTTTCCGTAG	121	2.00
<i>HPRT1</i>	CCTGGCGTCGTGATTAGTGAT	AGACGTTTCAGTCCTGTCCATAA	131	1.78
<i>TBP</i>	GCACAGGAGCCAAGAGTGA	GTTGGTGGGTGAGCACAAG	174	2.10

PCR, polymerase chain reaction; *GAPDH*, glyceraldehyde3-phosphate dehydrogenase; *RPLP0*, ribosomal protein large P0; *ACTB*,  $\beta$ -actin; *PPIA*, peptidylprolylisomerase A; *PGK1*, phosphoglycerate kinase-1; *B2M*,  $\beta$ -2 microglobulin; *RPL13A*, ribosomal protein L13a; *HPRT1*, hypoxanthine phosphoribosyl transferase-1; *TBP*, TATA box binding protein.

by 40 cycles of denaturation at 95°C for 5 sec, annealing at 58°C for 60 sec and elongation at 72°C for 30 sec. Each assay was performed three times. The RT-PCR products were then subjected to 1% agarose gel electrophoresis containing ethidium bromide.

**Statistical analysis.** Data analysis was performed using ABI 7500 SDS system software (version 1.4; Applied Biosystems). All biological replicates were used to calculate the average threshold cycle (Ct) values. The stability of the 10 candidate reference genes was comprehensively evaluated using NormFinder (version 0953; <http://moma.dk/normfinder-software>) (16) algorithms, geNorm (version 3.4; <http://medgen.ugent.be/~jvdesomp/genorm/>) (19) and BestKeeper (version 1; <http://www.gene-quantification.com/bestkeeper.html>) (20,21). In order to enter the Ct values into geNorm and NormFinder software, the (Ct) values were converted into relative quantities using the following formula:  $2^{-\Delta Ct}$  ( $\Delta Ct = Ct - \text{lowest Ct}$ ). The raw data was entered into the BestKeeper program and RT-qPCR efficiency was determined for each primer pair using slope analysis with a linear regression model. Relative standard curves for transcripts were performed with serial dilutions of cDNA at 1/2.5, 1/5, 1/10 and 1/20 ng. The corresponding RT-qPCR efficiencies (E) were calculated according to the following equation:  $E = 2^{-1/\text{slope}}$ .

## Results

**RNA purity and concentration.** All RNA samples were examined for their purity and concentration. The absorbance ratio at 260/280 nm was 1.85-1.95 for each RNA sample group, reflecting high purity and concentration.

**RT-qPCR efficiency of each primer pair.** The RT-qPCR efficiency of each primer pair was determined by serial dilution. The results demonstrated that the efficiencies of the HKGs of interest ranged between 1.78 (*HPRT1*) and 2.74 (*RPLP0*) for each primer pair (Table II).




Figure 1. Mean Ct values of the candidate reference genes among experimental samples. Ct, threshold cycle; *ACTB*,  $\beta$ -actin; *GAPDH*, glyceraldehyde-3-phosphate dehydrogenase; *RPL13A*, ribosomal protein L13a; *PPIA*, peptidylprolyl isomerase A; *RPLP0*, ribosomal protein large P0; *B2M*,  $\beta$ -2-microglobulin; *PGK1*, phosphoglycerate kinase-1; *HPRT1*, hypoxanthine phosphoribosyl transferase-1; *TBP*, TATA box binding protein.

### Candidate reference gene expression levels and ranges.

In general, the 10 candidate reference genes revealed wide expression levels with mean Ct values in a range that is usually covered by HKGs, varying between 17.5 (*ACTB*) and 25.5 (*TBP*) among the three groups of lung cancer cell lines. 18S revealed the lowest variability of Ct among all groups of the three cell lines (Fig. 1). The dissociation curve of each target reference gene demonstrated one single peak, which confirmed the specific amplification of the target reference gene (Fig. 2A). The gel electrophoresis results demonstrated one single band which further confirmed the specific RT-qPCR amplification of the target reference gene (Fig. 2B).

**Candidate HKG expression stability.** The expression stability of each of the 10 reference genes was analyzed using three commonly used software programs, geNorm, NormFinder and BestKeeper.

**geNorm analysis.** The geNorm software program is an Excel based program that calculates and compares the gene expression stability measure (M) of all candidate genes, and excludes genes with an M-value >1.5. The lower the M value the higher




Figure 2. Reverse transcription-quantitative polymerase chain reaction product of each of the 10 target reference genes. (A) Dissociation curves of each of the 10 target reference genes. (B) 1% agarose gel electrophoresis of amplified fragments. Bp, base pairs; *ACTB*,  $\beta$ -actin; *GADPH*, glyceraldehyde-3-phosphate dehydrogenase; *RPL13A*, ribosomal protein LI3a; *PPIA*, peptidylprolyl isomerase A; *RPLP0*, ribosomal protein large P0; *B2M*,  $\beta$ -2-microglobulin; *PGK1*, phosphoglycerate kinase-1; *HPRT1*, hypoxanthine phosphoribosyl transferase-1; *TBP*, TATA box binding protein.




Figure 3. Stability values of the candidate control genes analyzed by geNorm. (A) Determination of the optimal number of control genes. The software calculates the normalization factor from the least two stable genes at which the variable defines the pairwise variation between two sequential normalization factors. (B) M value of the 10 candidate reference genes analyzed by geNorm software. The x-axis from left to right indicates the ranking of the genes according to their stability; lower M values indicate higher stability. M, gene expression stability measure; V, variation; *ACTB*,  $\beta$ -actin; *GADPH*, glyceraldehyde-3-phosphate dehydrogenase; *RPL13A*, ribosomal protein LI3a; *PPIA*, peptidylprolyl isomerase A; *RPLP0*, ribosomal protein large P0; *B2M*,  $\beta$ -2-microglobulin; *PGK1*, phosphoglycerate kinase-1; *HPRT1*, hypoxanthine phosphoribosyl transferase-1; *TBP*, TATA box binding protein.

the gene expression stability and repeats the calculations until there are two genes remaining. The M value indicates the average pairwise variation of a gene compared with all the

Figure 4. Stability values of the candidate reference genes calculated by NormFinder software. The lower stability values indicate higher stability. *ACTB*,  $\beta$ -actin; *GADPH*, glyceraldehyde-3-phosphate dehydrogenase; *RPL13A*, ribosomal protein LI3a; *PPIA*, peptidylprolyl isomerase A; *RPLP0*, ribosomal protein large P0; *B2M*,  $\beta$ -2-microglobulin; *PGK1*, phosphoglycerate kinase-1; *HPRT1*, hypoxanthine phosphoribosyl transferase-1; *TBP*, TATA box binding protein.

other candidate genes. In order to determine the maximum number of genes necessary for adequate normalization in each panel of the experiment, geNorm determines pairwise variation ( $V$ )  $V_n/V_{n+1} = 0.15$  which is used as a cut-off value. A value  $<0.15$  indicates the number of control genes that is sufficient for valid normalization (Fig. 3A). The results demonstrated that the M-value for each of the 10 reference genes was  $<1.5$ , thus there was no exclusion. The most stable genes were *ACTB* and *PGK1*, followed by *PPIA*, while the gene with the least expression stability was *B2M* followed by *RPLP0* (Fig. 3A).

**NormFinder analysis.** The NormFinder software calculates the stability value based on an estimation of intra and intergroup variation for the analyzed genes. A low stability value has a low gene expression variance and indicates high stability expression.



Table III. Descriptive statistical analysis of candidate reference genes analyzed by BestKeeper software.

	18S	ACTB	GAPDH	RPL13A	PPIA	RPLP0	B2M	PGK1	HPRT1	TBP
N=12	12	12	12	12	12	12	12	12	12	12
Geo mean (CP)	7.29	16.65	17.18	17.22	17.31	17.58	17.96	19.81	21.60	24.56
Ar mean (CP)	7.32	16.65	17.19	17.23	17.31	17.60	18.00	19.82	21.61	24.57
Min (CP)	6.24	16.06	16.48	16.39	17.01	16.50	16.50	19.04	20.80	24.07
Max (CP)	7.98	17.14	18.14	18.24	17.70	18.63	19.75	20.29	22.34	25.73
SD ( $\pm$ CP)	0.55	0.35	0.41	0.38	0.21	0.69	0.99	0.38	0.53	0.52
CV (% CP)	7.49	2.09	2.38	2.23	1.22	3.94	5.53	1.94	2.44	2.13
Min (x-fold)	-2.07	-1.50	-1.62	-1.78	-1.23	-2.12	-2.74	-1.71	-1.74	-1.41
Max (x-fold)	1.60	1.41	1.95	2.03	1.31	2.07	3.47	1.39	1.67	2.25
SD ( $\pm$ x-fold)	1.46	1.27	1.33	1.3	1.16	1.62	1.99	1.30	1.44	1.44
Coeff. of corr. (r)	0.73	0.64	0.76	-0.51	0.10	0.14	0.72	0.62	-0.05	-0.49

<sup>a</sup>SD>1.00, was excluded from further analysis. CP, crossing point; SD, standard deviation; CV, coefficient of variance; ACTB,  $\beta$ -actin; GAPDH, glyceraldehyde3-phosphate dehydrogenase; RPL13A, ribosomal protein LI3a; PPIA, peptidylprolyl isomerase A; RPLP0, ribosomal protein large P0; B2M,  $\beta$ -2 microglobulin; PGK1, phosphoglycerate kinase-1; HPRT1, hypoxanthine phospho-ribosyl transferase-1; TBP, TATA box binding protein.




Figure 5. BestKeeper results for candidate reference genes. (A) SD was plotted on the y-axis. Genes with an SD >1.00 were excluded from further analysis. (B) Coefficient of correlation was plotted on the y-axis. A higher coefficient of correlation indicates a more stably expressed gene. SD, standard deviation; ACTB,  $\beta$ -actin; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; RPL13A, ribosomal protein LI3a; PPIA, peptidylprolyl isomerase A; RPLP0, ribosomal protein large P0; B2M,  $\beta$ -2-microglobulin; PGK1, phosphoglycerate kinase-1; HPRT1, hypoxanthine phosphoribosyl transferase-1; TBP, TATA box binding protein.

The output of this software analysis revealed that the most stable gene was PPIA followed by ACTB and PGK1. The least stable gene expression was B2M followed by RPLP0 (Fig. 4). These results were consistent with the geNorm analysis output.

**BestKeeper analysis.** BestKeeper is an excel based tool that assesses the stability of candidate HKGs based on

Table IV. Ranking of candidate control genes using BestKeeper, NormFinder and geNorm software programs.

geNorm	NormFinder	BestKeeper
ACTB, PGK1	PPIA	GAPDH
	ACTB	18S
PPIA	PGK1	B2M
HPRT1	GAPDH	ACTB
TBP	18S	PGK1
RPL13A	RPL13A	RPL13A
GAPDH	HPRT1	TBP
18S	TBP	RPLP0
RPLP0	RPLP0	PPIA
B2M	B2M	HPRT1

ACTB,  $\beta$ -actin; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; RPL13A, ribosomal protein LI3a; PPIA, peptidylprolyl isomerase A; RPLP0, ribosomal protein large P0; B2M,  $\beta$ -2-microglobulin; PGK1, phosphoglycerate kinase-1; HPRT1, hypoxanthine phosphoribosyl transferase-1; TBP, TATA box binding protein.

the inspection of calculated variation, including the standard deviation (SD) (22) and the coefficient of variance values (Table III). According to the BestKeeper program, the lowest variations revealed the highest stability. Genes with an SD >1 are considered to have an unacceptable range of variation (Fig. 5A). The analysis demonstrated that all 10 candidate HKGs had an SD  $\leq$ 1. GAPDH was the most stable, followed by 18S. HPRT1 was the least stable, followed by PPIA (Fig. 5B). The results from the BestKeeper software were therefore inconsistent with those of the geNorm and NormFinder software. A summary of the rankings produced by each of the three software programs is exhibited in Table IV.

## Discussion

Lung cancer is the most common type of cancer and the most common cause of cancer-related mortality worldwide (23). NSCLC is a highly fatal disease with a poor prognosis and low survival rate (24). To increase the survival rate of patients with NSCLC, the disease must be diagnosed as early as possible. Lung tumor cell lines have been widely dispersed to and used in experimental studies, including DNA sequencing (25), microRNA and microarray analyses (26,27) and detection of genome-wide methylated sequences (28,29). Previous RT-qPCR has been demonstrated to be useful for early NSCLC diagnosis, prognosis, prediction and gene expression analysis (30). The use of RT-qPCR technology to study gene expression levels requires reliable normalization of data to avoid unspecific variability caused by the differences in cDNA quantity and/or quality, incorrect interpretation of experimental results and mistaken analyses. Although diverse methods are employed to normalize RT-qPCR, it remains one of the main challenges in the efficacy of this technique (31). The identification of internal control gene(s) is therefore essential for accurate quantification of target mRNA by RT-qPCR in a given set of experimental samples (32). Statistical software, including NormFinder, BestKeeper and geNorm has been developed to identify the stability of reference genes in a given set of biological samples. Several studies have used these software programs in the assessment of diverse HKGs to ascertain their suitability as reference genes for normalization of qPCR data (17,33). The present study examined the RNA transcription levels of 10 common housekeeping genes, including 18S, *GAPDH*, *RPLP0*, *ACTB*, *PPIA*, *PGK1*, *B2M*, *RPL13A*, *HPRT1* and *TBP* (Table I) in the NSCLC cell lines NCI-H A549, NCI-H446 and NCI-H460. The three statistical softwares NormFinder, BestKeeper and geNorm (34) were used to assess the expression level stabilities of candidate reference genes. These programs use different calculation algorithms and therefore may provide different results (35,36). The present study demonstrated the following i) the purity and concentration of total RNA extracted from the abovementioned cell lines using TRIzol reagent; ii) the expression levels of the 10 reference genes determined in the above cell lines using qPCR and iii) the expression stability of the candidate reference genes in the above cell lines using geNorm, NormFinder and BestKeeper programs. In general, the present study demonstrated that almost all 10 candidate reference genes analyzed by the three independent programs could be used for future studies using lung cancer cell lines. This finding was somewhat in concordance with a previous study by Jacob *et al* (34). The analysis result of NormFinder was consistent with geNorm analysis output; both identified that *ACTB*, *PGK1* and *PPIA* were the most stable reference genes. By contrast, *B2M* and *RPLP0* were the least stable. BestKeeper analysis revealed that *GAPDH*, 18S and *B2M* were the most stable and *RPLP0*, *PPIA* and *HPRT1* were the least stable reference genes. This was consistent with previous studies demonstrating that *GADPH* and 18S were the most stable reference genes in NSCLC (31,37). By contrast, another previous study using lung tissue samples demonstrated that *GADPH* and *HPRT1* were the least stable reference genes (38). Variations obtained from these three programs were expected

given their distinct statistical algorithms. NormFinder and geNorm use relative quantities transformed from Ct values for stability calculation whereas BestKeeper uses Ct values directly, which may explain the different outputs from these three software programs (38). Several previous studies on reference gene selection for lung cancer also identified discrepancies between these programs (20,37) and there was no agreement regarding which was the best method. Few experimental studies have analyzed the stability of potential reference genes in lung cancer cell lines. To the best of our knowledge, no previous study has analyzed 10 reference genes in lung cancer cell lines using three different statistical software programs. The present study concluded that *ACTB*, *PPIA* and *PGK1* were the most stable reference genes analyzed by the three statistical programs geNorm, NormFinder and BestKeeper. These findings were somewhat inconsistent with those of previous studies and it was not possible to determine a single universal reference gene. Therefore, it is suggested that appropriate reference genes require selection on the basis of specific requirements and study conditions and in consideration of the characteristics of target genes in practical applications.

## Acknowledgements

This study was supported in part by grants from the Scientific Research Foundation of Jilin Province (nos. 20100942, 20110740 and 2013727038YY) and a grant from the Scientific Research Foundation of Jilin Province Development and Reform Commission (no. 2013c014-4).

## References

1. Koutsokera A, Kiagia M, Saif MW, Souliotis K and Syrigos KN: Nutrition habits, physical activity and lung cancer: an authoritative review. *Clin Lung Cancer* 14: 342-350, 2013.
2. Perroud MW, Jr, Honma HN, Barbeiro AS, *et al*: Mature autologous dendritic cell vaccines in advanced non-small cell lung cancer: a phase I pilot study. *J Exp Clin Cancer Res*: 30: 65, 2011.
3. Travis WD: Pathology of lung cancer. *Clin Chest Med* 32: 669-692, 2011.
4. Watanabe T, Miura T, Degawa Y, *et al*: Comparison of lung cancer cell lines representing four histopathological subtypes with gene expression profiling using quantitative real-time PCR. *Cancer Cell Int* 10: 2, 2010.
5. Guenin S, Mauriat M, Pelloux J, Van Wuytswinkel O, Bellini C and Gutierrez L: Normalization of qRT-PCR data: the necessity of adopting a systematic, experimental conditions-specific, validation of references. *J Exp Bot* 60: 487-493, 2009.
6. Dheda K, Huggett JF, Bustin SA, Johnson MA, Rook G and Zumla A: Validation of housekeeping genes for normalizing RNA expression in real-time PCR. *Biotechniques* 37: 112-114, 2004.
7. Hendriks-Balk MC, Michel MC and Alewijnse AE: Pitfalls in the normalization of real-time polymerase chain reaction data. *Basic Res Cardiol* 102: 195-197, 2007.
8. Eisenberg E and Levanon EY: Human housekeeping genes, revisited. *Trends Genet* 29: 569-574, 2013.
9. Arenas-Hernandez M and Vega-Sanchez R: Housekeeping gene expression stability in reproductive tissues after mitogen stimulation. *BMC Res Notes* 6: 285, 2013.
10. Huggett J, Dheda K, Bustin S and Zumla A: Real-time RT-PCR normalisation; strategies and considerations. *Genes Immun* 6: 279-284, 2005.
11. Barber RD, Harmer DW, Coleman RA and Clark BJ: GAPDH as a housekeeping gene: analysis of GAPDH mRNA expression in a panel of 72 human tissues. *Physiol Genomics* 21: 389-395, 2005.

12. Bas A, Forsberg G, Hammarstrom S and Hammarstrom ML: Utility of the housekeeping genes 18S rRNA, beta-actin and glyceraldehyde-3-phosphate-dehydrogenase for normalization in real-time quantitative reverse transcriptase-polymerase chain reaction analysis of gene expression in human T lymphocytes. *Scand J Immunol* 59: 566-573, 2004.
13. Selvey S, Thompson EW, Matthaei K, Lea RA, Irving MG and Griffiths LR: Beta-actin - an unsuitable internal control for RT-PCR. *Mol Cell Probes* 15: 307-311, 2001.
14. Andersen CL, Jensen JL and Orntoft TF: Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Can Res* 64: 5245-5250, 2004.
15. Rodriguez-Mulero S and Montanya E: Selection of a suitable internal control gene for expression studies in pancreatic islet grafts. *Transplantation* 80: 650-652, 2005.
16. Sorby LA, Andersen SN, Bukholm IR and Jacobsen MB: Evaluation of suitable reference genes for normalization of real-time reverse transcription PCR analysis in colon cancer. *J Exp Clinical Cancer Res* 29: 144, 2010.
17. Chua SL, See Too WC, Khoo BY and Few LL: UBC and YWHAZ as suitable reference genes for accurate normalisation of gene expression using MCF7, HCT116 and HepG2 cell lines. *Cytotechnology* 63: 645-654, 2011.
18. Kim B, Lee HJ, Choi HY, *et al*: Clinical validity of the lung cancer biomarkers identified by bioinformatics analysis of public expression data. *Cancer Res* 67: 7431-7438, 2007.
19. Schlotter YM, Veenhof EZ, Brinkhof B, *et al*: A GeNorm algorithm-based selection of reference genes for quantitative real-time PCR in skin biopsies of healthy dogs and dogs with atopic dermatitis. *Vet Immunol Immunopathol* 129: 115-118, 2009.
20. Wang Q, Ishikawa T, Michiue T, Zhu BL, Guan DW and Maeda H: Stability of endogenous reference genes in post-mortem human brains for normalization of quantitative real-time PCR data: comprehensive evaluation using geNorm, NormFinder and BestKeeper. *Int J Legal Med* 126: 943-952, 2012.
21. Pfaffl MW, Tichopad A, Prgomet C and Neuvians TP: Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper - Excel-based tool using pair-wise correlations. *Biotechnol Lett* 26: 509-515, 2004.
22. de Oliveira LA, Breton MC, Bastolla FM, *et al*: Reference genes for the normalization of gene expression in eucalyptus species. *Plant Cell Physiol* 53: 405-422, 2012.
23. Bunn PA, Jr: Worldwide overview of the current status of lung cancer diagnosis and treatment. *Arch Pathology Lab Med* 136: 1478-1481, 2012.
24. Tsuboi M, Ohira T, Saji H, *et al*: The present status of postoperative adjuvant chemotherapy for completely resected non-small cell lung cancer. *Ann Thoracic Cardiovasc Surg* 13: 73-77, 2007.
25. Liu J, Lee W, Jiang Z, *et al*: Genome and transcriptome sequencing of lung cancers reveal diverse mutational and splicing events. *Genome Res* 22: 2315-2327, 2012.
26. Sokilde R, Kaczowski B, Podolska A, *et al*: Global microRNA analysis of the NCI-60 cancer cell panel. *Mol Cancer The* 10: 375-384, 2011.
27. Che CL, Zhang YM, Zhang HH, *et al*: DNA microarray reveals different pathways responding to paclitaxel and docetaxel in non-small cell lung cancer cell line. *Int J Clin Exp Path* 6: 1538-1548, 2013.
28. Gazdar AF, Girard L, Lockwood WW, Lam WL and Minna JD: Lung cancer cell lines as tools for biomedical discovery and research. *J Natl Cancer Inst* 102: 1310-1321, 2010.
29. Gazdar AF, Gao B and Minna JD: Lung cancer cell lines: Useless artifacts or invaluable tools for medical science? *Lung Cancer* 68: 309-318, 2010.
30. van Eijk R, Licht J, Schrupf M, *et al*: Rapid KRAS, EGFR, BRAF and PIK3CA mutation analysis of fine needle aspirates from non-small-cell lung cancer using allele-specific qPCR. *PloS One* 6: e17791, 2011.
31. Saviozzi S, Cordero F, Lo Iacono M, Novello S, Scagliotti GV and Calogero RA: Selection of suitable reference genes for accurate normalization of gene expression profile studies in non-small cell lung cancer. *BMC Cancer* 6: 200, 2006.
32. Ho-Pun-Cheung A, Bascoul-Mollevi C, Assenat E, *et al*: Validation of an appropriate reference gene for normalization of reverse transcription-quantitative polymerase chain reaction data from rectal cancer biopsies. *Anal Biochem* 388: 348-350, 2009.
33. Ohl F, Jung M, Xu C, *et al*: Gene expression studies in prostate cancer tissue: which reference gene should be selected for normalization? *J Mol Medicine* 83: 1014-1024, 2005.
34. Jacob F, Guertler R, Naim S, *et al*: Careful selection of reference genes is required for reliable performance of RT-qPCR in human normal and cancer cell lines. *PloS One* 8: e59180, 2013.
35. Mafra V, Kubo KS, Alves-Ferreira M, *et al*: Reference genes for accurate transcript normalization in citrus genotypes under different experimental conditions. *PloS One* 7: e31263, 2012.
36. Mallona I, Lischewski S, Weiss J, Hause B and Egea-Cortines M: Validation of reference genes for quantitative real-time PCR during leaf and flower development in *Petunia hybrida*. *BMC Plant Biol* 10: 4, 2010.
37. Liu DW, Chen ST and Liu HP: Choice of endogenous control for gene expression in nonsmall cell lung cancer. *Eur Respir J* 26: 1002-1008, 2005.
38. Gresner P, Gromadzinska J and Wasowicz W: Reference genes for gene expression studies on non-small cell lung cancer. *Acta Biochim Pol* 56: 307-316, 2009.