

β -adrenergic stimulation activates protein kinase C ϵ and induces extracellular signal-regulated kinase phosphorylation and cardiomyocyte hypertrophy

LIN LI^{1*}, HONGYAN CAI^{1*}, HUA LIU² and TAO GUO¹

Departments of ¹Cardiology and ²Clinical Laboratory, First Affiliated Hospital of Kunming Medical University, Kunming, Yunnan 650032, P.R. China

Received March 24, 2014; Accepted January 7, 2015

DOI: 10.3892/mmr.2015.3316

Abstract. The cardiac adrenergic signaling pathway is important in the induction of cardiac hypertrophy. The cardiac adrenergic pathway involves two main branches, phospholipase C (PLC)/protein kinase C (PKC) and the adenylyl cyclase (cAMPase)/protein kinase A (PKA) signaling pathways. It is hypothesized that PLC/PKC and cAMPase/PKA are activated by the α -adrenergic receptor (α AR) and the β -adrenergic receptor (β AR), respectively. Previous studies have demonstrated that exchange protein directly activated by cAMP (Epac), a guanine exchange factor, activates phospholipase C ϵ . It is possible that there are β AR-activated PKC pathways mediated by Epac and PLC. In the present study, the role of Epac and PLC in β AR activated PKC pathways in cardiomyocytes was investigated. It was found that PKC ϵ activation and translocation were induced by the β AR agonist, isoproterenol (Iso). Epac agonist 8-CPT-2'OMe-cAMP also enhanced PKC ϵ activation. β AR stimulation activated PKC ϵ in the cardiomyocytes and was regulated by Epac. Iso-induced change in PKC ϵ was not affected in the cardiomyocytes, which were infected with adenovirus coding rabbit muscle cAMP-dependent protein kinase inhibitor. However, Iso-induced PKC ϵ activation was inhibited by the PLC inhibitor, U73122. The results suggested that Iso-induced PKC ϵ activation was independent of PKA, but was regulated by PLC. To further investigate the downstream signal target of PKC ϵ activation, the expression of phosphorylated extracellular signal-regulated kinase (pERK)1/2 and the levels of ERK phosphorylation was

analyzed. The results revealed that Iso-induced PKC ϵ activation led to an increase in the expression of pERK1/2. ERK phosphorylation was inhibited by the PKC ϵ inhibitor peptide. Taken together, these data demonstrated that the β AR is able to activate PKC ϵ dependent on Epac and PLC, but independent of PKA.

Introduction

Myocardial hypertrophy, which is a reaction of cardiomyocytes to a variety of pathological stimuli, is a common complication of hypertension, coronary heart disease, valvular heart disease, congenital heart disease, and other cardiovascular diseases. However, when stressors persist, the compensatory hypertrophy can evolve into a decompensated state, with profound changes in the gene expression program, contractile dysfunction and extracellular remodeling (1,2). Pathological hypertrophy increases myocardial oxygen consumption and reduces myocardial compliance, ultimately leading to heart failure, arrhythmia and sudden death (3-5). Cardiac hypertrophy is an independent risk factor of cardiovascular morbidity and mortality (3,4).

The β -adrenergic receptor (β AR) and its signal transduction pathway are important factors leading to cardiac hypertrophy (6,7). Although the acute stimulation of β AR may have a beneficial effect on cardiac function, evidence suggests that the long-term activation of human cardiac β AR may lead to heart dysfunction, apoptosis and cardiac remodeling (8,9). Studies have revealed that the chronic stimulation of β AR may lead to cardiomyocyte hypertrophy (10) and inappropriate cardiac hypertrophy may develop into heart failure (11). The clarification of the signal transduction mechanism of cardiomyocyte hypertrophy by β AR stimulation may contribute to improving the prevention and treatment of cardiac hypertrophy and heart failure.

Since the adrenergic receptor (AR) was categorized into α AR and β AR, it has been suggested that the protein kinase (PK)C signal transduction pathway is mediated by α AR, and that PKA is a β AR-exclusive downstream effector. A new family of proteins, designated exchange protein directly activated by cAMP (Epac), has been reported, which makes the cAMP-mediated signaling mechanism

Correspondence to: Professor Tao Guo, Department of Cardiology, First Affiliated Hospital of Kunming Medical University, 295 Xichang Road, Kunming, Yunnan 650032, P.R. China
E-mail: guotao20@hotmail.com

*Contributed equally

Key words: β -adrenergic receptor, protein kinase C ϵ , cardiomyocyte hypertrophy, phospholipase C, extracellular signal-regulated kinase

more complex. These proteins have been identified as the nucleotide exchange factors for small GTPases from the Rap family (12,13). A previous study by Schmidt *et al* (14) demonstrated that the activated adenylate cyclase (AC)-coupled β 2AR in HEK-293 cells can stimulate a novel phospholipase C (PLC) ϵ subtype of PLC independently of PKA, but is mediated by Epac, suggesting that Epac-mediated signal transduction pathways may exist between β AR and PKC (15). An investigation into the inflammatory pain mechanism for nociceptive neurons by Hucho *et al* (16) revealed that β AR stimulation in neurons may lead to PKC ϵ activation, and that Epac mediates signal transduction between cAMP and PKC ϵ . Studies by Oestreich *et al* (17,18) demonstrated that Epac and PLC ϵ can mediate the β AR regulation of calcium ion release in cardiomyocytes. These studies suggest that there is a possible association between PKC and β AR, with the exception of α AR signaling. It is possible that Epac and PLC mediate the interconnection between the β -adrenergic and PKC pathways.

PKC ϵ , one of the major PKC isozymes expressed in the heart, is important in cardiac cell signal transduction and function, including involvement in ischemic preconditioning, cardiac hypertrophy, myocardial fibrosis, heart failure and other signal transduction pathways (19,20). Upon activation, PKC ϵ undergoes translocation from the cytoplasm to the cell membrane (particulate fractions) to phosphorylate its targets, including extracellular signal-regulated kinases (ERK) (19-21). ERK1/2 are considered the downstream kinases of PKC and are implicated in a wide range of cellular processes, including cell growth, proliferation to apoptosis (22,23). The ERK pathway is involved in cardiac development and hypertrophy (24). The β AR may activate ERK and ERK may also be activated in a PKC-dependent way (7,22,24). It remains to be elucidated whether β AR activates PKC ϵ , mediated by Epac and PLC, to induce cardiac hypertrophy and the phosphorylation of ERK1/2 in cardiomyocytes.

The aim of the present study was to investigate PKC ϵ translocation by stimulation with the β AR agonist, isoproterenol (Iso), in isolated cardiomyocytes, to clarify the role of Epac and PLC in the cross-talk between β AR and PKC ϵ , and to investigate the effect of this signaling pathway on the phosphorylation of ERK and cardiomyocyte hypertrophy.

Materials and methods

Materials. Iso, phorbol 12-myristate 13-acetate (PMA), and PLC inhibitor (U73122) were obtained from Sigma-Aldrich (St. Louis, MI, USA). Rabbit polyclonal anti-PKC ϵ (1:2,000; sc-214), rabbit polyclonal ERK2 antibody (1:1,000; sc-154), mouse monoclonal anti-pERK 1/2 (1:2,000; sc-81492), mouse monoclonal β -actin antibody (1:5,000; sc-47778), goat anti-rabbit horseradish peroxidase (HRP)-conjugated immunoglobulin G (IgG) antibody [1:4,000(ERK2)-1:6,000(PKC ϵ); sc-2004], goat anti-mouse IgG-HRP [1:5,000(p-ERK 1/2)- 1:7,500(β -actin); sc-2005] and the PKC ϵ translocation inhibitor peptide (sc-3095) were obtained from Santa Cruz Biotechnology Inc. (Santa Cruz, CA, USA). The Epac agonist 8-CPT-2'OMe-cAMP (8-CPT) was obtained from Biolog Life Science Institute (Bremen, Germany). The green fluorescent protein (GFP) and human mutant Epac R279 K dominant negative (DN) in GFP structure were obtained from Genethon Center (Evry, France),

and the adenovirus coding for rabbit muscle cAMP-dependent protein kinase inhibitor (Ad.PKI) was obtained from Rush University Medical Center (Chicago, IL, USA). The electrochemiluminescence (ECL) kit was obtained from GE Healthcare (Amersham, UK).

Primary culture of cardiomyocytes. Cardiomyocyte culture was performed, as previously described, with a minor modification (25). Briefly, 20 pregnant female rats (Shanghai SLAC Laboratory Animal Co. Ltd., Shanghai, China) were maintained in a controlled environment (22-24°C; 12 h light/12 h dark cycle) with *ad libitum* access to food and water. Subsequent to sacrifice by decapitation, the hearts were removed from the new-born Wistar rats (1-3 days old) and placed into pre-cooled 1X Ads buffer solution (NaCl, 17 g; Hepes, 11.9 g; NaH₂PO₄, 0.3 g; KCl, 1 g; glucose, 2.5 g; MgSO₄, 0.25 g; in 250 ml distilled water; Sigma-Aldrich), the ventricular muscle was cut into 1 mm³ sections and a solution of pancreatin and collagenase (0.05%; Sigma-Aldrich) was added. Subsequently, 20 min digestion in pancreatin and collagenase at 37°C and a centrifugation (500x g, 6 min) for supernatant removal was performed, 2 ml of newborn calf serum (HyClone Laboratories, Inc., Logan, UT, USA) was added and mixed evenly for subsequent use. The digested products were collected for all cells and subjected to a 6 min centrifugation at 500 x g. The ventricular myocytes were then purified with discontinuous Percoll density gradient centrifugation (2,000 x g, 30 min). The ventricular muscle cell layers were collected, washed with Ads two times (6 min centrifugation at 500 x g), the supernatant and culture medium [4:1 Dulbecco's modified Eagle's medium/M199 (Gibco Life Technologies, Carlsbad, CA, USA) containing 10% horse serum (HyClone Laboratories, Inc.), 5% newborn calf serum and 1% penicillin/streptomycin (Sigma-Aldrich)] was added. This was then mixed evenly, inoculated in a petri dish (5x10⁴ cells/cm²), and cultivated at 37°C under conditions of 5% CO₂. After 48 h cultivation, the medium was replaced with serum-free medium and the subsequent experiments were carried out after 24 h. The study was approved by the Ethics Committee of Kunming Medical University (Kunming, China) and conformed to the standards set by the Yunnan Experimental Animal Management Board.

Cell treatment. The β AR agonist, Iso (1 μ mol/l, 1 min), Epac agonist, 8-CPT (1 μ mol/l, 10 min), PLC inhibitor, U73122 (2 μ mol/l, 30 min), and PKC agonist, PMA (1 μ mol/l, 5 min), were provided for cell treatment. Following cell infection (5x10²/cm²) by green fluorescent protein (GFP), Epac R279 K and Ad.PKI adenovirus, and the cell transfection by specific PKC ϵ translocation inhibition peptide and scramble peptide, Iso (1 μ mol/l, 1 or 10 min) was then provided for cell treatment and PKC ϵ and pERK1/2 protein expression were detected.

Cardiomyocyte infection. The role of Epac was analyzed using Epac R279 K (dominant negative) to downregulate or inhibit Epac. The PKA-dependent activation of PKC was analyzed by overexpression of a specific PKA inhibitor peptide using Ad.PKI. Following infection with adenovirus coding for GFP, Epac R279 K with a multiplicity of infection (MOI) of 100 and Ad.PKI with an MOI of 100, the cells were treated with Iso (1 μ mol/l) for 1 min.

Peptide transfection. It is hypothesized that the mechanism of activation of PKC involves translocation from a cytosolic fraction to a membrane-rich fraction. The translocation occurs when PKC in the cytoplasm binds to an isozyme-specific membrane-bound anchor protein or Receptor for Activated C Kinase (RACK). The RACK then transports the activated PKC isozyme to its target protein. Recently, small peptides of six to eight amino acids have been used to inhibit specific isozymes of PKC from binding to the specific RACK (26,27). Therefore, the specific PKC ϵ inhibitor peptide was used in the present study to investigate the role of PKC ϵ in ERK phosphorylation and cardiomyocyte hypertrophy induced by Iso stimulation (1 μ mol/l for 10 min and 10 μ mol/l for 48 h). The primary cardiomyocytes (5x10²/cm²) were exposed at room temperature to sterile phosphate buffered saline (PBS; HyClone Laboratories, Inc.) for 2 min and then exposed at room temperature to permeabilization buffer (20 mmol/l HEPES pH 7.4, 10 mmol/l EGTA, 140 mmol/l KCl, 50 μ l/ml saponin, 5 mmol/l NaN₃ and 5 mmol/l oxalic acid dipotassium salt) containing either the PKC ϵ translocation inhibitor peptide (22 μ g/ml) or a control of scramble peptide (22 μ g/ml) containing the same amino acids as the inhibitor peptide, but in a different sequence. This was removed after 2 min and the cells were washed twice with PBS at room temperature. The cells were then exposed to sterile PBS for 2 min at 37°C prior to the addition of a fresh culture medium at 37°C. The cells were allowed to recover for at least 1 h prior to the initiation of experiments, during which the cells began beating normally.

Immunocytochemical staining of cardiomyocytes. The cardiomyocytes (5x10²/cm²) were inoculated onto a Lab-Tec Chamber Slide system (Nunc®, Sigma-Aldrich) coated with 0.2% gelatine (Sigma-Aldrich) for culture and fixed with 4% paraformaldehyde (Sigma-Aldrich) (5 min, room temperature). They were then washed with NH₄Cl/PBS (Sigma-Aldrich) (pH 7.4, 0.5 mol/l) twice for 5 min each at room temperature and permeabilization treatment was performed using 1% Triton X-100/PBS (5 min, room temperature). Following washing with PBS and blocking of unspecific sites in 5% bovine serum albumin (BSA; Roche Diagnostics, Basel, Switzerland)/PBS, cells were incubated with a rabbit anti-PKC ϵ (1:200) primary antibody in 1% BSA/PBS for 1 h at room temperature. After 1 h incubation with Alexa-488 conjugated goat anti-rabbit IgG antibody (Invitrogen Life Technologies, Carlsbad, CA, USA) at room temperature, the cells were washed with PBS, fixed with Vectashield mounting agent (Vector Laboratories, Burlingame, CA, USA), covered with a slide and sealed with nail polish. A confocal laser scanning microscope (Zeiss LSM510; Carl Zeiss Lase Optics GmbH, Oberkochen, Germany) was used to detect the translocation of PKC ϵ .

Detection of the expression of PKC ϵ and pERK1/2 by western blot analysis. Cell lysis buffer (Tris 12.5 mmol/l, pH 7.4, EDTA 1 mmol/l, EGTA 2.5 mmol/l, NaF 100 mmol/l) containing protease inhibitors was used to dissolve the cells and lysates were obtained following 15 min (4°C) of centrifugation at 14,000 x g. The supernatant consisted of the cytosolic fractions. The centrifuged precipitate under a

second suspension of lysis buffer containing 1% Triton X-100 consisted of the particulate fractions. The obtained cell particulate fractions were subject to protein quantification using Bradford's method and western blot analysis. Following cell lysis with radioimmunoprecipitation assay lysis buffer, containing Tris-HCl 25 mmol/l, pH 7.5, NaCl 150 mmol/l, 1% Triton X-100, 1% sodium deoxycholate, 0.1% SDS and protease inhibitor cocktail (Roche Diagnostics), the lysates were collected, subject to a 10 min centrifugation (4°C) at 10,000 x g, and the supernatant was collected for pERK1/2 protein expression detection by western blot analysis.

Western blot analysis. A total of 20 μ g protein was used for 10% SDS-PAGE and then transferred onto a polyvinylidene difluoride membrane following electrophoresis at 35 V overnight. Non-specific sites were blocked (room temperature, 1 h) with Tween-tris-buffered saline (TTBS 0.1%) containing 5% nonfat dried milk, and PKC ϵ antibody (1:2,000) and pERK1/2 (1:2,000) added to incubate at room temperature for 90 min. Following washing the membrane with TTBS, the membrane was incubated with horseradish peroxidase-conjugated IgG secondary antibody at room temperature for 1 h. All antibodies were visualized with chemiluminescence detection (ECL). β -actin (1:5,000) and ERK2 antibody (1:1,000) were used as the internal reference. The image signals were quantified by densitometric analysis of digitized films using the Scion Image (4.0.3) analysis system (National Institutes of Health, Bethesda, MD, USA).

Myocardial cell protein content and surface area determination after 48 h treatment with Iso. Following transfection with the PKC ϵ inhibitor and scramble peptides, the cells (5x10²/cm²) were incubated with Iso (10 μ mol/l) for 48 h. The protein content, DNA concentration and surface areas of the cells were then measured. Each dish was rinsed three times with PBS. The cell layer was scraped with standard sodium citrate containing 0.25% (w/v) SDS and frozen at -20°C. Prior to use, the extracts were thawed and vortexed (Vortex-Genie 1; Scientific Industries, Inc., Bohemia, NY, USA). Total cell protein was assayed using Bradford's method. The results were normalized against the DNA content, which was measured fluorometrically using Hoechst staining, using calf thymus DNA as a standard. Hoechst 33258 and calf thymus DNA were obtained from Sigma-Aldrich. Hoechst 33258 dye was diluted to 1.5x10⁻⁴ mol/l in distilled water, and diluted five times with 20X saline-sodium citrate solution (SSC; 175.3 g NaCl and 88.2 g sodium citrate in 1 liter distilled water; Sigma-Aldrich) before testing. Calf thymus DNA was dissolved in SSC (1g/l). The cell extracts were mixed with Hoechst 33258 dye, and placed in the dark for 10 min. Absorbance was detected at 350 nm excitation wave and 450 nm emitting wave using a Fluorospectrophotometer 850 (Hitachi Ltd., Tokyo, Japan). The cell surface area measurements involved the use of microscope images (LV100POL; Nikon Corporation, Tokyo, Japan) (magnification, x10) captured with a digital camera (E5200; Nikon Corporation) and the cell perimeter was measured with an image analysis system (Scion Image). A total of five fields in each group were randomly selected, with 20 cells per field, to measure the cell surface area and to calculate the mean values (n=5).




Figure 1. Time-dependent translocation of PKC ϵ under Iso stimulation. The protein expression levels of PKC ϵ of cellular particulate fractions in cardiomyocytes treated with Iso for different durations were detected using western blot analysis and β -actin was used as an internal reference (n=5). Data are presented as the means \pm standard error of the mean of five independent experiments. *P<0.05, compared with the control. Iso, isoproterenol; PKC, protein kinase C.

Statistical analysis. The results are expressed as the mean \pm standard error of the mean. Statistical analysis was performed using Student's t-test for single comparisons. Comparisons among three or more groups were made by one-way analysis of variance followed by the Student-Newman-Keuls test. P<0.05 was considered to indicate a statistically significant difference.

Results

β AR stimulation activates PKC ϵ in cardiomyocytes. PKC ϵ activation is associated with its translocation from the cytoplasm to the cellular particulate fractions (28). For evaluating PKC ϵ translocation, cardiomyocytes were incubated with Iso (1 μ mol/l) for different durations (1-30 min). Following cell scraping, differential centrifugation was performed in order to separate the particulate and cytosolic fractions. As shown in Fig. 1, the PKC ϵ content in the particulate fractions was significantly increased following 1 and 5 min incubation with Iso (P<0.05). Compared with the control, this increase persisted for 15 min and decreased to the control level at 30 min. The precise sub-cellular PKC ϵ localization was examined by immunostaining under confocal microscopy. As shown in Fig. 2, PKC ϵ was translocated to the perinuclear area as early as 1 min after Iso incubation. Similarly, the western blot analysis revealed that this pattern persisted during the 15 min of Iso incubation, and homogeneous PKC ϵ localization emerged after 30 min incubation. These results suggested that the Iso-induced translocation of PKC ϵ was transient, with a translocation peak at 1 min. Therefore, this duration was used in the subsequent experiments to evaluate PKC ϵ translocation.

PKC ϵ activation by Epac agonist 8-CPT. Similar to that of Iso pre-incubation, following incubation of the cells with the Epac agonist (1 μ mol/l 8-CPT for 10 min), the PKC ϵ content in the cellular particulate fractions was significantly higher compared with those of the control (Fig. 3). The PKC ϵ content of the particulate fractions increased by 168 \pm 23% compared with the control (P<0.05).

PKA-independent PKC ϵ activation by Iso. Epac is activated by cAMP independently of PKA (29), therefore the present

study subsequently examined whether the PKC ϵ translocation produced by β AR stimulation was independent of PKA. Following infection with Ad.PKI, which specifically inactivates PKA activity, no inhibition of Iso-induced translocation of PKC ϵ was observed (Fig. 4).

Inhibition of Iso-induced PKC ϵ activation by Epac inhibitor Epac R279K. To further confirm whether Epac was involved in the Iso-induced PKC ϵ activation, a dominant negative (DN) form of Epac (Epac R279K) was used to downregulate or inhibit the expression of Epac. The results revealed that DN Epac inhibited Iso-induced PKC ϵ translocation (Fig. 5).

PLC-dependent PKC ϵ activation by Iso. To assess the involvement of PLC, the cardiomyocytes were pre-incubated with the PLC inhibitor U73122 prior to stimulation with Iso. The translocation of PKC ϵ was inhibited by this inhibitor (Fig. 6).

Iso-induced PKC ϵ activation increases the expression of pERK1/2. Following introduction of the PKC ϵ control scramble peptide into the cardiomyocytes and exposure of the cells to Iso (1 μ mol/l, 10 min), there was a significant increase of pERK1/2 (P<0.05). However, when the PKC ϵ inhibitor peptide was introduced into the cells prior to Iso exposure, no differences were identified in the expression of pERK1/2 compared with the PKC ϵ inhibitor peptide alone (Fig. 7).

Iso-induced PKC ϵ activation leads to hypertrophy in cardiomyocytes. The cells, which were exposed to the control scramble peptide and followed by Iso (10 μ mol/l, 48 h) revealed significant increases in the cell protein/DNA ratio and in the cell surface area compared with the cells, which were exposed to the scramble peptide alone. However, following transfection of the cells with the PKC ϵ inhibitor peptide and exposure to Iso, no significant differences were identified in the cell protein/DNA ratio or in the cell surface area compared with the PKC ϵ inhibitor peptide alone (Figs. 8 and 9). These findings suggested that Iso-induced cardiomyocyte hypertrophy was mediated by PKC ϵ and, following inhibition of PKC ϵ activation by the PKC ϵ -specific




Figure 2. Confocal microscope images of Iso stimulation-induced time-dependent PKC ϵ activation (magnification, $\times 1,000$; confocal immunofluorescence staining). PKC ϵ was distributed evenly in the cytoplasm without Iso treatment. PKC ϵ underwent translocation following incubation with Iso, with translocation around the nucleus. Translocation was most marked between 1 and 15 min, and was distributed in the cytoplasm at 30 min ($n=3$). Iso, isoproterenol; PKC, protein kinase C.

translocation inhibitor peptide, this hypertrophic effect was eliminated.

Discussion

At least 11 different isozymes have been identified as members of the serine/threonine PKC family, which can be divided into three categories based on their molecular structures and activation modes, comprising the classical or conventional PKC, the novel PKC and the atypical PKC (30). The G-proteins G α_q and G $\beta\gamma$, rather than G α_s , have been hypothesized to activate phospholipase for PKC activation (31). However, previous studies on nociceptors have suggested that G α_s may activate PKC (32,33). In the G α_s signaling pathway, G α_s /AC/cAMP may induce PKC ϵ activation, and this pathway is important in β AR signal transduction. Therefore, it was hypothesized that the β AR agonist Iso activates PKC ϵ in cardiomyocytes, resulting in a series of pathological and physiological reactions.

In the present study, western blot analysis and confocal laser scanning microscopy were used to detect PKC ϵ translocation following Iso stimulation in cardiomyocytes. The results revealed that Iso enhanced the activation of PKC ϵ . The Iso-stimulated PKC ϵ activation was relatively short and

time-dependent, peaking following 1 min of Iso stimulation and returning to the baseline level after 30 min. Iso also induced PKC ϵ to translocate to the perinuclear area. Since the subcellular localization of PKC was key in determining its function and specificity regulation, the localization of PKC ϵ around the nucleus may lead to the phosphorylation of target proteins, thus regulating intranuclear transcription and protein synthesis. The results demonstrated that β AR stimulation activated PKC ϵ in cardiomyocytes.

It has been reported that the PKA inhibitor CMIQ does not inhibit β 2-AR-induced translocation of PKC ϵ in neurons (16,18,34) and the signaling pathway from AC/cAMP to PKC does not involve PKA, indicating other upstream branches of the Gs/cAMP second-messenger signaling pathway at PKA prior to the activation of PKC ϵ (35). To verify whether the Iso-induced PKC ϵ activation passed through PKA in cardiomyocytes, Ad.PKI was used to infect cardiomyocytes. Following specific inhibition of PKA activity, Iso remained able to activate PKC ϵ , suggesting that PKA did not mediate Iso-induced PKC ϵ activation and that PKC ϵ activation was PKA-independent, possibly mediated by the novel PKA-independent signaling pathway factor, Epac.

Epac, a protein identified by Kawasaki *et al* (12) and de Rooij *et al* (13), may be directly activated by cAMP and




Figure 3. Effects of Iso and 8-CPT on PKC ϵ translocation in cardiomyocytes. Western blotting was used to determine the PKC ϵ translocation in cellular particulate fractions and β -actin was used as an internal reference (n=6). PMA was used as a positive control. Data are presented as the means \pm standard error of the mean of six independent experiments. *P<0.05, compared with the control. Iso, isoproterenol; PKC, protein kinase C; 8-CPT, 8-CPT-2'OMe-cAMP; PMA, phorbol 12-myristate 13-acetate; C, control.




Figure 4. Effect of the PKA inhibitor, Ad.PKI, on Iso-induced PKC ϵ activation. Following infection of the cardiomyocytes by Ad.PKI and GFP, Iso treatment (1 μ Mol/l for 1 min) was performed. Western blot analysis was used to determine the PKC ϵ content in the cellular particulate fractions, with β -actin as an internal reference (n=4). Data are presented as the means \pm standard error of the mean of four independent experiments. **P<0.01, compared with the GFP and PKI control group. Iso, isoproterenol; PK, protein kinase; Ad.PKI, adenovirus-coded rabbit muscle cAMP-dependent protein kinase inhibitor; GFP, green fluorescent protein; C, control.

mediate the cAMP signal transduction process. Epac is involved in numerous physiological processes, including cell division and differentiation, exocytosis, insulin secretion, cell adhesion and amyloid protein secretion (36,37). Studies have revealed that Epac may be a positive regulatory protein for cardiomyocyte hypertrophy (38). It has been demonstrated that Epac1 is involved in the hypertrophic effects of β AR in a PKA-independent manner in adult rat ventricular cardiomyocytes (39). In the present study, following incubation of cardiomyocytes with the Epac activator, 8-CPT, PKC ϵ translocation to the cellular particulate fractions was observed, suggesting that Epac activated PKC ϵ . The pattern




Figure 5. Effect of the Epac inhibitor, DN Epac (Epac R279K), on Iso-induced PKC ϵ activation and translocation. Following infection of the cardiomyocytes by adenoviruses carrying EpacR279K (DN) and GFP, Western blot analysis was used to determine the PKC ϵ content in the particulate fractions of the cardiomyocytes, with β -actin as an internal reference (n=5). Data are presented as the means \pm standard error of the mean of five independent experiments. **P<0.01, compared with the GFP control group. Iso, isoproterenol; PKC, protein kinase C; GFP, green fluorescent protein; Epac, exchange protein directly activated by cAMP; C, control.




Figure 6. Effect of the phospholipase C Inhibitor, U73122, on Iso-induced PKC ϵ activation. Following preincubation with U73122, western blot was used to determine the PKC ϵ content in the particulate fractions of the cardiomyocytes, with β -actin as an internal reference (n=6). Data are presented as the means \pm standard error of the mean of six independent experiments. *P<0.05, compared with the control. Iso, isoproterenol; PKC, protein kinase C; C, control.

of 8-CPT-induced PKC ϵ translocation was similar to that of Iso and it was hypothesized that there may be causal association between Epac and Iso. To further clarify the role of Epac in the Iso-activated PKC ϵ signaling pathway, the mutant Epac R279K (DN) was constructed to inhibit the action of the wild-type Epac protein. The results revealed that Iso-induced PKC ϵ activation was inhibited in the mutant, suggesting an Epac-mediated β AR/PKC ϵ signaling pathway in cardiomyocytes.

Epac in cardiomyocytes is considered to mediate the activation of PLC ϵ and the β AR-dependent Ca²⁺ release (17) and a signal transduction pathway, beginning with β 2AR




Figure 7. Effect of Iso-induced PKC ϵ activation on the expression of pERK1/2 in cardiomyocytes. Following transfection of cardiomyocytes with the PKC ϵ -specific inhibitor and scramble peptides, western blot analysis was used to determine the expression of pERK1/2 in the cardiomyocytes, with ERK2 as an internal reference (n=4). Data are presented as the means \pm standard error of the mean of six independent experiments. *P<0.05, compared with the scramble peptide group. Iso, isoproterenol; PKC, protein kinase C; ERK, extracellular signal-regulated kinase; Scramble, scramble peptide; PKC ϵ inh, PKC ϵ inhibitor peptide; C, control.

and followed by cAMP, Epac and Rap2B, leading to PLC ϵ activation and calcium ion release, has been suggested (15). The results of the present study demonstrated that β AR stimulation activated PKC ϵ in cardiomyocytes and that Epac mediated this signal transduction pathway. Therefore, it was hypothesized that Epac may activate PLC and subsequently activate PKC ϵ . Following the preincubation of cardiomyocytes with the PLC inhibitor, U73122, the effect of PLC on Iso-induced PKC ϵ activation suggested that PLC mediated Iso-induced PKC ϵ activation.

ERK signaling is closely associated with the pathological processes of cardiomyocyte hypertrophy and apoptosis (40-42). Iso-induced cardiac hypertrophy is associated with ERK by increasing ERK1/2 mRNA transcription in cardiomyocytes (43). Iso may also activate MAPK in cardiomyocytes and phosphorylate the Raf/MEK/ERK pathways (44,45). PKC ϵ may phosphorylate Ras/Raf and lead to activation of ERK1/2, inducing myocardial protein synthesis and cardiomyocyte hypertrophy. Although β AR activates ERK, it remains to be elucidated whether β AR acts on ERK by activating PKC. In the present study, the PKC ϵ -specific translocation inhibitor peptide was used to inhibit the activation of PKC ϵ to observe the effect of PKC ϵ on pERK1/2. The results revealed that the PKC ϵ -specific inhibitor peptide inhibited the Iso-induced pERK1/2 increase, suggesting that PKC ϵ mediated the Iso-induced phosphorylation of ERK1/2. The increase in pERK1/2 was one of the downstream effectors of Iso-induced PKC ϵ activation and Iso may induce ERK1/2 phosphorylation through the activation of PKC ϵ , resulting in cardiomyocyte hypertrophy.

There is considerable evidence demonstrating that chronic β AR activation may result in myocardial hypertrophy and heart failure, accompanied by cardiomyocyte injury, apoptosis, necrosis and cardiac remodeling (46). The results presented in the present study demonstrated an interaction between β AR and PKC ϵ in cardiomyocytes. The possible pathway of PKC ϵ




Figure 8. Effect of Iso-induced PKC ϵ activation on cardiomyocyte protein content. The relative ratio of cardiomyocyte protein/DNA following transfection of cells with the PKC ϵ inhibitor and scramble peptides followed by 48 h Iso treatment (n=6). Data are presented as the means \pm standard error of the mean of six independent experiments. **P<0.01, compared with the scramble peptide group; #P<0.05, compared with the scramble peptide+Iso group. Scramble, scramble peptide; PKC ϵ inh, PKC ϵ inhibitor peptide; Iso, isoproterenol; PKC, protein kinase C.




Figure 9. Effect of Iso-induced PKC ϵ activation on cardiomyocyte surface area. The cell surface area of cardiomyocytes following transfection with the PKC ϵ inhibitor and scramble peptide followed by 48 h Iso treatment (n=6). Data are presented as the means \pm standard error of the mean of six independent experiments. *P<0.05, compared with the scramble peptide group; #P<0.05, compared with the scramble peptide+Iso group. Scramble, scramble peptide; PKC ϵ inh, PKC ϵ inhibitor peptide.

activation by β AR suggested that β AR stimulation may activate the Epac guanine exchange protein, which in turn activated PLC, resulting in PKC ϵ translocation to the cellular particulate fractions. ERK phosphorylation and cardiomyocyte hypertrophy are effects of Iso-induced PKC ϵ activation. The clarification of signal transduction of cardiac hypertrophy resulting from β AR activation in cardiomyocytes assists in further understanding the molecular mechanisms underlying cardiac hypertrophy, providing a reference for examining new preventative methods and developing effective drugs against cardiac hypertrophy.

Acknowledgements

The authors would like to thank Professor Weimin Yang from the School of Pharmaceutical Science and Yunnan Key Laboratory of Pharmacology for Natural Products, Kunming Medical University, whose guidance and support enabled successful completion of the this study. The present study was supported by grants from the Yunnan Provincial Science and Technology Department (grant no. 2014FB037) and the National Natural Science Foundation of China (grant no.81260027).

References

1. Lorell BH and Carabello BA: Left ventricular hypertrophy: pathogenesis, detection, and prognosis. *Circulation* 102: 470-479, 2000.

2. Diwan A and Dorn GW II: Decompensation of cardiac hypertrophy: cellular mechanisms and novel therapeutic targets. *Physiology (Bethesda)* 22: 56-64, 2007.
3. Artham SM, Lavie CJ, Milani RV, Patel DA, Verma A and Ventura HO: Clinical impact of left ventricular hypertrophy and implications for regression. *Prog Cardiovasc Dis* 52: 153-167, 2009.
4. Vakili BA, Okin PM and Devereux RB: Prognostic implications of left ventricular hypertrophy. *Am Heart J* 141: 334-341, 2001.
5. Frey N, Katus HA, Olson EN and Hill JA: Hypertrophy of the heart a new therapeutic target? *Circulation* 109: 1580-1589, 2004.
6. Dorn GW II: Adrenergic pathways and left ventricular remodeling. *J Card Fail* 8 Suppl 6: 370-373, 2002.
7. Barki-Harrington L, Perrino C and Rockman HA: Network integration of the adrenergic system in cardiac hypertrophy. *Cardiovasc Res* 63: 391-402, 2004.
8. Wang W, Zhu W, Wang S, Yang D, Crow MT, Xiao RP and Cheng H: Sustained β -adrenergic stimulation modulates cardiac contractility by Ca^{2+} /calmodulin kinase signaling pathway. *Circ Res* 95: 798-806, 2004.
9. Movsesian MA and Bristow MR: Alterations in cAMP-mediated signaling and their role in the pathophysiology of dilated cardiomyopathy. *Curr Top Dev Biol* 68: 25-48, 2005.
10. Lohse MJ, Engelhardt S and Eschenhagen T: What is the role of β -adrenergic signaling in heart failure? *Circ Res* 93: 896-906, 2003.
11. Osadchii OE: Cardiac hypertrophy induced by sustained beta-adrenoreceptor activation: pathophysiological aspects. *Heart Fail Rev* 12: 66-86, 2007.
12. Kawasaki H, Springett GM, Mochizuki N, Toki S, Nakaya M, Matsuda M, Housman DE and Graybiel AM: A family of camp-binding proteins that directly activate rap1. *Science* 282: 2275-2279, 1998.
13. de Rooij J, Zwartkruis FJ, Verheijen MH, Cool RH, Nijman SM, Wittinghofer A and Bos JL: Epac is a rap1 guanine-nucleotide-exchange factor directly activated by cyclic AMP. *Nature* 396: 474-477, 1998.
14. Schmidt M, Evellin S, Weernink PA, Von Dorp F, Rehmann H, Lomasney JW and Jakobs KH: A new phospholipase C-calcium signalling pathway mediated by cyclic AMP and a rap GTPase. *Nat Cell Biol* 3: 1020-1024, 2001.
15. Schmidt M, Evellin S, Weernink PA, von Dorp F, Rehmann H, Lomasney JW and Jakobs KH: A new phospholipase-C-calcium signalling pathway mediated by cyclic AMP and a Rap GTPase. *Nat Cell Biol* 3: 1020-1024, 2001.
16. Huch TB, Dina OA and Levine JD: Epac mediates a cAMP-to-PKC signaling in inflammatory pain: an isolectin B4(+) neuron-specific mechanism. *J Neurosci* 25: 6119-6126, 2005.
17. Oestreich EA, Wang H, Malik S, Kaproth-Joslin KA, Blaxall BC, Kelley GG, Dirksen RT and Smrcka AV: Epac-mediated activation of phospholipase C ϵ plays a critical role in β -adrenergic receptor-dependent enhancement of Ca^{2+} mobilization in cardiac myocytes. *J Biol Chem* 282: 5488-5495, 2007.
18. Oestreich EA, Wang H, Malik S, Goonasekera SA, Blaxall BC, Kelley GG, Dirksen RT and Smrcka AV: Epac and phospholipase C ϵ regulate Ca^{2+} release in the heart by activation of protein kinase C ϵ and calcium-calmodulin kinase II. *J Biol Chem* 284: 1514-1522, 2009.
19. Churchill E, Budas G, Vallentin A, Koyanagi T and Mochly-Rosen D: PKC isozymes in chronic cardiac disease: possible therapeutic targets? *Annu Rev Pharmacol Toxicol* 48: 569-599, 2008.
20. Inagaki K, Churchill E and Mochly-Rosen D: Epsilon protein kinase C as a potential therapeutic target for the ischemic heart. *Cardiovasc Res* 70: 222-230, 2006.
21. Mackay K and Mochly-Rosen D: Localization, anchoring, and functions of protein kinase C isozymes in the heart. *J Mol Cell Cardiol* 33: 1301-1307, 2001.
22. Schonwasser DC, Marais RM, Marshall CJ and Parker PJ: Activation of the mitogen-activated protein kinase/extracellular signal-regulated kinase pathway by conventional, novel and atypical protein kinase C isotypes. *Mol Cell Biol* 18: 790-798, 1998.
23. Nilsson D, Gustafsson L, Wackenfors A, Gesslein B, Edvinsson L, Paulsson P, Ingemansson R and Malmsjo M: Up-regulation of endothelin type B receptors in the human internal mammary artery in culture is dependent on protein kinase C and mitogen-activated kinase signaling pathways. *BMC Cardiovasc Disord* 8: 21, 2008.
24. Wang Y: Mitogen-activated protein kinases in heart development and diseases. *Circulation* 116: 1413-1423, 2007.
25. Iwaki K, Sukhatme VP, Shubeita HE and Chien KR: Alpha- and beta-adrenergic stimulation induces distinct patterns of immediate early gene expression in neonatal rat myocardial cells. *fos/jun* expression is associated with sarcomere assembly; *egr-1* induction is primarily an alpha 1-mediated response. *J Biol Chem* 265: 13809-13817, 1990.
26. Mochly-Rosen D: Localization of protein kinases by anchoring proteins: a theme in signal transduction. *Science* 268: 247-251, 1995.
27. Mochly-Rosen D and Gordon AS: Anchoring proteins for protein kinase C: a means for isozyme selectivity. *FASEB J* 12: 35-42, 1998.
28. Vincent F, Duquesnes N, Christov C, Damy T, Samuel JL and Crozatier B: Dual level of interactions between calcineurin and PKC-epsilon in cardiomyocyte stretch. *Cardiovasc Res* 71: 97-107, 2006.
29. Bos JL: Epac: a new cAMP target and new avenues in cAMP research. *Nat Rev Mol Cell Biol* 4: 733-738, 2003.
30. Newton AC: Protein kinase C: structural and spatial regulation by phosphorylation, cofactors and macromolecular interactions. *Chem Rev* 101: 2353-2364, 2001.
31. Gudermann T, Schoneberg T and Schultz G: Functional and structural complexity of signal transduction via G-protein-coupled receptors. *Annu Rev Neurosci* 20: 399-427, 1997.
32. Parada CA, Reichling DB and Levine JD: Chronic hyperalgesic priming in the rat involves a novel interaction between cAMP and PKC ϵ second messenger pathways. *Pain* 113: 185-190, 2005.
33. Gold MS, Levine JD and Correa AM: Modulation of TTX-R INa by PKC and PKA and their role in PGE2-induced sensitization of rat sensory neurons in vitro. *J Neurosci* 18: 10345-10355, 1998.
34. Fan L, Ma J, Chen YH and Chen XQ: Antioxidant and antimicrobial phenolic compounds from *Setaria viridis*. *Chem Nat Comp* 50: 433-437, 2014.
35. Lu ZX, Quazi NH, Deady LW and Polya GM: Selective inhibition of cyclic AMP-dependent protein kinase by isoquinoline derivatives. *Biol Chem Hoppe Seyler* 377: 373-384, 1996.
36. Bos JL: Epac proteins: multi-purpose cAMP targets. *Trends Biochem Sci* 31: 680-686, 2006.
37. Holz GG, Kang G, Harbeck M, Roe MW and Chepurny OG: Cell physiology of cAMP sensor epac. *J Physiol* 577: 5-15, 2006.
38. Morel E, Marcantoni A, Gastineau M, Birkedal R, Rochais F, Garnier A, Lompree A, Vandecasteele G and Lezoualch F: CAMP-binding protein epac induces cardiomyocyte hypertrophy. *Circ Res* 97: 1296-1304, 2005.
39. Métrich M, Morel E, Berthouze M, Pereira L, Charron P, Gomez A and Lezoualch F: Functional characterization of the cAMP-binding proteins Epac in cardiac myocytes. *Pharmacol Rep* 61: 146-153, 2009.
40. Wang H, Oestreich EA, Maekawa N, Bullard TA, Vikstrom KL, Dirksen RT, Kelley GG, Blaxall BC and Smrcka AV: Phospholipase C epsilon modulates beta-adrenergic Receptor-dependent cardiac contraction and inhibits cardiac hypertrophy. *Circ Res* 97: 1305-1313, 2005.
41. Kehat I and Molkentin JD: Extracellular signal-regulated kinase 1/2 (ERK1/2) signaling in cardiac hypertrophy. *Ann N Y Acad Sci* 1188: 96-102, 2010.
42. Bueno OF and Molkentin JD: Involvement of extracellular signal-regulated kinases 1/2 in cardiac hypertrophy and cell death. *Circ Res* 91: 776-781, 2002.
43. House SL, House BE, Glascock B, Kimball T, Nusayr E, Schultz JE and Doetschman T: Fibroblast growth factor 2 mediates isoproterenol-induced cardiac hypertrophy through activation of the extracellular regulated kinase. *Mol Cell Pharmacol* 2: 143-154, 2010.
44. Zhang GX, Kimura S, Murao K, Yu X, Obata K, Matsuyoshi H and Takaki M: Effects of angiotensin type I receptor blockade on the cardiac RAF/MEK/ERK cascade activated via adrenergic receptors. *J Pharmacol Sci* 113: 224-233, 2010.
45. Takeishi Y, Ping P, Bolli R, Kirkpatrick DL, Hoit B and Walsh RA: Transgenic overexpression of constitutively active protein kinase $\text{c}\epsilon$ causes concentric cardiac hypertrophy. *Circ Res* 86: 1218-1223, 2000.
46. Morisco C, Marrone C, Galeotti J, Shao D, Vatner DE, Vatner SF and Sadoshima J: Endocytosis machinery is required for β 1-adrenergic receptor-induced hypertrophy in neonatal rat cardiac myocytes. *Cardiovasc Res* 78: 36-44, 2008.