

Synergistic hepatotoxicity by cadmium and chlorpyrifos: Disordered hepatic lipid homeostasis

WEI HE^{1,2}, WENLI GUO², YI QIAN², SHUPING ZHANG², DIFENG REN¹ and SIJIN LIU²

¹Beijing Key Laboratory of Forestry Food Processing and Safety, College of Biological Sciences and Biotechnology, Beijing Forestry University, Beijing 100083; ²State Key Laboratory of Environmental Chemistry and Ecotoxicology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, P.R. China

Received May 13, 2014; Accepted January 23, 2015

DOI: 10.3892/mmr.2015.3381

Abstract. Due to its extensive application, chlorpyrifos (CPF) has contaminated a diverse range of environmental substrates, fruits and vegetables. A number of studies have suggested that CPF may incur adverse effects on human health, including neurotoxicity, hepatotoxicity and endocrine disruption. Additionally, cadmium (Cd) is one of the most prevalent environmental heavy metals, as a result of considerable use in a wide spectrum of industrial fields. Exposure to Cd can cause several lesions in various organs, including the liver, kidneys and lungs. CPF and Cd often co-exist in the environment, food and crops, however, their joint exposure and potential synergistic toxicity are largely neglected and unrecognized. Our previous study characterized an interaction between CPF and Cd, which may occur via bonding between Cd²⁺ and the nitrogen atom in the pyridine ring of CPF, or the chelation between one Cd²⁺ and two CPF molecules. Our previous study also identified increased hepatotoxicity induced by CPF and Cd together compared with the individual compounds. In the present study, the effects of the concomitant exposure of CPF and Cd on lipid metabolism in hepatocytes was investigated.

The results demonstrated an accumulation of lipids in hepatocytes, induced by the CPF and Cd complex, which was fundamentally distinct from its parental chemicals. Notably, the molecular mechanism by which the CPF-Cd complex significantly induced hepatic lipogenesis was revealed, elevating the concentrations of sterol regulatory element-binding protein-1 and fatty acid synthase. These findings pave the way for future studies in recognizing synergistic biological effects between pollutants.

Introduction

Organophosphorus pesticides have been widely used in agricultural environments, since the ban of organochlorine insecticides, to protect crops against a range of pests due to their broad spectrum of insecticidal activity, effectiveness and their non-persistence in the environment (1). Despite the benefits of pesticides, residues may remain in crops, animal feed and environmental substrates, leading to contamination and potential human exposure through the food chain (2-4). Chlorpyrifos (CPF) is a major type of organophosphorus pesticide and is currently used in the production of >40 crops and fruits, including peaches, citrus fruits, almonds and grapes (1). CPF and its metabolites have been detected in farm animals, including cattle and sheep (1,5), and CPF residues or metabolites have also been detected in food and human urine (6). In addition, CPF has been found in the umbilical cord blood from female individuals living in urban environments (7). CPF has been confirmed as a neurotoxin by inhibiting acetylcholinesterase in the central nervous system (8). This can damage the nervous system, resulting in dizziness, nausea, confusion and, at a high concentration, respiratory paralysis and mortality. Additionally, a previous study demonstrated that CPF incurs developmental toxicity in rats at concentrations of 25 mg/kg/day, which was a maternally toxic dose (9). Similar to the majority of environmental pollutants, CPF elicits significant cytotoxicity through generating oxidative stress (10-12).

Cadmium (Cd) is one of the most prevalent environmental heavy metals, and human exposure to Cd is associated with a number of toxicities, including hepatotoxicity, DNA damage and cell apoptosis (13-15). Notably, CPF and Cd often coexist in the same environmental media and food chains, causing simultaneous exposure to organisms (16,17) and resulting in

Correspondence to: Professor Difeng Ren, Beijing Key Laboratory of Forestry Food Processing and Safety, College of Biological Sciences and Biotechnology, Beijing Forestry University, 35 Qinghua East Road, Haidian, Beijing 100083, P.R. China
E-mail: rendifeng@bjfu.edu.cn

Professor Sijin Liu, State Key Laboratory of Environmental Chemistry and Ecotoxicology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, 18 Shuangqing Road, Haidian, Beijing 100085, P.R. China
E-mail: sjliu@rcees.ac.cn

Abbreviations: CPF, chlorpyrifos; Cd, cadmium; OPPs, organophosphorus pesticides; FASN, fatty acid synthase; SREBP, sterol regulatory element-binding protein; DMSO, dimethyl sulfoxide; FBS, fetal bovine serum; ACC, acetyl-coenzyme A carboxylase; CoA, coenzyme A

Key words: chlorpyrifos, cadmium, synergistic toxicity, lipid metabolism

common toxicities, including carcinogenicity and hepatotoxicity (18-20). Our previous study characterized an interaction between Cd and CPF, and this interaction likely occurs due to bonding between Cd and nitrogen atoms in the pyridine ring of CPF, or the chelation between one Cd²⁺ and two CPF molecules (21). The joint hepatotoxicity of Cd and CPF to HepG2 cells was also demonstrated, with the Cd-CPF complex increasing the level of apoptosis by compared with its parental components (21).

Hepatic lipid metabolism is important in governing the whole body energy metabolism, as the liver is the major site for the storage and release of glucose and lipids (22,23). Lipid accumulation within liver has been suggested to cause obesity, insulin resistance and type II diabetes (22-24), and also predisposes individuals to nutritional stresses. Increasing evidence suggests that fatty acid synthase (FASN) is a critical regulator of hepatic lipid homeostasis, including fat and cholesterol synthesis (25,26). FASN encodes one of the key enzymes involved in fatty acid synthesis and is required for the *de novo* synthesis of fatty acids (27). Upregulated expression levels of FASN have been reported in various types of human cancer, and has been suggested to contribute to poor prognosis and recurrence of these types of cancer (28). Sterol regulatory element-binding protein (SREBP) is the predominant transcriptional activator of FASN (29,30), and previous studies have revealed that the increased expression of SREBP-1 is significantly associated with the extent of fatty liver in mouse models of diabetes mellitus (31-33). The present study aimed to investigate the synergistic effect of Cd and CPF on fat metabolism and fat accumulation in hepatocytes.

Materials and methods

Chemicals and reagents. CPF was purchased from Shuangma Fine Chemical Co., Ltd. (Nantong, China) and was dissolved in dimethyl sulfoxide (DMSO) (Solarbio Science & Technology Co., Ltd., Beijing, China). The final concentration of DMSO was <0.1% in the culture medium. CdCl₂ was purchased from Sigma-Aldrich (St. Louis, MO, USA). Sterile water was used to dissolve the CdCl₂, and the stock solution was filtered through a 0.45 mm membrane (Solarbio Science & Technology Co., Ltd.).

Cell culture. The HepG2 human hepatic carcinoma cell line was purchased from Shanghai Cell Bank of Type Culture Collection of the Chinese Academy of Sciences (Shanghai, China). The HepG2 cells were cultured at a concentration of 5.0x10³ cells/well in RPMI-1640 medium (Gibco Life Technologies, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum (FBS; HyClone, Logan, UT, USA) and 100 U/ml penicillin-streptomycin (HyClone) at 37°C in an atmosphere of 5% CO₂. The cell culture medium was changed every day and the cells were passed every other day.

Cytotoxicity assessment. The cytotoxicity was initially screened using an MTT assay (Invitrogen Life Technologies, Carlsbad, CA, USA). Briefly, the cells were inoculated and cultured overnight at 37°C in 96-well plates (Corning, Inc., New York, NY, USA) at a density of 8.0x10³ cells/well in RPMI-1640 medium supplemented with penicillin, streptomycin (100 mM) and 1% FBS. Following culture, the HepG2 cells were treated with

various concentrations of CPF (10, 50, 100, 500, 1,000, 1,500, 2,000 or 2,500 μM) or Cd (5, 10, 20, 40, 60, 80 or 100 μM) for a further 24 h at 37°C, and 20 μl MTT (5 mg/ml) was then added to each well. Following an additional 4 h incubation at 37°C, 100 μl DMSO was added to each well, followed by absorption assessment at 490 nm on a microplate reader (Multiskan MK3; Thermo Fisher Scientific, Co., Ltd., Waltham, MA, USA). The synergistic toxicity was further determined using an Alamar Blue assay (Invitrogen Life Technologies) and a bromodeoxyuridine (BrdU) assay (Roche Diagnostics, Mannheim, Germany). Similar to the MTT assay, following treatment with 10 μM CPF or Cd, 10 μl Alamar Blue reagent (Invitrogen Life Technologies) was added to each well and incubated for 2 h at 37°C, prior to the plates being read on a microplate reader with an emission wavelength of 590 nm and an excitation wavelength of 540 nm. The BrdU assay was performed according to the manufacturer's instructions, as described previously (34).

High glucose exposure. The cells were exposed to high glucose, as described previously (35). Briefly, the HepG2 cells were seeded into 6-well plates in RPMI-1640 medium supplemented with 10% FBS. The cells were grown to 70% confluence, prior to being maintained in serum-free RPMI-1640 medium, containing high concentrations of glucose (30 mmol/l) overnight. The cells were subsequently treated with Cd and CPF, as described above.

Determination of total cholesterol and triglyceride levels. The intracellular concentrations of total cholesterol and triglycerides were assayed in the HepG2 cell lysates, according to the manufacturer's instructions of the Triglycerides Assay and Total Cholesterol Assay kits (Nanjing Jiancheng Bioengineering Institute, Nanjing, China). The cell pellet was collected and 0.3 ml Triton-X 100 (1-2%) was added. It was lysed for 30 min prior to the start of the assays according to the manufacturer's instructions. The mixture was incubated at 37°C for 5 min, followed by reading with the microplate reader at 590 nm.

Western blotting. Following the treatment with Cd and CPF, described above, the HepG2 cells were collected and washed twice with phosphate-buffered saline (Solarbio Science & Technology Co., Ltd.). The harvested cells were lysed in radioimmunoprecipitation lysis buffer supplemented with protease inhibitor cocktail (Roche Diagnostics) on ice for 30 min. The cells were subsequently centrifuged at 12,000 x g for 10 min. The supernatants were collected and were subjected to 10% SDS-PAGE (gel, Beijing ComWin Biotech Co., Ltd., Beijing, China; SDS buffer, Solarbio Science & Technology Co., Ltd.) and were transferred onto nitrocellulose membranes (Solarbio Science & Technology Co., Ltd.). The membranes were incubated with primary antibodies against SREBP-1 (1:200; bs-1402R), FASN (1:200; bs-1498R-PE) and β-actin (1:1,000; bs-0061R) (Bioss, Beijing, China) in 5% milk overnight at 4°C. They were then washed 3 times in Tris-buffered saline with Tween-20 (Solarbio Science & Technology Co., Ltd.) for 5 min, and then incubated with anti-rabbit-horseradish peroxidase secondary antibodies (1:8,000; ComWin Biotech Co., Ltd.) for 1 h at 37°C in 5% milk, then washed again in a similar manner. The target proteins were colored by a chemical




Figure 1. Cytotoxicity assay of HepG2 cells treated with various concentrations of CPF or Cd²⁺. The cytotoxicity was assessed by measuring absorbance following an MTT assay of the HepG2 cells 24 h after treatment with 10-2,500 μM (A) CPF and (B) Cd²⁺ (n=5). CPF, chlorpyrifos; Cd, cadmium; Cont, untreated control.




Figure 2. Cell viability following concomitant exposure of CPF and Cd²⁺ in HepG2 cells. (A) Alamar Blue and (B) BrdU assays were performed on HepG2 cells treated with CPF and Cd²⁺ either alone (10 μM each) or concomitantly for 24 h (n=4-5). CPF, chlorpyrifos; Cd, cadmium; BrdU, bromodeoxyuridine; Cont, untreated control.

method. The intensity of the protein bands were assessed using Image J software, version 1.48 (NIH, Bethesda, MD, USA).

Statistical analysis. Using SPSS software, version 17.0 (SPSS, Inc., Chicago, IL, USA) Student's two-tailed t-test was performed to analyze the experimental data between two groups, and one-way analysis of variance was used to analyze the mean differences between groups relative to the control. The data are expressed as the mean ± standard deviation. P<0.05 was considered to indicate a statistically significant difference.

Results and Discussion

The toxicity of combinations of environmental pollutants is important when determining the health risks of each individually. Cd and CPF have been demonstrated to have common targets and elicit similar phenotypes, including hepatotoxicity (11,36-38). Our previous study revealed a novel interaction between Cd ions and CPF, and described the cytotoxicity induced by the Cd-CPF complex (21). The present study hypothesized that the synergistic cytotoxicity between CPF and Cd²⁺ was likely to cause metabolic disorders in hepatocytes.

Initially, a series of experiments were performed to confirm the synergistic toxicity, as previously discussed (21). As shown in Fig. 1A, the MTT assay revealed that CPF induced significant toxicity to the HepG2 cells following 24 h treatment with 0.5, 1, 1.5, 2 and 2.5 mM, in a dose-dependent manner, compared with the control (P<0.05). The half maximal inhibitory concentration (IC₅₀) was ~0.5 mM in the HepG2 cells (Fig. 1A). No cytotoxicity was observed in the HepG2 cells at CPF concentrations ≤100 μM (P>0.05; Fig. 1A). In addition, the cytotoxicity of Cd was determined in HepG2 cells treated with 5, 10, 20, 40, 60, 80 and 100 μM CdCl₂. Following treatment for 24 h, no toxicity was observed ≤10 μM (P>0.05; Fig. 1B), however, CdCl₂ induced significant cell death at Cd concentrations >10 μM (P<0.05). The IC₅₀ value for Cd²⁺ was ~20 μM (Fig. 1B).

Previous studies have suggested that the formation of a complex between chemicals alters their transport across the cell membrane and increases intracellular localization, resulting in enforced cytotoxicity, which may not occur with individual chemicals (39-41). A classical interaction of CPF was identified with methyl mercury, as the formation of this complex significantly increased the bioaccumulation of methyl mercury, coupled with increased toxicity (42). Our previous study demonstrated




Figure 3. Morphological changes of the HepG2 cells following concomitant exposure of CPF and Cd^{2+} . The phase-contrast images demonstrated morphological changes of the HepG2 cells following treatment with CPF and Cd^{2+} either alone or concomitantly (magnification, x200). CPF, chlorpyrifos; Cd, cadmium; Cont, untreated control.




Figure 4. Lipid alterations due to the synergistic toxicity of CPF and Cd^{2+} . The levels of intracellular (A) triglycerides and (B) total cholesterol in the HepG2 cells treated for 24 h with CPF and Cd^{2+} (10 μM each) either alone or concomitantly ($n=4-5$). CPF, chlorpyrifos; Cd, cadmium; Cont, untreated control.




Figure 5. Western blot analysis of the proteins responsible for lipid metabolism in HepG2 cells treated with CPF and Cd^{2+} . The cells were exposed to 10 μM CPF and/or Cd^{2+} for 24 h and were subsequently collected and subjected to western blotting to determine the expression levels of (A) SREBP-1 and (B) FASN. β -actin was used as a loading control. CPF, chlorpyrifos; Cd, cadmium; SERBP, sterol regulatory element-binding protein; FASN, fatty acid synthase; Cont, untreated control.

that the combined toxicity of Cd and CPF was attributable to the Cd-CPF complex-facilitated intracellular transport (21). In the present study the combined toxicity of CPF and Cd was further confirmed using Alamar Blue and BrdU assays, and by selecting non-toxic concentrations ($10\ \mu\text{M}$ each) of CPF and Cd. As shown in Fig. 2A, cell viability was significantly reduced by 21% upon concomitant exposure of Cd^{2+} and CPF ($P<0.05$) compared with the control or following individual treatment with either Cd^{2+} or CPF, as demonstrated using an Alamar Blue assay. Similarly, in the BrdU incorporation assay, the concomitant exposure of Cd^{2+} and CPF markedly inhibited cell proliferation by 23%, relative to the control or following individual treatment with Cd^{2+} or CPF ($P<0.05$; Fig. 2B). In addition, a marked morphological change, of smaller and rounder cells, was observed in the cells simultaneously treated with CPF and Cd compared with the cells in the control and individual treatment groups (Fig. 3). Consistent with the cytotoxicity results, as discussed above, these morphological changes were indicative of cell death (Fig. 3). Taken together, these results revealed a significant synergistic cytotoxic effect of Cd^{2+} and CPF on the HepG2 cells, which was distinct from the effects of the individual chemicals.

To investigate the potential disturbance to lipid metabolism by the synergistic exposure of CPF and Cd, the present study investigated the lipid concentrations in HepG2 cells post-treatment. To improve characterization of lipid metabolism, this was performed using a cell model treated with high glucose, as previously described (35). As shown in Fig. 4A, the concentration of triglycerides increased ~2-fold in cells exposed to simultaneous treatment of CPF and Cd compared with the untreated cells or single compound-treated cells. Similar to the changes in triglyceride levels, the concomitant exposure to CPF and Cd^{2+} ($10\ \mu\text{M}$ each) significantly increased the total cholesterol concentration by 80% ($P<0.05$) compared with the untreated cells or those treated with individual components ($P<0.05$; Fig. 4B). These results collectively suggested that the synergistic effect of CPF and Cd markedly altered hepatic lipid metabolism, associated with cholesterol and triglyceride accumulation, in the hepatocytes.

The present study subsequently aimed to investigate the molecular mechanism underlying CPF/Cd-mediated disorders in lipid metabolism. SREBP is a critical regulator of hepatic lipid metabolism, including glucose transport, gluconeogenesis and lipolysis (43). The SREBP family consists of SREBP-1a, SREBP-1c and SREBP-2 (44), and these members are essential for regulating the expression of lipogenic enzymes, including acetyl-coenzyme A (CoA) carboxylase and FASN (45,46). SREBP-1 is an important transcription factor, which stimulates lipogenic enzymes involved in liver fatty-acid synthesis, whereas SREBP-2 is relatively specific to the regulation of genes responsible for cholesterol synthesis and uptake, including low-density lipoprotein receptor and 3-hydroxy-3-methylglutaryl CoA reductase (44). The lipid accumulation, induced by exposure to CPF/Cd, may reside in the dysfunction of SREBP-1. The concentration of SREBP-1 was assessed in HepG2 cells following various treatments. Western blot analysis revealed that the expression of SREBP-1 was markedly increased by >2-fold in the cells following combined treatment with CPF and Cd compared with the untreated cells or those treated with CPF or Cd alone (Fig. 5A). The expression of

FASN was also markedly induced following treatment with CPF/Cd (>2-fold) compared with the untreated cells or those treated with the individual chemicals (Fig. 5B). These results demonstrated that the concomitant exposure of CPF and Cd induced hepatic fat accumulation through an increase in hepatic lipogenesis by increasing the expression levels of SREBP-1 and FASN.

Organophosphorus pesticides and heavy metals are amongst the most serious environmental pollutants. CPF, as a broad class of organophosphorus pesticides, is widely used throughout the world for agricultural and non-agricultural purposes (1-4,13). Cd is a toxic metal, commonly found in industrial workplaces and diverse environmental substrates, which contaminates food chains, including crops and fruits (47). Cd is toxic to various cell types, even at low concentrations, and has a long biological half-life in humans (10-30 years) (13,21). However, CPF and Cd are often detected in identical environmental substrates and food chains, and they elicit similar damage to organisms, including hepatotoxicity (16,17). Our previous study demonstrated the synergistic effect of CPF and Cd on reducing the viability of HepG2 cells (21). The present study demonstrated a novel HepG2 cell phenotype, caused by the CPF/Cd complex, with significant cholesterol and triglyceride accumulation in the hepatocytes. The molecular mechanism underlying hepatic lipogenesis elicited by joint exposure of CPF+Cd was found to occur through inducing the expression levels of SREBP-1 and FASN. Future investigations are required to determine the detailed mechanisms responsible for CPF/Cd-mediated action in lipogenesis.

Acknowledgements

This study was supported by grants from the Special Fund for Forest Scientific Research in the Public Welfare (no. 201304805), the National Natural Science Foundation of China (nos. 21377159 and 31201339), the Chinese Academy of Sciences (nos. KZCX2-EW-404 and XDB14000000) and the New Century Excellent Talents Program in University of the Ministry of Education (no. NCET-11-0587). The authors would like to thank the laboratory members for their assistance with experiments and reagents.

References

1. Chung SW and Chan BT: Validation and use of a fast sample preparation method and liquid chromatography-tandem mass spectrometry in analysis of ultra-trace levels of 98 organophosphorus pesticide and carbamate residues in a total diet study involving diversified food types. *J Chromatogr A* 1217: 4815-4824, 2010.
2. Štěpán R, Tichá J, Hajslová J, Kovalczuk T and Kocourek V: Baby food production chain: pesticide residues in fresh apples and products. *Food Addit Contam* 22: 1231-1242, 2005.
3. Szpyrka E, Kurdziel A, Slowik-Borowiec M, Grzegorzak M and Matyaszek A: Consumer exposure to pesticide residues in apples from the region of south-eastern Poland. *Environ Monit Assess* 185: 8873-8878, 2013.
4. Reiss R, Johnston J, Tucker K, DeSesso JM and Keen CL: Estimation of cancer risks and benefits associated with a potential increased consumption of fruits and vegetables. *Food Chem Toxicol* 50: 4421-4427, 2012.
5. Ivey MC and Palmer JS: Chlorpyrifos and 3,5,6-trichloro-2-pyridinol: residues in body tissues of swine treated with chlorpyrifos for hog louse and itch mite control. *J Econ Entomol* 72: 837-838, 1979.

6. Fenske RA, Lu C, Barr D and Needham L: Children's exposure to chlorpyrifos and parathion in an agricultural community in central Washington State. *Environ Health Perspect* 110: 549-553, 2002.
7. Whyatt RM, Rauh V, Barr DB, *et al*: Prenatal insecticide exposures and birth weight and length among an urban minority cohort. *Environ Health Perspect* 112: 1125-1132, 2004.
8. Caughlan A, Newhouse K, Namgung U and Xia Z: Chlorpyrifos induces apoptosis in rat cortical neurons that is regulated by a balance between p38 and ERK/JNK MAP kinases. *Toxicol Sci* 78: 125-134, 2004.
9. Farag A, El Okazy AM and El-Aswed AF: Developmental toxicity study of chlorpyrifos in rats. *Reprod Toxicol* 17: 203-208, 2003.
10. Bagchi D, Bagchi M, Hassoun EA and Stohs SJ: In vitro and in vivo generation of reactive oxygen species, DNA damage and lactate dehydrogenase leakage by selected pesticides. *Toxicology* 104: 129-140, 1995.
11. Das PC, Cao Y, Rose RL, Cherrington N and Hodgson E: Enzyme induction and cytotoxicity in human hepatocytes by chlorpyrifos and N,N-diethyl-m-toluamide (DEET). *Drug Metabol Drug Interact* 23: 237-260, 2008.
12. Ki YW, Park JH, Lee JE, Shin IC and Koh HC: JNK and p38 MAPK regulate oxidative stress and the inflammatory response in chlorpyrifos-induced apoptosis. *Toxicol Lett* 218: 235-245, 2013.
13. Son YO, Lee JC, Hitron JA, Pan J, Zhang Z and Shi X: Cadmium induces intracellular Ca^{2+} - and H_2O_2 -dependent apoptosis through JNK- and p53-mediated pathways in skin epidermal cell line. *Toxicol Sci* 113: 127-137, 2010.
14. Nzengue Y, Steiman R, Garrel C, Lefèbvre E and Guiraud P: Oxidative stress and DNA damage induced by cadmium in the human keratinocyte HaCaT cell line: role of glutathione in the resistance to cadmium. *Toxicology* 243: 193-206, 2008.
15. Zhou T, Zhou G, Song W, *et al*: Cadmium-induced apoptosis and changes in expression of p53, c-jun and MT-I genes in testes and ventral prostate of rats. *Toxicology* 142: 1-13, 1999.
16. Mansour SA, Belal MH, Abou-Arab AA and Gad MF: Monitoring of pesticides and heavy metals in cucumber fruits produced from different farming systems. *Chemosphere* 75: 601-609, 2009.
17. Fatta D, Canna-Michaelidou S, Michael C, *et al*: Organochlorine and organophosphoric insecticides, herbicides and heavy metals residue in industrial wastewaters in Cyprus. *J Hazard Mater* 145: 169-179, 2007.
18. Tuzmen N, Candan N, Kaya E and Demiryas N: Biochemical effects of chlorpyrifos and deltamethrin on altered antioxidative defense mechanisms and lipid peroxidation in rat liver. *Cell Biochem Funct* 26: 119-124, 2008.
19. Rai A, Maurya SK, Khare P, Srivastava A and Bandyopadhyay S: Characterization of developmental neurotoxicity of As, Cd, and Pb mixture: synergistic action of metal mixture in glial and neuronal functions. *Toxicol Sci* 118: 586-601, 2010.
20. Padilla S, Marshall RS, Hunter DL, *et al*: Neurochemical effects of chronic dietary and repeated high-level acute exposure to chlorpyrifos in rats. *Toxicol Sci* 88: 161-171, 2005.
21. Chen L, Qu G, Sun X, *et al*: Characterization of the interaction between cadmium and chlorpyrifos with integrative techniques in incurring synergistic hepatotoxicity. *PLoS One* 8: e59553, 2013.
22. Mingrone G, Rosa G, Greco AV, *et al*: Intramyocytic lipid accumulation and SREBP-1c expression are related to insulin resistance and cardiovascular risk in morbid obesity. *Atherosclerosis* 170: 155-161, 2003.
23. Xia C, Li R, Zhang S, *et al*: Lipid accumulation product is a powerful index for recognizing insulin resistance in non-diabetic individuals. *Eur J Clin Nutr* 66: 1035-1038, 2012.
24. Krssak M and Roden M: The role of lipid accumulation in liver and muscle for insulin resistance and type 2 diabetes mellitus in humans. *Rev Endocr Metab Disord* 5: 127-134, 2004.
25. Moreau A, Têrue C, Beylot M, *et al*: A novel pregnane X receptor and S14-mediated lipogenic pathway in human hepatocyte. *Hepatology* 49: 2068-2079, 2009.
26. Hao J, Zhu L, Zhao S, Liu S, Liu Q and Duan H: PTEN ameliorates high glucose-induced lipid deposits through regulating SREBP-1/FASN/ACC pathway in renal proximal tubular cells. *Exp Cell Res* 317: 1629-1639, 2011.
27. Vazquez-Martin A, Colomer R, Brunet J, Lupu R and Menendez JA: Overexpression of fatty acid synthase gene activates HER1/HER2 tyrosine kinase receptors in human breast epithelial cells. *Cell Prolif* 41: 59-85, 2008.
28. Wu X, Qin L, Fako V and Zhang JT: Molecular mechanisms of fatty acid synthase (FASN)-mediated resistance to anti-cancer treatments. *Adv Biol Regul* 54: 214-221, 2014.
29. Jeon BN, Kim YS, Choi WI, *et al*: Kr-pok increases FASN expression by modulating the DNA binding of SREBP-1c and S14 at the proximal promoter. *J Lipid Res* 53: 755-766, 2012.
30. Choi WI, Jeon BN, Park H, *et al*: Proto-oncogene FBI-1 (Pokemon) and SREBP-1 synergistically activate transcription of fatty-acid synthase gene (FASN). *J Biol Chem* 283: 29341-29354, 2008.
31. Horton JD, Goldstein JL and Brown MS: SREBPs: activators of the complete program of cholesterol and fatty acid synthesis in the liver. *J Clin Invest* 109: 1125-1131, 2002.
32. Yahagi N, Shimano H, Hasty AH, *et al*: Absence of sterol regulatory element-binding protein-1 (SREBP-1) ameliorates fatty livers but not obesity or insulin resistance in Lep(ob)/Lep(ob) mice. *J Biol Chem* 277: 19353-19357, 2002.
33. Goda Y, Shimizu T, Kato Y, *et al*: Two acylated anthocyanins from purple sweet potato. *Phytochemistry* 44: 183-186, 1997.
34. Qu G, Liu S, Zhang S, *et al*: Graphene oxide induces toll-like receptor 4 (TLR4)-dependent necrosis in macrophages. *ACS Nano* 7: 5732-5745, 2013.
35. Hwang YP, Choi JH, Han EH, *et al*: Purple sweet potato anthocyanins attenuate hepatic lipid accumulation through activating adenosine monophosphate-activated protein kinase in human HepG2 cells and obese mice. *Nutr Res* 31: 896-906, 2011.
36. Akhtar N, Srivastava MK and Raizada RB: Assessment of chlorpyrifos toxicity on certain organs in rat, *Rattus norvegicus*. *J Environ Biol* 30: 1047-1053, 2009.
37. Zhukalin M, Blanksma MK, Silva TD, *et al*: Characterization and in vitro cytotoxicity testing of ethanolamine-derived cadmium chelating agents. *Biomaterials* 20: 61-72, 2007.
38. Gottschalg E, Moore NE, Ryan AK, *et al*: Phenotypic anchoring of arsenic and cadmium toxicity in three hepatic-related cell systems reveals compound- and cell-specific selective up-regulation of stress protein expression: implications for fingerprint profiling of cytotoxicity. *Chem Biol Interact* 161: 251-261, 2006.
39. Goel A, Dani V and Dhawan DK: Protective effects of zinc on lipid peroxidation, antioxidant enzymes and hepatic histoarchitecture in chlorpyrifos-induced toxicity. *Chem Biol Interact* 156: 131-140, 2005.
40. Dondero F, Piacentini L, Banni M, Rebelo M, Burlando B and Viarengo A: Quantitative PCR analysis of two molluscan metallothionein genes unveils differential expression and regulation. *Gene* 345: 259-270, 2005.
41. Ediz L, Hiz O, Ozkol H, Gulcu E, Toprak M and Ceylan MF: Relationship between anti-CCP antibodies and oxidant and anti-oxidant activity in patients with rheumatoid arthritis. *Int J Med Sci* 8: 139-147, 2011.
42. Steevens JA and Benson WH: Interactions of chlorpyrifos and methyl mercury: a mechanistic approach to assess chemical mixtures. *Mar Environ Res* 50: 113-117, 2000.
43. Viollet B, Foretz M, Guigas B, *et al*: Activation of AMP-activated protein kinase in the liver: a new strategy for the management of metabolic hepatic disorders. *J Physiol* 574: 41-53, 2006.
44. Yuan H, Shyy JY and Martins-Green M: Second-hand smoke stimulates lipid accumulation in the liver by modulating AMPK and SREBP-1. *J Hepatol* 51: 535-547, 2009.
45. Lee MS, Kim D, Jo K and Hwang JK: Nordihydroguaiaretic acid protects against high-fat diet-induced fatty liver by activating AMP-activated protein kinase in obese mice. *Biochem Biophys Res Commun* 401: 92-97, 2010.
46. Rawson RB: Control of lipid metabolism by regulated intramembrane proteolysis of sterol regulatory element binding proteins (SREBPs). *Biochem Soc Symp* 2003 221-231, 2003.
47. Leach RM, Jr., Wang KW and Baker DE: Cadmium and the food chain: the effect of dietary cadmium on tissue composition in chicks and laying hens. *J Nutr* 109: 437-443, 1979.