

# Rho-associated protein kinase inhibitor, Y-27632, significantly enhances cell adhesion and induces a delay in G<sub>1</sub> to S phase transition in rabbit corneal endothelial cells

YU-MEI DIAO and JING HONG

Department of Ophthalmology, Peking University Third Hospital, Haidian, Beijing 100191, P.R. China

Received March 16, 2014; Accepted January 7, 2015

DOI: 10.3892/mmr.2015.3628

**Abstract.** Human corneal endothelial cells are a non-proliferative cell type. As a result of the increase in corneal endothelium disease, increasing numbers of studies have been conducted in order to promote corneal endothelial cell proliferation. The aim of the present study was to investigate the proliferative effects of Rho-associated protein kinase inhibitor, Y-27632, on rabbit corneal endothelial cells (rCECs). Y-27632 (1, 10 or 30  $\mu$ M) was added at two different time points to two groups of rCECs. The first group received Y-27632 when rCECs were initially plated, and the second following 72 h of cell growth. Cell morphology and cell adhesion ratios were subsequently observed using light microscopy. A cell counting kit was used to measure the number of viable cells that adhered to culture plates. Cell cycle transitions and levels of Annexin V-positive apoptotic cells were detected using flow cytometry. Cells treated with 1  $\mu$ M Y-27632 and 10  $\mu$ M Y-27632 retained their cell shape. At a concentration of 30  $\mu$ M Y-27632, the cell shape became irregular. Cell adhesion ratios, in 1  $\mu$ M Y-27632 (36.84%), 10  $\mu$ M Y-27632 (84.21%) and 30  $\mu$ M Y-27632 (84.21%) were higher than the adhesion ratio in the control group ( $P<0.01$ ). The optical densities of rCECs treated with 10  $\mu$ M or 30  $\mu$ M Y-27632 following 72 h of cell growth was less than that of the control cells ( $P<0.01$ ), but higher than that of cells which received Y-27632 at the time of plating ( $P<0.01$ ). Flow cytometry results also demonstrated that there was a delay in G<sub>1</sub> to S phase cell cycle progression in rCECs following administration of 10  $\mu$ M Y-27632 ( $P<0.01$ ). Cell apoptosis was inhibited when 10  $\mu$ M Y-27632 was added, at the time of cell plating, as well as when added following 72 h of cell growth ( $P<0.01$ ). At a concentration of 10  $\mu$ M Y-27632, there was an improvement in cell adhesion and an inhibition of the cell cycle in rabbit corneal endothelial cells. In conclusion,

Y-27632 has different effects on rCECs when administered at varying concentrations and at particular stages of cell growth.

## Introduction

Corneal endothelial cells (CECs) are crucial for the maintenance of corneal transparency. Human CECs (hCECs) are a non-proliferative monolayer of cells. This results in an average cell loss of 0.3-0.6% per year (1). The decrease may be accelerated as a result of accidental trauma; certain systemic diseases, such as diabetes (2); treatment for glaucoma (3); or endothelial dystrophies (4). When corneal endothelial cell density falls below a critical threshold, the functions of the endothelium are compromised, which leads to the formation of corneal edema and the loss of visual acuity (5).

In order to promote CEC proliferation, numerous mechanisms aiming to alter the non-proliferative status of these cells have been investigated. These include the release of cell-cell contact (6); growth factor-induced proliferation (7); the use of siRNA to reduce cyclin-dependent kinase inhibitors (8,9), SV40 large-T antigen (10); and induction of cell division by the human papilloma virus type 16 oncoproteins, E6/E7 (11). However, these results have not yet been transferred to clinical trials.

The Ras-related small GTPase, Rho, functions as a molecular switch of various cellular processes by shuttling between the inactive GDP-bound and active GTP-bound forms (12). Various cellular actions of Rho, include cell-to-substrate adhesion and motility, G<sub>1</sub>-S cell cycle progression and cell transformation, amongst other roles, by serum response factor (13). In addition, the microinjection of active RhoA into quiescent Swiss 3T3 fibroblasts (14) was found to induce the G<sub>1</sub>-S transition via downstream effectors, such as the Rho-associated coiled-coil forming protein serine/threonine kinase (ROCK) family of proteins. In particular, p160ROCK (ROCK-I) (15) and ROKa/Rho-kinase/ROCK-II (16-18). These studies verified that active RhoA induces G<sub>1</sub>-S transition.

In contrast to the findings from these studies, Y-27632, a specific inhibitor of the ROCK family of kinases (19), was used to promote cell cycle progression (20). However, these studies did not assess the cell cycle in detail; instead using changes in positive Ki67 and BrdU expression to measure cell cycle progression. Therefore, the present study was designed to further evaluate cell cycle G<sub>1</sub>-S phase progression and the effects of Y-27632 on rabbit CECs (rCECs).

*Correspondence to:* Professor Jing Hong, Department of Ophthalmology, Peking University Third Hospital, 49 North Garden Road, Haidian, Beijing 100191, P.R. China  
E-mail: hongjing1964@163.com

**Key words:** Y-27632, corneal endothelial cells, proliferation, cell cycle

## Materials and methods

**Animals.** A total of 15 New Zealand white rabbits (Peking University Health Science Center, Beijing, China; weight, 1.5-2.0 kg) were used in the present study. The experiments performed were in accordance with the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research and the Declaration of Helsinki. All animal care and experimental protocols complied with the Animal Management Rules of the Ministry of Health of the People's Republic of China and the guide for the Care and Use of the Laboratory Animals of Peking University. All animal protocols were approved by the Animal Research Committee of the Peking University Health Science Center. All efforts were made to minimize animal suffering.

**Primary culture.** Descemet membranes were stripped to obtain CECs. The cells were then incubated in 0.25% trypsin (Sigma-Aldrich, St. Louis, MO, USA) at 37°C in 5% CO<sub>2</sub>. After 15 min, the CECs were resuspended in culture medium and plated in one well of a 12-well plate. All primary cell cultures and passages of CECs were performed in Dulbecco's modified Eagle's medium (DMEM; cat no. 31600-034, Invitrogen Life Technologies, Carlsbad, CA, USA) supplemented with 15% fetal bovine serum (FBS; cat no. 16000-044, Gibco-BRL, Carlsbad, CA, USA), 50 U/ml penicillin and 50 µg/ml gentamicin. CECs were maintained in a humidified atmosphere at 37°C in 5% CO<sub>2</sub>. Culture medium was changed every two days. When cells reached confluence, they were trypsinized and passaged at ratios ranging from 1:2 to 1:4. rCECs from passage three were used in the present study.

**Effect of Y-27632 on cell shape.** Passage three rCECs were plated in 24-well plates (1.1875x10<sup>4</sup> cells/well). Different concentrations of Y-27632 (1, 10 or 30 µM) were added simultaneously when plating or after culture for 72 h. Cells without Y-27632 treatment were used as the control. After 12, 24 and 72 h, changes in cell morphology were observed using an inverted microscope (Eclipse Ti; Nikon Corp., Tokyo, Japan).

**Adhesion assay.** The adhesion assay was performed as reported previously (21). Briefly, rCECs (1.1875x10<sup>4</sup>) with Y-27632 (1, 10 and 30 µM) were plated in each well of 24-well plates. The cells without Y-27632 were used as controls. There were three wells, each at different concentrations. Following incubation at 37°C in 5% CO<sub>2</sub> for 2 h, the 24-well plate was washed with phosphate-buffered saline (PBS) and confirmed that there were no residual non-adherent cells. Subsequently, the non-adherent cells of each well were removed into centrifuge tubes. The ratios of adherent cells were calculated using the following formula: adherent ratio = (1.1875x10<sup>4</sup> - number of unattached cells) / 1.1875x10<sup>4</sup>.

**Cell counting kit-8 (CCK-8) assay to evaluate cell proliferation.** Passage three rCECs were seeded in 96-well plates (2,000 cells/well) and the medium was changed daily. rCECs that were treated with Y-27632, received this compound either at plating or after 72 h of cell culture. For rCECs that received Y-27632 during plating, the cells were cultured with Y-27632 (1, 10 or 30 µM) for 12, 24 and 72 h. At these time points, the

treatment medium was then replaced with fresh culture medium that did not contain Y-27632 and the treatment groups were cultured for a further 96 h. For rCECs that received Y-27632 following 72 h of cell growth, 1, 10 and 30 µM concentrations of Y-27632 were applied for 24 h. In order to assay the number of viable cells present, complete medium (100 µl) plus CCK-8 (10 µl) was added to each well. Untreated cells were used as controls and wells that did not contain cells were used as blank controls. After 2 h at 37°C, culture plates were agitated for 5 min prior to recording of optical density (OD) values at 450 nm. OD values for viable cells were calculated using the following formula: Fold difference in viable cell number = (OD<sub>sample</sub> - OD<sub>blank</sub>) / (OD<sub>control</sub> - OD<sub>blank</sub>).

**Analysis of cell cycle using flow cytometry.** For rCECs treated with Y-27632 at the time of plating and those treated after 72 h of cell growth, cells were harvested and resuspended in 2.1 ml absolute ethyl alcohol and 0.9 ml PBS. Cells were then stained for 1 h in darkness with PBS containing 50 µg/ml propidium iodide (PI) and 100 µg/ml ribonuclease A [included in the Annexin-fluorescein isothiocyanate (FITC) apoptosis detection kit; cat no. APOAF-50TST; Sigma-Aldrich]. Cell cycle progression was then evaluated using flow cytometry (FACSCalibur; BD Biosciences, Franklin Lakes, NJ, USA). The results shown are representative of three independent experiments.

**Apoptosis analysis.** Using an Annexin-FITC apoptosis detection kit, cells were washed twice with Dulbecco's PBS (Sigma-Aldrich) and resuspended in 1X binding buffer at a concentration of 1x10<sup>6</sup> cells/ml. For each sample, a 500 µl aliquot was combined with 5 µl Annexin V-FITC and 10 µl PI solution and incubated at room temperature in darkness for 10 min. The fluorescence for each sample was then detected and analyzed using flow cytometry (BD Biosciences).

**Statistical analysis.** Statistical analyses were performed using SPSS 16.0 statistical software (IBM, Armonk, NY, USA). A completely randomized design analysis of variance (ANOVA; one-factor ANOVA) and independent samples t-test were used to evaluate test data. Experimental data are expressed as the mean ± standard deviation (SD). P<0.05 was considered to indicate a statistically significant difference.

## Results

**Effects of Y-27632 on cell shape.** For cells treated with 1 µM (Fig. 1A) and 10 µM (Fig. 1B) Y-27632 at the time of plating, normal cellular morphology was observed at 24 h. However, for cells plated with 30 µM Y-27632 for 12 h, thin, long, neurite-like processes and cell extensions were observed (Fig. 1C). In addition, the cell shapes observed were not homogenous compared with the control group, or with cells treated with 1 and 10 µM Y-27632. At 24 h following removal of Y-27632, cells which had been treated with 30 µM Y-27632 remained irregular in shape, while the cells treated with 1 and 10 µM Y-27632 retained a generally normal shape. Furthermore, these changes were found to be concentration- and time-dependent (Fig. 1D). Parallel experiments with rCECs treated with Y-27632 following 72 h of cell growth were also performed and similar results were obtained.

Table I. Apoptosis assay of rCECs treated with Y-27632.

Treatment group	UR+LR (%)
A-12 h-con	61.8±0.92
A-12 h	47.91±0.37 <sup>a</sup>
B-24 h-con	46.41±0.98
B-24 h	8.48±0.98 <sup>a</sup>

Levels of apoptosis in groups A-12 h and B-24 h were lower than those of the corresponding controls. <sup>a</sup>P<0.01; n=3. rCECs, rabbit corneal epithelial cells; UR, Upper right quadrant of dot plot of apoptosis data; LR, lower right quadrant of dot plot of apoptosis data. A-12 h, Y-27632 was added at the time of plating; con, control; B-24 h, Y-27632 was added following 72 h of cell growth.

Table II. Cell cycle profiles for rCECs treated with Y-27632.

Treatment group	G <sub>1</sub> (%)	S/G <sub>2</sub> (%)
A-12 h-con	83.91±0.71	16.09±0.71
A-12 h	86.77±0.89 <sup>a</sup>	13.23±0.89 <sup>a</sup>
A-72 h-con	83.59±1.9	16.41±1.9
A-72 h	91.98±1.2 <sup>b</sup>	8.020±1.2 <sup>b</sup>
B-12 h-con	72.24±0.53	27.75±0.54
B-12 h	76.34±0.61 <sup>b</sup>	23.66±0.61 <sup>b</sup>
B-72 h-con	93.06±0.76	6.930±0.75
B-72 h	95.70±0.33 <sup>b</sup>	4.300±0.33 <sup>b</sup>

Cell cycle profiles were analyzed up to 12 and 72 h following treatment of rCECs with or without Y-27632 at the time of plating (A), as indicated. Cell cycle profiles were analyzed at the same time points following treatment of rCECs with or without Y-27632 after 72 h of cell growth (B), as indicated. In all experiments, the percentage of cells in the G<sub>1</sub> phase was higher in the treated cell groups than in the control cells. <sup>a</sup>P<0.05, <sup>b</sup>P<0.01; n=3. rCECs, rabbit corneal epithelial cells; con, control.

**Cell adhesion following Y-27632 treatment.** Cell adhesion assays were performed and cell adhesion ratios were calculated 2 h after the cells were plated. A ratio of 84.21% (Fig. 2C and D) was recorded for rCECs treated with 10 and 30  $\mu$ M Y-27632, while the ratios for control cells (19.33%; Fig. 2A) and the 1  $\mu$ M treatment group (36.84%; Fig. 2B) were lower. Furthermore, the differences in adhesion ratios between the treatment groups and the control group, as well as between the 10 and 30  $\mu$ M treatment groups and the 1  $\mu$ M group, were statistically significant (P<0.01 in each case) (Table III).

**Assessment of cell proliferation.** Using a CCK-8 assay, the number of viable cells present was assayed (Figs. 3 and 4). For rCECs that were treated with Y-27632 following 72 h of cell growth, the number of viable cells present was determined at 24 h following treatment with Y-27632. In these cells, a dose-dependent decrease in viability was observed. In addition, the proliferation of the control cells compared with the cells treated with 1  $\mu$ M Y-27632 did not differ (P>0.05). However,




Figure 1. Cell morphology examined using light microscopy for rCECs treated with Y-27632. (A) 1  $\mu$ M Y-27632 for 24 h, (B) 10  $\mu$ M Y-27632 for 24 h, (C) 30  $\mu$ M Y-27632 for 12 h and (D) 30  $\mu$ M Y-27632 for 24 h. Magnification, x100. rCECs, rabbit corneal epithelial cells.

Table III. Cell adhesion ratios under different Y-27632 concentrations (1  $\mu$ m, 10  $\mu$ m and 30  $\mu$ m).

Y-27632 concentrations	Cell adhesion ratios
Control	19.33%
1 $\mu$ m	36.84%
10 $\mu$ m	84.21%
30 $\mu$ m	84.21%

Cell adhesion ratios were calculated 2 h after the cells were plated. The treatment groups were statistically significant compared with the control group (P<0.01 in each case).

the proliferation of the control cells was higher than that of the cells treated with 10 and 30  $\mu$ M Y-27632 (P<0.01 in each case). Fig. 4 shows the results of the cell counting assays performed in parallel, with rCECs treated with Y-27632 at the time of plating. The number of viable cells was subsequently assayed at 12 and 24 h time points. The highest and lowest levels of proliferation were observed in the group that received treatment with 10  $\mu$ M Y-27632 (P<0.01) and in the control cells (P<0.01), respectively. However, at the 72 h time point, the highest proliferation was observed for the rCECs treated with 1  $\mu$ M Y-27632 (P<0.01). In addition, at each of the time points assayed, the proliferation levels of the cells treated with 30  $\mu$ M Y-27632 were lower than those for cells that received 1 or 10  $\mu$ M Y-27632. Proliferation was highest at the 72 h time point following treatment of rCECs with 1  $\mu$ M Y-27632 (P<0.01). A difference in cell proliferation was also observed at the 12 and 24 h time points, although the difference was not significant (P>0.05). Furthermore, for cells receiving 10 or 30  $\mu$ M Y-27632, a time-dependent decrease in proliferation was observed (P<0.01).




Figure 2. Cell adhesion ratios were determined using light microscopy. (A) Untreated rCECs (control) were compared with (B) rCECs treated with 1  $\mu$ M Y-27632 for 2 h, (C) rCECs treated with 10  $\mu$ M Y-27632 for 2 h and (D) rCECs treated with 30  $\mu$ M Y-27632 for 2 h. Magnification, x100. rCECs, rabbit corneal epithelial cells.




Figure 3. Comparison of cell proliferation data for rCECs treated with various concentrations of Y-27632 following 72 h of cell growth. Proliferation of the control cells and the cells treated with 1  $\mu$ M Y-27632 did not differ ( $P > 0.05$ ). In addition, proliferation of the cells treated with 10  $\mu$ M Y-27632 or 30  $\mu$ M Y-27632 was lower than that of the control cells (\* $P < 0.01$ ). The error bars represent the standard deviation for the experiments independently performed three times. rCECs, rabbit corneal endothelial cells.

**Apoptotic analysis.** Levels of apoptosis were assayed following the treatment of rCECs with 10  $\mu$ M Y-27632 (Table I and Fig. 5). Using Annexin V/PI staining, the percentage of cells undergoing apoptosis following treatment with Y-27632 during plating was  $47.91 \pm 0.37\%$  after 12 h. For rCECs treated with Y-27632 following 72 h of cell growth, the percentage of cells undergoing apoptosis at 24 h following the administration of Y-27632 was  $8.48 \pm 0.98\%$ . Cell apoptosis ratios were also lower for these two treatment groups compared with control cells, at  $61.8 \pm 0.92$  and  $46.41 \pm 0.98\%$ , respectively ( $P < 0.01$ ).

**Cell cycle progression.** The distribution of cells between various phases of the cell cycle was assayed using flow cytometry (Table II and Fig. 6). For rCECs treated with 10  $\mu$ M Y-27632 during plating, a reduced number of cells




Figure 4. Cell proliferation of rCECs assayed at different time points following the addition of various concentrations of Y-27632 at the time of plating. The number of viable cells increased with each treatment and time point assayed, compared with the control group. The highest fold increase was observed at 72 h after the 1  $\mu$ M Y-27632 treatment and 12 h after the 10  $\mu$ M Y-27632 treatment. The error bars represent the standard deviation for the experiments, which were performed independently three times. (\* $P < 0.01$ , \* $P < 0.05$ ). rCECs, rabbit corneal endothelial cells.




Figure 5. rCECs treated with Y-27632 were assayed for apoptosis. Apoptosis was detected for rCECs treated for 12 h, (A) without and (B) with Y-27632 administered at the time of plating. Apoptosis was also detected for rCECs treated for 24 h, (C) without and (D) with Y-27632 administered following 72 h of cell growth. Levels of apoptosis for B and D were lower than those for A and C ( $P < 0.01$ ;  $n = 3$ ). rCECs, rabbit corneal endothelial cells.

transitioning from  $G_1$  phase to S/ $G_2$  phase was observed at 12 h and 72 h, compared with control cells examined at the same time points. In addition, the differences at the two time points were significant, and the percentage of cells in the  $G_1$  phase at 72 h was greater than the number of cells in the  $G_1$  phase at 12 h ( $P < 0.01$ ). For cells that were treated with Y-27632 following 72 h of cell growth, 10  $\mu$ M Y-27632 induced the transition from S phase to  $G_1$  phase after 72 h ( $P < 0.01$ ).

## Discussion

In the present study, treatment of rCECs with 10  $\mu$ M Y-27632 maintained an improved cell shape and cell adhesion compared with treatment with 30 and 1  $\mu$ M Y-27632. Previously, it has




Figure 6. Cell cycle profiles for rCECs treated with Y-27632. rCECs treated without and with Y-27632 at the time of plating were analyzed for cell cycle distribution at 12 h (A and B, respectively) and 72 h (C and D, respectively) following the addition of Y-27632. rCECs were also treated without and with Y-27632 following completion of 72 h of cell growth. Cell cycle distribution was then assayed at 12 h (E and F, respectively) and 72 h (G and H, respectively) after Y-27632 was added. In all of the assays, the percentage of cells in the G<sub>1</sub> phase was higher for the treated cells than the control cells ( $P<0.01$ ;  $n=3$ ). rCECs, rabbit corneal endothelial cells.

been demonstrated that CECs, in particular hCECs, require the promotion of their attachment for cell growth (22). For example, extracellular matrices (ECM) plated on coated culture surfaces, including collagen (23), ECM derived from bovine CECs (24), a mixture of laminin and chondroitin sulfate (25), and a commercially available FNC coating mix containing fibronectin, collagen and albumin (26), have been reported to promote cell adhesion. Although the underlying mechanisms responsible for enhanced cell adhesion have not been well-characterized, it has been hypothesized that the actin cytoskeleton has a critical role in combination with integrins (27-29). Vinculin, which was involved in the linkage of the integrin adhesion complex to the actin cytoskeleton, was upregulated in ROCK inhibitor-treated CECs. However, additional studies are required to further elucidate the mechanisms that mediate enhanced adhesion induced by ROCK inhibitors.

In the CCK-8 assays performed 24 h after the addition of Y-27632, it was initially observed that rCECs treated with 10  $\mu$ M Y-27632 following 72 h of cell growth exhibited lower OD values than control cells. These results suggested that treatment with 10  $\mu$ M Y-27632 did not promote rCEC proliferation; and instead either induced cell apoptosis or delayed cell cycle progression. This did not coincide with the results of a previous study (30), which reported that rCECs and monkey CECs treated with Y-27632 exhibited an increased number of proliferating cells. It was inferred that treatment different time points may produced different results. Therefore, the effects of Y-27632 added at the time of plating were then evaluated. These results demonstrated that OD value folds of Y-27632 (1, 10 and 30  $\mu$ M) were markedly higher than that of the control groups. It appeared that Y-27632, when added at the same time as plating increased cell proliferation. However, due to the concomitant significant increase in cell adhesion, it was

hypothesized that the increase in OD values was a result of the increased number of adherent cells.

In order to test this hypothesis, the effects of Y-27632 on cell cycle and apoptosis in rCECs were assessed using flow cytometry. As shown in the present study, with regard to cell shape, cell adhesion and the results of the CCK-8 assay, as well as the results from other studies (5,30-32), 10  $\mu$ M Y-27632 was the optimal concentration for treatment of rCECs. Therefore, in the following study, 10  $\mu$ M Y-27632 was used to investigate its effects on the cell cycle and apoptosis. In cells treated at the time of plating, at 12 and 72 h a delay in G<sub>1</sub>-S cell cycle progression was observed, and the percentage of G<sub>1</sub>-S transition at 72 h was lower than it was at 12 h. The effects of Y-27632 on the cell cycle appeared to occur in a time-dependent manner.

In cells treated following 72 h of cell growth, at 12 and 24 h, reduced G<sub>1</sub>-S phase cell cycle progression was also observed. This indicated that 10  $\mu$ M Y-27632 did not promote G<sub>1</sub>-S cell cycle progression. Furthermore, the apoptosis assay revealed that 10  $\mu$ M Y-27632 inhibited cell apoptosis. These results suggested that Y-27632 significantly promoted cell adhesion and inhibited cell apoptosis, and did not induce proliferation of rCECs. It was inferred that in previous studies, the apparent increase in rCECs proliferation may have been due to an increase of the numbers of attached cells, leading to an increase in the numbers of Ki67- and Brdu-positive cells. The results of the present study, corresponded with the results of previous studies, which demonstrated the effects of Y-27632 on hCECs (5), vascular smooth muscle cell (33), Swiss 3T3 cells (12) and myofibroblasts (34). A previous study reported that Y-27632 significantly inhibited thrombin-induced DNA synthesis in cultured aortic smooth muscle cells at 10 mM (33). This may explain why Y-27632 delayed cell cycle progression.

Notably, in the CCK-8 assay, it was shown that the OD value folds of cells treated with 1  $\mu$ M Y-27632 increased with increasing time from the time of plating. However, in cells treated with 10 and 30  $\mu$ M, the OD value folds decreased with increasing time from the time of plating. Furthermore, the OD value folds in cells treated with 1  $\mu$ M at 72 h were higher than those in cells treated with 10  $\mu$ M at 12 h ( $P < 0.05$ ) and 24 h ( $P < 0.01$ ). It was also observed that the OD values of cells treated with 1  $\mu$ M Y-27632 following 72 h cell proliferation was increased in comparison with control. Therefore, 1  $\mu$ M Y-27632 may be the optimal dose.

In 2008, Yin and Yu (35) reported that Y-27632 induced a reduction of CEC proliferation. The beneficial effects of Y-27632 on the eye remained to be elucidated. Therefore, the present results suggest that the use of Y-27632, either as a cell culture additive or in the form of eye drops, should be thoroughly investigated.

In conclusion, in the present study, Y-27632 significantly enhanced the adhesion of rCECs, inhibited cell apoptosis and delayed cell cycle progression. The mechanisms underlying the effect of Y-27632 on CECs and the other tissues of the eye remain to be elucidated and warrant further investigation prior to progression to clinical trials.

## Acknowledgements

The present study was supported by the National Natural Science Foundation of China (grant no. 31271045).

## References

- Murphy C, Alvarado J, Juster R and Maglio M: Prenatal and postnatal cellularity of the human corneal endothelium. A quantitative histologic study. *Invest Ophthalmol Vis Sci* 25: 312-322, 1984.
- Schultz RO, Matsuda M, Yee RW, *et al*: Corneal endothelial changes in type I and type II diabetes mellitus. *Am J Ophthalmol* 98: 401-410, 1984.
- Bigar F and Witmer R: Corneal endothelial changes in primary acute angle-closure glaucoma. *Ophthalmol* 89: 596-599, 1982.
- Stainer GA, Akers PH, Binder PS and Zavala EY: Correlative microscopy and tissue culture of congenital hereditary endothelial dystrophy. *Am J Ophthalmol* 93: 456-465, 1982.
- Pipparelli A, Arsenijevic Y, Thuret G, Gain P, Nicolas M and Majo F: ROCK inhibitor enhances adhesion and wound healing of human corneal endothelial cells. *PLoS One* 8: e62095, 2013.
- Gerdes J, Schwab U, Lemke H and Stein H: Production of a mouse monoclonal antibody reactive with a human nuclear antigen associated with cell proliferation. *Int J Cancer* 31: 13-20, 1983.
- Haj FG, Markova B, Klamann LD, Bohmer FD and Neel BG: Regulation of receptor tyrosine kinase signaling by protein tyrosine phosphatase-1B. *J Biol Chem* 278: 739-744, 2003.
- Kikuchi M, Zhu C, Senoo T, Obara Y and Joyce NC: p27kip1 siRNA induces proliferation in corneal endothelial cells from young but not older donors. *Invest Ophthalmol Vis Sci* 47: 4803-4809, 2006.
- Joyce NC and Harris DL: Decreasing expression of the G1-phase inhibitors, p21Cip1 and p16INK4a, promotes division of corneal endothelial cells from older donors. *Mol Vis* 16: 897-906, 2010.
- Wilson SE, Lloyd SA, He YG and McCash CS: Extended life of human corneal endothelial cells transfected with the SV40 large T antigen. *Invest Ophthalmol Vis Sci* 34: 2112-2123, 1993.
- Wilson SE, Weng J, Blair S, He YG and Lloyd S: Expression of E6/E7 or SV40 large T antigen-coding oncogenes in human corneal endothelial cells indicates regulated high-proliferative capacity. *Invest Ophthalmol Vis Sci* 36: 32-40, 1995.
- Ishizaki T, Uehata M, Tamechika I, *et al*: Pharmacological properties of Y-27632, a specific inhibitor of Rho-associated kinases. *Mol Pharmacol* 57: 976-983, 2000.
- Narumiya S: The small GTPase Rho: cellular functions and signal transduction. *J Biochem* 120: 215-228, 1996.
- Olson MF, Ashworth A and Hall A: An essential role for Rho, Rac and Cdc42 GTPases in cell cycle progression through G1. *Science* 269: 1270-1272, 1995.
- Ishizaki T, Maekawa M, Fujisawa K, *et al*: The small GTP-binding protein Rho binds to and activates a 160 kDa Ser/Thr protein kinase homologous to myotonic dystrophy kinase. *EMBO J* 15: 1885-1893, 1996.
- Leung T, Manser E, Tan L and Lim L: A novel serine/threonine kinase binding the Ras-related RhoA GTPase which translocates the kinase to peripheral membranes. *J Biol Chem* 270: 29051-29054, 1995.
- Matsui T, Amano M, Yamamoto T, *et al*: Rho-associated kinase, a novel serine/threonine kinase, as a putative target for small GTP binding protein Rho. *EMBO J* 15: 2208-2216, 1996.
- Nakagawa O, Fujisawa K, Ishizaki T, Saito Y, Nakao K and Narumiya S: ROCK-I and ROCK-II, two isoforms of Rho-associated coiled-coil forming protein serine/threonine kinase in mice. *FEBS Lett* 392: 189-193, 1996.
- Uehata M, Ishizaki T, Satoh H, *et al*: Calcium sensitization of smooth muscle mediated by a Rho-associated protein kinase in hypertension. *Nature* 389: 990-994, 1997.
- Okumura N, Koizumi N, Ueno M, *et al*: The new therapeutic concept of using a rho kinase inhibitor for the treatment of corneal endothelial dysfunction. *Cornea* 30 Suppl 1: S4-S9, 2011.
- Bi YL, Zhou Q, Du F, Wu MF, Xu GT and Sui GQ: Regulation of functional corneal endothelial cells isolated from sphere colonies by Rho-associated protein kinase inhibitor. *Exp Ther Med* 5: 433-437, 2013.
- Peh GS, Beuerman RW, Colman A, Tan DT and Mehta JS: Human corneal endothelial cell expansion for corneal endothelium transplantation: an overview. *Transplantation* 91: 811-819, 2011.
- Li W, Sabater AL, Chen YT, *et al*: A novel method of isolation, preservation and expansion of human corneal endothelial cells. *Invest Ophthalmol Vis Sci* 48: 614-620, 2007.
- Blake DA, Yu H, Young DL and Caldwell DR: Matrix stimulates the proliferation of human corneal endothelial cells in culture. *Invest Ophthalmol Vis Sci* 38: 1119-1129, 1997.
- Engelmann K, Bohnke M and Friedl P: Isolation and long-term cultivation of human corneal endothelial cells. *Invest Ophthalmol Vis Sci* 29: 1656-1662, 1988.
- Zhu C and Joyce NC: Proliferative response of corneal endothelial cells from young and older donors. *Invest Ophthalmol Vis Sci* 45: 1743-1751, 2004.
- Sastry SK and Burridge K: Focal adhesions: a nexus for intracellular signaling and cytoskeletal dynamics. *Exp Cell Res* 261: 25-36, 2000.
- Worthylake RA, Lemoine S, Watson JM and Burridge K: RhoA is required for monocyte tail retraction during transendothelial migration. *J Cell Biol* 154: 147-160, 2001.
- Worthylake RA and Burridge K: RhoA and ROCK promote migration by limiting membrane protrusions. *J Biol Chem* 278: 13578-13584, 2003.
- Okumura N, Ueno M, Koizumi N, *et al*: Enhancement on primate corneal endothelial cell survival in vitro by a ROCK inhibitor. *Invest Ophthalmol Vis Sci* 50: 3680-3687, 2009.
- Okumura N, Koizumi N, Ueno M, *et al*: ROCK Inhibitor converts corneal endothelial cells into a phenotype capable of regenerating in vivo endothelial tissue. *Am J Pathol* 181: 268-277, 2012.
- Liu X, Ory V, Chapman S, *et al*: ROCK inhibitor and feeder cells induce the conditional reprogramming of epithelial cells. *Am J Pathol* 180: 599-607, 2012.
- Seasholtz TM, Majumdar M, Kaplan DD and Brown JH: Rho and Rho kinase mediate thrombin-stimulated vascular smooth muscle cell DNA synthesis and migration. *Circ Res* 84: 1186-1193, 1999.
- Porter KE, Turner NA, O'Regan DJ, Balmforth AJ and Ball SG: Simvastatin reduces human atrial myofibroblast proliferation independently of cholesterol lowering via inhibition of RhoA. *Cardiovasc Res* 61: 745-755, 2004.
- Yin J and Yu FS: Rho kinases regulate corneal epithelial wound healing. *Am J Physiol Cell Physiol* 295: C378-C387, 2008.