

Naringin ameliorates cognitive deficits via oxidative stress, proinflammatory factors and the PPAR γ signaling pathway in a type 2 diabetic rat model

ZHONGHUA QI¹, YINGHUI XU², ZHANHUA LIANG¹, SHENG LI¹, JIE WANG², YI WEI² and BIN DONG²

Departments of ¹Neurology and ²Neurosurgery, The First Affiliated Hospital of Dalian Medical University, Dalian, Liaoning 116011, P.R. China

Received October 22, 2014; Accepted July 17, 2015

DOI: 10.3892/mmr.2015.4232

Abstract. Naringenin is a flavonoid polyphenolic compound, which facilitates the removal of free radicals, oxidative stress and inflammation. The present study aimed to obtain a better understanding of the effects of Naringin on the regulation of diabetes-associated cognitive decline, and its underlying mechanisms. An experimental diabetes mellitus (DM) rat model was induced by streptozotocin (50 mg/kg). Following treatment with naringin (100 and 200 mg/kg) for 16 weeks, the body weight and blood glucose levels of the DM rats were measured. A morris water maze test was used to analyze the effects of naringin on the cognitive deficit of the DM rats. The levels of oxidative stress, proinflammatory factors, caspase-3 and caspase-9, and the protein expression of peroxisome proliferator-activated receptor γ (PPAR γ) were quantified in the DM rats using a commercially-available kit and western blot assay, respectively. In addition, a GW9662 PPAR γ inhibitor (0.3 mg/kg) was administered to the DM rats to determine whether PPAR γ affected the effects of naringin on the cognitive deficit of the DM rats. The results demonstrated that naringin increased the body weight, blood glucose levels, and cognitive deficits of the DM rats. The levels of oxidative stress and proinflammatory factors in the naringin-treated rats were significantly lower, compared with those of the DM rats. In addition, naringin activated the protein expression of PPAR γ , and administration of the PPAR γ inhibitor decreased the protein expression of PPAR γ , and attenuated the effects of naringin on cognitive deficit. The results also demonstrated that naringin decreased the expression levels of caspase-3 and caspase-9 in the DM rats. These results suggested that naringin ameliorated cognitive

deficits via oxidative stress, proinflammatory factors and the PPAR γ signaling pathway in the type 2 diabetic rat model. Furthermore, oxidative stress, proinflammatory factors and PPAR γ signaling may be involved in mediating these effects.

Introduction

With the increase in population age, senile dementia has become a social problem worldwide (1). The central nervous system damage caused by type 2 diabetes mellitus (T2DM) is attracting increasing attention (2). Several studies have demonstrated the association between T2DM and cognitive dysfunction, with the central nervous system damage secondary to T2DM termed 'diabetes-associated cognitive decline (DACD)' (3,4).

At present, the precise mechanism underlying DACD remains to be elucidated. A variety of factors, including abnormal glucose metabolism, oxidative stress, T2DM complications and the inflammatory response, are involved in DACD, and there is overlapping among the pathogenic factors (5). Several investigations have been performed on the association between oxidative stress and cognitive dysfunction. Free radicals are highly reactive, and are involved in oxidization, biomolecular damage and toxicity to nerve cells, thus leading to lipofuscin deposition, increased age spots and vacuolar degeneration (6). Malardé *et al* (7) demonstrated that fermented soy permeate exhibited anti-oxidant and anti-inflammatory properties in streptozotocin (STZ)-induced diabetic rats. In addition, Wang *et al* (8) reported that chronic treatment with oxymatrine alleviates DACD, which is associated with oxidative stress, inflammation and apoptosis in rats.

Previous evidence has demonstrated that inflammation is involved in pathological damage via multiple mechanisms, which damages vascular function integrity in DACD (9). In treating the pathological mechanisms, reducing the inflammatory reaction in the brain can significantly improves neuronal damage and nerve fiber degeneration, thereby improving learning and memory (10). Mao *et al* (11) reported that Huperzine A ameliorates DACD via oxidative stress, inflammation and apoptosis. In addition, Li *et al* (9) reported that chrysin markedly alleviates DACD via oxidative stress, inflammation and apoptosis (9).

Correspondence to: Mr. Bin Dong, Department of Neurosurgery, The First Affiliated Hospital of Dalian Medical University, 222 Zhongshan Road, Dalian, Liaoning 116011, P.R. China
E-mail: bindongmr@163.com

Key words: naringenin, diabetes-associated cognitive decline, oxidative stress, peroxisome proliferator-activated receptor γ , inflammation

Peroxisome proliferator-activated receptor γ (PPAR γ) belongs to the superfamily of nuclear receptors and is a ligand-dependent transcription factor (12). PPARs regulate the gene expression of various start regions containing PPAR response elements, intracellular transcription levels and fatty acid and glucose metabolism, and inhibit the inflammatory response (12). Tharahaswari *et al* (13) reported that trigonelline and diosgenin attenuate endoplasmic reticulum stress and enhance adipose tissue PPAR γ activity in T2DM rats. Capobianco *et al* (14) demonstrated that PPAR activation regulates nitric oxide production, lipid concentration and lipoperoxidation in the placenta of patients with T2DM. In addition, pioglitazone ameliorates memory deficits in diabetic mice via the activation of PPAR γ (15).

Naringenin is a natural dihydro flavonoid, which is widely distributed in grapefruit and other citrus fruits and may also be chemically synthesized (16). Previous studies have demonstrated that naringenin exhibits various pharmacological effects, including anti-tumor, anti-mutagenic and anti-atherosclerotic effects (17-19). Studies have suggested that naringenin may reverse the liver damage caused by drugs or toxic chemical compounds, including alcohol, cadmium, carbon tetrachloride, oxytetracycline and dimethyl nitrosamine (20-23). The present study aimed to investigate the effects of naringenin on DACD. In addition, the correlation between nuclear oxidative stress, proinflammatory factors, PPAR γ and DACD was investigated.

Materials and methods

Animals. A total of 32 male 6-week-old Sprague-Dawley rats (weight, 270 \pm 20 g) obtained from the experimental center of the Dalian Medical University (Dalian, China) were selected for the experimental procedures in the present study. All animal procedures were performed in accordance with the guidelines of the First Affiliated Hospital of the Dalian Medical University and were approved by the Ethics Committee of Dalian Medical University (Dalian, China). Prior to experimentation, the rats were provided with *ad libitum* access to food and water, and were housed in a laboratory animal room at 23 \pm 1°C in 50-70% humidity on a 12 h light/dark cycle.

Drugs and chemicals. Naringin (purity, \geq 98%) and STZ were purchased from Sigma-Aldrich (St. Louis, MO, USA). Glutathione peroxidase (GSH-Px; S0056), glutathione (GSH; S0053), superoxide dismutase (SOD; S0038), malondialdehyde (MDA; S0131), ELISA kits, bicinchoninic acid (BCA; P0006) protein assay kits and caspase-3 (C1115) and caspase-9 (C1157) activity test kits were purchased from Beyotime Institute of Biotechnology (Haimen, China), and tumor necrosis factor α (TNF- α ; RT029) and interleukin (IL)-6 (RI001) were purchased from Shanghai Gefan Biological Technology Co., Ltd. (Shanghai, China).

Establishment of the diabetic rat model. At the onset of the experiment, the body weights of the experimental rats were measured. Plasma glucose levels were then detected using an enzymatic glucose oxidase peroxidase diagnostic kit. Fasting blood glucose levels >250 mg/dl were considered diabetic and were used for further experimentation. The chemical structure




Figure 1. Chemical structure of naringin.

of naringin is shown in Fig. 1. After 1 week of acclimatization, the rats received a single intraperitoneal injection of STZ (50 mg/kg) to induce T2DM, with the exception of the normal healthy controls, as previously described (24). At the end of the experiment (48 h following injection), the body weight and blood glucose levels of the experimental rats were measured.

Experimental design. In total, eight normal rats were injected with physiological saline, defined as the control group; eight diabetic rats were injected with physiological saline, defined as the DM group; 24 diabetic rats were randomly divided into three groups, and were treated with naringin at doses of 100 mg/kg and 200 mg/kg into the caudal vein, as previously described (25). Rats from each group (4/group) were immediately sacrificed by cervical dislocation under 30 mg/kg pentobarbital (Sigma-Aldrich), and brain tissue and blood samples were collected under anesthesia (300 mg/kg intraperitoneally). The samples were stored at -80°C for further experimentation.

Morris water maze (MWM) assessment. Following treatment with naringin (100 and 200 mg/kg) for 16 weeks, MWM assessments were performed, as previously described (26,27). Rats from each group (4/group) were trained to swim freely with a circular Plexiglas platform (14 cm diameter) submerged 1.5 cm beneath the surface of the water prior to performing the MWM test. The platform was located in a fixed position, equidistant from the center and the wall of the tank. The rats were subjected to four training trials per day. The rats were placed into the tank at one of the four designated start points per day, in a pseudorandom order, and trained for as many days as required to reach the criterion of 25 sec to reach the platform. If the rats failed to find the platform within 60 sec, they were manually guided to the platform and allowed to remain there for 5 sec. Following the final training session, a probe trial was performed after 24 h, consisting of a 60 sec free swim in the pool in the absence of the platform. The MWM assessments were recorded via video capture and analyzed using a SMARTW system (PanLab, Barcelona, Spain).

Oxidative stress assessment. Tissue sections (~5 mg) of the cerebral cortex and hippocampus were added to 100 μ l tissue lysis buffer (Beijing Solarbio Science & Technology Co., Ltd., Beijing, China) on ice, and incubated for 10 min. The homogenates were then centrifuged at 20,000 \times g for 15 min

at 4°C. The clear supernatant was collected and analyzed for oxidative stress. This was achieved by quantifying the levels of GSH-Px, GSH, SOD and MDA in the cerebral cortex and hippocampus, using commercially-available kits (Beyotime Institute of Biotechnology), according to the manufacturer's instructions.

TNF- α and IL-6 assessment. Tissue sections (~5 mg) of the cerebral cortex and hippocampal samples were added to 100 μ l tissue lysis buffer on ice, and incubated for 10 min. The homogenates were then centrifuged at 20,000 x g for 15 min at 4°C. TNF- α and IL-6 ELISA kits (Beyotime Institute of Biotechnology) were used to quantify the protein levels, according to the manufacturer's instructions, and the samples were analyzed spectrophotometrically (Model 550; Bio-Rad Laboratories, Inc., Hercules, CA, USA) at 450 nm absorbance.

Western blot analysis. Tissue sections (~5 mg) of the cerebral cortex and hippocampal tissue samples were added to 100 μ l tissue lysis buffer on ice and incubated for 10 min. The homogenates were then centrifuged at 20,000 x g for 15 min at 4°C. The supernatant was collected and the protein concentration was measured using a BCA protein assay kit. The protein was separated by 8-12% SDS-PAGE (Sigma-Aldrich), and electrotransferred to polyvinylidene difluoride membranes (0.22 mm; Beijing Zhongshan Golden Bridge Biotechnology Co., Ltd., Beijing, China). The membranes were blocked with tris-buffered saline (TBS; Wuhan Boster Biological Technology, Ltd., Wuhan, China) containing 5% non-fat milk for 2 h. The membranes were subsequently incubated with monoclonal mouse anti-human PPAR γ (sc-7273; 1:2,000; Santa Cruz Biotechnology, Inc., Dallas, TX, USA) and polyclonal mouse anti-human β -actin (bs-0061R; 1:500; BIOSS, Beijing, China) primary antibodies, overnight at 4°C. The membranes were washed three times with TBS with Tween 20 (1%; Sigma-Aldrich) for 2 h, and then incubated with anti-mouse horseradish peroxidase-conjugated IgG (sc-2370; 1:3,000; Santa Cruz Biotechnology, Inc.), for 2 h. The bands were visualized using an ECL kit (Bio-Rad Laboratories, Inc.) and quantified using densitometry (Image Quant LAS 4000 software; GE Healthcare Life Sciences, Chalfont, UK).

Caspase-3 and caspase-9 activity level quantification. Tissue sections (~5 mg) of the cerebral cortex and hippocampal tissue samples were added to 100 μ l tissue lysis buffer on ice and incubated for 10 min. The homogenates were then centrifuged at 20,000 x g for 15 min at 4°C. The activity levels of caspase-3 and caspase-9 were measured using a caspase-3 and caspase-9 activity test kit, according to the manufacturer's instructions, and incubated at 37°C for 120 min. The levels of caspase-3 and caspase-9 were measured using the Model 550 spectrophotometer at 405 nm.

Statistical analysis. Statistical analyses were performed using one-way analysis of variance followed by Dunnett's test. Statistical analyses were performed using SPSS 17.0 (SPSS, Inc., Chicago, IL, USA), and the data are expressed as the mean \pm standard deviation. $P < 0.05$ was considered to indicated a statistically significant difference.




Figure 2. Effects of naringin on the (A) body weight and (B) blood glucose levels of DM rats. Data are expressed as the mean \pm standard deviation. ** $P < 0.01$, vs. the Con group; ## $P < 0.01$, vs. the DM group. Con, control group; DM, diabetes group; NAR (100), naringin (100 mg/kg)-treated group; NAR (200), naringin (200 mg/kg)-treated group.

Results

Effects of naringin on body weight and blood glucose levels. The chemical structure of naringin is shown in Fig. 1. The DM rats exhibited a significant decrease in body weight and an increase in blood glucose levels, compared with the normal control group. Following treatment with naringin (100 and 200 mg/kg) for 16 weeks, the body weights and blood glucose levels of the DM rats were significantly increased and decreased, respectively, compared with those of the untreated DM group (Fig. 2A-B).

Effects of naringin on cognitive deficit. Following 16 weeks treatment with naringin (100 and 200 mg/kg), a MWM test was used to assess cognitive function. The escape latency markedly increased in the DM group, compared with the control group (Fig. 3A). Following treatment with naringin, the escape latency was reduced, compared with that of the DM group (Fig. 3A). The mean path length significantly increased in the DM rats, compared with the control group. The mean path length following treatment with naringin was significantly reduced, compared with that of the DM group (Fig. 3B). The duration spent in the target quadrant and the number of times the rats crossed the former platform location were significantly reduced in the DM group, compared with those observed in the control group (Fig. 3C-D). Treatment with naringin markedly reversed these effects in the DM rats (Fig. 3C-D). However, the swimming speed of the rats in the control group was similar to those observed in the other groups (Fig. 3E).

Effects of naringin on diabetes-induced changes in oxidative stress. To examine the effects of naringin on oxidative stress in the brain tissue, the expression levels of GSH-Px, GSH, SOD




Figure 3. Effects of naringin on cognitive deficit. The effects of naringin on (A) escape latency, (B) mean path length, (C) percentage of time spent in the target quadrant, (D) number of times crossing the platform and (E) swimming speed in the rats. Data are expressed as the mean \pm standard deviation. ** $P < 0.01$, vs. Con group; ## $P < 0.01$, vs. DM group. Con, control; DM, diabetes group; NAR (100), naringin (100 mg/kg)-treatment; NAR (200), naringin (200 mg/kg) treatment.

and MDA were measured in the cerebral cortex and hippocampus tissue samples. As shown in Fig. 4A-C, the expression levels of GSH-Px, GSH and SOD were significantly decreased in the cerebral cortex and hippocampus of the DM group, compared with those of the control group. Treatment of the STZ-induced diabetic rats with naringin (100 and 200 mg/kg) significantly increased the expression levels of GSH-Px, GSH and SOD in the cerebral cortex and hippocampus (Fig. 4A-C). In addition, the expression levels of MDA in the DM group were significantly increased, compared with those of the control group (Fig. 4D). Following treatment with naringin (100 and 200 mg/kg) for 16 weeks, these expression levels were significantly reduced (Fig. 4D).

Effects of naringin on DM-induced changes in proinflammatory cytokines. To determine the effects of naringin on the brain proinflammatory cytokines in the DM rat, the expression levels of TNF- α and IL-6 were measured in the cerebral cortex and hippocampus. Compared with those of the control group, the expression levels of TNF- α and IL-6 were significantly increased in the cerebral cortex and hippocampus tissues of the STZ-induced DM rats (Fig. 5A and B). Treatment with naringin (100 and 200 mg/kg) significantly reversed the elevated expression levels of TNF- α and IL-6 in the cerebral cortex and hippocampus, compared with the DM group (Fig. 5A and B).

Effects of naringin on the expression levels of PPAR γ . To investigate whether naringin exerts its effects through upregulation of the expression of PPAR γ , the expression levels of PPAR γ were measured in the cerebral cortex and hippocampus tissues using western blotting. As shown in Fig. 6A and B, the expression levels of PPAR γ were significantly decreased in the cerebral cortex and hippocampus of the DM group, compared with those of the control group. Treatment of the STZ-induced DM rats with naringin (100 and 200 mg/kg) significantly increased the protein expression levels of PPAR γ in the cerebral cortex and hippocampus, compared with the DM rats without naringin (Fig. 6A-B).

PPAR γ inhibitor regulates the effects of naringin on DACD. To confirm the observed results that the PPAR γ GW9662 inhibitor (0.3 mg/kg) regulated the effects of naringin on DACD, a MWM test was performed. The PPAR γ inhibitor significantly decreased the protein expression levels of PPAR γ (Fig. 7A-B). In addition, PPAR γ inhibitor reversed the effects of naringin on DACD following treatment with naringin (100 and 200 mg/kg) for 16 weeks (Fig. 7C-G).

Effects of naringin on the expression levels of caspase-3 and caspase-9. To examine whether naringin affected the levels of caspase-3 and caspase-9, the expression levels of caspase-3




Figure 4. Effects of naringin on diabetes-induced changes in oxidative stress. The effects of naringin on the expression levels of (A) GSH-Px, (B) GSH, (C) SOD and (D) MDA in the rats. Data are expressed as the mean \pm standard deviation. ** $P < 0.01$, vs the Con group; ## $P < 0.01$, vs the DM group. Con, control group; DM, diabetes group; NAR (100), naringin (100 mg/kg)-treated group; NAR (200), naringin (200 mg/kg)-treated group; GSH, glutathione; GSH-Px, GSH peroxidase; SOD, superoxide dismutase; MDA, malondialdehyde.




Figure 5. Effects of naringin on diabetes-induced changes in the proinflammatory cytokines. Effects of naringin on the expression levels of (A) TNF- α and (B) IL-6 in the rats. Data are expressed as the mean \pm standard deviation. ** $P < 0.01$, vs the Con group; ## $P < 0.01$, vs the DM group. Con, control group; DM, diabetes group; NAR (100), naringin (100 mg/kg)-treated group; NAR (200), naringin (200 mg/kg)-treated group. TNF, tumor necrosis factor; IL, interleukin.




Figure 6. Effects of naringin on the expression levels of PPAR γ . (A) Representative western blotting analysis of the protein expression levels of PPAR γ in the cerebral cortex and hippocampus tissue samples. (B) Quantitative results of the protein expression levels of PPAR γ in the cerebral cortex and hippocampus tissues. Data are expressed as the mean \pm standard deviation. ** $P < 0.01$, vs the Con group; ## $P < 0.01$, vs the DM group. Con, control group; DM, diabetes group; NAR (100), naringin (100 mg/kg)-treated group; NAR (200), naringin (200 mg/kg)-treated group. PPAR γ , peroxisome proliferator-activated receptor γ .

and caspase-9 were measured in the cerebral cortex and hippocampus tissues using ELISA assays. As shown in Fig. 8A and B, naringin notably increased the expression levels of caspase-3 and caspase-9 in the cerebral cortex and hippocampus of the

DM group, compared with the control group. Treatment with naringin (100 and 200 mg/kg) decreased the expression levels of caspase-3 and caspase-9 in the cerebral cortex and hippocampus, compared with the control group (Fig. 8A and B).




Figure 7. PPAR γ inhibitor regulates the effects of naringin on DACD. (A) Representative western blots and (B) quantitative results of the protein expression of PPAR γ in the cerebral cortex and hippocampus. Effects of naringin on the (C) escape latency, (D) mean path length, (E) mean percentage of time spent in the target quadrant, (F) number of times of platform crossing, and (G) swimming speed in the rats. Data are expressed as the mean \pm standard deviation. ** $P < 0.01$, vs. the Con group; ## $P < 0.01$, vs. the DM group; and ### $P > 0.05$, vs. the DM group. Con, control group; DM, diabetes group; NAR (200), naringin (200 mg/kg)-treated group; NAR+GW (naringin, 200 mg/kg+GW9662, 0.3 mg/kg)-treated group. PPAR γ , peroxisome proliferator-activated receptor γ .




Figure 8. Effects of naringin on the expression levels of caspase-3 and caspase-9. (A and B) The effects of naringin on the activity levels of caspase-3 and caspase-9 in rats. Data are expressed as the mean \pm standard deviation. ** $P < 0.01$, vs. the Con group; ## $P < 0.01$, vs. the DM group. Con, control group; DM, diabetes group; NAR (100), naringin (100 mg/kg)-treated group; NAR (200), naringin (200 mg/kg)-treated group.

Discussion

With changes in lifestyles and aging of the population, T2DM incidence is increasing year by year (28). As a systemic disease, T2DM can cause a variety of organizational, structural and functional changes in organs, and the lesions can affect the whole body. Therefore, early recognition and treatment of DACD can delay and reduce the incidence of dementia, and improve the life quality of patients with T2DM (29). In the present study, naringin significantly increased the body weight and reduced the levels of blood glucose in DM rats. Oršolić *et al* (30) reported that the DNA-protective effects of naringenin increased the body weight of alloxan-induced diabetic mice. Naringenin has also been observed to markedly normalize the body weights of albino mice (31). Priscilla *et al* (32) demonstrated that naringenin reduces the postprandial glycemic response in DM rats. The results of the present study demonstrated that naringin effectively improved the cognitive deficit of DM rats.

The biological basis of DACD in cognitive impairment may be associated with abnormal brain development, in which a large number of reactive oxygen species produced by oxidative stress and secondary cell injury are important (33). Under normal circumstances, the body has an effective antioxidant defense mechanism, and SOD is one of the most important antioxidant enzymes (34). An increase in the levels of reactive oxygen species can cause lipid peroxidation, DNA damage and cell-associated molecule or gene regulation, thereby inducing the nerve cell damage characteristic of certain cognitive deficits (35). The compensatory activity of antioxidant enzyme SOD is then increased. Therefore, the possible biological basis of DACD may be associated with SOD. Oxidized low density lipoprotein is highly cytotoxic, which can lead to vascular endothelial cell necrosis, and it closely associated with the occurrence of atherosclerosis and other cardiovascular and cerebrovascular diseases (36). MDA is a major metabolite involved in the biological membrane damage by free radicals, and can cause disorders of protein synthesis, which results in memory and mental decline (36). The levels of MDA reflect the degree of lipid peroxidation in the body (36). The results of the present study further determined that the effects of naringin markedly increased the expression levels of GSH-Px, GSH and SOD in the cerebral cortex and hippocampus of the DM rats. In addition, the expression levels of MDA were significantly reduced following treatment with naringin in the DM rats. Previous studies have demonstrated that naringenin has a similar protective effects to L-arginine in monocrotaline-induced pulmonary hypertension through oxidative stress, inflammation and nitric oxide in rats (37). Jeon *et al* (38) reported that naringenin increases the expression levels of GSH-Px, GSH and SOD in rats fed a high-cholesterol diet. Hermenean *et al* (39) demonstrated that naringenin increases the expression of MDA and decreases the expression level of SOD, catalase, GSH and GSH-Px in the mouse kidney (39).

TNF- α is a type of cytokine produced by activated monocytes, which exhibits a wide range of activities (40). TNF- α may be involved in nerve damage, and a previous study on the brain neuroinflammatory response of patients with dementia demonstrated that the primary deposit of

TNF- α is β -amyloid, which is one of the causes of neurodegenerative dementia (41). IL-6 is produced by monocytes or macrophages, and is a stimulating factor produced by several cells *in vivo*; IL-6 exhibits a wide range of biological activities, and is an important member in the complex network of cytokines in the body, which are involved in various pathophysiological processes (42). An increase in the expression levels of IL-6 in the plasma of elderly patients is closely associated with cognitive impairment, which is a risk factor for cognitive dysfunction, and the rise of inflammatory factors in the plasma of patients occurs prior to the clinical diagnosis of dementia, suggesting that inflammatory cytokines of the peripheral blood may be involved in the pathophysiology of dementia (43). The results of the present study demonstrated that naringin decreased the expression levels of TNF- α and IL-6 in the cerebral cortex and hippocampus of DM rats. A previous study demonstrated that naringin significantly decreased the production and expression levels of IL-1 β and IL-6 in diabetic mice (44). The inhibition of TNF- α and IL-6 by naringenin may contribute to its anti-inflammatory activity in rats with ethanol-induced liver injury (45). Bodet *et al* (46) also reported that naringin exhibits anti-inflammatory properties in macrophages.

PPARs are a type of ligand-activated nuclear transcription factor that regulate the expression of several key genes, including those involved in glucose and lipid metabolism (47). There are three subtypes of PPARs: PPAR- α , PPAR- γ and PPAR- δ . PPAR- γ is an important transcription factor in lipogenesis, and promotes adipocyte differentiation, increased insulin sensitivity and lower blood sugar levels (48). PPAR- γ reduces the levels of blood fat and increases insulin sensitivity. Therefore, PPAR- γ has become the focus of investigations on anti-DM drugs (49). In the cerebral cortex and hippocampus, naringin activated the protein expression of PPAR γ in DM rats. Sharma *et al* (50) provided evidence that naringin ameliorates insulin resistance, hepatic steatosis and kidney damage in T2DM rats by regulating oxidative stress, inflammation and upregulation of PPAR γ . Naringin reduces ethanol intake and ethanol-conditioned place preference in mice (51). The present study demonstrated that the PPAR γ inhibitor, GW9662, decreased the protein expression of PPAR γ , as well as the effects of naringin on the cognitive deficit of the DM rats. These results suggested that naringin enhanced the cognitive deficit of DM rats via the upregulation of PPAR γ .

The results of the present study demonstrated that naringin reduced the expression levels of caspase-3 and caspase-9 in the cerebral cortex and hippocampus of DM rats, and decreased cell apoptosis in the brain tissues of the DM rats. Treatment with naringin improves functional recovery via the inhibition of caspase-3 following spinal cord injury in rats (52). In addition, naringin reduces the levels of 3-nitropropionic acid-induced apoptosis through decreased caspase-3 activation (53).

In conclusion, the present study demonstrated that naringin significantly ameliorated cognitive deficits via oxidative stress, and the proinflammatory and PPAR γ signaling pathways in T2DM rats. Future research will focus on the characterization of naringin, and aim to investigate the therapeutic effects of naringin on DACD *in vitro* and *in vivo*.

Acknowledgements

The current study was supported by the National Natural Scientific Foundation Project (grant no. 81371454; 2013); the Natural Scientific Foundation Project of Liaoning Province (grant no. 2014023021; 2014); the Dalian Natural Scientific Foundation Project (grant no. 2014E145F176); and the Natural Science Foundation of China (grant no. 30872665 to Mr. Bin Dong; 2008).

References

- Lin Z, Gu J, Xiu J, Mi T, Dong J and Tiwari JK: Traditional chinese medicine for senile dementia. *Evid Based Complement Alternat Med* 2012: 692621, 2012.
- Wang YB, Wang S, Bai R, Du JL, Xing Q, Ba Y, Yang Y, Zhang XY, Shi CH and Yao JJ: Efficacy of switching from premixed insulin to insulin glargine regimen in Type 2 diabetes mellitus patients with different islet functions. *Mol Med Rep* 10: 1096-1102, 2014.
- Tonoli C, Heyman E, Roelands B, Pattyn N, Buyse L, Piacentini MF, Berthoin S and Meeusen R: Type 1 diabetes-associated cognitive decline: A meta-analysis and update of the current literature. *J Diabetes* 6: 499-513, 2014.
- Chen J, Liang L, Zhan L, Zhou Y, Zheng L, Sun X, Gong J, Sui H, Jiang R, Zhang F and Zhang L: ZiBuPiYin recipe protects db/db mice from diabetes-associated cognitive decline through improving multiple pathological changes. *PLoS One* 9: e91680, 2014.
- Chang XH, Liang LN, Zhan LB, Lu XG, Shi X, Qi X, Feng ZL, Wu MJ, Sui H, Zheng LP, et al: The effect of Chinese Jinzhida recipe on the hippocampus in a rat model of diabetes-associated cognitive decline. *BMC Complement Altern Med* 13: 161, 2013.
- Kuhad A and Chopra K: Effect of sesamol on diabetes-associated cognitive decline in rats. *Exp Brain Res* 185: 411-420, 2008.
- Malardé L, Groussard C, Lefevre-Orfila L, Vincent S, Efstathiou T and Gratas-Delamarche A: Fermented soy permeate reduces cytokine level and oxidative stress in streptozotocin-induced diabetic rats. *J Med Food* 18: 67-75, 2015.
- Wang SB and Jia JP: Oxymatrine attenuates diabetes-associated cognitive deficits in rats. *Acta Pharmacol Sin* 35: 331-338, 2014.
- Li R, Zang A, Zhang L, Zhang H, Zhao L, Qi Z and Wang H: Chrysin ameliorates diabetes-associated cognitive deficits in Wistar rats. *Neurol Sci* 35: 1527-1532, 2014.
- Maczurek A, Hager K, Kenkles M, Sharman M, Martins R, Engel J, Carlson DA and Münch G: Lipoic acid as an anti-inflammatory and neuroprotective treatment for Alzheimer's disease. *Adv Drug Deliv Rev* 60: 1463-1470, 2008.
- Mao XY, Cao DF, Li X, Yin JY, Wang ZB, Zhang Y, Mao CX, Zhou HH and Liu ZQ: Huperzine A ameliorates cognitive deficits in streptozotocin-induced diabetic rats. *Int J Mol Sci* 15: 7667-7683, 2014.
- Umemoto T and Fujiki Y: Ligand-dependent nucleo-cytoplasmic shuttling of peroxisome proliferator-activated receptors, PPAR α and PPAR γ . *Genes Cells* 17: 576-596, 2012.
- Tharahaswari M, Jayachandra Reddy N, Kumar R, Varshney KC, Kannan M and Sudha Rani S: Trigonelline and diosgenin attenuate ER stress, oxidative stress-mediated damage in pancreas and enhance adipose tissue PPAR γ activity in type 2 diabetic rats. *Mol Cell Biochem* 396: 161-174, 2014.
- Capobianco E, Martinez N, Fornes D, Higa R, Di Marco I, Basualdo MN, Faingold MC and Jawerbaum A: PPAR activation as a regulator of lipid metabolism, nitric oxide production and lipid peroxidation in the placenta from type 2 diabetic patients. *Mol Cell Endocrinol* 377: 7-15, 2013.
- Liu LP, Yan TH, Jiang LY, Hu W, Hu M, Wang C, Zhang Q, Long Y, Wang JQ, Li YQ, et al: Pioglitazone ameliorates memory deficits in streptozotocin-induced diabetic mice by reducing brain β -amyloid through PPAR γ activation. *Acta Pharmacol Sin* 34: 455-463, 2013.
- Shi Y, Tan Y, Mao S and Gu W: Naringenin inhibits allergen-induced airway remodeling in a murine model of asthma. *Mol Med Rep* 9: 1204-1208, 2014.
- Arul D and Subramanian P: Naringenin (citrus flavonone) induces growth inhibition, cell cycle arrest and apoptosis in human hepatocellular carcinoma cells. *Pathol Oncol Res* 19: 763-770, 2013.
- Shimizu T, Lin F, Hasegawa M, Okada K, Nojiri H and Yamane H: Purification and identification of naringenin 7-O-methyltransferase, a key enzyme in biosynthesis of flavonoid phytoalexin sakuranetin in rice. *J Biol Chem* 287: 19315-19325, 2012.
- Lee S, Lee CH, Moon SS, Kim E, Kim CT, Kim BH, Bok SH and Jeong TS: Naringenin derivatives as anti-atherogenic agents. *Bioorg Med Chem Lett* 13: 3901-3903, 2003.
- Jayaraman J, Veerappan M and Namasivayam N: Potential beneficial effect of naringenin on lipid peroxidation and antioxidant status in rats with ethanol-induced hepatotoxicity. *J Pharm Pharmacol* 61: 1383-1390, 2009.
- Renugadevi J and Prabu SM: Cadmium-induced hepatotoxicity in rats and the protective effect of naringenin. *Exp Toxicol Pathol* 62: 171-181, 2010.
- Yen FL, Wu TH, Lin LT, Cham TM and Lin CC: Naringenin-loaded nanoparticles improve the physicochemical properties and the hepatoprotective effects of naringenin in orally-administered rats with CCl₄ (4)-induced acute liver failure. *Pharm Res* 26: 893-902, 2009.
- Pari L and Gnanasoundari M: Influence of naringenin on oxytetracycline mediated oxidative damage in rat liver. *Basic Clin Pharmacol Toxicol* 98: 456-461, 2006.
- Siddiqui O, Sun Y, Liu JC and Chien YW: Facilitated transdermal transport of insulin. *J Pharm Sci* 76: 341-345, 1987.
- Yu W, Wu J, Cai F, Xiang J, Zha W, Fan D, Guo S, Ming Z and Liu C: Curcumin alleviates diabetic cardiomyopathy in experimental diabetic rats. *PLoS One* 7: e25013, 2012.
- Wang X, Liu H, Zhang Y, Li J, Teng X, Liu A, Yu X, Shan Z and Teng W: Effects of isolated positive maternal thyroglobulin antibodies on brain development of offspring in an experimental autoimmune thyroiditis model. *Thyroid* 25: 551-558, 2015.
- Stemper BD, Shah AS, Pinter FA, McCrea M, Kurpad SN, Glavaski-Joksimovic A, Olsen C and Budde MD: Head rotational acceleration characteristics influence behavioral and diffusion tensor imaging outcomes following concussion. *Ann Biomed Eng* 43: 1071-1088, 2015.
- Uslu S, Kebapçı N, Kara M and Bal C: Relationship between adipocytokines and cardiovascular risk factors in patients with type 2 diabetes mellitus. *Exp Ther Med* 4: 113-120, 2012.
- Fang H, Luo X, Wang Y, Liu N, Fu C, Wang H, Fang Y, Shi W, Zhang Y, Zeng C and Wang X: Correlation between single nucleotide polymorphisms of the ACTA2 gene and coronary artery stenosis in patients with type 2 diabetes mellitus. *Exp Ther Med* 7: 970-976, 2014.
- Oršolić N, Gajski G, Garaj-Vrhovac V, Dikić D, Prskalo ZŠ and Sirovina D: DNA-protective effects of quercetin or naringenin in alloxan-induced diabetic mice. *Eur J Pharmacol* 656: 110-118, 2011.
- Roy A, Das A, Das R, Haldar S, Bhattacharya S and Haldar PK: Naringenin, a citrus flavonoid, ameliorates arsenic-induced toxicity in swiss albino mice. *J Environ Pathol Toxicol Oncol* 33: 195-204, 2014.
- Priscilla DH, Roy D, Suresh A, Kumar V and Thirumurugan K: Naringenin inhibits α -glucosidase activity: A promising strategy for the regulation of postprandial hyperglycemia in high fat diet fed streptozotocin induced diabetic rats. *Chem Biol Interact* 210: 77-85, 2014.
- Wang Y, He H, Li D, Zhu W, Duan K, Le Y, Liao Y and Ou Y: The role of the TLR4 signaling pathway in cognitive deficits following surgery in aged rats. *Mol Med Rep* 7: 1137-1142, 2013.
- Zhu X, Su B, Wang X, Smith MA and Perry G: Causes of oxidative stress in Alzheimer disease. *Cell Mol Life Sci* 64: 2202-2210, 2007.
- Kuhad A, Sethi R and Chopra K: Lycopene attenuates diabetes-associated cognitive decline in rats. *Life Sci* 83: 128-134, 2008.
- Liu YW, Zhu X, Yang QQ, Lu Q, Wang JY, Li HP, Wei YQ, Yin JL and Yin XX: Suppression of methylglyoxal hyperactivity by mangiferin can prevent diabetes-associated cognitive decline in rats. *Psychopharmacology (Berl)* 228: 585-594, 2013.
- Ahmed LA, Obaid AA, Zaki HF and Agha AM: Naringenin adds to the protective effect of L-arginine in monocrotaline-induced pulmonary hypertension in rats: Favorable modulation of oxidative stress, inflammation and nitric oxide. *Eur J Pharm Sci* 62: 161-170, 2014.
- Jeon SM, Kim HK, Kim HJ, Do GM, Jeong TS, Park YB and Choi MS: Hypocholesterolemic and antioxidative effects of naringenin and its two metabolites in high-cholesterol fed rats. *Transl Res* 149: 15-21, 2007.

39. Hermenean A, Ardelean A, Stan M, Herman H, Mihali CV, Costache M and Dinischiotu A: Protective effects of naringenin on carbon tetrachloride-induced acute nephrotoxicity in mouse kidney. *Chem Biol Interact* 205: 138-147, 2013.
40. Lichte P, Grigoleit JS, Steiner EM, Kullmann JS, Schedlowski M, Oberbeck R, Kobbe P: Low dose LPS does not increase TLR4 expression on monocytes in a human in vivo model. *Cytokine* 63: 74-80, 2013.
41. Li J, Wang Y, Zhou Y and Liu J: Gastric bypass surgery alters the mechanisms of insulin resistance in the adipose tissue of GK rats. *Mol Med Rep* 6: 1111-1116, 2012.
42. Ross JH, Hardy DC, Schuyler CA, Slate EH, Mize TW and Huang Y: Expression of periodontal interleukin-6 protein is increased across patients with neither periodontal disease nor diabetes, patients with periodontal disease alone and patients with both diseases. *J Periodontol Res* 45: 688-694, 2010.
43. Serlin Y, Levy J and Shalev H: Vascular pathology and blood-brain barrier disruption in cognitive and psychiatric complications of type 2 diabetes mellitus. *Cardiovasc Psychiatry Neurol* 2011: 609202, 2011.
44. Tsai SJ, Huang CS, Mong MC, Kam WY, Huang HY and Yin MC: Anti-inflammatory and antifibrotic effects of naringenin in diabetic mice. *J Agric Food Chem* 60: 514-521, 2012.
45. Jayaraman J, Jesudoss VA, Menon VP and Namasivayam N: Anti-inflammatory role of naringenin in rats with ethanol induced liver injury. *Toxicol Mech Methods* 22: 568-576, 2012.
46. Bodet C, La VD, Epifano F and Grenier D: Naringenin has anti-inflammatory properties in macrophage and ex vivo human whole-blood models. *J Periodontol Res* 43: 400-407, 2008.
47. Pejčić T, Stanković I, Petković TR, Borovac DN, Djordjević I and Jeftović-Stoimenov T: Peroxisome proliferator-activated receptor gamma as modulator of inflammation in pulmonary sarcoidosis. *Srp Arh Celok Lek* 141: 705-709, 2013.
48. Grygiel-Gorniak B: Peroxisome proliferator-activated receptors and their ligands: Nutritional and clinical implications - a review. *Nutr J* 13: 17, 2014.
49. Liu Q, Chen L, Hu L, Guo Y and Shen X: Small molecules from natural sources, targeting signaling pathways in diabetes. *Biochim Biophys Acta* 1799: 854-865, 2010.
50. Sharma AK, Bharti S, Ojha S, Bhatia J, Kumar N, Ray R, Kumari S and Arya DS: Up-regulation of PPAR γ , heat shock protein-27 and -72 by naringin attenuates insulin resistance, β -cell dysfunction, hepatic steatosis and kidney damage in a rat model of type 2 diabetes. *Br J Nutr* 106: 1713-1723, 2011.
51. Bahi A, Nurulain SM and Ojha S: Ethanol intake and ethanol-conditioned place preference are reduced in mice treated with the bioflavonoid agent naringin. *Alcohol* 48: 677-685, 2014.
52. Rong W, Wang J, Liu X, Jiang L, Wei F, Hu X, Han X and Liu Z: Naringin treatment improves functional recovery by increasing BDNF and VEGF expression, inhibiting neuronal apoptosis after spinal cord injury. *Neurochem Res* 37: 1615-1623, 2012.
53. Gopinath K, Prakash D and Sudhandiran G: Neuroprotective effect of naringin, a dietary flavonoid against 3-nitropropionic acid-induced neuronal apoptosis. *Neurochem Int* 59: 1066-1073, 2011.