

# Bioinformatics analysis of molecular mechanisms involved in intervertebral disc degeneration induced by TNF- $\alpha$ and IL-1 $\beta$

FENG XU<sup>1</sup>, FENG GAO<sup>2</sup>, YADONG LIU<sup>1</sup>, ZHENYU WANG<sup>1</sup>, XINMING ZHUANG<sup>1</sup>, ZHIGANG QU<sup>1</sup>,  
HUI MA<sup>3</sup>, YI LIU<sup>1</sup>, CHANGFENG FU<sup>1</sup>, QI ZHANG<sup>1</sup> and XIAOYING DUAN<sup>4</sup>

<sup>1</sup>Department of Spine Surgery, The First Hospital of Jilin University, Changchun, Jilin 130021;

<sup>2</sup>Department of Orthopaedics, The Second Hospital of Jilin University, Changchun, Jilin 130041;

<sup>3</sup>Teaching and Research Department of Health Service Management, Changchun University of Chinese Medicine, Changchun, Jilin 130117; <sup>4</sup>Department of Acupuncture and Moxibustion, The Second Hospital of Jilin University,

Changchun, Jilin 130041, P.R. China

Received March 12, 2015; Accepted January 12, 2016

DOI: 10.3892/mmr.2016.4861

**Abstract.** The present study aimed to explore the molecular mechanisms associated with intervertebral disc degeneration (IDD) induced by tumor necrosis factor (TNF)- $\alpha$  and interleukin (IL)-1 $\beta$ . The microarray dataset no. GSE42611 was downloaded from the Gene Expression Omnibus database. The differentially expressed genes (DEGs) between four experimental nucleus pulposus samples and four control nucleus pulposus samples were analyzed. Subsequently, Gene Ontology (GO) and pathway enrichment analyses of DEGs were performed, followed by protein-protein interaction (PPI) network construction and prediction of a regulatory network of transcription factor (TFs). Finally, the transcriptional regulatory network was integrated into the PPI network to analyze the network modules. A total of 246 upregulated and 290 downregulated DEGs were identified. The upregulated DEGs were mainly associated with GO terms linked with inflammatory response and apoptotic pathways, while the downregulated DEGs were mainly associated with GO terms linked with cell adhesion and pathways of extracellular matrix - receptor interaction. In the PPI network, *IL6*, *COL1A1*, *NFKB1* and *HIF1A* were hub genes, and in addition, *NFKB1* and *HIF1A* were TFs. Pathways of apoptosis and extracellular matrix - receptor interaction may have important roles in IDD

progression. *IL6*, *COL1A1* and the TFs *NFKB1* and *HIF1A* may be used as biomarkers for IDD diagnosis and treatment.

## Introduction

Intervertebral discs lie between adjacent vertebrae in the spine, forming a fibrocartilaginous joint to allow slight movement of the vertebrae (1). The layers of fibrocartilage contain the nucleus pulposus, which functions as a shock absorber dissipating compressive forces outward from its center to the surrounding annulus fibrosus (2). However, the annulus fibrosus becomes weaker with increasing age (3), resulting in intervertebral disc degeneration (IDD). Beyond age 40, >60% of individuals show symptoms of IDD; furthermore, IDD is the most common cause of disability among workers aged 18-64 years (4). IDD is characterized by decreases in intervertebral disc function and height due to cell loss through apoptosis, increased breakdown of matrix or altered matrix synthesis, with the underlying pathological processes being complex (5-7).

Numerous studies have intended to explore the molecular mechanisms involved in IDD. Vo *et al* (4) reported that IDD was a consequence of increased catabolism of the extracellular matrix (ECM), since the proteolytic degradation of ECM macromolecules led to marked structural changes of the intervertebral disc. These catabolic processes are mediated by a number of cytokines in the nucleus pulposus, among which interleukin (IL)-1 $\beta$  and tumor necrosis factor (TNF)- $\alpha$  have been suggested to have crucial roles in the development of IDD. However, the involvement of IL-1 $\beta$  and TNF- $\alpha$  in IDD has remained to be fully elucidated.

Markova *et al* (8) cultured rat intervertebral discs in the presence of IL-1 $\beta$ , TNF- $\alpha$  and serum-limiting conditions to mimic a degenerative insult to identify the differentially expressed genes (DEGs) between experimental and control groups. Their microarray data (no. GSE42611) have been deposited at the National Center of Biotechnology Information Gene Expression Omnibus (GEO) database (<http://www.ncbi.nlm.nih.gov/geo/>) and were used in the present study for a bioinformatics analysis. DEGs between experimental and control

**Correspondence to:** Dr Yi Liu, Department of Spine Surgery, The First Hospital of Jilin University, 71 Xinmin Street, Chaoyang, Changchun, Jilin 130021, P.R. China  
E-mail: liuyigk@hotmail.com

Dr Feng Gao, Department of Orthopaedics, The Second Hospital of Jilin University, 218 Ziqiang Street, Changchun, Jilin 130041, P.R. China  
E-mail: highfeng@yeah.net

**Key words:** intervertebral disc degeneration, differentially expressed genes, pathway enrichment analysis, protein-protein interaction network, transcription factor

samples were identified, Gene Ontology (GO) and pathway enrichment analyses were performed and a protein-protein interaction (PPI) network was constructed. The present study aimed to identify the DEGs and pathways associated with IDD caused by the presence of IL-1 $\beta$  and TNF- $\alpha$ .

## Materials and methods

**Affymetrix microarray data.** The microarray expression profile dataset GSE42611 (8) was downloaded from the GEO (<http://www.ncbi.nlm.nih.gov/geo/>) database based on the platform of Affymetrix Rat Gene 1.0 ST Array [transcript (gene) version; Affymetrix Inc., Santa Clara, CA, USA]. The dataset contained four experimental nucleus pulposus (ENP) samples and four control nucleus pulposus (CNP) samples. The experimental lumbar discs had been cultured in Dulbecco's modified Eagle's medium (DMEM; R&D Systems, Inc., Minneapolis, MN, USA) containing 10 ng IL-1 $\beta$  (R&D Systems, Inc.), 100 ng/ml TNF- $\alpha$  (R&D Systems, Inc.), 1% fetal bovine serum (FBS; R&D Systems, Inc.), 50  $\mu$ g/ml L-ascorbate (Cellgro; Corning Incorporated, Corning, NY, USA), 40 mM NaCl (Cellgro; Corning Incorporated), antibiotics and antimycotics (Cellgro; Corning Incorporated), while the control discs were cultured in DMEM containing 10% FBS, 50  $\mu$ g/ml L-ascorbate, 40 mM NaCl, antibiotics and antimycotics.

**Data preprocessing and differential expression analysis.** The original array data were converted into gene symbols and then subjected to background correction and quartile data normalization using the robust multiarray average (9) algorithm in the oligo package (10), which is available through BioConductor (<http://www.bioconductor.org>).

The paired Student's t-test based on the Limma package (11) in R was used to identify DEGs between ENP and CNP samples. Multiple testing correction was performed using the Benjamini-Hochberg method (12) to obtain the adjusted P-value. Subsequently, the log<sub>2</sub>-fold change (log<sub>2</sub>FC) was calculated. Only genes with an adjusted P<0.05 and a |log<sub>2</sub>FC|>1.0 were regarded as DEGs.

**GO and pathway enrichment analyses.** GO (<http://www.geneontology.org/>) (13) is a database used for unification of biological data, which comprises a structured, defined and controlled vocabulary for large-scale gene annotation. The Kyoto Encyclopedia of Genes and Genomes (KEGG; <http://www.genome.jp/>) (14) database is a collection of online databases of genomes, enzymatic pathways and biological chemicals.

The present study performed GO and KEGG pathway enrichment analyses to determine the function of DEGs using the Database for Annotation, Visualization and Integrated Discovery (<http://david.abcc.ncifcrf.gov/>) (15) online tool, which is a comprehensive functional annotation tool for associating functional terms with gene lists using a clustering algorithm. P<0.05 and a gene count >2 were set as thresholds.

**PPI network construction.** The Search Tool for the Retrieval of Interacting Genes (STRING; <http://string-db.org/>) (16) database is a pre-computed global resource, which was designed to predict functional associations between proteins. In the

present study, the STRING online tool was applied to analyze the PPIs of DEGs with experimentally validated interactions with a combined score of >0.4 considered significant.

The previously obtained biological networks indicated that the majority of the biological networks are characterized as scale-free (17). Thus, the connectivity degree was assessed by statistical analysis of networks to identify important nodes, which are being referred to as hub proteins (18).

**Prediction of regulatory networks of transcription factors (TFs).** The TRANSFAC<sup>®</sup> database (19), which is available at <http://www.gene-regulation.com>, consists of TFs, their target genes and regulatory binding sites. In the present study, the transcriptional regulatory network within the PPI network was predicted based on the TRANSFAC<sup>®</sup> database.

**Network module identification and functional enrichment analysis.** After integrating the transcriptional regulatory network into the PPI network, the modular complex detection plugin (20) in cytoscape (<http://www.cytoscape.org/>) (21) was used to identify the network modules. The selected modules with high node scores and connectivity degrees were the subjected to GO and KEGG pathway enrichment analyses.

## Results

**Identification of DEGs.** After data pre-processing, a total of 536 DEGs were obtained between ENP and CNP samples. Among these DEGs, 246 were upregulated and 290 were downregulated.

**GO and pathway enrichment analyses.** The top two clustering groups obtained by GO enrichment analysis are shown in Table I. The upregulated DEGs were mainly enriched in biological processes (BPs) associated with inflammatory response and responses to organic substances. The downregulated DEGs were enriched in the BP terms associated with cell adhesion and collagen fibril organization.

The results of the pathway enrichment analysis are shown in Table II. The upregulated DEGs were enriched in nine pathways, which included apoptotic pathways and the NOD-like receptor signaling pathway. The downregulated DEGs were also enriched in nine pathways, including pathways of ECM-receptor interaction and focal adhesion.

**PPI network construction.** Based on the STRING database, 1,345 PPI pairs were obtained and the PPI network was constructed (Fig. 1). The PPI network contained 36 DEGs with a degree of >25, referred to as hub genes, which included *IL6* (degree, 67), *COL1A1* (degree, 36), *NFKB1* (degree, 31) and *HIF1A* (degree, 26).

**Regulatory networks of TFs.** Based on the PPI network and the TRANSFAC<sup>®</sup> database, 81 PPI pairs of transcriptional regulatory interactions were obtained. In addition, seven TFs were obtained, namely *RAR- $\alpha$* , *ANPEP*, *ETS2*, *ATF3*, *EGRI*, *HIF1A* and *NFKB1*.

**Network module identification and functional enrichment analyses.** Using the MCODE plugin in cytoscape, four modules were

Table I. GO functional enrichment analysis for the upregulated and downregulated differentially expressed genes.

GO Term	Biological process	Count	P-value
Enrichment score, 7.038			
GO:0009611	Response to wounding	24	2.10x10 <sup>-8</sup>
GO:0006954	Inflammatory response	17	5.50x10 <sup>-8</sup>
GO:0006952	Defense response	21	6.67x10 <sup>-7</sup>
Enrichment score, 5.439			
GO:0010033	Response to organic substance	34	4.24x10 <sup>-7</sup>
GO:0009719	Response to endogenous stimulus	24	4.27x10 <sup>-6</sup>
GO:0009725	Response to hormone stimulus	21	2.66x10 <sup>-5</sup>
Enrichment score, 6.414			
GO:0007155	Cell adhesion	28	7.91x10 <sup>-9</sup>
GO:0022610	Biological adhesion	28	7.91x10 <sup>-9</sup>
GO:0016337	Cell-cell adhesion	12	9.22x10 <sup>-4</sup>
Enrichment score, 6.331			
GO:0030199	Collagen fibril organization	8	3.80x10 <sup>-8</sup>
GO:0030198	Extracellular matrix organization	12	1.74x10 <sup>-7</sup>
GO:0043062	Extracellular structure organization	12	1.54x10 <sup>-5</sup>

GO, gene ontology.


Table II. KEGG pathway enrichment analysis for the upregulated and downregulated DEGs.

Term	Biological process/pathway	Count	P-value
Upregulated DEGs			
rno04210	Apoptosis	9	4.15x10 <sup>-5</sup>
rno04621	NOD-like receptor signaling pathway	7	3.43x10 <sup>-4</sup>
rno04060	Cytokine-cytokine receptor interaction	11	8.06x10 <sup>-4</sup>
rno04062	Chemokine signaling pathway	9	4.53x10 <sup>-3</sup>
rno04630	Jak-STAT signaling pathway	8	5.29x10 <sup>-3</sup>
rno05222	Small cell lung cancer	6	8.70x10 <sup>-3</sup>
rno05200	Pathways in cancer	12	8.81x10 <sup>-3</sup>
rno04620	Toll-like receptor signaling pathway	6	1.21x10 <sup>-2</sup>
rno00230	Purine metabolism	7	3.06x10 <sup>-2</sup>
Downregulated DEGs			
rno04512	Extracellular matrix - receptor interaction	16	7.56x10 <sup>-12</sup>
rno04510	Focal adhesion	20	1.13x10 <sup>-9</sup>
rno05200	Pathways in cancer	14	4.34x10 <sup>-3</sup>
rno04640	Hematopoietic cell lineage	6	1.30x10 <sup>-2</sup>
rno04670	Leukocyte transendothelial migration	7	1.73x10 <sup>-2</sup>
rno05410	Hypertrophic cardiomyopathy	6	1.75x10 <sup>-2</sup>
rno04110	Cell cycle	7	2.59x10 <sup>-2</sup>
rno05219	Bladder cancer	4	2.66x10 <sup>-2</sup>
rno04115	p53 signaling pathway	5	3.08x10 <sup>-2</sup>

KEGG, Kyoto Encyclopedia of Genes and Genomes; DEG, differentially expressed gene.

obtained. Module 1 contained 29 nodes and 376 edges; however, it only contained downregulated DEGs and no TFs. Module 2 contained 20 nodes and 74 edges, and included the upregulated TFs *HIF1A* and *NFKB1* (Fig. 2). Module 3 contained 10 nodes

and 33 edges while module 4 contained 14 nodes and 42 edges. Module 2 was further subjected to GO and pathway enrichment analyses, and the results were shown in Table III. The DEGs in module 2 were predominantly enriched in GO terms associated



with the regulation of cell proliferation and wound healing. In addition, they were also enriched in pathways associated with cancer and focal adhesion.

## Discussion

leading to a decrease of life quality or even disability (22). To date, the molecular mechanisms involved in the pathology of IDD have not been fully elucidated. In the present study, a total of 246 upregulated and 290 downregulated DEGs were identified between ENP and CNP samples. Several upregulated DEGs, including *NFKB1*, were enriched in apoptotic pathways, while downregulated DEGs, including *COL1A1*, were enriched in the pathway of ECM-receptor interaction. In






Figure 2. Protein-protein interaction network of module 2. Circles and squares indicate DEGs and triangles signify transcription factors (upregulated). Circular nodes indicate upregulated DEGs and square nodes signify downregulated DEG. Arrows indicate transcriptional regulation, while lines signify protein-protein interactions. DEG, differentially expressed gene.

addition, in the PPI network, *IL6*, *COL1A1*, *NFKB1* and *HIF1A* were hub genes with a high connectivity degree. Furthermore, module analysis indicated that the DEGs in module 2 were predominantly enriched in GO terms associated with cell proliferation and pathways associated with cancer. The results suggested that these genes and pathways may be candidates for IDD diagnosis and treatment.

A central feature of IDD is loss of tissue cellularity, which may be the result of programmed cell death (23). The results of the pathway enrichment analysis indicated that apoptosis was the most significant pathway. Apoptosis is a type of programmed cell death and is characterized by chromosomal concentration, DNA degradation and cell shrinkage (24). Apoptosis acts as a quality control mechanism for the maintenance of tissue homeostasis by eliminating defective cells (25). Apoptosis not only exists in certain physiological processes, but is also involved in numerous pathological degenerative diseases, including neurodegeneration and IDD (26,27). Boos *et al* (28) reported that the initiation of IDD is associated with the changes of intervertebral disc cell behavior, including increased cell death. Gruber and Hanley (29) suggested that a large proportion of intervertebral disc cells underwent apoptosis in patients with degenerated intervertebral discs. Therefore, pathways of apoptosis may be involved in the progression of IDD in the ENP.

In addition, the present study found that the TF *NFKB1* was upregulated in the pathway of apoptosis. In particular, *NFKB1*

was a hub gene in the PPI network. *NFKB1* encodes the nuclear factor kappa B p105/p50 isoforms (30). NFKB proteins are a family of transcription factors that have critical roles in the regulation of various biological defense processes, including immune responses, cell-growth control and apoptosis (31). Inappropriate activation of NFKB has been associated with numerous inflammatory diseases (30). Wuertz *et al* (32) have reported that IDD is characterized not only by an imbalance between anabolic and catabolic processes, but also by inflammatory mechanisms. Of note, IL-1 $\beta$  and TNF- $\alpha$  are pro-inflammatory cytokines, which have been detected in the degenerated disc (33). In conclusion, the upregulation of *NFKB1* may have been stimulated by IL-1 $\beta$  and TNF- $\alpha$ , which suggests that the TF *NFKB1* may be an important biomarker for IDD.

In the present study, several downregulated DEGs, including *COL1A1*, were enriched in the pathway of ECM - receptor interaction; furthermore, *COL1A1* was a hub gene in the PPI network. Over the previous decade, IDD research has focused on elucidating the mechanisms of ECM degradation, as it causes marked structural changes, including dehydration and fibrosis of the nucleus pulposus, disorganization of the annulus fibrosus and calcification of the cartilaginous end plates (28). These changes ultimately lead to structural failure in IDD. Of note, the protein collagen type I, alpha 1, encoded by *COL1A1*, is a major ECM component (34). Col1a1 has been

Table III. GO and KEGG pathway enrichment analyses for the differentially expressed genes in module 2.

Term	Biological process/pathway	Count	P-value
GO			
GO:0042127	Regulation of cell proliferation	12	6.47x10 <sup>-10</sup>
GO:0009611	Response to wounding	10	4.98x10 <sup>-9</sup>
GO:0042060	Wound healing	7	1.89x10 <sup>-7</sup>
GO:0001666	Response to hypoxia	7	3.28x10 <sup>-7</sup>
GO:0070482	Response to oxygen levels	7	4.70x10 <sup>-7</sup>
GO:0009725	Response to hormone stimulus	9	4.75x10 <sup>-7</sup>
GO:0031099	Regeneration	6	8.12x10 <sup>-7</sup>
GO:0009719	Response to endogenous stimulus	9	1.15x10 <sup>-6</sup>
GO:0008284	Positive regulation of cell proliferation	8	1.20x10 <sup>-6</sup>
GO:0001817	Regulation of cytokine production	6	3.26x10 <sup>-6</sup>
KEGG			
rno05200	Pathways in cancer	8	7.66x10 <sup>-6</sup>
rno05212	Pancreatic cancer	4	7.37x10 <sup>-5</sup>
rno04510	Focal adhesion	5	1.45x10 <sup>-3</sup>
rno04150	mTOR signaling pathway	3	8.56x10 <sup>-3</sup>
rno04621	NOD-like receptor signaling pathway	3	1.16x10 <sup>-2</sup>
rno04115	p53 signaling pathway	3	1.31x10 <sup>-2</sup>
rno05211	Renal cell carcinoma	3	1.42x10 <sup>-2</sup>
rno04060	Cytokine-cytokine receptor interaction	4	1.43x10 <sup>-2</sup>
rno05220	Chronic myeloid leukemia	3	1.67x10 <sup>-2</sup>
rno05222	Small cell lung cancer	3	2.02x10 <sup>-2</sup>
rno05215	Prostate cancer	3	2.35x10 <sup>-2</sup>

KEGG, Kyoto Encyclopedia of Genes and Genomes; GO, gene ontology.

found in most connective tissues and is abundant in bone (35). Several studies have demonstrated that the expression of genes encoding connective tissue components, particularly collagen, can be affected by numerous cytokines, including IL-1 and TNF- $\alpha$  (36,37). Mori *et al* (38) found that TNF- $\alpha$  decreased *Colla1* expression through suppressing promoter activity of *Colla1*. Feng *et al* (39) suggested that *Colla1* can provide tensile strength in the annulus fibrosus of the intervertebral disc. Thus, the downregulated expression of *COL1A1* in the present study may be regulated by IL-1 $\beta$  and TNF- $\alpha$ , resulting in the initiation of IDD.

In the PPI network, *IL6* and *HIF1A* were hub genes with a high connectivity degree, while *HIF1A* was also a TF. *IL6* together with IL-1 $\alpha$ , IL-1 $\beta$  and TNF- $\alpha$  are among the most potent catabolic cytokines and pro-inflammatory mediators as mentioned above (40). Noponen-Hietala *et al* (41) reported that the features of pain, tissue destruction and inflammation in IDD were linked with the functions of IL6, and IL6 was also associated with IDD-associated radiculopathy. Therefore, IL6 is not the sole pro-inflammatory cytokine in the pathogenesis of IDD and appears to be important in the mediation of pain in IDD.

In the present study, the TF of *HIF1A* was also found in module 2 and was involved in GO terms associated with cell proliferation. HIF1 functions as a transcriptional regulator of the adaptive response to hypoxia and activates transcription

of certain genes involved in energy metabolism, angiogenesis and apoptosis (42). Wu *et al* (43) found that Hif1 $\alpha$  has an important role in the metabolism and synthesis of ECM and is a pivotal contributor to the survival of nucleus pulposus cells. Therefore, *HIF1A* may be involved in the development of IDD.

In conclusion, the present study provided a comprehensive bioinformatics analysis of the molecular mechanisms of IDD induced by IL-1 $\beta$  and TNF- $\alpha$ . Pathways of apoptosis and ECM - receptor interaction as well as their enriched DEGs may have important roles in the development and progression of IDD. In addition, the TFs *NFKB1* and *HIF1A* and the DEGs *COL1A1* and *IL6* are hypothesized to interact with IL-1 $\beta$  and TNF- $\alpha$  to participate in the development of IDD. However, further bioinformatics and experimental studies with larger sample sizes are required to confirm the results of the present study.

### Acknowledgements

The present study was supported by the National Natural Youth Science Foundation of China (grant no. 31300778).

### References

1. Bolger C: Spine disorders (medical and surgical management). The Surgeon 8: 185, 2010.

2. McCann MR, Tamplin OJ, Rossant J and Séguin CA: Tracing notochord-derived cells using a Noto-cre mouse: Implications for intervertebral disc development. *Dis Model Mech* 5: 73-82, 2012.
3. Roberts S, Evans H, Trivedi J and Menage J: Histology and pathology of the human intervertebral disc. *J Bone Joint Surg Am* 88 (Suppl 2): 10-14, 2006.
4. Vo NV, Hartman RA, Yurube T, Jacobs LJ, Sowa GA and Kang JD: Expression and regulation of metalloproteinases and their inhibitors in intervertebral disc aging and degeneration. *Spine J* 13: 331-341, 2013.
5. Adams MA and Roughley PJ: What is intervertebral disc degeneration and what causes it? *Spine (Phila Pa 1976)* 31: 2151-2161, 2006.
6. Johnson WE and Roberts S: 'Rumours of my death may have been greatly exaggerated': A brief review of cell death in human intervertebral disc disease and implications for cell transplantation therapy. *Biochem Soc Trans* 35: 680-682, 2007.
7. Le Maitre C, Pockert A, Buttle D, Freemont A and Hoyland J: Matrix synthesis and degradation in human intervertebral disc degeneration. *Biochem Soc Trans* 35: 652-655, 2007.
8. Markova DZ, Kepler CK, Addya S, Murray HB, Vaccaro AR, Shapiro IM, Anderson DG, Albert TJ and Risbud MV: An organ culture system to model early degenerative changes of the intervertebral disc II: Profiling global gene expression changes. *Arthritis Res Ther* 15: R121, 2013.
9. Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U and Speed TP: Exploration, normalization and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4: 249-264, 2003.
10. Carvalho BS and Irizarry RA: A framework for oligonucleotide microarray preprocessing. *Bioinformatics* 26: 2363-2367, 2010.
11. Diboun I, Wernisch L, Orenge CA and Koltzenburg M: Microarray analysis after RNA amplification can detect pronounced differences in gene expression using limma. *BMC Genomics* 7: 252, 2006.
12. Benjamini Y and Hochberg Y: Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc B (Methodological)* 57: 289-300, 1995.
13. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, *et al*: Gene ontology: Tool for the unification of biology. The gene ontology consortium. *Nat Genet* 25: 25-29, 2000.
14. Kanehisa M and Goto S: KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Res* 28: 27-30, 2000.
15. Huang DW, Sherman BT, Tan Q, Collins JR, Alvord WG, Roayaei J, Stephens R, Baseler MW, Lane HC and Lempicki RA: The DAVID Gene functional classification tool: A novel biological module-centric algorithm to functionally analyze large gene lists. *Genome Biol* 8: R183, 2007.
16. Von Mering C, Huynen M, Jaeggi D, Schmidt S, Bork P and Snel B: STRING: A database of predicted functional associations between proteins. *Nucleic Acids Res* 31: 258-261, 2003.
17. Albert R and Barabási AL: Statistical mechanics of complex networks. *Rev Mod Phys* 74: 47-97, 2002.
18. He X and Zhang J: Why do hubs tend to be essential in protein networks? *PLoS Genet* 2: e88, 2006.
19. Matys V, Fricke E, Geffers R, Gössling E, Haubrock M, Hehl R, Hornischer K, Karas D, Kel AE, Kel-Margoulis OV, *et al*: TRANSFAC: Transcriptional regulation, from patterns to profiles. *Nucleic Acids Res* 31: 374-378, 2003.
20. Bader GD and Hogue CW: An automated method for finding molecular complexes in large protein interaction networks. *BMC Bioinformatics* 4: 2, 2003.
21. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B and Ideker T: Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Res* 13: 2498-2504, 2003.
22. Akhatib B, Onnerfjord P, Gawri R, Ouellet J, Jarzem P, Heinegård D, Mort J, Roughley P and Haglund L: Chondroadherin fragmentation mediated by the protease HTRA1 distinguishes human intervertebral disc degeneration from normal aging. *J Biol Chem* 288: 19280-19287, 2013.
23. Ariga K, Miyamoto S, Nakase T, Okuda S, Meng W, Yonenobu K and Yoshikawa H: The relationship between apoptosis of endplate chondrocytes and aging and degeneration of the intervertebral disc. *Spine (Phila Pa 1976)* 26: 2414-2420, 2001.
24. Lockshin RA and Zakeri Z: Apoptosis, autophagy and more. *Int J Biochem Cell Biol* 36: 2405-2419, 2004.
25. Fuchs Y and Steller H: Programmed cell death in animal development and disease. *Cell* 147: 742-758, 2011.
26. Moreno S, Imbrogliini V, Ferraro E, Bernardi C, Romagnoli A, Berrebi AS and Cecconi F: Apoptosome impairment during development results in activation of an autophagy program in cerebral cortex. *Apoptosis* 11: 1595-1602, 2006.
27. Ye W, Xu K, Huang D, Liang A, Peng Y, Zhu W and Li C: Age-related increases of macroautophagy and chaperone-mediated autophagy in rat nucleus pulposus. *Connect Tissue Res* 52: 472-478, 2011.
28. Boos N, Weissbach S, Rohrbach H, Weiler C, Spratt KF and Nerlich AG: Classification of age-related changes in lumbar intervertebral discs: 2002 Volvo Award in basic science. *Spine (Phila Pa 1976)* 27: 2631-2644, 2002.
29. Gruber HE and Hanley EN Jr: Analysis of aging and degeneration of the human intervertebral disc: Comparison of surgical specimens with normal controls. *Spine (Phila Pa 1976)* 23: 751-757, 1998.
30. Karban AS, Okazaki T, Panhuysen CI, Gallegos T, Potter JJ, Bailey-Wilson JE, Silverberg MS, Duerr RH, Cho JH, Gregersen PK, *et al*: Functional annotation of a novel NFKB1 promoter polymorphism that increases risk for ulcerative colitis. *Hum Mol Genet* 13: 35-45, 2004.
31. Baldwin AS: Series introduction: The transcription factor NF-kappaB and human disease. *J Clin Invest* 107: 3-6, 2001.
32. Wuertz K, Vo N, Kleitsas D and Boos N: Inflammatory and catabolic signalling in intervertebral discs: The roles of NF-κB and MAP kinases. *Eur Cell Mater* 23: 103-119, 2012.
33. Akeda K, An H, Gemba T, Okuma M, Miyamoto K, Chujo T, Kitahara S and Masuda K: A new gene therapy approach: *In vivo* transfection of naked NFκB decoy oligonucleotide restored disc degeneration in the rabbit annular needle puncture model. *Trans Orthop Res Soc* 30: 45, 2005.
34. Wu F and Chakravarti S: Differential expression of inflammatory and fibrogenic genes and their regulation by NF-kappaB inhibition in a mouse model of chronic colitis. *J Immunol* 179: 6988-7000, 2007.
35. Alberts B, Johnson A, Lewis J, Raff M, Roberts K and Walter P: The extracellular matrix of animals, 2002.
36. Postlethwaite AE, Raghow R, Stricklin GP, Poppleton H, Seyer JM and Kang AH: Modulation of fibroblast functions by interleukin 1: Increased steady-state accumulation of type I procollagen messenger RNAs and stimulation of other functions but not chemotaxis by human recombinant interleukin 1 alpha and beta. *J Cell Biol* 106: 311-318, 1988.
37. Scharffetter K, Heckmann M, Hatamochi A, Mauch C, Stein B, Riethmüller G, Ziegler-Heitbrock HW and Krieg T: Synergistic effect of tumor necrosis factor-alpha and interferon-gamma on collagen synthesis of human skin fibroblasts in vitro. *Expe Cell Res* 181: 409-419, 1989.
38. Mori K, Hatamochi A, Ueki H, Olsen A and Jimenez S: The transcription of human alpha 1 (I) procollagen gene (COL1A1) is suppressed by tumour necrosis factor-alpha through proximal short promoter elements: Evidence for suppression mechanisms mediated by two nuclear-factor binding sites. *Biochem J* 319: 811-816, 1996.
39. Feng H, Danfelter M, Strömquist B and Heinegård D: Extracellular matrix in disc degeneration. *J Bone Joint Surg Am* 88 (Suppl 2): 25-29, 2006.
40. Benoist M: The natural history of lumbar disc herniation and radiculopathy. *Joint Bone Spine* 69: 155-160, 2002.
41. Noponen-Hietala N, Virtanen I, Karttunen R, Schwenke S, Jakkula E, Li H, Merikivi R, Barral S, Ott J, Karppinen J and Ala-Kokko L: Genetic variations in IL6 associate with intervertebral disc disease characterized by sciatica. *Pain* 114: 186-194, 2005.
42. Zou H, Kang X, Pang LJ, Hu W, Zhao J, Qi Y, Hu J, Liu C, Li H, Liang W, *et al*: Xp11 translocation renal cell carcinoma in adults: A clinicopathological and comparative genomic hybridization study. *Int J Clin Exp Pathol* 7: 236-245. eCollection 2014, 2013.
43. Wu WJ, Zhang XK, Zheng XF, Yang YH, Jiang SD and Jiang LS: SHH-dependent knockout of HIF-1 alpha accelerates the degenerative process in mouse intervertebral disc. *Int J Immunopathol Pharmacol* 26: 601-609, 2013.