

Increased expression of dermatopontin and its implications for testicular dysfunction in mice

JUN CAI^{1,2*}, WEIJIA LIU^{1*}, JIE HAO³, MAOXIN CHEN¹ and GANG LI¹

¹Institute of Life Sciences, Chongqing Medical University; ²Department of Clinical Laboratory, Chongqing Provincial Crops Hospital of PAP; ³Experimental Research Center, The First Affiliated Hospital, Chongqing Medical University, Chongqing 400016, P.R. China

Received March 13, 2015; Accepted January 11, 2016

DOI: 10.3892/mmr.2016.4879

Abstract. An array of specific and non-specific molecules, which are expressed in the testis, have been demonstrated to be responsible for testicular function. Our previous study revealed that dermatopontin (DPT) is expressed in Sertoli cells of the testis, however, its roles in testicular function remains somewhat elusive. In the present study, CdCl₂- and busulfan-induced testicular dysfunction models were used to investigate the implications of DPT expression for testicular function. The mRNA and protein expression levels of DPT were detected using reverse transcription-quantitative polymerase chain reaction and western blotting, respectively. A negative correlation was observed between testicular damage and the expression of DPT, which suggested that an increase in DPT expression may be a marker for testicular dysfunction. This result was corroborated by the finding that transgenic mice exhibiting Sertoli cell-specific overexpression of DPT exhibited damage to their testicular morphology. Additionally, DPT overexpression in the testis affected the expression levels of claudin-11 and zonula occludens-1, which indicated that DPT may affect testicular function by affecting the integrity of the blood-testis barrier (BTB). In conclusion, the present study provided evidence to suggest that DPT may be indicative of mouse testicular dysfunction, since increased expression may be associated with damage to the BTB.

Introduction

Infertility is a growing problem worldwide. The term 'infertility' refers to the failure of a couple to conceive

following 12 months of regular intercourse. Infertility affects 10-15% of couples, and a male factor is responsible for ~50% of infertility cases (1,2). Numerous reports have shown that greater exposure to electromagnetic radiation (3), smoking (4), heat (5), the consumption of certain drugs (6), exposure to heavy metals (7-9), exposure to pesticides (10,11), and biological hazards may partly explain the mechanisms underlying male infertility.

The majority of causes of male infertility are considered to be associated with abnormal spermatogenesis and failures in sperm function (12). Mammalian spermatogenesis, which is a complex and highly regulated process of germ cell proliferation and differentiation, which occurs within the seminiferous tubules of the testes, involves dynamic interactions between the developing germ cells and the Sertoli cells. The Sertoli cells provide structural and nutritional support for the germ cells. An array of specific and non-specific molecules that are expressed by the Sertoli cells, including proteases, protease inhibitors, signaling molecules, growth factors and cell adhesion molecules, have been identified, which are responsible for the regulation of spermatogenesis (13-15). Numerous harmful factors can target the molecules that are expressed by the Sertoli cells, and disruptions to these molecules may arrest germ cell production and maturation or dysfunctional spermatozoa may be produced, resulting in male infertility (16,17). For example, the dibutyl phthalate-induced collapse of Sertoli cell vimentin filaments may lead to the detachment of spermatogenic cells, and the detached cells may undergo apoptosis since they have lost support from the Sertoli cells (18). Zhang and Lui (19) showed that nectin-2 is a direct molecular target for cadmium and proposed that the dysregulation of nectin-2 in the Sertoli cells may explain cadmium-induced male infertility (19).

Dermatopontin (DPT), a 22 kDa non-collagenous extracellular matrix protein, which promotes cellular adhesion and extracellular matrix assembly, initially co-purifies with decorin from bovine dermal extracts (20-22). Western blotting and northern blot analysis indicates that DPT is expressed in the skin, skeletal muscle, bone, cartilage and other tissue types (22,23). DPT reportedly modifies the behavior of transforming growth factor (TGF)- β on mink lung epithelial cells via its interaction with decorin within the extracellular matrix *in vivo* (24). Furthermore, a reduction in the expression of DPT is a molecular link between uterine leiomyomas and keloids.

Correspondence to: Professor Gang Li, Institute of Life Sciences, Chongqing Medical University, 1 Yixueyuan Road, Yuzhong, Chongqing 400016, P.R. China
E-mail: lg168sn@yahoo.com

*Contributed equally

Key words: dermatopontin, busulfan, CdCl₂, testicular dysfunction, blood-testis barrier

Light microscopy has demonstrated that *Dpt*-null corneas from 2-month-old mice exhibit a 24% reduction in the average stromal thickness compared with the corneas from wild-type mice, which suggests that DPT is important in collagen fibril organization (25). Additionally, DPT may be involved in the pathogenesis and growth of prostate cancer cells (26).

Our previous study demonstrated that DPT is expressed in the Sertoli cells within the testis (27). An *in vitro* study showed that DPT is a novel regulator of the CdCl₂-induced reduction in claudin-11 expression, which suggests that DPT may be associated with testicular dysfunction. Therefore, the present study aimed to further investigate the implications of DPT expression for testicular function.

Materials and methods

Materials. The protein extraction kit and the Beyo-ECL Plus western blotting reagents were purchased from Beyotime Institute of Biotechnology (Jiangsu, China). The mouse monoclonal DPT antibody (cat. no. sc-376863) was purchased from Santa Cruz Biotechnology, Inc. (Dallas, TX, USA). The anti-claudin-11 (cat. no. 36-4500) and anti-zonula occludens (ZO-1; cat. no. 61-7300) antibodies were purchased from Thermo Fisher Scientific, Inc. (Waltham, MA, USA). All secondary antibodies were purchased from ZsBio (Beijing, China). The FLAG-tag antibody (cat. no. 20543-1-AP) was purchased from Proteintech (Wuhan, China). The SYBR[®] PrimeScript[®] RT-PCR kit (Perfect Real Time) was purchased from Takara Bio, Inc. (Liaoning, China). CdCl₂ and busulfan were purchased from Sigma-Aldrich (St. Louis, MO, USA). The remaining chemicals were purchased from Sangon Biotech (Shanghai, China).

Animal experiments. A total of 90 C57BL mice, aged between 3 and 4 months (weight, 25–32 g) were obtained from the Experimental Animal Centre at Chongqing Medical University (Chongqing, China). The animals were maintained on a 12 h light/dark cycle, with free access to food and water. The mice were randomly divided into groups containing 6 mice. The mice received 30 mg/kg⁻¹ busulfan or 3.5 mg/kg⁻¹ CdCl₂, which were administered via intraperitoneal injection. Mice were sacrificed 5, 15, 25, 35 or 70 days following treatment with an overdose of 6% chloral hydrate. The left testes were removed and were fixed in a formaldehyde solution at room temperature. The right testes were removed for protein and RNA extraction. All procedures were approved by the Animal Care and Use Committee of Chongqing Medical University (Chongqing, China).

Total RNA extraction and the reverse transcription-quantitative polymerase chain reaction (RT-qPCR). The total RNA was extracted from the testes using TRIzol[®] reagent (Thermo Fisher Scientific, Inc.), according to the manufacturer's protocol. RNA concentrations were quantified using a Nanodrop 2000 spectrophotometer (Nanodrop Technologies, Wilmington, DE, USA). Aliquots of the total RNA (1 µg) from each sample were reverse transcribed into cDNA using the PrimeScript[®] RT Enzyme (Takara Bio, Inc.), according to the manufacturer's protocol.

The RT-qPCR was performed in a thermal cycler (CFX96; Bio-Rad Laboratories, Inc. Hercules, CA, USA) using the SYBR[®] PrimeScript[®] RT-PCR kit (Perfect Real

Time), according to the manufacturer's protocol. β-actin served as the internal control. The primers used were as follows: DPT, forward: 5'-GGTGGCTACGGGTACCCA TA-3' and reverse: 5'-GTCAGAGCCTTCCTTCTTGC-3'; β-actin, forward: 5'-TCGTGCGTGACATCAAAGAG-3' and reverse: 5'-CAAGAAGGAAGGCTGGAAAA-3'. The primers were synthesized by Sangon Biotech. The reaction solution (25 µl) contained 1 µl cDNA, 10.5 µl water, 12.5 µl SYBR[®] Premix Ex Taq[™] and 0.5 µl of 10 µM forward and reverse primers. The PCR thermal cycling parameters were 1 min at 94°C, followed by 40 cycles of 10 sec at 94°C, 15 sec at 58°C and 15 sec at 72°C. The relative expression of DPT was normalized against β-actin and was calculated using the 2^{-ΔΔC_t} method.

Western blotting. The total protein was extracted from the testes using radioimmunoprecipitation assay lysis buffer, and the concentrations were determined using the bicinchoninic acid assay. Protein samples (40 µg) were separated on a 10% sodium dodecyl sulfate-polyacrylamide gel and were transferred onto polyvinylidene difluoride membranes. The membranes were blocked with 5% non-fat milk for 2 h at 37°C, then incubated with the primary antibodies against DPT (1:500, mouse monoclonal), claudin-11 (1:300; rabbit polyclonal), ZO-1 (1:300, rabbit polyclonal) and FLAG (1:1000, rabbit polyclonal) overnight at 4°C. Following incubation, the membranes were washed briefly with Tris-buffered saline containing 0.1% Tween-20 (100mM Tris, 0.9% NaCl, 0.1% Tween-20; pH 7.4), and were subsequently incubated for 2 h at room temperature with the following corresponding secondary antibody: Horseradish peroxidase (HRP) conjugated-anti-rabbit immunoglobulin G (IgG; 1:3,000; cat. no. ZB-2301) and HRP-anti-mouse IgG (1:3,000; cat. no. ZB-2305). The proteins were detected using the Beyo-ECL kit, according to the manufacturer's protocol. The densities of bands were analyzed using Quantity One (version 4.6.2; Bio-Rad Laboratories, Inc.) and were normalized against β-actin, which served as the loading control.

Histopathological examination. Routine histology was performed to assess the statuses of the testes. The formalin-fixed testes were embedded in paraffin and 4 µm sections were cut using a microtome (Leica RM2135; Wetzlar, Germany). The sections were de-waxed and were re-hydrated through a descending series of alcohol concentrations to distilled water. The tissue sections were subsequently stained with hematoxylin and eosin. The samples were analyzed for changes in testicular morphology and structure using light microscopy (Olympus BX51; Olympus, Tokyo, Japan).

Vector constructs. To generate transgenic mice with a Sertoli cell-specific overexpression of DPT, a fragment of the Müllerian-inhibiting substance (MIS) upstream promoter, which was previously validated to be sufficient for the expression of growth hormone (28), nuclear receptor subfamily 0, group B, member 1 (29) and Smad4 (30) in Sertoli cells, was used. The transgenic construct contained the following components: The upstream promoter region (-180 to +1) of the MIS, FLAG tags, the complete open reading frame of mouse DPT and the simian virus (SV)40 polyadenylation poly (A) sequence. The MIS promoter fragment was amplified from the




Figure 1. Hematoxylin and eosin staining of paraffin embedded testis tissue. Histology of the testes from mice treated with (A-F) 30 mg/kg⁻¹ busulfan or (G-L) 3.5 mg/kg⁻¹ CdCl₂ at (B and H) 5, (C and I) 15, (D and J) 25, (E and K) 35 and (F and L) 70 days following treatment (scale bar, 50 μ m). (A and G) Samples from 4 month old control mice.

C57BL mice genomic deoxyribonucleic acid (DNA) using the primers MIS-FLAG-F: 5'-GTCGACCTCAGGCCTCTGCAGTTATGGG-3' and MIS-FLAG-R: 5'-GAGGTCCATCTTGTCATCGTCGTCCTTGTAGTCCATGGTGGTACAGCAAG-3'. To clone DPT from the testes of C57BL mice and to add the FLAG epitope (DYKDDDDK) to the DPT protein, the primers FLAG-Dpt-F: 5'-CCACCATGGACTACAAGGACGACGATGACAAGATGGACCTCACTCTTC TG-3' and FLAG-Dpt-R: 5'-TGGCTGGCAACTAGAAGGCACAGCTAAACGTTTTCGAATTCGAGTCGTA-3' were used. The SV40 fragment was amplified from the pcDNA3.1 vector using the primers Dpt-SV40-F1: 5'-GAATTCGAAAACGTTTAGTCTAGATAAGTAATGAT-3' and Dpt-SV40-R1: 5'-GATCCTCTGGAGATACAGACATGATAAGATACATTG-3'. Subsequently, the transgenic cassette, MIS-FLAG-DPT-SV40, was generated by in-fusion PCR using the primers MIS-Dpt-F1: 5'-ATAATCAATGTCAACCCTCAGGCCTCTGCAGTTA-3' and Dpt-SV40-R1: 5'-GATCCTCTGGAGATACAGACATGATAAGATACATTG-3'. The produced fragment was cloned into an 18-T simple vector (Takara Bio, Inc.) for sequencing.

Generation of mice with overexpression of DPT. The purified transgenic constructs were microinjected into the pronuclei of

fertilized oocytes from C57BL mice. To identify the founder transgenic mice, tail biopsies were collected for genomic DNA isolation using the Universal Genomic DNA Extraction kit (Takara Bio, Inc.) when the mice were 2 weeks of age. The resulting genomic DNA samples were screened by PCR using the primers, MIS-FLAG-F and Dpt-SV40-R1. The amplified products were resolved using 1.5% agarose gel electrophoresis.

Immunohistochemical detection of the FLAG-tagged DPT protein. Testicular sections (5 μ m) were deparaffinized in xylene and were hydrated in a descending series of ethanol concentrations, which was followed by antigen retrieval in 0.01 mol/l citrate buffer (pH 6.0). Non-specific background was eliminated by blocking with goat serum buffer (ZsBio) at room temperature for 1 h, followed by incubation of the slides with anti-FLAG antibody (1:100; rabbit polyclonal) overnight at 4°C. The samples were subsequently incubated for 1 h at room temperature with goat-anti-rabbit immunoglobulin G secondary antibody (1:1,000; polyclonal; cat. no. ZB-2301). Following washing with phosphate-buffered saline, containing 0.1% Tween-20, the sections were incubated with diaminobenzidine and were counterstained with hematoxylin. The tissue sections were examined and images were captured using a light microscope (Olympus).




Figure 2. Effect of 30 mg/kg⁻¹ busulfan or 3.5 mg/kg⁻¹ CdCl₂ on the expression of DPT in the testes of mice. The mice were administered either 30 mg/kg⁻¹ busulfan or 3.5 mg/kg⁻¹ CdCl₂ via intraperitoneal injection and were sacrificed 5, 15, 25, 35 and 70 days later (n=6 per time point). Testicular lysates were obtained for immunoblotting, and RNA was extracted for RT-qPCR. (A and B) The mRNA expression levels of DPT were determined using RT-qPCR. The expression levels were normalized against that of β-actin and the mRNA expression within the control group was set to 1. (C and D) The protein expression levels of DPT were determined by western blotting. β-actin was used as a loading control. (E and F) The relative protein expression levels of DPT in the mouse testicular tissue was normalized against β-actin and the protein expression level within the control group was set to 1. The results are presented as the mean ± standard error of the mean (*P<0.05; ^bP<0.01, compared with the control group). DPT, dermatopontin; RT-qPCR, reverse transcription-quantitative polymerase chain reaction; d, days.

Statistical analysis. The data are expressed as the mean ± standard error. The statistical differences were evaluated using one-way analysis of variance and Newman-Keuls test. P<0.05 was considered to indicate a statistically significant difference.

Results

CdCl₂- or busulfan-induced testicular morphology damage. Routine histology was performed to assess spermatogenesis in the mice following 30 mg/kg⁻¹ busulfan or 3.5 mg/kg⁻¹ CdCl₂ treatment (Fig. 1). No histopathological changes were observed in the testes of the mice in the control groups (Fig. 1A and G). After 5 days of treatment with busulfan, the mice exhibited no obvious changes in the morphology of the epithelium of the seminiferous tubules compared with the control mice (Fig. 1B). After 15 days treatment with busulfan, the seminiferous tubule epithelium had become thinner, and exhibited the detachment of the spermatogenic cells (Fig. 1C). At 25 days following treatment with busulfan, 15% of the seminiferous tubules contained no germ cells (Fig. 1D) and 35 days following treatment with busulfan, 40% of the seminiferous tubules contained no germ cells (Fig. 1E). At 70 days following treatment with busulfan, a

histological analysis determined that the majority of the testes contained germ cells and that regeneration was occurring in the seminiferous tubules (Fig. 1F).

In the CdCl₂-treated mice, the histological analysis determined that the basement membrane of the seminiferous tubules was discontinuous, and that secondary spermatocytes and round spermatids were present in the tubules' lumens at 5 days post-treatment (Fig. 1H). At 15 days following treatment with CdCl₂, severe necrosis and degeneration of the seminiferous tubules were apparent, with 20% of the seminiferous tubules exhibiting germ cell losses (Fig. 1I). At 25 days following treatment with CdCl₂, the majority of the seminiferous tubules exhibited severe histopathological changes (Fig. 1J). Between 35 and 70 days following treatment with CdCl₂, the structure of the testes exhibited more severe disorganization and the degree of fibrosis within the testes appeared to increase in a time-dependent manner (Fig. 1K and L).

Effects of busulfan or CdCl₂ treatment on the expression of DPT in the testes of mice. The effects of busulfan or CdCl₂ treatment on the expression levels of DPT in the testes of mice were determined using PCR and western blotting. At 5 days following treatment, the busulfan-treated group exhibited no change in the expression of DPT (P>0.05). Between 15 and 25 days




Figure 3. Generation and analysis of the DPT-TG mice. (A) Polymerase chain reaction analysis of the genomic integration of the mouse DPT transgene. (B) The protein expression levels of DPT, FLAG-DPT and β -actin in the testes of DPT-TG mice at the age of 4 months were assessed by western blotting. (C) The relative expression levels of DPT and FLAG-DPT in the testes of the mice were normalized against β -actin from five different samples, and are presented as the mean \pm standard error or the mean. The protein level within the control group was set to 1 ($^bP < 0.01$, compared with the control group). (D and E) Immunolocalization of the FLAG-DPT protein in the testes of adult mice was determined. The sections of testicular tissue were incubated with (D) an anti-FLAG antibody and (E) the control immunoglobulin G1, and the nuclei were counterstained with hematoxylin. Specific signals were identified in the Sertoli cells. Magnification x400. DPT, dermatopontin; N, negative control; P, positive control; TG, transgenic.




Figure 4. Overexpression of DPT in the Sertoli cells led to the disruption of the testicular architecture, which that was visualized as germ cell loss and the formation of vacuoles. Hematoxylin and eosin-stained sections of the testicular tissue from 4-month-old (A) control mice and (B) DPT transgenic mice (scale bar, 50 μ m). The red arrows indicate the desquamation of the spermatogenic cells in the seminiferous tubules and the green arrows indicate the formation of vacuoles. DPT, dermatopontin.

following treatment with busulfan, the DPT levels gradually increased with time. Between 35 and 70 days following treatment with busulfan, the DPT levels gradually normalized (Fig. 2A, C, and E).

The pattern of DPT expression in the mice administered CdCl_2 differed from that observed in the busulfan-treated mice. Time-dependent changes in the expression of DPT mRNA (Fig. 2B) and protein (Fig. 2D and F) were observed in the mouse testes in response to treatment with CdCl_2 .

Establishment of a transgenic mouse with DPT overexpression.

Two males (lines 37 and 45) positive for the mouse DPT transgene (Fig. 3A) were selected for further breeding. The 4-month-old mice, originally derived from the two male founder mice, exhibited an upregulation of the protein expression of DPT in their testes, however, the control litter mate did not (Fig. 3B and C). No apparent differences were observed in the expression of testicular DPT between the transgenic lines. Since the transgenic construct contains a FLAG tag,




Figure 5. Overexpression of DPT in the Sertoli cells reduces the expression levels of ZO-1 and claudin-11 in the testes. (A) Representative western blotting analyses of ZO-1 and claudin-11 in the control mice and the mice with DPT overexpression. (B) The relative expression levels of ZO-1 and claudin-11 were determined using densitometric analysis of the data normalized against the β -actin from five different samples. The data are presented as the mean \pm standard error of the mean ($^*P < 0.05$, compared with the control group). The protein expression in the control group was set to 1. DPT, dermatopontin; TG, transgenic; ZO-1, zonula occludens-1.

the present study also tested for the expression of the FLAG epitope in the testes, and the results revealed that while no expression of the FLAG epitope occurred in the control mice, its protein was detected in the testes from the transgenic animals at a molecular weight that was appropriate for DPT (Fig. 3B). Histological analysis also immunolocalized the FLAG epitope in the Sertoli cells that were within the sections of the testicular tissue from the DPT overexpression mice (Fig. 3D), which further substantiated the expression of the transgenic construct within the testes.

Overexpression of DPT in the Sertoli cells results in germ cell losses and vacuole formation in the seminiferous tubules. Testes from the 4-month-old transgenic mice exhibited histopathological changes in the seminiferous tubules, which included the desquamation of the spermatogenic cells (Fig. 4). In addition, a significant proportion of the tubules within the testes of the 4-month-old transgenic mice exhibited impairments of the spermatogenesis process, which included the formation of vacuoles, suggesting that endogenous DPT overexpression in the Sertoli cells also negatively impacts upon reproductive function.

Overexpression of DPT in the Sertoli cells leads to a decrease in the expression levels of ZO-1 and claudin-11 in the testes. To determine whether the integrity of the blood-testis barrier (BTB) is affected by DPT overexpression *in vivo*, the present study assessed the steady-state levels of claudin-11 and ZO-1 by immunoblotting. As shown in Fig. 5, the relative expression levels of ZO-1 and claudin-11 in the DPT overexpression mice were significantly reduced as compared with the control mice (Fig. 5), which suggested that DPT may be associated with injury to the BTB.

Discussion

DPT is a gene with unknown function in male reproduction, which was previously shown to be expressed in the testis. The present study demonstrated that increased expression of DPT has an affect on testicular dysfunction in mice. To investigate changes in the expression of DPT in mice with testicular dysfunction, an animal model of testicular dysfunction was

established by treating mice with a single dose of busulfan or CdCl_2 . Findings from previous research have demonstrated that busulfan or CdCl_2 can damage reproductive function in males by disrupting the endogenous balance within the testicular microenvironment, which leads to germ cell losses and somatic cell damage (31-33). In the present study, busulfan or CdCl_2 injections damaged the structure of the testicles of the mice, which concurs with results reported previously. The histological data suggested that the degree of damage to the testes was considerably lower in the busulfan-treated mice compared with the CdCl_2 -treated mice. At 70 days following busulfan treatment, the testicular histology appeared to be recovering to an almost normal profile, while exposure to CdCl_2 resulted in progressive testicular injury. Subsequently, the expression pattern of DPT in the testis with regard to busulfan or CdCl_2 treatment was characterized. In busulfan treated groups, the expression of DPT was normal up to day 5, however, was significantly increased on days 15, 25 and 35, and with recovery up to day 70. Taking the results from the histological analyses of the testes following busulfan treatment together with the levels of DPT expression in the testes at different time points following busulfan treatment, which were determined using PCR and western blotting, suggested that an increase in the expression of DPT may be a good marker for male testicular dysfunction. Therefore, a high level of DPT expression may indicate a low level of reproductive function.

Time-dependent changes in the expression of DPT were observed in the mouse testes in response to treatment with CdCl_2 . The pattern of DPT expression in the mice administered CdCl_2 differed from that observed in the busulfan-treated mice, which further supports the concept that an increase in the expression of DPT may be indicative of reproductive dysfunction in male mice.

Since the increases in the expression of DPT in testis were induced by exogenous detrimental factors, namely CdCl_2 or busulfan administration, the present study hypothesized that endogenous increases in the levels of DPT expression were directly associated with testicular function. Our previous study revealed that DPT is expressed in the Sertoli cells (27), after which DPT transgenic mice, which had a Sertoli cell-specific overexpression of DPT, were established to facilitate studies into the effects of endogenous DPT

overexpression on testicular function in mice. A significant proportion of the tubules within the testes of the 4-month-old transgenic mice exhibited impairments of the spermatogenesis process suggested that just as exogenous factors induce increases in the levels of DPT expression, endogenous DPT overexpression in the Sertoli cells also implicates testicular dysfunction.

The present study further investigated the possible mechanisms underlying testicular injury in mice with Sertoli cell-specific DPT overexpression.

Alterations in the BTB is associated with male fertility (34). Sertoli cell tight junctions (TJ) are an essential component of the BTB, which create an immunologically balanced microenvironment, and they provide structural support for spermatogenesis. TJ are multimolecular membrane specializations comprising several integral membrane proteins, including occludin, claudins, and ZO-1. The findings from previous *in vitro* and *in vivo* investigations show that particular factors can affect the integrity of the BTB by targeting TJ (35). For example, TGF- β 3 and tumor necrosis factor- α , which are secreted by the Sertoli cells and the germ cells, can induce reversible disruption to the BTB *in vivo* by reducing the steady-state levels of occludin and ZO-1 (36).

Our previous data from *in vitro* investigations showed that CdCl₂-induced downregulation of claudin-11 is partially counteracted by DPT silencing in 15P-1 Sertoli cells, which suggests that DPT is a novel regulator of CdCl₂-induced BTB damage *in vitro* (27). In the present study, the steady-state levels of claudin-11 and ZO-1 were assessed by immunoblotting to determine whether the integrity of the BTB is affected by DPT overexpression *in vivo*. It was revealed that compared with the control mice, the relative expression levels of ZO-1 and claudin-11 in the DPT overexpression mice were significantly reduced, which suggested that DPT overexpression may be associated with injury to the BTB, and, in turn, testicular dysfunction.

In conclusion, the expression patterns of DPT in mice treated with CdCl₂ or busulfan indicated that an increase in DPT expression has implications for testicular dysfunction in mice. This finding was corroborated by the findings in the testicles of mice exhibiting endogenous overexpression of DPT, rather than increases in DPT expression induced by busulfan or CdCl₂. The increase in the expression of DPT may harm the integrity of the BTB, which may, in turn, be injurious to testicular function.

Acknowledgements

The present study was supported by the National Nature Science Foundation of China (nos. 81270687 and 30901062).

References

- Irvine DS: Epidemiology and aetiology of male infertility. *Hum Reprod* 13 (Suppl 1): 33-44, 1998.
- Gnoth C, Godehardt E, Frank-Herrmann P, Friol K, Tigges J and Freundl G: Definition and prevalence of subfertility and infertility. *Hum Reprod* 20: 1144-1147, 2005.
- Sepehrimanesh M, Kazemipour N, Saeb M and Nazifi S: Analysis of rat testicular proteome following 30-day exposure to 900 MHz electromagnetic field radiation. *Electrophoresis* 35: 3331-3338, 2014.
- Fowler PA, Cassie S, Rhind SM, Brewer MJ, Collinson JM, Lea RG, Baker PJ, Bhattacharya S and O'Shaughnessy PJ: Maternal smoking during pregnancy specifically reduces human fetal desert hedgehog gene expression during testis development. *J Clin Endocrinol Metab* 93: 619-626, 2008.
- Thonneau P, Bujan L, Multigner L and Mieuisset R: Occupational heat exposure and male fertility: A review. *Hum Reprod* 13: 2122-2125, 1998.
- Benavides-Garcia R, Joachim R, Pina NA, Mutoji KN, Reilly MA and Hermann BP: Granulocyte colony-stimulating factor prevents loss of spermatogenesis after sterilizing busulfan chemotherapy. *Fertil Steril* 103: 270.e8-280.e8, 2015.
- Spiazzi CC, Manfredini V, Barcellos da Silva FE, Flores EM, Izaguirry AP, Vargas LM, Soares MB and Santos FW: γ -Oryzanol protects against acute cadmium-induced oxidative damage in mice testes. *Food Chem Toxicol* 55: 526-532, 2013.
- Houem S, Najjar MF, El Hani A and Sakly R: Accumulation of cadmium and its effects on testis function in rats given diet containing cadmium-polluted radish bulb. *Exp Toxicol Pathol* 59: 307-311, 2008.
- Mabrouk A and Cheikh HB: Thymoquinone supplementation ameliorates lead-induced testis function impairment in adult rats. *Toxicol Ind Health*, 2014 (Epub ahead of print).
- Sharma P, Huq AU and Singh R: Cypermethrin-induced reproductive toxicity in the rat is prevented by resveratrol. *J Hum Reprod Sci* 7: 99-106, 2014.
- Ismail MF and Mohamed HM: Deltamethrin-induced genotoxicity and testicular injury in rats: Comparison with biopesticide. *Food Chem Toxicol* 50: 3421-3425, 2012.
- Walczak-Jedrzejowska R, Wolski JK and Slowikowska-Hilczner J: The role of oxidative stress and antioxidants in male fertility. *Cent European J Urol* 66: 60-67, 2013.
- Desai SS, Roy BS and Mahale SD: Mutations and polymorphisms in FSH receptor: Functional implications in human reproduction. *Reproduction* 146: R235-R248, 2013.
- Smith LB and Walker WH: The regulation of spermatogenesis by androgens. *Semin Cell Dev Biol* 30: 2-13, 2014.
- Xiao X, Mruk DD and Cheng CY: Intercellular adhesion molecules (ICAMs) and spermatogenesis. *Hum Reprod Update* 19: 167-186, 2013.
- Eid N, Ito Y and Otsuki Y: Enhanced mitophagy in Sertoli cells of ethanol-treated rats: Morphological evidence and clinical relevance. *J Mol Histol* 43: 71-80, 2012.
- Li MW, Mruk DD, Lee WM and Cheng CY: Disruption of the blood-testis barrier integrity by bisphenol A *in vitro*: Is this a suitable model for studying blood-testis barrier dynamics? *Int J Biochem Cell Biol* 41: 2302-2314, 2009.
- Alam MS, Ohsako S, Tay TW, Tsunekawa N, Kanai Y and Kurohmaru M: Di (n-butyl) phthalate induces vimentin filaments disruption in rat Sertoli cells: A possible relation with spermatogenic cell apoptosis. *Anat Histol Embryol* 39: 186-193, 2010.
- Zhang X and Lui WY: Dysregulation of nectin-2 in the testicular cells: An explanation of cadmium-induced male infertility. *Biochim Biophys Acta* 1839: 873-884, 2014.
- Neame PJ, Choi HU and Rosenberg LC: The isolation and primary structure of a 22-kDa extracellular matrix protein from bovine skin. *J Biol Chem* 264: 5474-5479, 1989.
- Choi HU, Johnson TL, Pal S, Tang LH, Rosenberg L and Neame PJ: Characterization of the dermatan sulfate proteoglycans, DS-PGI and DS-PGII, from bovine articular cartilage and skin isolated by octyl-sepharose chromatography. *J Biol Chem* 264: 2876-2884, 1989.
- Superti-Furga A, Rocchi M, Schäfer BW and Gitzelmann R: Complementary DNA sequence and chromosomal mapping of a human proteoglycan-binding cell-adhesion protein (dermatopontin). *Genomics* 17: 463-467, 1993.
- Okamoto O, Suzuki Y, Kimura S and Shinkai H: Extracellular matrix 22-kDa protein interacts with decorin core protein and is expressed in cutaneous fibrosis. *J Biochem* 119: 106-114, 1996.
- Okamoto O, Fujiwara S, Abe M and Sato Y: Dermatopontin interacts with transforming growth factor beta and enhances its biological activity. *Biochem J* 337: 537-541, 1999.
- Cooper LJ, Bentley AJ, Nieduszynski IA, Talabani S, Thomson A, Utani A, Shinkai H, Fullwood NJ and Brown GM: The role of dermatopontin in the stromal organization of the cornea. *Invest Ophthalmol Vis Sci* 47: 3303-3310, 2006.
- Takeuchi T, Suzuki M, Kumagai J, Kamijo T, Sakai M and Kitamura T: Extracellular matrix dermatopontin modulates prostate cell growth *in vivo*. *J Endocrinol* 190: 351-361, 2006.

27. Yang Q, Hao J, Chen M and Li G: Dermatopontin is a novel regulator of the CdCl₂-induced decrease in claudin-11 expression. *Toxicol In Vitro* 28: 1158-1164, 2014.
28. Giuli G, Shen WH and Ingraham HA: The nuclear receptor SF-1 mediates sexually dimorphic expression of mullerian inhibiting substance, *in vivo*. *Development* 124: 1799-1807, 1997.
29. Jeffs B, Ito M, Yu RN, Martinson FA, Wang ZJ, Doglio LT and Jameson JL: Sertoli cell-specific rescue of fertility, but not testicular pathology, in Dax1 (Ahch)-deficient male mice. *Endocrinology* 142: 2481-2488, 2001.
30. Narula A, Kilen S, Ma E, Kroeger J, Goldberg E and Woodruff TK: Smad4 overexpression causes germ cell ablation and leydig cell hyperplasia in transgenic mice. *Am J Pathol* 161: 1723-1734, 2002.
31. Zohni K, Zhang X, Tan SL, Chan P and Nagano MC: The efficiency of male fertility restoration is dependent on the recovery kinetics of spermatogonial stem cells after cytotoxic treatment with busulfan in mice. *Hum Reprod* 27: 44-53, 2012.
32. Kanatsu-Shinohara M, Toyokuni S, Morimoto T, Matsui S, Honjo T and Shinohara T: Functional assessment of self-renewal activity of male germline stem cells following cytotoxic damage and serial transplantation. *Biol Reprod* 68: 1801-1807, 2003.
33. Siu ER, Mruk DD, Porto CS and Cheng CY: Cadmium-induced testicular injury. *Toxicol Appl Pharmacol* 238: 240-249, 2009.
34. Koksall IT, Ishak Y, Usta M, Danisman A, Guntekin E, Bassorgun IC and Ciftcioglu A: Varicocele-induced testicular dysfunction may be associated with disruption of blood-testis barrier. *Arch Androl* 53: 43-48, 2007.
35. Erkanlı Şentürk G, Ersoy Canillioglu Y, Umay C, Demiralp-Eksioglu E and Ercan F: Distribution of Zonula Occludens-1 and Occludin and alterations of testicular morphology after *in utero* radiation and postnatal hyperthermia in rats. *Int J Exp Pathol* 93: 438-449, 2012.
36. Xia W, Wong EW, Mruk DD and Cheng CY: TGF-beta3 and TNFalpha perturb blood-testis barrier (BTB) dynamics by accelerating the clathrin-mediated endocytosis of integral membrane proteins: A new concept of BTB regulation during spermatogenesis. *Dev Biol* 327: 48-61, 2009.