

Characterization of human bone morphogenetic protein gene variants for possible roles in congenital heart disease

FEI-FENG LI^{1,2}, XIA DENG^{1,2}, JING ZHOU³, PENG YAN⁴, ER-YING ZHAO¹ and SHU-LIN LIU^{1,2,5}

¹Genomics Research Center (One of The State-Key Laboratory of Biopharmaceutical Engineering, China), Harbin Medical University; ²Translational Medicine Research and Cooperation Center of Northern China,

Heilongjiang Academy of Medical Sciences, Harbin, Heilongjiang 150081; ³Intensive Care Unit;

⁴Department of Colorectal Surgery, The Second Affiliated Hospital of Harbin Medical University, Harbin, Heilongjiang 150086, P.R. China; ⁵Department of Microbiology and Infectious Diseases,

University of Calgary, Calgary, AB T2N 4N1, Canada

Received July 3, 2015; Accepted April 25, 2016

DOI: 10.3892/mmr.2016.5428

Abstract. Congenital heart disease (CHD) is a complex illness with high rates of morbidity and mortality. In embryonic development, the heart is the first formed organ, which is strictly controlled by gene regulatory networks, including transcription factors, signaling pathways, epigenetic factors and microRNAs. Bone morphogenetic protein (BMP)-2 and -4 are essential in cardiogenesis as they can induce the expression of transcription factors, NKX2-5 and GATA binding protein 4, which are important in the development of the heart. The inhibition of BMP-2 and -4 inhibits the late expression of NKX2-5 and affects cardiac differentiation. The aim of the present study was to investigate whether BMP-2 and -4 variations may be associated with CHD in Chinese Han populations. The rs1049007, rs235768 and rs17563 single nucleotide polymorphisms (SNPs), which are genetic variations located within the translated region of the BMP-2 and -4, were evaluated in 230 patients with CHD from the Chinese Han population and 160 non-CHD control individuals. Statistical analyses were performed using the χ^2 test, implemented using SPSS software (version 13.0). The Hardy-Weinberg equilibrium test was performed on the population using online Online Encyclopedia for Genetic Epidemiology studies software, and multiple-sequence alignments of the BMP proteins were performed using Vector NTI software. No statistically significant associations were identified between these genetic variations and the risk of CHD (rs1049007, P-value=0.560;

rs235768, P-value=0.972; rs17563, P-value=0.787). In addition, no correlation was found between the patients with CHD and the non-CHD control individuals. Therefore, the rs1049007, rs235768 and rs17563 genetic variations of BMP-2 were not associated with CHD in the Chinese Han population.

Introduction

Congenital heart disease (CHD) is a common, complex illness with high rates of morbidity and mortality, the genetic etiology of which remains to be fully elucidated (1,2). Worldwide, the incidence of moderate and severe CHD is ~6/1,000 live births, however, if small muscular ventricular septal defects (VSDs) and other minor lesions are included, the incidence is ~75/1,000 live births (3). Appropriately, 1% of patients with CHD require intervention (4), and ~13% of patients with CHD show recognizable chromosomal variants (5,6). The majority of adult patients with show a variety of cardiac complications, including coronary heart disease, arrhythmias and heart failure (7). Although extensive genetic investigations and high-resolution technologies have revealed subtle defects in familial and sporadic cases of CHD (2,8), the subtle genetic causes and molecular mechanisms of CHD remain to be fully elucidated.

In embryonic development, the heart is the first formed organ, which is strictly controlled by gene regulatory networks, including transcription factors, signaling pathways, epigenetic factors and microRNAs (2,9,10). During the last few decades, several cardiac-enriched transcription factors have been identified, and a variety of CHD-causing mutations have been identified in those factor genes, which provide molecular markers and models for the analysis of heart development and the molecular mechanisms underlying CHD (6,11). For example, NKX2-5 is a member of the NK-2 class homeodomain proteins, and is among the earliest markers of cardiac specification. *Drosophila* embryos fail to form a heart in the presence of a mutant NKX gene (12) and, if the functions of the NKX family members are inhibited, vertebrate embryos also fail to form a heart (13). The zinc finger binding proteins, GATA binding protein (GATA)-4, -5 and -6, are also members

Correspondence to: Dr Shu-Lin Liu, Genomics Research Center (One of The State-Key Laboratory of Biopharmaceutical Engineering, China), Harbin Medical University, 157 Baojian Road, Harbin, Heilongjiang 150081, P.R. China
E-mail: slliu@ucalgary.ca

Key words: congenital heart disease, Hardy-Weinberg equilibrium test, single nucleotide polymorphism, bone morphogenetic protein 2, bone morphogenetic protein 4

of cardiac-enriched transcription factors, and are expressed early in heart development, being involved in the initial steps of cardiomyocyte differentiation (14). Mutations in GATA-4 have been identified to be associated with CHD (15), and a variety of other CHD-causing gene mutations have been identified, including CITED2 (16), CFC1 (17) and TBX1 (18). These genes are critical in cardiac development; mutations in these genes lead to cardiovascular malformations and contribute to CHD (19).

In non-precardiac mesoderm, the expression of the heart-specific transcription factors, NKX2-5 and GATA-4, can be induced by BMP-2 and -4, and inhibition of BMP-2 and -4 signaling inhibits the late expression of NKX2-5 and affects the cardiac differentiation (20). In zebrafish, the BMPs also have a direct role in development of heart tube looping (21), and in *Xenopus*, BMP-2 and -4 are broadly expressed in the anterior endoderm and cardiac mesoderm, prior to cardiac differentiation in heart development (22-24). The inhibition of BMP signaling in mice lacking Sma6, leads to the development of hyperplasia in heart valves and defects in the aorticopulmonary septum (25).

BMP-2 and -4 have important functions at multiple stages of cardiogenesis, particularly in the initial induction of heart development. However, the exact nature of these roles, and the association between BMP-2 and -4, with CHD remains to be elucidated. To determine whether there is an association between BMP-2 and -4, and CHD, the gene sequences of BMP-2 and -4 were compared between 230 Chinese Han patients with CHD and 160 control individuals, focusing on the rs1049007, rs235768 and rs17563 SNP genetic variations. The results revealed no correlation between these candidate SNPs and the risk of CHD, and the BMP-2 rs1049007, rs235768 and BMP-4 rs17563 genetic variations may not be risk factors for CHD.

Materials and methods

Study population. A total of 230 patients with CHD and 160 control subjects with no reported cardiac phenotypes were recruited for the present study from the Second Affiliated Hospital of Harbin Medical University (Harbin, China), as shown in Table I. Written informed consent was obtained from each participant and the study was reviewed by the Ethics Committee of Harbin Medical University, consistent with the 1975 Declaration of Helsinki (26,27). The Ethics Committee approved the study. Detailed records on the medical history, physical examination and chest X-ray examination, electrocardiogram and ultrasonic echocardiogram were obtained.

BMP-2 rs1049007 and rs235768, and BMP-4 rs17563 SNP genotyping analysis and statistical methods. Whole genomic DNA was extracted from peripheral blood leukocytes using a QIAamp DNA Blood Mini Kit (cat. no. 51104; Qiagen, Hilden, Germany) (28). The genotypes for the rs1049007, rs235768 and rs17563 SNPs associated with the BMP-2 and -4 genes, respectively (Fig. 1), were determined using a two-stage method. First, rs1049007, rs235768 and rs17563 (Table II) were amplified using standard procedures (2,6,29), following which the PCR products were sequenced

Table I. Clinical characteristics of the study populations.

Parameter	CHD	Control
Sample (n)	230	160
Male/Female (n)	142/88	105/55
Age (years)	16.18±10.22	7.88±11.96

Data are presented as the mean ± standard deviation; CHD, congenital heart disease.




Figure 1. Schematic diagrams of rs1049007, rs235768 and rs17563 locations within the translated regions of the BMP-2 and -4 genes. BMP, bone morphogenetic protein.

(Genewiz, Inc., South Plainfield, NJ, USA) to determine the genotype (Fig. 2). The statistical analyses were performed using χ^2 tests (descriptive statistic crosstalk) to calculate the odds ratios and P-values, implemented using SPSS software (version 13.0; SPSS, Inc., Chicago, IL, USA). In addition, Online Encyclopedia for Genetic Epidemiology studies (OEGE; <http://www.oege.org/software>) online software was used to perform the Hardy-Weinberg equilibrium test for the CHD and control populations.

Multiple sequence alignment. From the National Center for Biotechnology Information website (<http://www.ncbi.nlm.nih.gov/>), the BMP-2 and -4 protein sequences of various species were obtained. Using Vector NTI software (Suite 9; Thermo Fisher Scientific, Inc., Waltham, MA, USA), multiple-sequence alignments of these proteins were performed.

Results

Patients. Clinical diagnosis of the recruited patients was confirmed at the Second Affiliated Hospital of Harbin Medical University. There were no histories of other systemic abnormalities in the patients with CHD, and their mothers had no history of taking medicines or contracting infections during pregnancy (30).

BMP genotyping and statistical analysis. To confirm the hypothesis that there are possible associations between BMPs, which inhibit the late expression of NKX2-5 and affects cardiac differentiation (20), and CHD, the present study performed SNP analyses. No significant differences

Table II. Polymerase chain reaction primers used for BMP genotyping sequence analysis.

Gene	Single nucleotide polymorphism	Primer	Size (bp)	Temperature (°C)
BMP2	rs1049007	Forward CGGGACCCGCTGTCTTCT Reverse TGGAAACGTCCGCTGGTG	455	60.5
	rs235768	Forward CCCACGGAGGAGTTTATC Reverse GCCACTTCCACCACGAAT	275	52.5
BMP4	rs17563	Forward CCCCACCTTATCTGCTCCT Reverse AGTTTGGCTGCTTCTCCC	500	52.8

BMP, bone morphogenetic protein.

Table III. Genotype and allele frequencies of the rs1049007, rs235768 and rs17563 SNPs in 230 Chinese Han patients with CHD and 160 non-CHD control individuals.

SNP	Group	n	Genotype frequency, n (%)			Allele frequency, n (%)	
rs1049007			G/G	A/G	A/A	G	A
	CHD	230	155 (67.4)	68 (29.6)	7 (3.0)	378 (82.2)	82 (17.8)
	Control	160	103 (64.4)	54 (33.8)	3 (1.9)	260 (81.3)	60 (18.8)
rs235768			T/T	T/A	A/A	T	A
	CHD	230	142 (61.7)	80 (34.8)	8 (3.5)	364 (79.1)	96 (20.9)
	Control	160	97 (60.6)	57 (35.6)	6 (3.8)	251 (78.4)	69 (21.6)
rs17563			T/T	T/C	C/C	T	C
	CHD	230	114 (49.6)	96 (41.7)	20 (8.7)	324 (70.4)	136 (29.6)
	Control	160	85 (53.1)	62 (38.8)	13 (8.1)	232 (72.5)	88 (27.5)

SNP, single nucleotide polymorphism; CHD, congestive heart disease.

Table IV. rs1049007, rs235768 and rs17563 SNPs within BMP-2 and -4 are not associated with risk of congenital heart disease.

Genotyped SNP	Gene	Genotype/allele	Pearson's χ^2				Pearson's R			
			χ^2	Min count ^a	df	Asymp. P-value (2-sided)	Pearson's R-value	Asymp. SE ^b	Approx. T-value ^c	Approx. P-value ^d
rs1049007	BMP2	Genotype	1.160	4.10	2	0.560	0.017	0.050	0.337	0.737 ^d
		Allele	0.108	58.26	1	0.742	0.012	0.036	0.329	0.743 ^d
rs235768	BMP2	Genotype	0.568	5.74	2	0.972	0.012	0.051	0.239	0.811 ^d
		Allele	0.054	67.69	1	0.816	0.008	0.036	0.233	0.816 ^d
rs17563	BMP4	Genotype	0.479	13.54	2	0.787	-0.032	0.051	-0.623	0.534 ^d
		Allele	0.393	91.90	1	0.531	-0.022	0.036	-0.626	0.531 ^d

^aMinimum expected count; ^bNot assuming the null hypothesis; ^cUsing the asymptotic standard error assuming the null hypothesis; ^dBased on normal approximation. df, degrees of freedom; Asymp, asymptomatic; Approx, approximate; SNP, single nucleotide polymorphism; BMP, none morphogenetic protein.

were found between the patients with CHD and the 160 CHD-free control individuals in relation to the risk of CHD (Tables III and IV). The present study also performed Hardy-Weinberg equilibrium tests for the patients with CHD

and control individuals, and found that the χ^2 value was 3.06 for the patients with CHD and 2.86 for the CHD-free control individuals. The results were concordant with the Hardy-Weinberg equilibrium.




Figure 2. DNA sequence chromatograms of the rs1049007, rs235768 and rs17563 single nucleotide polymorphisms. (A) rs1049007; (B) rs235768; (C) rs17563. Arrows indicate the sites of variation.




Figure 3. Multiple-sequence alignment of BMP-2 and -4 from birds, fish and mammals, including *Homo sapiens*, *Pan troglodytes* and *Macaca mulatta*. Conservation of 190Ser and 152Val residues in BMP-2 and -4, respectively, were high and in a highly conserved regions. BMP, bone morphogenetic protein.

Conservation of proteins in evolution. The comparison of the BMP-2 and -4 protein sequences between different species included birds, fishes, rodents and mammals, including *Homo sapiens*, *Pan troglodytes* and *Macaca mulatta*. The results of the multiple-sequence alignment analysis showed that the conservation of the rs235768 and rs17563 variations were high; the rs1049007 variation did not alter the protein sequence. The conservation of the 190Ser and 152Val residues in BMP-2 and -4, respectively, were high and were located in highly conserved regions of the proteins (Fig. 3).

Discussion

BMP4, a member of the transforming growth factor (TGF)- β family, is capable of causing human embryonic stem cells (hESCs) to differentiate, and this differentiation can occur without extensive generation of mesoderm and endoderm (31). When inhibiting the fibroblast growth factor (FGF)2 pathway and maximizing BMP4 signaling, BMP4 can direct the hESCs to differentiate towards syncytiotrophoblasts (32,33). The differentiation program induced by BMP4 involves rapid induction, and occurs prior to the expression of caudal type homeobox 2 and several other mesoderm marker genes (32,34). BMP2 is a homodimeric disulfide-bonded protein and is also a member of the TGF- β family (35-37). Despite substantial differences in amino acid sequences with cystine-knot growth factors, it has a similar monomer structure with the factor (38), and has a similar dimer structure with the TGF- β family (39,40).

Several previous studies have shown that BMP-2 and -4 are implicated in the formation of the heart from the overlying mesoderm (41). The expression of the early cardiac markers, NKX2-5 and GATA-4, can be induced by BMP-2 within anterior mesodermal cells, which are located close to the heart forming region (20,42). Following pretreatment with FGFs, BMP-4 can increase the activity of the posterior mesoderm cells (20,43). If the expression of BMP-2 and -4 are inhibited, cardiac differentiation can be inhibited (20,44). Therefore, BMP-2 and -4 are essential in cardiogenesis, and the inductive functions of BMP-2 and -4 may result from the interactions between BMP and other extracellular signaling molecules (45). BMP signaling is important in the formation of different early cell types from hESCs, including the mesoderm (46), endoderm (18) and trophoblasts (47). However, BMP signaling is not required for the expression of early markers of cardiac specification, including NKX2-5, and GATA-4, -5 and -6 (41,48). Therefore, the present study focussed on the rs1049007, rs235768 and rs17563 SNP genetic variations, the aim of which was to analyze the association between BMP-2 and -4, and CHD.

Previously, Qian *et al* found that the rs762642 polymorphism in BMP4 may increase susceptibility to sporadic CHD, and that the polymorphism contributes to the susceptibilities to atrial septal defects and VSDs (49). In addition, it was found that the rs17563 SNP in BMP4 was not associated with the risk of CHD or types of CHD (49). These previous results do not conflict with the results of the present study. The rs762642 SNP is located within the intron between the first and second exon of BMP4, which may affect the function of the promoter and enhancer regions of the gene. Although the rs17563 SNP within the fourth exon resulted in an amino

acid change within the CDS region of the gene, the SNP may not be as important as the rs762642 SNP for the function of BMP4.

In conclusion, the present study compared the gene sequences of BMP-2 and -4 between 230 Chinese Han patients with CHD and 160 control individuals, focusing on the rs1049007, rs235768 and rs17563 SNP genetic variations. The meta-analysis assisted in clarifying the associations between these SNPs with CHD, the results of which revealed that there were no correlations between these candidate SNPs and the risk of CHD in the Chinese population. The risk of developing CHD in individuals with these variants of the BMP-2 and -4 genes may be low; the results of the present study demonstrated that these variations in the BMP-2 and -4 genes were not associated with CHD in the Chinese Han population.

Acknowledgements

This study was supported by a grant from the Heilongjiang Innovation Research Foundation for Graduate Studies (grant no. YJSCX2014-10HYD) to Dr Fei-Feng Li and grants from the National Natural Science Foundation of China (grant nos. NSFC81271786, 81110378, 30970119 and 81030029) to Dr Shu-Lin Liu.

References

1. Zomer AC, Uiterwaal CS, van der Velde ET, Tijssen JG, Mariman EC, Verheugt CL, Vaartjes I, Pieper PG, Meijboom FJ, Grobbee DE and Mulder BJ: Mortality in adult congenital heart disease: Are national registries reliable for cause of death? *Int J Cardiol* 152: 212-217, 2011.
2. Li FF, Zhou J, Zhao DD, Yan P, Li X, Han Y, Li XS, Wang GY, Yu KJ and Liu SL: Characterization of SMAD3 gene variants for possible roles in ventricular septal defects and other congenital heart diseases. *PLoS One* 10: e0131542, 2015.
3. Hoffman JI and Kaplan S: The incidence of congenital heart disease. *J Am Coll Cardiol* 39: 1890-1900, 2002.
4. Hoffman JI, Kaplan S and Liberthson RR: Prevalence of congenital heart disease. *Am Heart J* 147: 425-439, 2004.
5. Pierpont ME, Basson CT, Benson DW Jr, Gelb BD, Giglia TM, Goldmuntz E, McGee G, Sable CA, Srivastava D and Webb CL: American Heart Association Congenital Cardiac Defects Committee, Council on Cardiovascular Disease in the Young: Genetic basis for congenital heart defects: Current knowledge: A scientific statement from the American heart association congenital cardiac defects committee, council on cardiovascular disease in the young: Endorsed by the American academy of pediatrics. *Circulation* 115: 3015-3038, 2007.
6. Deng X, Zhou J, Li FF, Yan P, Zhao EY, Hao L, Yu KJ and Liu SL: Characterization of nodal/TGF-lefty signaling pathway gene variants for possible roles in congenital heart diseases. *PLoS One* 9: e104535, 2014.
7. van der Bom T, Zomer AC, Zwinderman AH, Meijboom FJ, Bouma BJ and Mulder BJ: The changing epidemiology of congenital heart disease. *Nat Rev Cardiol* 8: 50-60, 2011.
8. Richards AA and Garg V: Genetics of congenital heart disease. *Curr Cardiol Rev* 6: 91-97, 2010.
9. Buckingham M, Meilhac S and Zaffran S: Building the mammalian heart from two sources of myocardial cells. *Nat Rev Genet* 6: 826-835, 2005.
10. van Weerd JH, Koshida-Takeuchi K, Kwon C and Takeuchi JK: Epigenetic factors and cardiac development. *Cardiovasc Res* 91: 203-211, 2011.
11. Fishman MC and Olson EN: Parsing the heart: Genetic modules for organ assembly. *Cell* 91: 153-156, 1997.
12. Azpiazu N and Frasch M: Tinman and bagpipe: Two homeobox genes that determine cell fates in the dorsal mesoderm of *Drosophila*. *Genes Dev* 7: 1325-1340, 1993.

13. Grow MW and Krieg PA: Tinman function is essential for vertebrate heart development: Elimination of cardiac differentiation by dominant inhibitory mutants of the tinman-related genes, *XNkx2-3* and *XNkx2-5*. *Dev Biol* 204: 187-196, 1998.
14. Charron F and Nemer M: GATA transcription factors and cardiac development. *Semin Cell Dev Biol* 10: 85-91, 1999.
15. Butler TL, Esposito G, Blue GM, Cole AD, Costa MW, Waddell LB, Walizada G, Sholler GF, Kirk EP, Feneley M, *et al*: GATA4 mutations in 357 unrelated patients with congenital heart malformation. *Genet Test Mol Biomarkers* 14: 797-802, 2010.
16. Sperling S, Grimm CH, Dunkel I, Mebus S, Sperling HP, Ebner A, Galli R, Lehrach H, Fusch C, Berger F and Hammer S: Identification and functional analysis of CITED2 mutations in patients with congenital heart defects. *Hum Mutat* 26: 575-582, 2005.
17. Wang B, Wang J, Liu S, Han X, Xie X, Tao Y, Yan J and Ma X: CFC1 mutations in Chinese children with congenital heart disease. *Int J Cardiol* 146: 86-88, 2011.
18. Teo AK, Ali Y, Wong KY, Chipperfield H, Sadasivam A, Poobalan Y, Tan EK, Wang ST, Abraham S, Tsuneyoshi N, *et al*: Activin and BMP4 synergistically promote formation of definitive endoderm in human embryonic stem cells. *Stem Cells* 30: 631-642, 2012.
19. Gong W, Gottlieb S, Collins J, Blescia A, Dietz H, Goldmuntz E, McDonald-McGinn DM, Zackai EH, Emanuel BS, Driscoll DA and Budarf ML: Mutation analysis of TBX1 in non-deleted patients with features of DGS/VCFS or isolated cardiovascular defects. *J Med Genet* 38: E45, 2001.
20. Ladd AN, Yatskiyevych TA and Antin PB: Regulation of avian cardiac myogenesis by activin/TGFbeta and bone morphogenetic proteins. *Dev Biol* 204: 407-419, 1998.
21. Chen JN, van Eeden FJ, Warren KS, Chin A, Nüsslein-Volhard C, Haffter P and Fishman MC: Left-right pattern of cardiac BMP4 may drive asymmetry of the heart in zebrafish. *Development* 124: 4373-4382, 1997.
22. Wang S, Krinks M, Kleinwaks L and Moos M Jr: A novel *Xenopus* homologue of bone morphogenetic protein-7 (BMP-7). *Genes Funct* 1: 259-271, 1997.
23. Hawley SH, Wunnenberg-Stapleton K, Hashimoto C, Laurent MN, Watabe T, Blumberg BW and Cho KW: Disruption of BMP signals in embryonic *Xenopus* ectoderm leads to direct neural induction. *Genes Dev* 9: 2923-2935, 1995.
24. Hemmati-Brivanlou A and Thomsen GH: Ventral mesodermal patterning in *Xenopus* embryos: Expression patterns and activities of BMP-2 and BMP-4. *Dev Genet* 17: 78-89, 1995.
25. Galvin KM, Donovan MJ, Lynch CA, Meyer RI, Paul RJ, Lorenz JN, Fairchild-Huntress V, Dixon KL, Dunmore JH, Gimbrone MA Jr, *et al*: A role for smad6 in development and homeostasis of the cardiovascular system. *Nat Genet* 24: 171-174, 2000.
26. Tyebkhan G: Declaration of Helsinki: The ethical cornerstone of human clinical research. *Indian J Dermatol Venereol Leprol* 69: 245-247, 2003.
27. Palacios R: Post-trial access and the new version of the Declaration of Helsinki. *Colomb Med (Cali)* 44: 206-207, 2013.
28. Gebril OH and Meguid NA: HFE gene polymorphisms and the risk for autism in Egyptian children and impact on the effect of oxidative stress. *Dis Markers* 31: 289-294, 2011.
29. Abanmi A, Al Harthi F, Zouman A, Kudwah A, Jamal MA, Arfin M and Tariq M: Association of Interleukin-10 gene promoter polymorphisms in Saudi patients with vitiligo. *Dis Markers* 24: 51-57, 2008.
30. Fung A, Manlhiot C, Naik S, Rosenberg H, Smythe J, Loughheed J, Mondal T, Chitayat D, McCrindle BW and Mital S: Impact of prenatal risk factors on congenital heart disease in the current era. *J Am Heart Assoc* 2: e000064, 2013.
31. Amita M, Adachi K, Alexenko AP, Sinha S, Schust DJ, Schulz LC, Roberts RM and Ezashi T: Complete and unidirectional conversion of human embryonic stem cells to trophoblast by BMP4. *Proc Natl Acad Sci USA* 110: E1212-E1221, 2013.
32. Sudheer S, Bhushan R, Fauler B, Lehrach H and Adjaye J: FGF inhibition directs BMP4-mediated differentiation of human embryonic stem cells to syncytiotrophoblast. *Stem Cells Dev* 21: 2987-3000, 2012.
33. Kraushaar DC, Rai S, Condac E, Nairn A, Zhang S, Yamaguchi Y, Moremen K, Dalton S and Wang L: Heparan sulfate facilitates FGF and BMP signaling to drive mesoderm differentiation of mouse embryonic stem cells. *J Biol Chem* 287: 22691-22700, 2012.
34. Greber B: When BMP meets FGF. *Cell Stem Cell* 9: 91-92, 2011.
35. Isaacs NW: Cystine knots. *Curr Opin Struct Biol* 5: 391-395, 1995.
36. Griffith DL, Keck PC, Sampath TK, Rueger DC and Carlson WD: Three-dimensional structure of recombinant human osteogenic protein 1: Structural paradigm for the transforming growth factor beta superfamily. *Proc Natl Acad Sci USA* 93: 878-883, 1996.
37. Vallejo LF and Rinas U: Folding and dimerization kinetics of bone morphogenetic protein-2, a member of the transforming growth factor-beta family. *FEBS J* 280: 83-92, 2013.
38. McDonald NQ and Hendrickson WA: A structural superfamily of growth factors containing a cystine knot motif. *Cell* 73: 421-424, 1993.
39. Scheufler C, Sebald W and Hülsmeier M: Crystal structure of human bone morphogenetic protein-2 at 2.7 Å resolution. *J Mol Biol* 287: 103-115, 1999.
40. Shibata Y, Suzuki D, Wurihan, Yamada A, Maruyama N, Fujisawa N, Kamijo R and Miyazaki T: Lysyl oxidase like-2 reinforces unsatisfactory ossification induced by bone morphogenetic protein-2: Relating nanomechanical properties and molecular changes. *Nanomedicine* 9: 1036-1047, 2013.
41. Walters MJ, Wayman GA and Christian JL: Bone morphogenetic protein function is required for terminal differentiation of the heart but not for early expression of cardiac marker genes. *Mech Dev* 100: 263-273, 2001.
42. Waldo KL, Kumiski DH, Wallis KT, Stadt HA, Hutson MR, Platt DH and Kirby ML: Conotruncal myocardium arises from a secondary heart field. *Development* 128: 3179-3188, 2001.
43. Lough J, Barron M, Brogley M, Sugi Y, Bolender DL and Zhu X: Combined BMP-2 and FGF-4, but neither factor alone, induces cardiogenesis in non-precordial embryonic mesoderm. *Dev Biol* 178: 198-202, 1996.
44. Borges L, Iacovino M, Mayerhofer T, Koyano-Nakagawa N, Baik J, Garry DJ, Kyba M, Letarte M and Perlingeiro RC: A critical role for endoglin in the emergence of blood during embryonic development. *Blood* 119: 5417-5428, 2012.
45. Leung AW, Kent Morest D and Li JY: Differential BMP signaling controls formation and differentiation of multipotent preplacodal ectoderm progenitors from human embryonic stem cells. *Dev Biol* 379: 208-220, 2013.
46. Zhang P, Li J, Tan Z, Wang C, Liu T, Chen L, Yong J, Jiang W, Sun X, Du L, *et al*: Short-term BMP-4 treatment initiates mesoderm induction in human embryonic stem cells. *Blood* 111: 1933-1941, 2008.
47. Yu P, Pan G, Yu J and Thomson JA: FGF2 sustains NANOG and switches the outcome of BMP4-induced human embryonic stem cell differentiation. *Cell Stem Cell* 8: 326-334, 2011.
48. Shi Y, Katsev S, Cai C and Evans S: BMP signaling is required for heart formation in vertebrates. *Dev Biol* 224: 226-237, 2000.
49. Qian B, Mo R, Da M, Peng W, Hu Y and Mo X: Common variations in BMP4 confer genetic susceptibility to sporadic congenital heart disease in a han Chinese population. *Pediatr Cardiol* 35: 1442-1447, 2014.