

Resistant mechanisms and molecular epidemiology of imipenem-resistant *Acinetobacter baumannii*

SHU-ZHEN XIAO^{1*}, HAI-QING CHU², LI-ZHONG HAN^{1*},
ZHE-MIN ZHANG^{2*}, BING LI², LAN ZHAO² and LIYUN XU²

¹Department of Clinical Microbiology, Ruijin Hospital, Shanghai Jiaotong University School of Medicine, Shanghai 200025; ²Department of Respiratory Medicine, Shanghai Pulmonary Hospital, Tongji University School of Medicine, Shanghai 200433, P.R. China

Received January 5, 2016; Accepted July 22, 2016

DOI: 10.3892/mmr.2016.5538

Abstract. The aim of the study was to investigate the resistant mechanisms and homology of imipenem-resistant *Acinetobacter baumannii* (*A. baumannii*). A total of 46 non-duplicate imipenem-resistant *A. baumannii* clinical isolates were collected from three tertiary hospitals between July, 2011 and June, 2012. The minimal inhibitory concentrations (MICs) of antimicrobial agents were determined using the agar dilution method. Phenylalanine-arginine β -naphthylamide was used to detect the presence of the efflux pump-mediated resistant mechanism. Polymerase chain reaction was employed to amplify genes associated with drug resistance, including β -lactamase genes, efflux pump genes and outer membrane protein gene *CarO*. A few amplicons were randomly selected and sequenced. Multilocus sequence analysis (MLST) was employed in typing *A. baumannii*. *A. baumannii* was resistant to imipenem, simultaneously showing resistance to several other antimicrobials. In addition, 13 *A. baumannii* were found to mediate drug resistance through operation of the efflux pump. Of the various drug resistance genes tested, *bla*_{OXA-51} was present in 46 isolates, *bla*_{OXA-23} gene was present in 44 isolates and *bla*_{NDM} gene was found in only one strain. Other drug resistant-associated genes, including *bla*_{KPC}, *bla*_{IMP}, *bla*_{OXA-24}, *bla*_{OXA-58}, *bla*_{SHV}, *bla*_{GIM} and *bla*_{VIM} were not detected. Mutation of *adeS* and outer membrane protein gene *CarO* were found in a few of the imipenem-resistant isolates. The MLST analysis revealed that all 46 clinical isolates were clustered into 11 genotypes and the most frequent genotype

was ST208. In conclusion, β -lactamase genes, genes involved in efflux pump and mutation of outer membrane protein encoding gene may be important in mediating imipenem resistance in *A. baumannii*. Of the 11 different genotypes, ST11 was shared by the majority of *A. baumannii*, which may be due to horizontal transfer of patients from hospitals.

Introduction

Acinetobacter baumannii (*A. baumannii*) has emerged as a major pathogen of nosocomial infections and is associated with high rates of morbidity and mortality in recent years (1,2). A nationwide surveillance program, including hospitals from 14 geographically different regions in China revealed that the ratio of *A. baumannii* is on the increase annually (3).

Carbapenem has good antibacterial activity against *A. baumannii* and was the first choice in treatment of infection caused by *A. baumannii* in the past years (4). However, the emergence of resistance to carbapenem was reported in 1991 (5), followed by similar reports from different parts of the world (6,7). In China, 57 and 61% of *Acinetobacter* spp. (*A. baumannii* accounted for 89.6%) showed resistance to imipenem and meropenem, respectively (3). International studies in China as well as in other parts of the world focused only on evaluating the resistance of *A. baumannii* to various antimicrobials (8-10). However, to the best of our knowledge, few studies have investigated the molecular mechanism underlying drug resistance. Additionally, no data are available on the epidemiological characteristics of imipenem-resistant *A. baumannii* in Shanghai.

Thus, *A. baumannii* clinical isolates were collected from three tertiary hospitals in Shanghai and their drug resistance pattern to a spectrum of antimicrobials, molecular mechanisms (including carbapenemase, efflux pumps and membrane proteins) behind their resistance and multilocus sequence analysis (MLST) were analyzed to assess their molecular epidemiology.

Materials and methods

Bacterial strains. During the period July, 2011 to June, 2012, 46 non-duplicate imipenem-resistant *A. baumannii* strains

Correspondence to: Dr Hai-Qing Chu, Department of Respiratory Medicine, Shanghai Pulmonary Hospital, Tongji University School of Medicine, 507 ZhengMin Road, Shanghai 200433, P.R. China
E-mail: nqy320@163.com

*Contributed equally

Key words: *Acinetobacter baumannii*, imipenem, efflux pump, minimal inhibitory concentration, multilocus sequence analysis, molecular epidemiology

Table I. Gene-specific primers used in this study.

Genes	Primer sequences
<i>recA</i>	F: CCTGAATCTTCTGGTAAAAC R: GTTTCTGGGCTGCCAAACATTAC
<i>ITS</i>	F: CATTATCACGGTAATTAGTG R: AGAGCACTGTGCACTTAAG
<i>bla_{KPC}</i>	F: TTACTGCCCCGTTGACGCCCAATCC R: TCGCTAAACTCGAACAGG
<i>bla_{IMP}</i>	F: AACCAGTTTTGCCTTACCAT R: CTACCGCAGCAGAGTCTTTG
<i>bla_{NDM}</i>	F: CCGCCAGATCCTCAACT R: ATCAGGCAGCCACCAAAA
<i>bla_{OXA-51}</i>	F: TAATGCTTTGATCGGCCTTG R: TGGATTGCACTTCATCTTGG
<i>bla_{OXA-23}</i>	F: GATCGGATTGGAGAACCAGA R: ATTTCTGACCCATTTCCAT
<i>bla_{OXA-24}</i>	F: GGTTAGTTGGCCCCCTTAAA R: AGTTGAGCGAAAAGGGGATT
<i>bla_{OXA-58}</i>	F: AAGTATTGGGGCTTGTGCTG R: CCCCTCTGCGCTCTACATAC
<i>bla_{SHV}</i>	F: GGTTATGCGTTATATTGCCC R: TTAGCGTTGCCAGTGCTC
<i>bla_{GIM}</i>	F: AGAACCTTGACCGAACGCAG R: ACTCATGACTCCTCACGAGG
<i>bla_{VIM}</i>	F: TCCACGCACTTTCATGACGA R: AGACGTGCGTGACAACCTCAT
<i>adeA</i>	F: GAAATCCGTCCGCAAGTC R: ACACGCACATACATACCC
<i>adeB</i>	F: AAAGACTTCAAAGAGCGG R: TCACGCATTGCTTCACCC
<i>adeC</i>	F: ATTTCAAGTCGTAGCATT R: CTTGATAAGTAGAGTAGGGATT
<i>adeS</i>	F: ACTGTTATCTTCTGTGGCTGTA R: GTGGACGTTAGGTCAAGTTCTG
<i>adeR</i>	F: AAACGGTTGGGAAGTATTA R: ATGGCTATCTACGGTTCCG
<i>CarO</i>	F: AAGGAGAAAACGATGA R: TTATTACGTGGTTATGG
<i>gltA</i>	F: AATTACAGTGGCACATTAGGTCCC R: GCAGAGATACCAGCAGAGATACACG
<i>gyrB</i>	F: TGAAGGCGGCTTATCTGAGT R: GCTGGGTCTTTTTCCTGACA
<i>gdhB</i>	F: ACCACATGCTTTGTTATG R: GTTGGCGTATGTTGTGC
<i>recA</i>	F: CCTGAATCTTCYGGTAAAAC R: GTTTCTGGGCTGCCAAACATTAC
<i>cpn60</i>	F: GGTGCTCAACTTGTTTCGTGA R: CACCGAAACCAGGAGCTTTA
<i>Gpi</i>	F: GAAATTTCCGGAGCTCACAA R: TCAGGAGCAATACCCCACTC
<i>rpoD</i>	F: ACCCGTGAAGGTGAAATCAG R: TTCAGCTGGAGCTTTAGCAAT

were collected from three tertiary hospitals located in Shanghai, China.

Reconfirmation of strains. The collected strains were subjected to gram staining, biochemical tests, and *recA* gene and 16S-23S rRNA gene intergenic spacer region to reconfirm them as *A. baumannii* (7).

Antimicrobial susceptibility and efflux phenotype tests. The collected *A. baumannii* isolates were subjected to an antimicrobial susceptibility test against imipenem, meropenem, amikacin, piperacillin, ceftazidime, cefotaxime, minocycline, ciprofloxacin, ampicillin/sulbactam, sulbactam, cefoperazone/sulbactam, piperacillin/tazobactam, colistin, tigecycline and trimethoprim/sulfamethoxazole using agar dilution method. *Escherichia coli* strain ATCC25922 and *Pseudomonas aeruginosa* (*P. aeruginosa*) strain ATCC27853 were used as reference strains.

Strains in which efflux pump operation was detected by agar dilution method where imipenem- and meropenem-resistant isolates were cultured in Mueller-Hinton agar contained the efflux pump inhibitor phenylalanine-arginine β -naphthylamide (PA β N) at a final concentration of 20 mg/l (11,12). A ≥ 4 -fold reduction of imipenem or meropenem minimal inhibitory concentrations (MICs) in the presence of PA β N possessed an operating drug efflux pump.

Analysis of genes responsible for drug resistance, drug efflux and outer membrane protein. Polymerase chain reaction (PCR) was performed for the genes, *bla_{KPC}*, *bla_{IMP}*, *bla_{NDM}*, *bla_{OXA-51}*, *bla_{OXA-23}*, *bla_{OXA-24}*, *bla_{OXA-58}*, *bla_{SHV}*, *bla_{GIM}* and *bla_{VIM}*, *CarO*, *adeA*, *adeB*, *adeC*, *adeS* and *adeR*. Thus, obtained amplicons were subjected to sequencing analysis.

A fresh and pure bacterial colony was suspended in distilled water and boiled at 100°C for 15 min. After centrifugation at 8,000 \times g for 15 min, 1 μ l of the supernatant was used for PCR analysis with the primers (Table I). PCR was performed in a total volume of 50 μ l containing 0.25 μ l Taq DNA polymerase (Takara Bio, Inc., Tokyo, Japan), 5 μ l 10X PCR buffer (Mg²⁺ Plus), 4 μ l dNTP mixture (2.5 mM each), 2.5 μ l DNA template, 1 μ l of each primer (20 μ M), and 36.25 μ l ddH₂O. The PCR thermal cycle consisted of initial denaturation at 94°C for 5 min, followed by 30 cycles of 94°C for 30 sec, annealing 55°C for 1 min and 72°C for 1 min and a final extension at 72°C for 7 min. The PCR products were electrophoresed in 1% agarose gel and visualized under ultraviolet light, and subsequently sequenced (Sangon Biotech Co., Ltd., Shanghai, China).

MLST. Seven housekeeping genes including homologous recombination factor (*recA*), citrate synthase (*gltA*), DNA gyrase subunit (*gyrB*), glucose-6-phosphate isomerase (*gpi*), glucose dehydrogenase B (*gdhB*), 60-kDa chaperonin (*cpn60*), and RNA polymerase 70 factor (*rpoD*) were amplified in PCR using relevant primers (Table I) and appropriate thermal conditions. The amplicons were sequenced and the sequences were submitted to the MLST database (<http://pubmlst.org.net>) to compare them with sequences submitted from other parts of the world. Each strain was then characterized by a pattern of numbers defining its allelic profile.

A

Query	61	aTTAGTCACGGCGACCTCTCTGCTAGAGCTTACGATAATAGAATTCACCTCCGCCGAAATG	120
Sbjct	2041963	ATTAGTCACGGCGACCTCTCTGCTAGAGCTTACGATAACCGAATTCACCTCCGCCGAAATG	2042022
Query	121	TCGGAGCTTTTATATAATTTAATGATATGGCTCAAAAGCTAGAGGTTTCTGTTAAAAAT	180
Sbjct	2042023	TCGGAGCTTTTATATAATTTAATGATATGGCTCAAAAGCTAGAGGTTTCCGTTAAAAAT	2042082
Query	181	GCGCAGGTTTGGAAATCGACCATCGCACATGAGTTAAGAAGCCTATAACGATATTACAA	240
Sbjct	2042083	GCGCAGGTTTGGAAATCGACCATCGCACATGAGTTAAGAAGCCTATAACGATATTACAA	2042142
Query	241	GGTCGTTTACAAGGCATCATCGACGGTGTITTTAAACCTGATGAAGTCCTATTTAAAAAGC	300
Sbjct	2042143	GGTCGTTTACAAGGAATTAATTGATGGCGTITTTAAACTTGATGAAGTCTATTTAAAAAGT	2042202
Query	301	CTTTTAAATCAAGTTGAAGTTTATCTCATTAGTCGAAGACTTACGGACTTTAAGCTTA	360
Sbjct	2042203	CTTTTAAATCAAGTTGAAGTTTATCTCATTAGTCGAAGACTTACGGACTTTAAGCTTA	2042262
Query	361	GTAGAGAACCAGCAACTCCGGTTAAATTATGAATTGTTTGACTTTAAGCGCGTAGITGAA	420
Sbjct	2042263	GTAGAGAACCAGCAACTCCGGTTAAATTATGAATTGTTTGACTTTAAGCGCGTAGITGAA	2042322
Query	421	AAAGTTCTTAAAGCATTGGAAGATCGTTTGGATGAAGCTAAGCTAGTACCAGAACCTTGAC	480
Sbjct	2042323	AAAGTTCTTAAAGCATTGGAAGATCGTTTGGATGAAGCTAAGCTAGTACCAGAACCTTGAC	2042382

B

Query	1	MRLAKRFIVPINFLAEAAKIKSHGDLARAYDNRIHSAEMSELLYNFNDMAQKLEVSVKV	60
Sbjct	1	MRLAKRFIVPINFLAEAAKIKSHGDLARAYDNRIHSAEMSELLYNFNDMAQKLEVSVKV	60
Query	61	AQVWNAIAHELRTPTITLQGRLLQGIIDGVFKPDEVLFKSLNQVEVLSHLVEDLRTLSL	120
Sbjct	61	AQVWNAIAHELRTPTITLQGRLLQGIIDGVFKPDEVLFKSLNQVEVLSHLVEDLRTLSL	120
Query	121	VENQQLRLNYELDFKAVVEKVLKAFEDRLDEAKLVPELDLTSTPVYCDRRRIEQVLIAL	180
Sbjct	121	VENQQLRLNYELDFKAVVEKVLKAFEDRLDQAKLVPELDLTSTPVYCDRRRIEQVLIAL	180
Query	181	IDNAIRYSNAGKLKISSEVVSQNWILKIEDEGPGIATEFQDDLYKFFRLEESRNKEFGG	240
Sbjct	181	IDNAIRYSNAGKLKISSEVVSQNWILKIEDEGPGIATEFQDDLYKFFRLEESRNKEFGG	240
Query	241	TGLGLAVVHAIIVALKGTIQYSNQGSVFTIKISMGHHEIG	282
Sbjct	241	TGLGLAVVHAIIVALKGTIQYSNQGSVFTIKISMGHHEIG	282

Figure 1. (A) Comparison of nucleotide sequence of *adeS* gene between resistant strain and reference strain ATCC17978. (B) Comparison of amino acid sequence of *adeS* gene between resistant strain and reference strain ATCC17978.

Results

Antimicrobial susceptibility. *A. baumannii* resistant to imipenem simultaneously showed resistance to several other common antimicrobials. The resistance rate was >80% for all the antimicrobials except minocycline and colistin. Antibiotic susceptibility of the 46 clinical isolates is shown in Table II. Thirteen imipenem-resistant *A. baumannii* isolates were positive for efflux pump.

Detection of genes involved in drug resistance, drug efflux and outer membrane protein. Of the various drug resistance genes tested, *bla*_{OXA-51} was present in 46 isolates, *bla*_{OXA-23} gene was present in 44 isolates and *bla*_{NDM} gene was found in only one strain. Other drug-resistant genes including *bla*_{KPC}, *bla*_{IMP}, *bla*_{OXA-24}, *bla*_{OXA-58}, *bla*_{SHV}, *bla*_{GIM} and *bla*_{VIM} were not detected in the isolates.

Of the five genes associated with the drug efflux pump tested, all five were found to be present in the isolates. Several mutations were found in the sequences of *adeS* gene in isolates with efflux phenotype. Differences were observed at three places when nucleotide sequences were translated into an amino acid sequence. This amino acid sequence was then

Table II. The drug-resistant rates of imipenem-resistant *Acinetobacter baumannii*.

Drug	Resistance rate No. of resistant strains (%)
Meropenem	41 (89)
Amikacin	38 (83)
Piperacillin	46 (100)
Ceftazidime	46 (100)
Minocycline	34 (74)
Ciprofloxacin	45 (98)
Ampicillin/sulbactam	43 (93)
Piperacillin/tazobactam	46 (100)
Colistin	1 (2)
Trimethoprim/sulfamethoxazole	43 (93)
Cefotaxime	45 (98)

compared to the amino acid sequence of the reference strain ATCC17978 (Fig. 1).

A				
Query	1	TTACCAGAAGAAGTTCACACCAACTTTACCAACTGGCAACCATTTGTATTTATCATCGTT	60	
Sbjct	2936379	TTACCAGTAGAAGTTTACACCAACTTTACCAACTGGGAACCATTTGTATTTATCATCATT	2936438	
Query	61	AGCAATTTTACGAGCTTCCGCATTTACTGCCTCTTCAAGTGATTGATCA-C--CAGTTGT	117	
Sbjct	2936439	ACGGATTTTGTCTTCTCAGCACGTAAGTCAGCATC-CG-CATT-ACCACCTGCAGCATT	2936495	
Query	118	AACGTCAGAACCTGACGAAACAAGCTTAAGTGTGGTTACCAAGTATAGTAAGCGCTAC	177	
Sbjct	2936496	AACAAAAGTACCTTGTTTATCTAAGTCTACAGTGGTTACCAAGTGTAGTAAGCGCTAC	2936555	
Query	178	TTCACCGAATACGCCCCAGTTTATTGATTTTAGGTGCAAAACCAAACTAAGTATGG	237	
Sbjct	2936556	TTCACCAAATACACCCAGTTTATTGATTTTAGGTGCGAAACCAAACTAAGTACGG	2936615	
Query	238	AGCAATATCATTTTATA--TGACATTGACCGTTAATTTGACACCATCGGCACCCGCA	295	
Sbjct	2936616	AGCGATATCATTTTGTAACTTA-A-CTGACCATT--TACAGA-ACCAT---TATAAG-A	2936666	
Query	296	ATAAAGTCTT-GGTTATTTACACGGAATGAACGAGTCGCATCAACGTTACGAGTTAAATC	354	
Sbjct	2936667	ATAATTTGTTCCGTTAATTT---TGATAGTACCA-TCTGATGAACGTT--TAGTTAGGTC	2936720	
Query	355	ATAATCGTTATCAAGGTAAGCCGCCAGCAGCTACATATAAGCCTTGAGCCCAACGGTT	414	
Sbjct	2936721	ATAATCGTTATCTAAATACCCGCCACCTGCAGCTACATATAAGCCTTGAGCCCAACGGTT	2936780	
Query	415	AGTGCTTGACCCCATGGACGAATCTCAGCATTTAAATAAACGTTGTTATTATCCATATC	474	
Sbjct	2936781	AGTACTTGACCCCATGGCGAATCTCAGCATTTAAATATACGTTGTTATTATCCATATC	2936840	
Query	475	AAGGTCATAAGTTGATCCATTGACTTTTACATCATCAGACCAAGAAATGTCACCGCGGT	534	
Sbjct	2936841	AACGTCATATTTAGTACCATTAATTGATAAGTCATCTGCCAAGAAATGTCACCAACGGT	2936900	
Query	535	ATAACCAATGCTAAACCTACATATGGGTTTGCTTGCCATAACAAAGCACCACCGTAACC	594	
Sbjct	2936901	ATAACCAATGCTAAACCTACATATGGGTTTGCTTGCCATAACAAAGCACCACCGTAACC	2936960	
Query	595	TGTAGTACCTACTTCAGCACGAGCGCCTACTGGAATTAATTGGTTTTATCGAATGCATA	654	
Sbjct	2936961	TGTAGTACCTACTTCAGCACGAGCGCCTACTGGAATTAATTGGTTTTATCGAATGCATA	2937020	
Query	655	GCTGTCATGAACAACAGCTTCATCCGCCAT	684	
Sbjct	2937021	ACTGTCATGAACAACAGCTTCATCAGCCAT	2937050	
B				
Query	6	VVHDSYAFDKQLIPVGARAEVGTGTYGGALLWQANFYVGLALGYNGDISWSDDVKVNG	65	
Sbjct	25	VVHDSYAFDKQLIPVGARAEVGTGTYGGALLWQANFYVGLALGYNGDISW DD+ +NG	84	
Query	66	STYDLMDNNNVYLNAEIRPWGASTNRWAQGLYVAAGAAAYLDNDYDLT-RNVDA TRSFRV	124	
Sbjct	85	+ YD+DMDNNNVYLNAEIRPWGASTNRWAQGLYVAAGAAAYLDNDYDLT R+ D T ++	142	
Query	125	NNQDFIAGDGVKINGMSYKNDIAPYLGFGFAPKINKNWGVFGEVGAYITGNPTVKLV	184	
Sbjct	143	N NGTNYSYNG---SVNGQLSYKNDIAPYLGFGFAPKINKNWGVFGEVGAYITGNPTVELDK	199	
Query	185	SGSAVITGDQSLEEA VNAEARKIANDDKYKWLFPVGVGVNFVW	227	
Sbjct	200	G+ V + + + AE KI NDDKYK W PVGVGVNF+ W	242	

Figure 2. (A) Comparison of nucleotide sequence of *CarO* gene between resistant and reference strain ATCC17978. (B) Comparison of amino acid sequence of *CarO* gene between resistant and reference strain ATCC17978.

Similarly, the nucleotide sequence of the outer membrane protein encoding gene *CarO*, when compared with the nucleotide sequence of reference strain ATCC17978, harbored mutations that were reflected in the amino acid sequence (Fig. 2).

Genotyping of isolates by MLST. The MLST analysis revealed that the isolates were clustered in 11 different genotypes or STs. The ST208 genotype was shared by the majority of isolates (58.7%, 27/46), followed by ST191 (10.9%, 5/46) and ST451 (6.5%, 3/46). We also detected some other STs shared by certain isolates such as ST75 (2.1%, 1/46), ST90 (4.2%,

2/46), ST92 (2.1%, 1/46), ST108 (2.1%, 31/46), ST109 (2.1%, 1/46), ST172 (2.1%, 1/46), ST368 (4.2%, 2/46) and ST69 (4.2%, 2/46). These STs were grouped into the three clonal complexes, CC92, CC109 and CC28.

Discussion

A. baumannii develops resistance to imipenem through a variety of mechanisms. Carbapenemase is an important factor responsible for imipenem resistance. In the present study, common carbapenemases were detected in the

isolates, including *bla*_{OXA-51}, *bla*_{OXA-23}, *bla*_{OXA-24}, *bla*_{OXA-58}, *bla*_{KPC}, *bla*_{IMP}, *bla*_{SHV}, *bla*_{GIM}, *bla*_{NDM} and *bla*_{VIM}. OXA-type enzymes are naturally present in *Acinetobacter spp.* and are usually expressed in small amounts (13). The expression of such genes is markedly higher under the effect of a strong promoter (insertion sequence ISAbal is the most shared) and induce drug resistance only when combined with a reduction in outer membrane permeability and/or activation of the efflux pump (14). In the present study, *bla*_{OXA-51} and *bla*_{OXA-23} genes were prevalent among the isolates, results that are consistent with other reports (15-17). Carbapenemases that are different from OXA, such as KPC, IMP, SHV, GIM, NDM and VIM have strong carbapenem-hydrolysing activity (14). However, such types of carbapenemases were rarely detected in *A. baumannii*. The *bla*_{NDM} gene was identified in only one strain while the remaining resistant genes were not detected. NDM was first identified in *Escherichia coli* and *Klebsiella pneumoniae* in 2008 in India (18). This finding was followed by reports on NDM-producing *P. aeruginosa*, *Enterobacter cloacae*, *Citrobacter freundii* and *Enterococcus faecium* (19-23). In China, NDM-producing *A. baumannii* was first reported in 2011. Of the antimicrobials tested one NDM-positive isolate in the present study was identified that was multidrug-resistant, and only susceptible to amikacin, colistin and minocycline.

Drug efflux systems including AdeABC, AdeIJK, AdeDE and AdeXYZ (RND family) have been found in *A. baumannii*. Of these, the AdeABC efflux system is common in *A. baumannii* (24). This efflux pump, together with other resistant mechanisms, can lead to high-level imipenem resistance. Although mediated by the substrate, its expression may increase when a single point mutation occurs in the *adeR* or *adeS* gene (25). PAβN was proven to be an effective inhibitor of drug efflux. In the present study, *adeA*, *adeB* and *adeC* were present in all of the isolates because when PAβN was added the MICs inherent to imipenem in 13 isolates were decreased. The *adeS* gene differed from the *adeS* of standard strain and this is the possible reason for increased drug efflux associated with drug resistance.

Few studies concerning the impact of changes on membrane proteins in *A. baumannii* are available. In 2002, a laboratory in Argentina advocated for the first time that inducible resistance by imipenem can trigger loss of a 29-kDa membrane protein. In 2005, the same laboratory furthering their study, demonstrated that the outer membrane protein is encoded by the *CarO* gene and when there is an insertion mutation or any other mutation in the *CarO* gene makes it off and thus the strain become resistant to certain drugs (26). In the present study, the sequence of *CarO* gene had nucleotide insertions, deletions and point mutations in comparison with the standard strains and there were also differences in their nucleotide and amino acid sequences.

In summary, for a global epidemiologic analysis, a comparison of the results between different laboratories is required. MLST is a powerful tool used to transfer typing data and compare results via relevant databases. The MLST analysis revealed that the major epidemic clone of *A. baumannii* in Shanghai was ST208 (CC92 clone complex), which differed from the results obtained in other regions in China (27).

Acknowledgements

The project was supported by a grant from the Natural Science Foundation of Shanghai Science and Technology Committee (no. 12ZR1426200), the Medical Guide Program of Shanghai Science and Technology Committee (no. 14411962900), Key project of Shanghai Municipal Health and Family Planning Commission (no. 201540367) and Central Universities Basic Research Program (no. 1511219024).

References

1. Perez F, Hujer AM, Hujer KM, Decker BK, Rather PN and Bonomo RA: Global challenge of multidrug-resistant *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 51: 3471-3484, 2007.
2. Giamarellou H, Antoniadou A and Kanellakopoulou K: *Acinetobacter baumannii*: A universal threat to public health? *Int J Antimicrob Agents* 32: 106-119, 2008.
3. Wang F, Zhu D, Hu F, Jiang X, Hu Z, Li Q, Sun Z, Chen Z, Xu Y, Zhang X, et al: CHINET 2012 surveillance of bacterial resistance in China. *China J Infect Chemother* 13: 321-330, 2013 (In Chinese).
4. Kim YJ, Kim SI, Hong KW, Kim YR, Park YJ and Kang MW: Risk factors for mortality in patients with carbapenem-resistant *Acinetobacter baumannii* bacteremia: Impact of appropriate antimicrobial therapy. *J Korean Med Sci* 27: 471-475, 2012.
5. Go ES, Urban C, Burns J, Kreiswirth B, Eisner W, Mariano N, Mosinka-Snipas K and Rahal JJ: Clinical and molecular epidemiology of acinetobacter infections sensitive only to polymyxin B and sulbactam. *Lancet* 344: 1329-1332, 1994.
6. Pournaras S, Markogiannakis A, Ikonomidis A, Kondyli L, Bethimouti K, Maniatis AN, Legakis NJ and Tsakris A: Outbreak of multiple clones of imipenem-resistant *Acinetobacter baumannii* isolates expressing OXA-58 carbapenemase in an intensive care unit. *J Antimicrob Chemother* 57: 557-561, 2006.
7. Kuo LC, Teng LJ, Yu CJ, Ho SW and Hsueh PR: Dissemination of a clone of unusual phenotype of pandrug-resistant *Acinetobacter baumannii* at a university hospital in Taiwan. *J Clin Microbiol* 42: 1759-1763, 2004.
8. Xu J, Sun Z, Li Y and Zhou Q: Surveillance and correlation of antibiotic consumption and resistance of *Acinetobacter baumannii* complex in a tertiary care hospital in northeast China, 2003-2011. *Int J Environ Res Public Health* 10: 1462-1473, 2013.
9. Kwon NY, Kim JD and Pai HJ: The resistance mechanisms of b-lactam antimicrobials in clinical isolates of *Acinetobacter baumannii*. *Korean J Intern Med* 17: 94-99, 2002.
10. García-Quintanilla M, Pulido MR, Moreno-Martínez P, Martín-Peña R, López-Rojas R, Pachón J and McConnell MJ: Activity of host antimicrobials against multidrug-resistant *Acinetobacter baumannii* acquiring colistin resistance through loss of lipopolysaccharide. *Antimicrob Agents Chemother* 58: 2972-2975, 2014.
11. Chen TL, Siu LK, Wu RC, Shaio MF, Huang LY, Fung CP, Lee CM and Cho WL: Comparison of one-tube multiplex PCR, automated ribotyping and intergenic spacer (ITS) sequencing for rapid identification of *Acinetobacter baumannii*. *Clin Microbiol Infect* 13: 801-806, 2007.
12. Clinical Laboratory Standards Institute (CLSI): Performance standards for antimicrobial susceptibility testing: Twenty-second informational supplement. CLSI document M100-S22. CLSI, Wayne, PA, 2012.
13. Zhang JP, Zhu W, Tian SF, Chu YZ and Chen BY: Molecular characteristics and resistant mechanisms of imipenem-resistant *Acinetobacter baumannii* isolates in Shenyang, China. *J Microbiol* 48: 689-694, 2010.
14. Merckier AK and Centrón D: bla(OXA-51)-type beta-lactamase genes are ubiquitous and vary within a strain in *Acinetobacter baumannii*. *Int J Antimicrob Agents* 28: 110-113, 2006.
15. He C, Xie Y, Zhang L, Kang M, Tao C, Chen Z, Lu X, Guo L, Xiao Y, Duo L, et al: Increasing imipenem resistance and dissemination of the ISAbal-associated blaOXA-23 gene among *Acinetobacter baumannii* isolates in an intensive care unit. *J Med Microbiol* 60: 337-341, 2011.

16. Martínez P and Mattar S: Imipenem-resistant *Acinetobacter baumannii* carrying the ISAbal-bla OXA-23,51 and ISAbal-bla ADC-7 genes in Monteria, Colombia. *Braz J Microbiol* 43: 1274-1280, 2012.
17. Corvec S, Poirel L, Naas T, Drugeon H and Nordmann P: Genetics and expression of the carbapenem-hydrolyzing oxacillinase gene blaOXA-23 in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 51: 1530-1533, 2007.
18. Yong D, Toleman MA, Giske CG, Cho HS, Sundman K, Lee K and Walsh TR: Characterization of a new metallo-beta-lactamase gene, bla(NDM-1), and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother* 53: 5046-5054, 2009.
19. Cabanes F, Lemant J, Picot S, Simac C, Cousty J, Jalin L, Naze F, Boisson V, Cresta MP, André H, *et al*: Emergence of *Klebsiella pneumoniae* and *Salmonella* metallo-beta-lactamase (NDM-1) producers on reunion island. *J Clin Microbiol* 50: 3812, 2012.
20. Savard P, Gopinath R, Zhu W, Kitchel B, Rasheed JK, Tekle T, Roberts A, Ross T, Razeq J, Landrum BM, *et al*: First NDM-positive *Salmonella* sp. strain identified in the United States. *Antimicrob Agents Chemother* 55: 5957-5958, 2011.
21. Jovicic B, Lepsanovic Z, Suljagic V, Rackov G, Begovic J, Topisirovic L and Kojic M: Emergence of NDM-1 metallo- β -lactamase in *Pseudomonas aeruginosa* clinical isolates from Serbia. *Antimicrob Agents Chemother* 55: 3929-3931, 2011.
22. Poirel L, Dortet L, Bernabeu S and Nordmann P: Genetic features of blaNDM-1-positive *Enterobacteriaceae*. *Antimicrob Agents Chemother* 55: 5403-5407, 2011.
23. Ho PL, Lo WU, Yeung MK, Lin CH, Chow KH, Ang I, Tong AH, Bao JY, Lok S and Lo JY: Complete sequencing of pNDM-HK encoding NDM-1 carbapenemase from a multidrug-resistant *Escherichia coli* strain isolated in Hong Kong. *PLoS One* 6: e17989, 2011.
24. Chu YW, Chau SL and Houang ET: Presence of active efflux systems AdeABC, AdeDE and AdeXYZ in different AbeM (MATE) Ade ABC *Acinetobacter* genomic DNA groups. *J Med Microbiol* 55: 477-478, 2006.
25. Marchand I, Damier-Piolle L, Courvalin P and Lambert T: Expression of the RND-type efflux pump AdeABC in *Acinetobacter baumannii* is regulated by the AdeRS two-component system. *Antimicrob Agents Chemother* 48: 3298-3304, 2004.
26. Mussi MA, Limansky AS and Viale AM: Acquisition of resistance to carbapenems in multidrug-resistant clinical strains of *Acinetobacter baumannii*: Natural insertional inactivation of a gene encoding a member of a novel family of β -barrel outer membrane proteins. *Antimicrob Agents Chemother* 49: 1432-1440, 2005.
27. Zhou Z, Du X, Wang L, Yang Q, Fu Y and Yu Y: Clinical carbapenem-resistant *Acinetobacter baylyi* strain coharboring blaSIM-1 and blaOXA-23 from China. *Antimicrob Agents Chemother* 55: 5347-5349, 2011.