Hypoxia-induced hsa-miR-101 promotes glycolysis by targeting TIGAR mRNA in clear cell renal cell carcinoma

XIAOCHAO XU¹, CHAO LIU² and JINKU BAO³

¹College of Bioindustry, Chengdu University, Chengdu, Sichuan 610106; ²College of Life Science, Leshan Normal University, Leshan, Sichuan 614000; ³State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, Chengdu, Sichuan 610041, P.R. China

Received August 22, 2015; Accepted August 12, 2016

DOI: 10.3892/mmr.2017.6139

Abstract. Increasing evidence suggests that microRNAs (miRNAs) are essential in carcinogenesis, therefore, the present study investigated the role of hsa-miR-101 in renal tumorigenesis and cancer development. On identification of its expression pattern, it may serve as a diagnostic biomarker for clear cell renal cell carcinoma (ccRCC). In the present study, 10 pairs of ccRCC and noncancerous tissue samples were obtained to examine whether the expression of hsa-miR-101 is linked to cancer. The data obtained were validated using reverse transcription-quantitative polymerase chain reaction analysis. The levels of hsa-miR-101 were examined following exposure to hypoxia in ACHN and HK-2 cells. As a predicted target, the mRNA and protein levels of TP53-induced glycolysis and apoptosis regulator (TIGAR) were then assessed. A pcDNA-GFP-miR-101 plasmid was stably transfected into ACHN and HK-2 cells, following which the effects of hsa-miR-101 on the expression of TIGAR and inhibition of glycolysis were investigated. The present study also examined the association between the level of hsa-miR-101 and kidney tumors. It was identified that the expression level was significantly higher in the ccRCC tissues, compared with that in the corresponding noncancerous tissues. The expression values for the upregulated miRNA ranged between 4.6- and 67.9-fold. On demonstrating the functional link between hypoxia and the expression of miRNAs changes in the expression of hsa-miR-101 were examined following hypoxia exposure in kidney tumor and non-tumor cell lines. It was shown that hypoxia exposure significantly induced hsa-miR-101. The hypoxia-induced upregulation of hsa-miR-101 repressed the activity of TIGAR by targeting TIGAR mRNA and promoting glycolysis. The results showed that the upregulation of hsa-miR-101 in ccRCC was induced by hypoxia. Its expression decreased the protein expression of

TIGAR and promoted glycolysis. This regulatory pathway may represent a novel mechanism of carcinogenesis and requires further investigation.

Introduction

Renal cell carcinoma (RCC) is a common urologic malignancy, which accounts for 2-3% of adult malignancies, and clear cell RCC (ccRCC) is the most frequent of all types of RCC, which accounts for 85-95% of cases (1). The mortality rate of ccRCC is >90,000/year worldwide, and particularly high in China. Histological subtype detection shows ccRCC represents ~75-80% of all cases of RCC (1). Over the past decade, the incidence of ccRCC has increased significantly with an mortality rate of ~40% (2). In previous years, studies investigating the biological mechanisms, which cause ccRCC carcinogenesis focussed on genomic mutations, expression of protein-coding genes and epigenetic changes (3). However, there is increasing evidence suggesting that the microRNAs (miRNAs) are closely associated with cancer, and that the changes in their expression profiles may be considered as biomarkers for early detection, accounting for early diagnosis. Therefore, an improved understanding of miRNAs associated with the progression of ccRCC may lead to more accurate prognosis and therapeutic strategies (4).

miRNAs are a class of small noncoding RNAs, which have been shown to regulate gene expression by binding through partial sequence homology to the 3' untranslated region (3' UTR), causing target mRNA degradation or translational inhibition (5). The majority of miRNAs are transcribed as long monocistronic, bicistronic or polycistronic primary transcription units (pri-miRNAs) by RNA polymerase II and, following a series of cellular processing events, mature miRNAs are synthesized (6). The mature miRNAs are incorporated into the RNA-induced silencing complex and drives the selection of mRNAs of interest containing antisense sequences matching those of the miRNAs (7). As a result of their gene expression-associated activity, miRNAs have been identified as key regulators of several biological processes, including development, differentiation, apoptosis, proliferation and carcinogenesis (8-10). miRNAs are found aberrantly expressed or mutated in cancer, suggesting their role as a novel class of factors involved in promoting or inhibiting carcinogenesis (11,12). In

Correspondence to: Professor Jinku Bao, State Key Laboratory of Biotherapy and Cancer Center, West China Hospital, Sichuan University, 4 Keyuan Road, Chengdu, Sichuan 610041, P.R. China
E-mail: baojinku@scu.edu.cn

Key words: clear cell renal cell carcinoma, hsa-miR-101, TP53-induced glycolysis and apoptosis regulator, glycolysis, hypoxia

several types of cancer, miRNAs show significant differences, including prostate, lung, breast, colon and kidney malignancies (13), and evidence supports the promising use of miRNAs for diagnostic, prognostic and predictive purposes (14).

Hypoxia is an essential feature of the neoplastic micro-environment. The oxygen concentration in tumor tissue is substantially lower, compared with that in the surrounding non-tumor tissue. For example, in solid tumors of the breast, the O₂ pressure is ~10 mmHg, whereas normal breast tissue has an O₂ pressure of >60 mmHg (15). The regulation of hypoxia in cells occurs through affecting the expression of certain genes, which are sensitive to O₂ concentration. The molecular mechanisms responsible for the hypoxic survival of cancer cells remains to be fully elucidated, and additional information on this process may lead to novel strategies for pharmacological intervention.

In order to determine the possible role of hsa-miR-101 in hypoxic gene regulation and to examine this hypoxically-regulated miRNA, which may promote glycolysis by inhibiting TP53-induced glycolysis and apoptosis regulator (TIGAR), the present study examined changes in miRNA expression levels in response to hypoxia. The results obtained characterized in detail the hypoxic regulation of hsa-miR-101 and the mechanism underlying its regulation of TIGAR and promotion of glycolysis. The data also revealed the association between the level of hsa-miR-101 and the pathogenesis and prognosis of ccRCC.

Materials and methods

Patients and tissue samples. In total, 15 pairs of ccRCC tumor tissue and normal tissue samples were collected between July 2012 and May 2013 at the department of Urological Surgery, First People's Hospital of Beijing (Beijing, China). All patients who donated tissue were diagnosed with ccRCC via post-operative pathology. The current study was approved by the Ethics Committee of Xutong Hospital (Xutong, China).

Of the 15 tissue samples from the patients with ccRCC, seven were from men and eight were from women, with a mean age of 58 years (range, 36-71 years). Of these patients, nine were in clinical stage I and six were in stage II.

Cell culture and induction of hypoxia. In the present study, the normal HK-2 epithelial cell line (frozen in the laboratory) and the ACHN ccRCC cell line were used, obtained from American Type Culture Collection (ATCC; Manassas, VA, USA). The cells were grown in Eagle's minimum essential medium (Life Technologies; Thermo Fisher Scientific, Inc., Waltham, MA, USA) supplemented with 10% fetal bovine serum, 100 U/ml penicillin and 100 mg/ml streptomycin (Life Technologies; Thermo Fisher Scientific, Inc.) in a 5% CO₂ atmosphere at 37°C according to standard procedures. Hypoxia exposure was established by changing the medium, which had been gassed to equilibrium with 0.1% O₂, decreasing CO₂ flow rates to 10 cm³/sec in specifically designed chambers obtained from Oxoid (Adelaide, Australia).

Transfection. To generate transiently-expressing constructs, the genomic region surrounding the pri-miRNA sequence of hsa-miR-101 was amplified using primers the following primers: Forward 5'-CTCAACTGGTGTCTGGAGTCGGC

AATTCAGTTGAGTTCAGTTAT-3' and reverse 5'-ACACTC CAGCTGGGTACAGTACTGTGATAA-3'. The thermocycling protocol was as follows: 95°C for 5 min, then 30 cycles of amplification, each cycle consisted of 94°C for 30 sec, 55°C for 45 sec, and 72°C for 1 min. This generated a polymerase chain reaction (PCR) product of ~500 bp, which was directionally cloned into pLNCX (Clontech Laboratories, Mountain View, CA, USA) using *NotI* and *XhoI* (Takara Bio, Inc., Shiga, Japan). The ligated construct was transformed into the DH5α *E. coli* strain (ATCC) chemically by preparing endotoxic-free plasmid using Maxipreps (Qiagen, Hilden, Germany), packed with Lipofectamine 2000 (Thermo Fisher Scientific, Inc.) and mixed with Phoenix packaging cells (Orbigen, Inc., San Diego, CA, USA). Sequencing was performed by Qingkezixi (Chengdu, China) to confirm the insertion of the target sequence into the plasmid following the cytomegalovirus promoter. The supernatant was obtained following centrifugation at 800 x g for 10 min at 4°C was collected and used to infect the target cells, which were incubated in a 5% CO₂ atmosphere at 37°C, and these cells were harvested 48 h later, for further analysis.

RNA extraction and cDNA synthesis. Total RNA was extracted from the tissues by homogenizing 100 mg of the frozen tissue with a homogenizer (Thermo Fisher Scientific, Inc.), followed by isolation using the mir-VANA miRNA Isolation kit (Ambion, Austin, TX, USA). For reverse transcription (RT), 1 μg of the isolated miRNAs were incubated with RT mixture from Reverse-Transcriptional Fast kit (RiboBio Co., Ltd., Guangzhou, China) to simultaneously convert all small RNAs into detectable cDNAs for PCR analysis.

TaqMan RT-qPCR and quantitative (q)PCR. The method was optimized for miRNA, and all reagents, primers and probes were obtained from Applied Biosystems; Thermo Fisher Scientific, Inc. Human 28S rRNA was used to normalize all RNA samples. The RT reactions and qPCR analyses were performed according to the manufacturer's protocols. The previously prepared cDNA samples were run in duplicate in an Applied Biosystems 9700 thermocycler. The Taqman probe for hsa-miR-101 was 5'-CGGCGGTACAGTACTGTGATAA-3'; miR-93 5'-CGGCGGTGGAGTGTGACAATGG-3'. The thermocycling conditions were: 95° for 5 min, followed by 35 cycles at 95°C for 10 sec, 55°C for 10 sec and 72°C for 1 min. The gene expression levels were quantified using the ABI Prism 7900HT sequence detection system (Applied Biosystems; Thermo Fisher Scientific, Inc.). Analysis was performed in duplicate, including controls containing no template. The relative expression was calculated using the comparative quantification cycle method (16).

Total RNA was extracted from the cells using TRIzol reagent (Life Technologies; Thermo Fisher Scientific, Inc.), and a 1 μg from each cell line were used for RT (Life Technologies; Thermo Fisher Scientific, Inc.) according to the manufacturer's protocol. Subsequent qPCR was performed in the Bio-Rad CFX qPCR system (Bio-Rad Laboratories, Inc., Hercules, CA, USA) according to the manufacturer's protocol. The protocol used was as follows: 98°C 5 min, followed by 40 cycles of 98°C for 10 sec and 60°C for 1 min. The prediction of potential target for microRNAs was performed by using TargetScan (www.targetscan.org). The primers used

Figure 1. Detection of expression levels of hsa-miR-101 in ccRCC and adjacent tissue samples. (A) RT-qPCR analysis of the expression of hsa-miR-101 in 15 pairs of ccRCC and corresponding noncancerous tissue samples. (B) Conventional stem-loop semi-quantitative PCR analysis of the expression of hsa-miR-101 in ccRCC and corresponding noncancerous tissue samples, and ACHN and HK-2 kidney cell lines. U6 RNA was used as an internal control. All experiments were performed in triplicate. * $P < 0.01$. ccRCC, clear cell renal cell carcinoma; T, tumor tissue; N, noncancerous tissue; miR, microRNA; RT-qPCR, reverse transcription-quantitative polymerase chain reaction.

for qPCR were as follows: TIGAR, forward 5'-GTGAGGACT ACGCAGCATCA-3' and reverse 5'-GCATCAGAACCGTGA TATATTCT-3'; β -actin, forward 5'-GCGCGTGCCTTCATC AC-3' and reverse 5'-TCTGCGCCATAAGGTGGTAG-3'.

Semi-quantitative western blot analysis. To prepare the total protein samples, the cells were trypsinized and homogenized on ice in lysis buffer containing 10 mM Tris-HCl (pH 7.4), 1% Triton X-100, 150 mM NaCl and 100 mM KCl. The crude lysate was centrifuged at 20,000 $\times$ g for 45 min at 4°C and ~200 μ l of clear supernatant was collected. A bicinchoninic acid assay was performed to identify the total protein concentration of the collected supernatant. A final concentration of 2 μ g/ μ l total protein was adjusted with the addition of ddH₂O and 1X laemmli buffer and boiled at 98°C for 10 min. To fractionate the proteins, 4-12% gradient SDS-PAGE gels were used. Immunoblotting was performed with specific antibodies, including mouse anti-TIGAR antibody (cat. no. ab64622; 1:1,000; Abcam, Cambridge, UK), anti- β -actin antibody (cat. no. ab8226; 1:1,000; Abcam, Cambridge, UK) at 4°C overnight. The PVDF membranes were then incubated with goat anti-mouse horseradish peroxidase-labeled secondary antibody (cat. no. ab97040; 1:5,000; Abcam, Cambridge, UK) incubated at room temperature.

Cell proliferation assay. Cell proliferation was detected using 3-(4, 5-dimethylthazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT). The cells were seeded into 24-well plates (1.2 $\times$ 10⁴ cells/well) and allowed to attach overnight in a 5% CO₂ atmosphere at 37°C. After 24, 48, 72 and 96 h, cell viability was assessed using an MTT assay. The absorbance at 490 nm (A490) of each well was read on a spectrophotometer. Three independent experiments were performed in quadruplicate.

Metabolite determination. To measure the levels of fructose (Fru)-2,6-P₂, the treated cells were homogenized in

homogenizing buffer (100 mM NaOH and 0.1% Triton X-100), following which the crude lysate was heated to 80°C for 10 min and centrifuged at 20,000 $\times$ g for 10 min at 4°C. The supernatant was transferred to a fresh 1.5 ml tube. Fru-2,6-P₂ was identified in the supernatants by its ability to activate pyrophosphate-dependent phosphofructokinase-1 from potato tubers, as described previously (17). Standard enzymatic methods for measuring lactate spectrophotometrically in the neutralized perchloric extracts were performed according to the protocol described previously (18).

Statistical analysis. All statistical analyses were performed using SPSS 13.0 software (SPSS Inc., Chicago, IL, USA). The differences between the tumor and adjacent noncancerous tissues were assessed using the paired samples *t*-test. The differences between the tissues of patients with ccRCC were assessed using an independent samples *t*-test or non-parametric test. The clinical correlation analysis was performed using the variance test or Kruskal-Wallis test. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

hsa-miR-101 is significantly downregulated in human ccRCC tissues. In the present study, a stem-loop RT-qPCR assay was performed to identify the expression level of hsa-miR-101 in 15 pairs of matched ccRCC and noncancerous kidney tissue samples. As shown in Fig. 1A, significantly upregulated expression levels of hsa-miR-101 were detected in the ccRCC tissues, compared with the noncancerous tissue. The expression levels of hsa-miR-101 in 12 ccRCC tissue samples were upregulated between 4.6- and 17.9-fold, whereas the levels in three ccRCC tissue samples were downregulated between 0.2- and 0.7-fold. Two pairs of the matched ccRCC and noncancerous tissue samples, and the normal HK-2 epithelial cell line and ACHN

Figure 2. Time course and oxygen sensitivity of hsa-miR-101 following the induction of hypoxia. (A) ACHN and HK-2 cells were exposed to 1% oxygen for 0, 1, 3, 6, 12, 24 and 48 h. All time points were performed in triplicate. (B) ACHN and HK-2 cells were exposed for 16 h to oxygen concentrations of 0.1, 1, 3, 5 and 21%. All concentrations were performed in triplicate. hsa-miR-93 was considered a negative control, which is not affected by hypoxia. The expression levels of hsa-miR-101 and hsa-miR-93 were measured using reverse transcription-quantitative polymerase chain reaction analysis. The data are expressed as the mean fold difference in microRNA levels between the treated and untreated samples. * $P < 0.05$ and ** $P < 0.01$. miR, microRNA.

ccRCC cell line were used for further conventional RT-qPCR analyses, and the results were analyzed by gel electrophoresis. The results showed that the expression of hsa-miR-101 was upregulated in the ccRCC tissues (Fig. 1B). Of note, no significant differences were found in the expression of hsa-miR-101 in the ACHN cells, compared with the HK-2 cells, indicating the induction of hsa-miR-101 by other factors.

Confirmation of the hypoxic upregulation of hsa-miR-101 in ccRCC cells using RT-qPCR analysis. The present study hypothesized that hsa-miR-101 is involved in the hypoxic response, and this was investigated by examining the level of hsa-miR-101 following hypoxic exposure. The time course of the induction of hsa-miR-101 by hypoxia was examined. The ACHN and HK-2 cells were cultured in hypoxia (1% O₂) for 1, 3, 6, 12, 24 and 48 h. The expression of hsa-miR-101 was analyzed using TaqMan RT-qPCR analysis. Following incubation for 6 h, the upregulation of hsa-miR-101 by hypoxia was detected and showed a progressive increase in expression (Fig. 2A). Following hypoxia exposure for 12 h, the upregulation of hsa-miR-101 was significant (ACHN, 4.19 ± 0.17 ; HK-2, 2.67 ± 0.35 ; $P < 0.001$) and was maximal at the 48 h time point (ACHN, 5.79 ± 0.21 ; HK-2, 3.43 ± 0.25 ; $P < 0.001$). As a negative control, the expression of hsa-miR-93, which is not affected by hypoxic incubation (19), was completely unaffected (data not shown). To investigate the oxygen-dependent regulation of hsa-miR-101, a range of oxygen concentrations (0.1, 1, 3, 5 and 21% oxygen) were used for cell maintenance. The induction of hsa-miR-101 was most marked at 1% oxygen, with more modest regulation at 0.1, 3 and 5% oxygen, following hypoxic exposure for 24 h (Fig. 2B). No detectable change in the expression of hsa-miR-93 was observed following hypoxia exposure.

Overexpression of hsa-miR-101 knocks down the expression of TIGAR, and affects kidney cell metabolism and growth rate. According to the stem-loop character of hsa-miR-101 and cross-species comparison, computational algorithms have been developed. TargetScan was used to generate a potential miRNA with a high probability of binding to the 3'UTR of TIGAR. To further confirm the predicted result, a hsa-miR-101-expressing plasmid was constructed and transfected into ACHN and HK-2 cells. Total RNA was extracted from the stably transfected cells, and RT-qPCR analysis was performed. The data showed that the overexpression of hsa-miR-101 reduced the expression of TIGAR at the mRNA and protein levels (Fig. 3A and B).

TIGAR is important as a regulator of oxidative stress, therefore, the present study next examined the biological consequences of the specific knockdown of TIGAR by hsa-miR-101 in ACHN and HK-2 cells. Metabolic parameters associated with the function of TIGAR were determined, and the levels of Fru-2, 6-P₂ and lactate were significantly increased in the hsa-miR-101-overexpressing cells following transfection (Fig. 3C). The present study also evaluated the effects of the overexpression of hsa-miR-101 on the growth rate of the transfected ACHN and HK-2 cells. As shown in Fig. 3D, the ACHN/hsa-miR-101 and HK-2/hsa-miR-101 cell lines had significantly increased cell viability, compared with the mock-transfected ACHN and HK-2 cells ($P < 0.05$).

Discussion

Increasing reports have shown the close association between disturbances in the expression levels of miRNAs and the carcinogenic process (20-22). These miRNAs, the mutation or

Figure 3. Overexpression of hsa-miR-101 affects the metabolism and growth rate of clear cell renal cell carcinoma and normal kidney cell lines. (A) Reverse transcription-quantitative polymerase chain reaction and (B) western blot analyses of the expression of TIGAR in mock-, hsa-miR-101- or hsa-miR-93-stably transfected ACHN and HK-2 cells. (C) Intracellular Fru-2, 6-P₂ and lactate were detected. (D) Proliferation rates of cells were assayed by performing a 3-(4, 5-dimethylthazol-2-yl)-2, 5-diphenyltetrazolium bromide assay. All experiments were performed in triplicate. *P<0.01 vs. untreated cells. microRNA; TIGAR, TP53-induced glycolysis and apoptosis regulator; OD, optical density.

misexpression of which correlate with various types of human cancer, are referred to as oncomiRs (11). They can be involved in tumor suppression or formation.

The expression pattern of miRNAs may be altered during the progressive development of ccRCC. Gowrishankar *et al* (23) examined the differences in expression levels of >700 human miRNAs in a series of 94 ccRCC samples, and found that the accumulation of miR-21-5p and miR-142-3p were reduced. The overexpression of these miRNAs leads to proliferation and decreased cell death, suggesting their role as oncogenes. Previously, Cheng *et al* (24) showed that miR-34a and miR-224 were upregulated and had an anti-apoptotic effect in ccRCC. This suggests that these miRNAs may be involved in tumor suppression.

The effects of TIGAR on cell proliferation and glycolysis are considered to be cell- and context-dependent. Following moderate levels of stress, TIGAR is induced for repairing DNA damage and inhibiting glycolysis. Peña-Rico *et al* (25) showed that the silencing of TIGAR in glioblastoma cell lines causes higher levels of Fru-2,6-P₂ in cells, consequently increasing glycolysis and leading to the accumulation of

reactive oxygen species, which promotes cell death. In tumor tissues, TIGAR has been found to be markedly downregulated (26), which is predominantly caused by p53 dysfunction. However, the mechanism underlying the downregulation of TIGAR in tumors with functioning p53 remains to be elucidated, however, it is hypothesized to be involved with miRNAs.

The aim of the present study was to evaluate whether there is an association between hypoxia exposure and altered expression patterns of hsa-miR-101 in ccRCC. The results showed that hsa-miR-101 was frequently downregulated in human ccRCC tissues, compared with corresponding noncancerous kidney tissues, which is consistent with the results of Wotschovsky *et al* (27). In addition, hsa-miR-101 has been reported to be upregulated in HepG2 cells (28) and periodontal ligament cells (29), indicating its close association with different types of tumors. However, the mechanism causing the upregulation of hsa-miR-101 and the effect of the higher level of this miRNA remains to be elucidated.

Due to the close association between the induction of miRNA with hypoxia, the present study investigated whether

hsa-miR-101 is induced by hypoxia. The resulting data showed that hsa-miR-101 was induced by hypoxia *in vitro*. Consistent with previous results, hsa-miR-93 showed no responsiveness to hypoxia *in vitro*. The effects of alterations in the levels of hsa-miR-101 remain to be fully elucidated. Resultant modifications in target gene expression are possible, however, the relative effects on transcription, mRNA post-transcriptional regulation and translation remain to be elucidated. The computational prediction of mRNA targets for hsa-miR-101 binding and action generates a number of potential targeted mRNA sequences (www.MicroRNAs.org/microRNAs/home.do and www.targetscan.org). As TIGAR mRNA showed the highest binding affinity, it was selected for further experiments. The data obtained showed that the overexpression of hsa-miR-101 decreased the mRNA and protein levels of TIGAR, causing the stimulation of glycolysis and proliferation. The correlation between the levels of hsa-miR-101 and ccRCC was found to be marked. This may be due to the direct effect of hsa-miR-101 on tumor biology through its induction by tumor hypoxia. Of note, as a feature of a solid tumors, hypoxia induces hsa-miR-101 to target TIGAR, causing changes in metabolism. This indicates one mechanism involved in the regulation responses of miRNAs to microenvironmental factors.

The results of the present study showed that a significantly high proportion of hsa-miR-101 is overexpressed in human tumor tissues, and the alteration of this miRNA is caused by hypoxia *in vitro* and *in vivo*. By regulating its target gene, TIGAR, the overexpression of hsa-miR-101 stimulates glycolysis and increases proliferation. In conclusion, the present study may provide a novel therapeutic target site for ccRCC through targeting hsa-miR-101.

Acknowledgements

The authors would like to thank Dr Ziyi Zhao (Sichuan University, Chengdu, China) for their English editing, Dr Changjin Chen (Sichuan University, Chengdu, China) for his technical support and Miss Jiao Lv (Sichuan University, Chengdu, China) for her technical assistance.

References

- Jemal A, Siegel R, Ward E, Murray T, Xu J, Smigal C and Thun MJ: Cancer statistics, 2006. *CA Cancer J Clin* 65: 106-130, 2006.
- Gottardo F, Liu CG, Ferracin M, Calin GA, Fassan M, Bassi P, Seviquani C, Byrne D, Negrini M, Pagano F, *et al*: Micro-RNA profiling in kidney and bladder cancers. *Urol Oncol* 25: 387-392, 2007.
- Brugarolas J: Molecular genetics of clear-cell renal cell carcinoma. *J Clin Oncol* 32: 1986-1976, 2014.
- Cummins JM and Velculescu VE: Implications of micro-RNA profiling for cancer diagnosis. *Oncogene* 25: 6220-6227, 2006.
- Garzon R, Pichiorri F, Palumbo T, Visentini M, Ageilan R, Cimmino A, Wang H, Sun H, Volinia S, Alder H, *et al*: MicroRNA gene expression during retinoic acid-induced differentiation of human acute promyelocytic leukemia. *Oncogene* 26: 4148-4157, 2007.
- Siomi H and Siomi MC: On the road to reading the RNA-interference code. *Nature* 457: 396-404, 2009.
- Cullen BR: Transcription and processing of human microRNA precursors. *Mol Cell* 16: 861-865, 2004.
- Bhaskaran M and Mohan M: MicroRNAs: History, biogenesis, and their evolving role in animal development and disease. *Vet Pathol* 51: 759-774, 2014.
- Beroli G, Cava C and Castiglioni I: MicroRNAs: New biomarkers for diagnosis, prognosis, therapy prediction and therapeutic tools for breast cancer. *Theranostics* 5: 1122-1143, 2015.
- Su Z, Yang Z, Xu Y, Chen Y and Yu Q: MicroRNAs in apoptosis, autophagy and necroptosis. *Oncotarget* 6: 8474-8490, 2015.
- Esquela-Kerscher A and Slack FJ: Oncomirs-microRNAs with a role in cancer. *Nat Rev Cancer* 6: 259-269, 2006.
- Fulci V, Chiaretti S, Goldoni M, Azzalin G, Carucci N, Tavoraro S, Castellano L, Magrelli A, Citarella F, Messina M, *et al*: Quantitative technologies establish a novel microRNA profile of chronic lymphocytic leukemia. *Blood* 109: 4944-4951, 2007.
- Volinia S, Calin GA, Liu CG, Ambs S, Cimmino A, Petrocca F, Visone R, Iorio M, Roldo C, Ferracin M, *et al*: A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci USA* 103: 2257-2261, 2006.
- Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, Peck D, Sweet-Cordero A, Ebert BL, Mak RH, Ferrando AA, *et al*: MicroRNA expression profiles classify human cancers. *Nature* 435: 834-838, 2005.
- Vaupel P, Mayer A and Höckel M: Tumor hypoxia and malignant progression. *Methods Enzymol* 381: 335-354, 2004.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
- Van Schaftingen E, Lederer B, Bartrons R and Hers HG: A kinetic study of pyrophosphate: Fructose-6-phosphate phosphotransferase from potato tubers. Application to a microassay of fructose 2,6-bisphosphate. *Eur J Biochem* 129: 191-195, 1982.
- Gutmann I and Wahlefeld AW: *Methods of Enzymatic Analysis* (Bergmeyer, H.U., ed.), Academic Press, London, 2nd ed. v3: 1464-1468, 1974.
- Camps C, Buffa FM, Colella S, Moore J, Sotiriou C, Sheldon H, Harris AL, Gleadle JM and Ragoussis J: hsa-miR-210 is induced by hypoxia and is an independent prognostic factor in breast cancer. *Clin Cancer Res* 14: 1340-1348, 2008.
- Hung CH, Chiu YC, Chen CH, Hu TH: MicroRNAs in hepatocellular carcinoma: Carcinogenesis, progression, and therapeutic target. *Biomed Res Int* 486407, 2014.
- Lan H, Lu H, Wang X and Jin H: MicroRNAs as potential biomarkers in cancer: Opportunities and challenges. *Biomed Res Int* 125094, 2015.
- Tan G, Tang X and Tang F: The role of microRNAs in nasopharyngeal carcinoma. *Tumor Biol* 36: 69-79, 2015.
- Gowrishankar B, Ibragimova I, Zhou Y, Slifker M, Devarajan K, Alsalem T, Uzzo R and Cairns P: MicroRNA expression signatures of stage, grade, and progression in clear cell RCC. *Cancer Biol Ther* 15: 329-341, 2014.
- Cheng T, Wang L, Li Y, Huang C, Zeng L and Yang J: Differential microRNA expression in renal cell carcinoma. *Oncol Lett* 6: 769-776, 2013.
- Peña-Rico MA, Calvo-Vidal MN, Villalonga-Planells R, Martínez-Soler F, Giménez-Bonafé P, Navarro-Sabaté A, Tortosa A, Bartrons R and Manzano A: TP53 induced glycolysis and apoptosis regulator (TIGAR) knockdown results in radiosensitization of glioma cells. *Radiother Oncol* 101: 132-139, 2011.
- Bensaad K, Tsuruta A, Selak MA, Vidal MN, Nakano K, Bartrons R, Gottlieb E and Vousden KH: TIGAR, a p53-inducible regulator of glycolysis and apoptosis. *Cell* 126: 107-120, 2006.
- Wotschovsky Z, Liep J, Meyer HA, Jung M, Wagner I, Disch AC, Schaser KD, Melcher I, Kilic E, Busch J, *et al*: Identification of metastasis-associated microRNAs in clear cell renal cell carcinomas. *Int J Biol Sci* 8: 1363-1374, 2012.
- Li C, Li C, Yue J, Huang X, Chen M, Gao J and Wu B: miR-21 and miR-101 regulate PLAP-1 expression in periodontal ligament cells. *Mol Med Rep* 5: 1340-1346, 2012.
- Chiang CW, Huang Y, Leong KW, Chen LC, Chen HC, Chen SJ and Chou CK: PKCalpha mediated induction of miR-101 in human hepatoma HepG2 cells. *J Biomed Sci* 17: 35, 2010.