Overexpression of Tim-3 reduces *Helicobacter pylori*-associated inflammation through TLR4/NFκB signaling *in vitro*

FUCAI WANG^{1,2*}, ZHIRONG MAO^{1*}, DONGSHENG LIU^{1*}, JING YU¹, YOUHUA WANG¹, WEN YE^{1,2}, DONGJIA LIN², NANJIN ZHOU³ and YONG XIE¹

¹Department of Gastroenterology, The First Affiliated Hospital of Nanchang University, Gastroenterology Institute of Jiangxi, Key Laboratory of Digestive Diseases of Jiangxi; ²Department of Immunology, Medical College of Nanchang University; ³Institute of Immunology and Biological Therapy, Jiangxi Academy of Medical Sciences, Nanchang, Jiangxi 330006, P.R. China

Received January 10, 2016; Accepted January 19, 2017

DOI: 10.3892/mmr.2017.6346

Abstract. The present study aimed to investigate the interaction between T-cell immunoglobulin and mucin-domain-containing molecule-3 (Tim-3) and Toll-like receptor 4 (TLR4)/nuclear factor KB (NF-KB) signaling in Helicobacter pylori-infected RAW264.7 macrophage cells. RAW264.7 cells were co-cultured with H. pylori SS1 at different bacteria/cell ratios, and subsequently the mRNA expression of Tim-3, TLR4, and myeloid differentiation factor 88 (MyD88) was measured by reverse transcription-quantitative polymerase chain reaction (RT-qPCR). Furthermore, the effect of Tim-3 overexpression was examined by transfection of RAW264.7 with pLVX-IRES-ZsGreen-Tim-3 and co-culturing with H. pylori. mRNA and protein expression levels were then analyzed for Tim-3, TLR4, MyD88, and phosphorylated (p-) NF-KB by RT-qPCR and western blot analysis respectively. The concentrations of pro-inflammatory cytokines [tumor necrosis factor-a (TNF-a), interleukin 6 (IL-6), interferon- γ (IFN- γ) and interleukin 10 (IL-10)] released in the culture supernatants were measured by ELISA. H. pylori stimulation resulted in a significant increase of Tim-3, TLR4, and MyD88 mRNA expression in RAW264.7 cells. H.

Correspondence to: Professor Yong Xie, Department of Gastroenterology, The First Affiliated Hospital of Nanchang University, Gastroenterology Institute of Jiangxi, Key Laboratory of Digestive Diseases of Jiangxi, 17 Yongwaizheng Street, Nanchang, Jiangxi 330006, P.R. China E-mail: xieyong_med@163.com

Professor Nanjin Zhou, Institute of Immunology and Biological Therapy, Jiangxi Academy of Medical Sciences, 461 Bayi Road,

Nanchang, Jiangxi 330006, P.R. China

E-mail: jxznj@163.com

*Contributed equally

Key words: T-cell immunoglobulin and mucin-domain-containing molecule-3, toll-like receptor 4, *Helicobacter pylori*, inflammation

pylori stimulation upregulated Tim-3 expression even in the Tim-3-overexpressing RAW264.7 cells compared with unstimulated cells. TLR4, MyD88, and pNF-κB protein expression and pro-inflammatory cytokines (TNF- α , IL-6, and IFN- γ) release levels were increased in the control RAW264.7 cells following *H. pylori* infection, but not in the Tim-3-overexpressing RAW264.7 cells. By contrast, IL-10 levels were decreased following *H. pylori* infection in both control and Tim-3-overexpressing RAW264.7 cells. Overexpression of Tim-3 reduced *H. pylori*-associated inflammation in RAW264.7 macrophages, by downregulating expression of proteins in the TLR4 pathway and release of pro-inflammatory cytokines. These findings suggest that Tim-3 serves a crucial role in the negative regulation of *H. pylori*-associated inflammation and may be a novel therapeutic target for *H. pylori* infection.

Introduction

Chronic infection by *Helicobacter pylori* is achieved through the colonization of an almost exclusive niche and through evading detection by the host's cellular immune defense mechanisms (1). *H. pylori* is the only microorganism known to colonize the human stomach and to inhabit gastric mucosal cells. To achieve this colonization, *H. pylori* must escape detection by both innate and adaptive immune responses (2). Macrophages, phagocytic cells that are part of the innate immune system, are located in various tissues. A great number of cytokines assist macrophages in their role as custodians of the innate to adaptive immunity (3). Therefore, macrophages may serve an important role in *H. pylori* infection.

Toll-like receptor (TLR) 4, a pattern recognition receptor, recognizes pathogen-associated molecular patterns. It activates macrophages to secrete cytokines, which initiate and regulate the immune response (4). T-cell immunoglobulin and mucin-domain-containing molecule 3 (Tim-3) is an important member of the TIM family, a recently discovered family of trans-membrane proteins (5). Previous studies have demonstrated that Tim-3 serves a role in the differentiation of T helper (Th) 1 cells from Th2 cells; upon activation, Tim-3

downregulates Th1 cell function (5-8). Several studies have also demonstrated that TLR4 and Tim-3 are expressed on macrophages and that they participate in the regulation of cytokine secretion from macrophages (9-11), suggesting that Tim-3 impacts macrophage function by interacting with the TLR4 signaling pathway. However, it is unclear to date how *H. pylori* infection impacts the interaction of Tim-3 and TLR4 in macrophages.

Based on these previous studies, an investigation of the interaction between Tim-3 and TLR4 signaling pathways in H. *pylori*-associated inflammation was undertaken. The present study aimed to provide theoretical and experimental evidence to support Tim-3 as a potential therapeutic target for the control and prevention of H. *pylori* infection-related diseases.

Materials and methods

Cell lines and bacteria. The murine macrophage RAW264.7 cell line was obtained from the Gastroenterology Institute of Jiangxi Province (Nanchang, China) and was cultured in DMEM with 10% FBS, 100 U/ml penicillin (Gibco, Thermo Fisher Scientific, Inc., Waltham, MA, USA) and 100 μ g/ml streptomycin (Gibco, Thermo Fisher Scientific, Inc.) under 5% CO₂ at 37°C in a humidified atmosphere. The *H. pylori* standard strain SS1 (CagA⁺, VacA⁺) was obtained from the Chinese Center for Disease Control (Beijing, China) and was grown on a Campylobacter agar base from the Chinese Research Center for Diarrhoeal Disease Control (Shanghai, China), containing 10% sheep blood, under microaerobic conditions (5% O₂, 10% CO₂ and 85% N₂) at 37°C for 2-3 days.

Cell transfection. Murine Tim-3 genes was amplified by polymerase chain reaction (PCR) using the following primer: Forward, 5'-CAACTCGAGATGTTTTCAGGT CTTACCCT-3' and reverse, 5'-AACGGATCCTCAGGA TGGCTGCTGGCTGT-3'. The PCR products were purified to generate the plasmid pLVX-IRES-ZsGreen1 (Clontech Laboratories, Inc., Mountainview, CA, USA). The ligated products were transformed into TOP10 chemically competent Escherichia coli (Takara Bio, Inc., Otsu, Japan) and incubated on Luria-Bertani plates containing 100 μ g/ml ampicillin at 37°C overnight. Subsequently, eight putative ampicillin-resistant positive clones were selectedå to amplify, extract and purify for PCR amplification and electrophoresis detection. The pLVX-IRES-ZsGreen1-Tim-3 plasmid was constructed and the sequencing was completed by Shanghai ShengGong Biotechnology Co., Ltd. (Shanghai, China). Then the RAW264.7 cells were plated at a density of $2x10^5$ cells/ml in a 6-well plate, grown overnight, and transferred to serum-free medium prior to Tim-3 transfection. The RAW264.7 cells were transfected with 2.5 µg pLVX-IRES-ZsGreen1-Tim-3 plasmid or empty vector using Lipofectamine® LTX and Plus Reagent (15,338,100; Invitrogen; Thermo Fisher Scientific, Inc.) according to the manufacturer's protocols. Cells were observed for green fluorescent protein using fluorescence microscopy after 42 h transfection. The control cells were grown in serum-free medium alone. Tim-3 overexpression was confirmed by dual-endonuclease digestion and sequencing.

Reverse transcription (RT)-PCR. Total RNA was extracted from RAW264.7 cells in 6-well plate using an E.Z.N.A. Total RNA kit II (Omega BioTek, Inc., Norcross, GA, USA) according to the manufacturer's protocols. RNA quality was evaluated by 1% agarose gel electrophoresis. Using a Quantiscript RT Kit (Tiangen Biotech Co., Beijing, China), single-strand cDNA was synthesized and then used as a template for PCR amplification of Tim-3, TLR4, myeloid differentiation factor 88 (MyD88) and β -actin (used for normalization). The $2^{-\Delta\Delta Cq}$ method (12) was used to determine the relative quantities of products. Primer sequences were as follows: Tim-3, forward 5'-ACTCTACCTACATCTGGG ACACT-3' and reverse 5'-TAGGTCCCATGGTCATCCAG-3'; TLR4, forward 5'-AGAAACGGCAACTTGGACCT-3' and reverse 5'-GGCCTTAGCCTCTTCTCCT-3'; MyD88, forward 5'-CTGGCCTTGTTAGACCGTGA-3' and reverse 5'-TCG AAAAGTTCCGGCGTTTG-3'; and β-actin, forward 5'-GAG ACCTTCAACACCCCAGC-3' and reverse 5'-ATGTCACGC ACGATTTCCC-3'. Each 25 µl PCR consisted of 10 pmol of each primer, 10 ul 2xTaq Master Mix, 3 ul template, and 10 μ l ddH₂O. β-actin and MyD88 were amplified at 94°C for 90 sec (1 cycle); 94°C for 30 sec, 61°C for 30 sec and 72°C for 1 min (30 cycles); and 72°C for 5 min (1 cycle). Tim3 and TLR4 were amplified at 94°C for 90 sec (1 cycle); 94°C for 30 sec, 60°C for 30 sec and 72°C for 1 min (30 cycles); 72°C for 5 min (1 cycle). The PCR products were visualized following electrophoresis on 2% agarose gels. Densitometric analysis of the bands was carried out using a ChemiDoc MP System with Image LabTM software version 5.1 (Bio-Rad Laboratories, Inc., Hercules, CA, USA). Results are expressed as the ratio of the intensity of the band for Tim-3, TLR4, and MyD88 to the intensity of the band for β -actin.

Western blot analysis. Total protein was extracted from cells using cell lysis buffer (cat no. P0013; Beyotime Biotechnology, Shanghai, China). Protein concentrations were measured using a bicinchoninic acid (BCA) protein assay (Generay Biotech Co., Ltd., Shanghai, China), according to the manufacturer's protocols. Equal amounts of proteins (10 μ g) of cell lysates were separated on a 10% SDS-PAGE and electrotransferred onto a nitrocellulose membrane (EMD Millipore, Billerica, MA, USA). The membranes were blocked in Tris buffered saline Tween containing 5% fat-free dry milk and incubated overnight at 4°C with primary antibodies, including rabbit anti-Tim-3 antibody (M-171; sc-292390; 1:200; Santa Cruz Biotechnology, Inc., Dallas, TX, USA), mouse anti-TLR4 antibody (ab22048; 1:1,000; Abcam, Cambridge, UK), rabbit anti-MyD88 antibody (ab2068; 1:1,000; Abcam), rabbit anti-phosphorylated (p-) nuclear factor κB (NF- κB) p65 antibody (3033; 1:1,000; Cell Signaling Technology, Inc., Danvers, MA, USA) and mouse anti- β -actin antibody (ab1801; 1:1,000; Abcam). Goat anti-rabbit horseradish peroxidase (HRP)-conjugated immunoglobulin G (IgG; ZDR-5306; 1:2,000; ZSGB-BIO, Beijing, China) or rabbit anti-mouse HRP-IgG (ZDR-5109; 1:2,000; ZSJB-BIO) were used as secondary antibodies. Proteins were detected using a ChemiDoc MP System with Image Lab[™] Software version 5.1 (Bio-Rad Laboratories, Inc.). Data were normalized to β -actin levels.

ELISA. The concentrations of interleukin 6 (IL-6), tumor necrosis factor- α (TNF- α), interferon- γ (IFN- γ) and interleukin

10 (IL-10) in the cell culture supernatants of RAW264.7 cells were determined by ELISA, using a commercial human multiplex kit (IL-6, TNF- α , IFN- γ and IL-10; 88-5083; Aushon Biosystems, Inc., Wuxi, China) according to the manufacturer's protocol.

Statistical analysis. Data are expressed as the means \pm standard deviation of three independent experiments. Data were analyzed by paired or unpaired t-tests when comparing differences between paired samples or two independent groups respectively. One-way analysis of variance with the LSD post hoc test was used for multiple comparisons. Data analysis was performed using SPSS 17.0 (SPSS, Inc., Chicago, IL, USA). P<0.05 was considered to indicate a statistically significant difference.

Results

Effect of H. pylori infection on Tim-3, TLR4 and MyD88 mRNA expression in macrophages. H. pylori infection of macrophages was performed by co-culturing H. pylori with RAW264.7 cells for 12 h at various bacteria/cell ratios, i.e. multiplicities of infection (MOI). H. pylori infection significantly elevated Tim-3, TLR4 and MyD88 mRNA expression levels in RAW264.7 macrophages compared with control uninfected macrophages (P<0.01 compared with control; Fig. 1). Tim-3, TLR4 and MyD88 mRNA levels increased in a MOI-dependent manner (MOI range tested 25-200; Fig. 1). Tim-3 mRNA expression was significantly higher in the infected cells compared with the control cells in all MOI tested (P<0.01; Fig. 1), and no significant difference was observed between different MOI effects (P>0.05; Fig. 1). TLR4 mRNA expression was significantly higher at MOI ratios 50-200 compared with control uninfected cells (P<0.01; Fig. 1). Lastly, MyD88 mRNA expression was significantly higher at MOI ratios 100 and 200 compared with control uninfected cells (P<0.01; Fig. 1).

Construction of Tim-3-overexpressing macrophages. RAW264.7 cells were either left untransfected (control) or were transfected with the pLVX-IRES-ZsGreen1-Tim-3 expression plasmid, in order to establish Tim-3 overexpression in macrophages. Tim-3 mRNA (Fig. 2A) and protein (Fig. 2B) expression levels were significantly higher in transfected RAW264.7 cells than in the control cells (P<0.05; Fig. 2), suggesting that RAW264.7 cells were successfully transfected.

Effect of H. pylori infection on Tim-3 expression and TLR4 pathway proteins in Tim-3-overexpressing macrophages. RAW264.7 cells were either left untransfected (control), or were transfected with empty plasmid (mock) or Tim-3 overexpression plasmid (Tim-3), then cultured for an additional 12 h either alone or with H. pylori at an MOI of 100. In the control macrophages, Tim-3, TLR4, MyD88 mRNA and protein levels, and p-NF-kBp65 protein levels, were significantly higher in the H. pylori-infected cells compared with the H. pylori-negative cells (P<0.05; Fig. 3). In the Tim-3 overexpressing macrophages, TLR4 and MyD88 mRNA expression was increased following infection (P<0.01; Fig. 3A), but



Figure 1. Effect of *Helicobacter pylori* infection on Tim-3, TLR4 and MyD88 mRNA expression. RAW264.7 macrophages were cocultured with *H. pylori* at 0, 25, 50, 100, or 200 MOI for 12 h, following which mRNA expression for Tim-3, TLR4 and MyD88 was analyzed in the RAW264.7 cells by reverse transcription-quantitative polymerase chain reaction. n=4 independent experiments performed in duplicate. *P<0.05 and **P<0.01 vs. 0 MOI. Tim-3, T-cell immunoglobulin and mucin-domain-containing molecule-3; TLR4, Toll-like receptor 4; MyD88, myeloid differentiation factor 88; MOI, multiplicity of infection; NS, not significant.

no significant change was observed in TLR4, MyD88 and p-NF- κ Bp65 protein levels between the *H. pylori*-infected and uninfected cells (P>0.05; Fig. 3B). *H. pylori* infection increased Tim-3 mRNA and protein expression in the Tim-3-overexpressing cells compared with the untransfected cells (P<0.05, Fig. 3). In addition, infected by *H. pylori*, TLR4 and MyD88 mRNA expression levels were significantly lower in the Tim-3-overexpressing cells compared with the untransfected cells (P<0.01; Fig. 3A), but no significant change was observed in TLR4, MyD88 and p-NF- κ B p65 protein levels (P>0.05; Fig. 3B).

Effect of H. pylori infection on inflammatory cytokine secretion in Tim-3-overexpressing macrophages. RAW264.7 cells were either left untransfected (control), or were transfected with empty plasmid (mock) or Tim-3 overexpression plasmid (Tim-3), then either cultured alone or infected with H. pylori at MOI 100 for 12 h. The culture supernatants were subsequently assayed by ELISA for levels of the inflammatory cytokines TNF- α , IL-6, IFN- γ and IL-10. H. pylori infection increased TNF- α , IL-6 and IFN- γ secretion in the control and mock-transfected cells compared with the H. pylori-negative



Figure 2. Tim-3 overexpression in RAW264.7 cells. RAW264.7 cells were either left untransfected (control) or were transfected with the pLVX-IRES-ZsGreen1-Tim-3 overexpression plasmid (Tim-3) for 24 h. (A) Tim-3 mRNA expression levels, measured by reverse transcription-quantitative polymerase chain reaction. (B) Representative images and densitometric quantification from western blot analysis, with quantification relative to β -actin. Results reported from 3 independent experiments performed in duplicate. *P<0.05 and **P<0.01 vs. control. Tim-3, T-cell immunoglobulin and mucin-domain-containing molecule-3.

cells (P<0.01; Fig. 4A-C), but no significant differences were observed in the levels of these cytokines with or without infection in the Tim-3-transfected cells (Fig. 4A-C). By contrast, *H. pylori* infection significantly decreased IL-10 secretion in all cells compared with the *H. pylori*-negative cells, independent of whether they were control, or mock or Tim-3-transfected (P<0.01; Fig. 4D). When comparing the *H. pylori*-infected cells, TNF- α , IL-6, IFN- γ and IL-10 demonstrated significantly lower secretion in the Tim-3-transfected cells compared with the control or mock-transfected cells (P<0.01; Fig. 4).

Discussion

H. pylori infection leads to the accumulation of a large number of inflammatory cells in the gastric mucosa, including macrophages (13,14). Previous studies have demonstrated that TLR4 is a type of receptor that recognizes conserved microbial products in macrophages (15,16). Therefore, TLR4 is a signaling pathway that has important effects on the inflammatory and immune responses to *H. pylori* infections (17,18).

It has been reported that *H. pylori* lipopolysaccharide (LPS) activates NF-KB through the TLR4 signaling pathway, which leads to the release of pro-inflammatory cytokines, including IL-6, interleukin 1 β (IL-1 β) and TNF- α , and to the promotion of inflammatory reactions (19). However, its regulation and the exact mechanism by which it acts remain unclear. To determine the effect of *H. pylori* infection on the TLR4 signaling pathway, the expression of various proteins downstream of TLR4 was examined in macrophages that were infected with different H. pylori concentrations. In addition, the release of inflammatory cytokines from the macrophages was also evaluated. The results revealed that H. pylori infection at MOIs 25-100 significantly increased TLR4 and MyD88 mRNA expression levels. When cells were infected with H. pylori at MOI 100 for 12 h, the protein expression levels of TLR4, MyD88 and p-NF-kBp65 were significantly upregulated. H. pylori infection also significantly increased the secretion of pro-inflammatory cytokines, including IL-6, IFN-y and TNF- α , but reduced the secretion of the anti-inflammatory cytokine IL-10. The present results suggested that H. pylori infection activated the TLR4-mediated and MyD88-dependent signaling pathway, activated NF-KB, and promoted inflammatory reactions. Pathak et al (20) reported that H. pylori 0175 proteins come directly in contact with the extracellular domain of TLR4 and that they activate the MAPK and NF-ĸB signaling pathways, thus leading to IL-6 secretion from macrophages. Maeda et al (21) reported that H. pylori activated the NF- κ B signaling pathway and increased TNF- α release from human monocytic THP-1 cells but failed to activate the NF-KB signaling pathway in macrophages derived from TLR4 mutant mice. Mandell et al (22) reported that H. pylori infection increases the secretion of TNF- α and IL-6 from THP-1 cells in a concentration-dependent manner. However, macrophages lacking TLR4 expression do not respond to H. pylori LPS stimulation. In summary, TLR4 signaling pathway activation in macrophages is important in the abnormal inflammatory reaction that is caused by *H. pylori* infection, but the exact regulatory mechanism remains unclear.

Tim-3 was the first, and is presently the only, surface molecule shown to specifically identify Th1 cells in mice and humans (23). Numerous studies have demonstrated that Tim-3 is also expressed in innate immune cells, including dendritic cells, macrophages and natural killer cells, but the effect of Tim-3 on immune cells is different than on other cells (5,23). Tim-3 has been demonstrated to be important in innate immune cells, including macrophages/monocytes (24). The present study revealed that H. pylori infection upregulated the mRNA and protein expression of Tim-3 in macrophages in a concentration-dependent manner. In a previous study by this group, it was demonstrated that following 12 h of H. pylori stimulation, there was a marked increase in Tim-3 production by mouse spleen lymphocytes (25). Tim-3 expression was also confirmed in vivo to be higher in the gastric mucosa of H. pylori-infected mice than of uninfected mice (25). However, the function of Tim-3 in *H. pylori* infection remains unclear.

TLR4 and Tim-3 are commonly expressed in various immune cells, and their interaction is closely related to the development of many diseases (10,11). However, no report to date has examined the effect of Tim-3 overexpression on the activation of the TLR4 signaling pathway and the



Figure 3. Effect of *Helicobacter pylori* infection on expression of Tim-3 and TLR4 signaling pathway proteins in macrophages. RAW264.7 cells were either left untransfected (control), or were transfected with empty plasmid (Mock) or Tim-3 overexpression plasmid (Tim-3), and then they were cultured for 12 h either alone (*H. pylori*-) or with *H. pylori* at MOI 100 (*H. pylori*+). (A) mRNA expression was measured for Tim-3, TLR4 and MyD88 by reverse transcription-quantitative polymerase chain reaction and gel electrophoresis. (B) Protein expression was measured for Tim-3, TLR4, MyD88 and p-NF κ Bp65 mRNA by western blot analysis. Densitometric quantification and representative images of the blots are shown. Quantification for mRNA and protein was relative to β -actin. n=3 independent experiments performed in duplicate. *P<0.05 and **P<0.01, with comparisons indicated by lines. Tim-3, T-cell immunoglobulin and mucin-domain-containing molecule-3; TLR4, Toll-like receptor 4; MOI, multiplicity of infection; MyD88, myeloid differentiation factor 88; p-NF κ Bp65, phosphorylated nuclear factor κ B p65 subunit.



Figure 4. Effect of *Helicobacter pylori* infection on cytokine secretion in Tim-3-overexpressing macrophages. RAW264.7 cells were either left untransfected (control), or were transfected with empty plasmid (Mock) or Tim-3 overexpression plasmid (Tim-3), then cultured for 12 h either alone (*H. pylori*-) or with *H. pylori* at MOI 100 (*H. pylori*+). Levels of the inflammatory cytokines (A) TNF- α , (B) IFN- γ , (C) IL-6 and (D) IL-10 were detected in the culture supernatants by ELISA. n=3 independent experiments performed in duplicate. *P<0.05 and **P<0.01, with comparisons indicated by lines. Tim-3, T-cell immunoglobulin and mucin-domain-containing molecule-3; MOI, multiplicity of infection; TNF- α , tumor necrosis factor- α ; IFN- γ , interferon- γ ; IL-6, interleukin 6; IL-10, interleukin 10.

H. pylori-induced inflammatory reactions in macrophages. The present study revealed that mRNA expression of TLR4 and MyD88, and secretion of IL-6, IFN- γ and TNF- α , in Tim-3-overexpressing macrophages infected with H. pylori were significantly lower than control macrophages. These results suggest that Tim-3 inhibited the activation of macrophages by negatively regulating the TLR4 signaling pathway. A previous study from this group (26) demonstrated that blocking Tim-3 with an inhibitory antibody upregulated TLR4 and MyD88 expression, promoted NF-κB activation, decreased the number of CD4+CD25+Foxp3+ Treg cells, and upregulated Th1 immune responses, resulting in intensified inflammation of the gastric mucosa of *H. pylori*-infected mice. Frisancho et al (11) reported that, in a mouse model of inflammatory heart disease, blocking Tim-3 significantly increases TLR4 expression and cardiac inflammation, indicating that Tim-3 is a negative regulator of the TLR4 signaling pathway. The present study also demonstrated that following H. pylori infection, Tim-3 overexpression in macrophages inhibited the TLR4 signaling pathway and reduced pro-inflammatory cytokine secretion. However, these results were not statistically significant in the absence of H. pylori infection, indicating that Tim-3 strongly and negatively regulated TLR4 signaling only in the presence of pathology, and not under normal physiological conditions. The mechanism by which Tim-3 negatively regulated the TLR4 signaling pathway remains unclear. Yang et al (9) reported that Tim-3 overexpression in macrophages significantly suppresses TLR-mediated pro-inflammatory cytokine production. Li et al (27) reported that the H. pylori cytotoxin-associated gene A activates the phosphatidylinositol 3-kinase (PI3K)/AKT serine/threonine kinase 1 (Akt1) pathway, which inhibits interleukin 8 release in H. pylori-infected gastric carcinoma AGS cells. However, further studies are needed to determine how H. pylori infection induces Tim-3 overexpression and downregulates TLR4-mediated inflammatory cytokine production through the PI3K-Akt pathway. The present study also demonstrated that following H. pylori infection, Tim-3 overexpression in macrophages not only inhibited the release of IL-6, IFN- γ and TNF- α , but also suppressed the release of IL-10. Zhang et al (24) reported that blocking or silencing Tim-3 in macrophages not only increased the release of IL-12 and IL-6, but also increased the release of IL-10. Therefore, the role of Tim-3 in regulating inflammatory and immune responses may be influenced not only by Tim-3 expression levels but also by the state of the macrophages and the molecular balance between inhibition and stimulation. Anderson et al (28) reported that Tim-3, by virtue of its differential expression on cells of the innate and adaptive immune systems, can both promote inflammation and terminate Th1 immunity. Therefore, Tim-3 serves opposing roles, influenced by multiple factors, including variations in stimuli, cell type and immune activation state. Further research will focus on the mechanism by which Tim-3 downregulates the TLR4 signaling pathway during H. pylori infection.

In conclusion, *H. pylori* infection in RAW264.7 macrophages activated the TLR4 signaling pathway, upregulated Tim-3 expression and increased secretion of pro-inflammatory cytokines (TNF- α , IL-6 and IFN- γ), while decreasing secretion of an anti-inflammatory cytokine (IL-10). Under conditions of *H. pylori* infection, however, Tim-3 overexpression inhibited the TLR4 pathway activation and the secretion of pro-inflammatory cytokines. Taken together, the present study indicated that Tim-3 may represent a novel therapeutic target for treating *H. pylori* infection-related diseases.

Acknowledgements

The present study was supported by the Major Research Program Foundation of China (Prevention and control of major chronic non-communicable diseases, grant no. 2016YFC1302201) and the National Natural Science Foundation of China (grant no. 81260076), and was edited for proper English language, grammar, punctuation, spelling, and overall style by American Journal Experts.

References

- Koch M, Meyer TF and Moss SF: Inflammation, immunity, vaccines for Helicobacter pylori infection. Helicobacter 18 (Suppl 1): S18-S23, 2013.
- Demiray E and Bekmen N: Helicobacter pylori infection and phagocytosis. Mikrobiyol Bul 42: 177-184, 2008 (In Turkish).
- 3. Arango Duque G and Descoteaux A: Macrophage cytokines: Involvement in immunity and infectious diseases. Front Immunol 5: 491, 2014.
- Rad R, Ballhorn W, Voland P, Eisenächer K, Mages J, Rad L, Ferstl R, Lang R, Wagner H, Schmid RM, *et al*: Extracellular and intracellular pattern recognition receptors cooperate in the recognition of Helicobacter pylori. Gastroenterology 136: 2247-2257, 2009.
- 5. Freeman GJ, Casasnovas JM, Umetsu DT and DeKruyff RH: TIM genes: A family of cell surface phosphatidylserine receptors that regulate innate and adaptive immunity. Immunol Rev 235: 172-189, 2010.
- Mariat C, Sánchez-Fueyo A, Alexopoulos SP, Kenny J, Strom TB and Zheng XX: Regulation of T cell dependent immune responses by TIM family members. Philos Trans R Soc Lond B Biol Sci 360: 1681-1685, 2005.
- Monney L, Sabatos CA, Gaglia JL, Ryu A, Waldner H, Chernova T, Manning S, Greenfield EA, Coyle AJ, Sobel RA, *et al*: Th1-specific cell surface protein Tim-3 regulates macrophage activation and severity of an autoimmune disease. Nature 415: 536-541, 2002.
- Khademi M, Illés Z, Gielen AW, Marta M, Takazawa N, Baecher-Allan C, Brundin L, Hannerz J, Martin C, Harris RA, *et al*: T Cell Ig- and mucin-domain-containing molecule-3 (TIM-3) and TIM-1 molecules are differentially expressed on human Th1 and Th2 cells and in cerebrospinal fluid-derived mononuclear cells in multiple sclerosis. J Immunol 172: 7169-7176, 2004.
- 9. Yang X, Jiang X, Chen G, Xiao Y, Geng S, Kang C, Zhou T, Li Y, Guo X, Xiao H, *et al*: T cell Ig mucin-3 promotes homeostasis of sepsis by negatively regulating the TLR response. J Immunol 190: 2068-2079, 2013.
- Uchida Y, Ke B, Freitas MC, Yagita H, Akiba H, Busuttil RW, Najafian N and Kupiec-Weglinski JW: T-cell immunoglobulin mucin-3 determines severity of liver ischemia/reperfusion injury in mice in a TLR4-dependent manner. Gastroenterology 139: 2195-2206, 2010.
- 11. Frisancho-Kiss S, Davis SE, Nyland JF, Frisancho JA, Cihakova D, Barrett MA, Rose NR and Fairweather D: Cutting edge: Cross-regulation by TLR4 and T cell Ig mucin-3 determines sex differences in inflammatory heart disease. J Immunol 178: 6710-6714, 2007.

- 12. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. Methods 25: 402-408, 2001.
- 13. Schumacher MA, Donnelly JM, Engevik AC, Xiao C, Yang L, Kenny S, Varro A, Hollande F, Samuelson LC and Zavros Y: Gastric sonic hedgehog acts as a macrophage chemoattractant during the immune response to Helicobacter pylori. Gastroenterology 142: 1150-1159.e6, 2012.
- Deen NS, Gong L, Naderer T, Devenish RJ and Kwok T: Analysis of the relative contribution of phagocytosis, LC3-associated phagocytosis, and canonical autophagy during helicobacter pylori infection of macrophages. Helicobacter 20: 449-459, 2015.
 Zhang Y, Lu Y, Ma L, Cao X, Xiao J, Chen J, Jiao S, Gao Y, Liu C,
- Zhang Y, Lu Y, Ma L, Cao X, Xiao J, Chen J, Jiao S, Gao Y, Liu C, Duan Z, *et al*: Activation of vascular endothelial growth factor receptor-3 in macrophages restrains TLR4-NF-κB signaling and protects against endotoxin shock. Immunity 40: 501-514, 2014.
- 16. Byun EB, Sung NY, Byun EH, Song DŠ, Kim JK, Park JH, Song BS, Park SH, Lee JW, Byun MW and Kim JH: The procyanidin trimer C1 inhibits LPS-induced MAPK and NF-κB signaling through TLR4 in macrophages. Int Immunopharmacol 15: 450-456, 2013.
- 17. Käbisch R, Mejias-Luque R, Gerhard M and Prinz C: Involvement of toll-like receptors on Helicobacter pylori-induced immunity. PLoS One 9: e104804, 2014.
- Su YL, Yang JC, Lee H, Sheu F, Hsu CH, Lin SL and Chow LP: The C-terminal disulfide bonds of Helicobacter pylori GroES are critical for IL-8 secretion via the TLR4-dependent pathway in gastric epithelial cells. J Immunol 194: 3997-4007, 2015.
- Fujihara M, Muroi M, Tanamoto K, Suzuki T, Azuma H and Ikeda H: Molecular mechanisms of macrophage activation and deactivation by lipopolysaccharide: Roles of the receptor complex. Pharmacol Ther 100: 171-194, 2003.
- 20. Pathak SK, Basu S, Bhattacharyya A, Pathak S, Banerjee A, Basu J and Kundu M: TLR4-dependent NF-kappaB activation and mitogen- and stress-activated protein kinase 1-triggered phosphorylation events are central to Helicobacter pylori peptidyl prolyl cis-, trans-isomerase (HP0175)-mediated induction of IL-6 release from macrophages. J Immunol 177: 7950-7958, 2006.
- Maeda S, Akanuma M, Mitsuno Y, Hirata Y, Ogura K, Yoshida H, Shiratori Y and Omata M: Distinct mechanism of Helicobacter pylori-mediated NF-kappa B activation between gastric cancer cells and monocytic cells. J Biol Chem 276: 44856-44864, 2001.
- 22. Mandell L, Moran AP, Cocchiarella A, Houghton J, Taylor N, Fox JG, Wang TC and Kurt-Jones EA: Intact gram-negative Helicobacter pylori, Helicobacter felis, and Helicobacter hepaticus bacteria activate innate immunity via toll-like receptor 2 but not toll-like receptor 4. Infect Immun 72: 6446-6454, 2004.
- Anderson DE: TIM-3 as a therapeutic target in human inflammatory diseases. Expert Opin Ther Targets 11: 1005-1009, 2007.
 Zhang Y, Ma CJ, Wang JM, Ji XJ, Wu XY, Moorman JP and
- 24. Zhang Y, Ma CJ, Wang JM, Ji XJ, Wu XY, Moorman JP and Yao ZQ: Tim-3 regulates pro- and anti-inflammatory cytokine expression in human CD14+ monocytes. J Leukoc Biol 91: 189-196, 2012.
- Hu S, Xie Y, Zhou N, Jin L, Tan Y, Liu D, Gong Y, Liu L, Liu J, Liu W, *et al*: Expression of T-cell immunoglobulin- and mucin-domain-containing molecules-1 and -3 (Tim-1 and Tim-3) in Helicobacter pylori infection. Helicobacter 16: 373-381, 2011.
 Gong Y, Tao L, Wang F, Liu W, Jing L, Liu D, Hu S, Xie Y and
- Gong Y, Tao L, Wang F, Liu W, Jing L, Liu D, Hu S, Xie Y and Zhou N: Chitosan as an adjuvant for a Helicobacter pylori therapeutic vaccine. Mol Med Rep 12: 4123-4132, 2015. v
 Li SP, Chen XJ, Sun AH, Zhao JF and Yan J: CagA(+)
- Li SP, Chen XJ, Sun AH, Zhao JF and Yan J: CagA(+) H. pylori induces Akt1 phosphorylation and inhibits transcription of p21(WAF1/CIP1) and p27(KIP1) via PI3K/Akt1 pathway. Biomed Environ Sci 23: 273-278, 2010.
- 28. Anderson AC, Anderson DE, Bregoli L, Hastings WD, Kassam N, Lei C, Chandwaskar R, Karman J, Su EW, Hirashima M, et al: Promotion of tissue inflammation by the immune receptor Tim-3 expressed on innate immune cells. Science 318: 1141-1143, 2007.