

The transfection of A20 gene prevents kidney from ischemia reperfusion injury in rats

LIXIN WEI¹, XIANGHUI ZHANG², QIUPING YE¹, YUEER YANG¹ and XIAOWEN CHEN¹

¹Department of Nephropathy, Union Hospital, Fujian Medical University, Fuzhou, Fujian 350001; ²Department of Nephropathy, The Fifth Affiliated Hospital of Zunyi Medical University, Zhuhai, Guangdong 519000, P.R. China

Received April 22, 2016; Accepted March 24, 2017

DOI: 10.3892/mmr.2017.6725

Abstract. Ischemia/reperfusion may induce inflammation and cell death through the nuclear factor (NF)- κ B signaling pathway. As a negative regulator of NF- κ B, zinc finger A20 exhibits anti-apoptotic and anti-inflammatory effects *in vitro*. The present study was designed to upregulate A20 expression using an A20 transfection approach to investigate the *in vivo* protective effects of the A20 gene on renal ischemia reperfusion injury. The A20 gene was cloned into a pcDNA3.1 vector to construct the expression plasmid pcDNA3.1-A20. The plasmid was wrapped with a liposome and injected intravenously into rats 48 h prior to establishing the models of renal ischemia reperfusion injury. Saline and the empty plasmid pcDNA3.1 were used as controls. Following 24 h post-operation, A20 expression was determined using reverse transcription-quantitative polymerase chain reaction and western blotting. The renal function and structure were assessed by analyzing the concentrations of serum creatinine (Scr), blood urea nitrogen (BUN) and histological features. Renal tissues were additionally examined for renal tubular cell apoptosis and NF- κ B activity. The results demonstrated that *in vivo* transfection of pcDNA3.1-A20 induced renal A20 expression in rats. A20 overexpression *in vivo* significantly reduced renal injury as demonstrated by the improved levels of Scr and BUN and the reduction in histological damage. These improvements were accompanied by a suppression of renal proximal tubular epithelial cell apoptosis and an inhibition of NF- κ B activity. These results indicated that transfection of the A20 gene upregulates the expression of A20 *in vivo* and protects the kidneys from ischemia reperfusion injury via inhibition of the NF- κ B signal transduction pathway.

Introduction

Renal ischemia/reperfusion injury (IRI) is a primary cause of acute renal failure resulting from tubular dysfunction following shock, sepsis or renal transplantation (1). Despite various advances in vascular surgery, the rates of morbidity and mortality in patients with post-operative IRI remain particularly high (2). As with other pathological conditions of the kidney, renal ischemia may ultimately progress to chronic advanced kidney disease, which is characterized by tubule and capillary loss as well as regional interstitial fibrosis associated with chronic hypoxic stress (3-6).

In the ischemic kidney, and during the subsequent re-oxygenation, excessive reactive oxygen species (ROS) are generated in the reperfusion phase, initiating a cascade of deleterious cellular responses leading to inflammation, cell death and acute kidney failure (7). During the pathophysiological process, initiated in response to a broad spectrum of pro-inflammatory mediators including tumor necrosis factor (TNF)- α , interleukin-1 and ROS (8,9), nuclear factor (NF)- κ B is activated, mediating a vicious cycle of further inflammatory responses, which contribute to cell apoptosis. Therefore, targeting the NF- κ B signaling pathway is a promising novel therapeutic strategy to inhibit the inflammatory and innate immune cascades responsible for the amplification of injury, which may produce effective treatments or prevention strategies for IRI. This concept was supported by a previous study that performed treatments with *in vivo* transfection of NF- κ B decoy oligodeoxynucleotides, which significantly reduced the renal dysfunction and damage associated with ischemic acute renal failure (10).

Zinc finger protein A20 has been recognized as a central regulator of inflammation and apoptosis due to its ubiquitous inhibitory effect on NF- κ B (8,11,12). As an NF- κ B dependent gene, A20 forms a negative feedback loop, limiting NF- κ B activation and inflammation in cells exposed to hypoxia and re-oxygenation (8,13). Lee *et al* (14) reported that mice with A20 deficiency develop severe inflammation and cachexia as a result of failure to terminate the TNF-induced responses to NF- κ B and cell death. Overexpression of A20 in rat kidneys using recombinant adenovirus-mediated gene transfer, protects rat kidneys from acute tubular necrosis following renal ischemia (13).

Therefore, the authors hypothesized that A20 may prevent the kidneys from IRI via a combination of anti-apoptotic and

Correspondence to: Dr Lixin Wei, Department of Nephropathy, Union Hospital, Fujian Medical University, 29 Xinquan Road, Gulou, Fuzhou, Fujian 350001, P.R. China
E-mail: lixinwei66@126.com

Key words: kidney, ischemia reperfusion injury, zinc finger protein A20, cell apoptosis, nuclear factor- κ B signaling pathway

anti-inflammatory effects. The aim of the present study was to establish the A20 transfection model in rats and to evaluate the protective effect of *in vivo* A20 transfection on renal IRI by observing renal function, histopathological alterations, cell apoptosis and the NF- κ B signaling pathway.

Materials and methods

Construction of recombinant human A20 expression plasmid. Human A20 cDNA was cloned into a pcDNA3.1 vector (Invitrogen; Thermo Fisher Scientific, Inc., Waltham, MA, USA) to construct the expression plasmid, which was termed pcDNA3.1-A20. The preparation of A20 cDNA and construction of pcDNA3.1-A20 were performed by the Invitrogen Cn Service (Project No: KL120515006; Shanghai, China). The plasmid DNA was transformed into *E coli* DH5 α competent cells (Invitrogen; Thermo Fisher Scientific, Inc.) and grown in Luria Broth medium for propagation, according to the manufacturer's protocol. The plasmid DNA was subsequently extracted using the PureLink[®] HiPure Plasmid DNA Purification kit (Invitrogen; Thermo Fisher Scientific, Inc.) according to the manufacturer's protocol. The inserted sequences were digested with the restriction endonucleases *Nhe*I and *Xba*I (Promega Corporation, Madison, WI, USA) according to manufacturer's protocol, and verified by DNA sequencing, performed by the Invitrogen Cn Service, and separation by 1% gel electrophoresis.

Experimental animals. The animal experiments were approved by the Animal Ethics Committee of the Fujian University of Traditional Chinese Medicine (permit no. SCXK Shanghai 2012-0002; Fuzhou, China). Sprague-Dawley male rats (8-10 weeks in age and, 250-300 g in weight) were provided by the Laboratory Animal Center of Fujian University of Traditional Chinese Medicine. The rats received a standard diet, and were housed individually with free access to tap water. A 12 h light-dark cycle was provided. Room temperature was maintained at 21-23°C with 30-60% humidity. Rats (n=24) were randomly assigned into three groups: Saline (control), plasmid control and A20 groups (n=8/group). The three groups were given intravenous injections via the tail vein with saline (250 μ l), empty pcDNA3.1 plasmid (15 μ l of Lipofectamine 2000 (Invitrogen; Thermo Fisher Scientific, Inc.) + 10 μ g of pcDNA3.1 mixed with saline to a final volume of 250 μ l) or pcDNA3.1-A20 (15 μ l of Lipofectamine 2000 + 10 μ g of pcDNA3.1-A20 mixed with saline to a final volume of 250 μ l), respectively. Following 48 h post-injection, rats were anesthetized with sodium pentobarbital (50 mg/kg; Pitman-Moore Inc., Washington Crossing, NJ, USA) intraperitoneally. A ventral midline longitudinal incision (8-10 cm) was used to fully expose the left kidney, renal artery, abdominal aorta and vena cava. The renal ischemia reperfusion injury model was established by clamping the left renal artery and vein for 45 min, then cutting the right kidney prior to reperfusion of the left renal vasculature. At 24 h post-operation, at the time of euthanasia, 2 ml blood samples were drawn from the mesenteric vein and centrifuged at 1,500 \times g for 10 min to obtain sera. For paraffin histologic assessments and cell apoptosis analysis, one part of the left kidney was fixed in 10% buffered formalin for 24 h, the remaining parts of the kidneys were

snap-frozen in liquid nitrogen for western blot, gene expression and electrophoretic mobility shift assay analyses.

Western blot analysis. Cytoplasmic and nuclear proteins were extracted using the Nucleoprotein Cytoplasm Protein Extraction kit (Nanjing KeyGen Biotech Co., Ltd., Nanjing, China) according to manufacturer's protocol. The protein concentration was measured using a Bicinchoninic Protein Assay (Beyotime Institute of Biotechnology, Shanghai, China) according to manufacturer's protocol. Proteins (50 μ g) were separated by 10% SDS-PAGE and transferred to nitrocellulose membranes. Following blocking in phosphate-buffered saline (PBS) containing 5% skimmed milk overnight at 4°C, the membranes were incubated with primary antibodies against A20 (dilution, 1:1,000; rabbit; 5630; Cell Signaling Technology, Inc., Danvers, MA, USA) and β -actin (dilution, 1:2,000; rabbit; 4970; Cell Signaling Technology, Inc.) for 2 h at room temperature. Bound antibodies were detected using an anti-rabbit horseradish peroxidase-conjugated secondary antibody (dilution, 1:5,000; goat; HS101-01; Beijing TransGen Biotech Co., Ltd.) for 1 h at room temperature. A chemiluminescent substrate (Thermo Fisher Scientific, Inc.) was used to detect the immunoreactive protein signals. Computerized image analysis was performed using Image Lab[™] version 5.1 and a Molecular Imager[®] ChemiDoc[™] XRS System (both from Bio-Rad Laboratories, Inc., Hercules, CA, USA). The values for the A20 protein were quantified and normalized to β -actin abundance. All samples were performed in duplicate.

RNA isolation and reverse transcription-quantitative polymerase chain reaction (RT-qPCR). Total RNA was isolated from rat kidneys using TRIzol[®] (Invitrogen; Thermo Fisher Scientific, Inc.) and 5 μ g RNA was reverse-transcribed using the PrimeScript[™] II First strand cDNA Synthesis kit (Takara Bio, Inc., Otsu, Japan) according to the manufacturer's protocol. The following primers for A20 were designed to target the rat-specific mRNA region of the A20 gene: Forward, 5'-CGG CGTAGATTCTACGTCTTCA-3' and reverse, 5'-CTGTCT CGGATACTGCTTTGTCCA-3'. RT-qPCR was performed with the SYBR[®] Premix Ex Taq[™] (Takara Bio, Inc.) following the manufacturer's protocol. Transcript levels were quantified by qPCR using the following cycling conditions: 3 sec at 94°C, then 40 cycles of 3 sec at 95°C and 20 sec at 60°C. The expression levels of A20 were normalized to the housekeeping gene (β -actin: Forward, 5'-ACTTCGAGCAAGAGATGGC-3' and reverse, 5'-ACGTCACACTTCATGATGGA-3') and the initial amount of RNA used was calculated using the $2^{-\Delta\Delta C_q}$ method as described previously (15). All samples were performed in duplicate.

Renal function assay. Renal function was detected by measuring serum creatinine (Scr) and blood urea nitrogen (BUN) using Raichem[™] Enzymatic Creatinine Reagent and BUN Slow Rate Reagent (Cliniqa Corporation, San Marcos, CA, USA) according to the manufacturer's protocol, and measured on a Roche Automatic Biochemical Analyzer (Roche Diagnostics, Basel, Switzerland).

Histological analysis. One portion of the renal tissue was fixed in 10% buffered formalin for 24 h. Following fixation

and dehydration, sections (4 μ m) were cut and stained with haematoxylin and eosin for 1 min respectively at room temperature, followed by examination with a light microscope. Renal injury scores were determined by evaluating the degree of tubulointerstitial injury according to tubular necrosis, tubular dilatation and/or atrophy, inflammatory cell infiltration or cellular edema, as previously described (16): 0, normal kidney; 1, minimal necrosis (<5% involvement of the cortex or outer medulla); 2, mild necrosis (5-25% involvement of the cortex or outer medulla); 3, moderate necrosis (25-75% involvement of the cortex or outer medulla); 4, severe necrosis (>75% involvement of the cortex or outer medulla). Histologic examination was performed in a blinded manner by a renal pathologist using a light microscope (Zeiss AG, Oberkochen, Germany).

Evaluation of cell apoptosis. Cell apoptosis in the renal tissue sections (5- μ m), was quantified using a DeadEnd™ Colorimetric TUNEL System (Promega Corporation), according to the manufacturer's protocol. For all staining procedures, the terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL)-positive cells were counted in 8 random high-power fields using a light microscope by two individual evaluators. Disagreements in results were resolved by a third evaluator.

Electrophoretic mobility shift assay (EMSA). The DNA binding activity of NF- κ B was assessed on nuclear extracts of kidney tissue, extracted as aforementioned, using a Chemiluminescent EMSA kit (Beyotime Institute of Biotechnology) according to the manufacturer's protocol. Radio-labeled NF- κ B consensus oligonucleotide was incubated with 5 μ g nuclear extract in 10 μ l binding buffer at room temperature for 20 min. The negative control reaction, cold probes competitive reaction and the mutant cold probe competitive reaction were also incubated with the same amount of samples as the controls and using the exact same experimental conditions. Band shifts were resolved on a non-denaturing 6.5% polyacrylamide gel. Radiographic signals were quantified by densitometry using the Molecular imager ChemiDoc™ XRS+ system with Image Lab™ software (Version 5.1, Bio-Rad Laboratories, Inc.).

Statistical analysis. The data were analyzed using SPSS 18.0 (SPSS, Inc., Chicago, IL, USA). Differences among multiple groups (>2) were assessed using one-way analysis of variance, followed by the post hoc Tukey-Kramer test. The data differences between two groups were analyzed by an unpaired Student's t-test. $P < 0.05$ was considered to indicate a statistically significant difference. Data are presented as mean \pm standard deviation or as the median (range).

Results

A20 expression in renal tissue. The recombinant pcDNA3.1-A20 plasmid was constructed successfully as demonstrated by DNA sequence and restriction enzyme analyses (Fig. 1). Subsequently the plasmid DNA was wrapped by a liposome and transfected into the rats. As expected, transfection with the pcDNA3.1-A20 significantly increased




Figure 1. Plasmid DNA electrophoresis patterns. The plasmid DNA was verified by 1% agarose gel electrophoresis. M, DNA Ladder (1kb); lane 1 pcDNA3.1-A20 digested by *NheI* and *XbaI* restriction endonucleases; lane 2, pcDNA3.1-A20; lane 3, pcDNA3.1 digested by *NheI* and *XbaI* restriction endonucleases; lane 4, plasmid pcDNA3.1.




Figure 2. A20 protein expression in renal tissue. (A) Western blotting illustrated the expression of the A20 protein in cytosolic fractions using representative samples from each group. (B) A20 protein expression in the saline control, empty plasmid transfected and A20 transfected groups (n=7/group). Data are presented as the mean \pm standard deviation. ^{*} $P < 0.05$ vs. saline control; [#] $P < 0.05$ vs. plasmid control.

the cytoplasmic A20 protein expression *in vivo* (Fig. 2). When compared with the saline group, the A20 protein level increased 7-fold following 72 h of A20 transfection ($P < 0.05$). Although the concentration of A20 appeared to be elevated in the animals transfected with the plasmid control, when compared to the saline control the difference was not statistically significant ($P > 0.05$). The A20 mRNA level was further quantified in the three experimental groups, however, no significant differences were identified in mRNA expression of A20 between groups ($P > 0.05$; Fig. 3).




Figure 3. A20 gene expression in renal tissue. The A20 gene expression in the saline control (n=8), empty plasmid transfected (n=8) and A20 transfected groups (n=9). The box and whisker plots depict the median and the 10 and 90th centiles.




Figure 4. Serum Scr and BUN. The levels of (A) Scr and (B) BUN in the three groups. Data are presented as the mean \pm standard deviation (n=8/group). ***P<0.001 vs. saline control; ###P<0.001 vs. plasmid control. Scr, serum creatinine; BUN, blood urea nitrogen.

Renal injury. The present study determined the levels of the renal injury indicators, Scr and BUN following IRI. When compared with the animals pretreated with pcDNA3.1 or saline, overexpression of A20 induced an ~3-fold decrease in the levels of Scr and BUN (P<0.001; Fig. 4). The extent of renal tissue damage following IRI was evaluated using the histological approach. The saline and plasmid control groups exhibited a similar degree of pathological damage (Fig. 5), including, renal tubular dilatation, cracks in cells and cells flattening, injured tubular epithelial cells or loss of brush border, tubular epithelial cell shedding or necrosis and protein casts or cell debris observed in the tube cavity, and tubular epithelial cell shedding was observed to be the most serious feature. Conversely, in the A20 group, damage to the tubular epithelial cells brush border was the fundamental alteration. However, changes in renal glomerulus structures, including the endothelial cell lining and basement membrane, were not significantly different amongst the three groups. When analyzing the tubular damage scores, the A20 group was

significantly lower compared with the two control groups (P<0.05; Fig. 5).

Renal tubular epithelial cell apoptosis. The number of apoptotic cells in the kidney tissues of experimental rats was determined using a TUNEL assay. Apoptotic cells were primarily present in the renal tubules, particularly in the distal tubule, while few were identified in the renal interstitium and glomerular. The number of TUNEL-positive cells was significantly lower in the A20 group (56.25 \pm 26.47) when compared with the saline group (109.25 \pm 49.91) at 24 h of IRI (P<0.05; Fig. 6). In addition, the total number of apoptotic cells accumulated in the renal tubular epithelium was similar between the groups administrated saline or empty plasmid control.

Analysis of NF- κ B DNA-binding activity. A significant reduction in nuclear DNA binding activity of NF- κ B was observed in the experimental rats transfected with pcDNA3.1-A20. This




Figure 5. Histological characteristics in renal tissue. (A) Hematoxylin-eosin staining revealed the pathological alterations in the representative samples from each group. (B) The tubular damage scores in the three groups. Magnification, $\times 200$. Data are presented as the mean \pm standard deviation ($n=8$ /group). Arrows indicate injured renal tubular epithelial cells. ^{*} $P<0.05$ vs. saline control; [#] $P<0.05$ vs. plasmid control.




Figure 6. Detection of apoptotic cells in renal tissue. (A) Microscopic examination revealed the number of apoptotic cells present in renal tissues from the representative samples of each group. (B) The number of TUNEL-positive apoptotic cells in the three groups. Magnification, $\times 200$. Data are presented as the mean \pm standard deviation ($n=8$ /group). ^{*} $P<0.05$ vs. saline control.

activity was reduced in the A20 group by 53% when compared with the plasmid control group ($P=0.001$; Fig. 7).

Discussion

In the present study, a liposomal pcDNA3.1-A20 plasmid was successfully transfected into rats and a transient expression of A20 was produced in kidney tissues. Overexpression of




Figure 7. Nuclear NF- κ B activity in renal tissue. (A) The band shift assay determined the activity of the signaling molecule NF- κ B in nuclear fractions using representative samples from each group. (B) NF- κ B activity in the plasmid control and A20 groups. Data are presented as the mean \pm standard deviation, expressed as the fold-change relative to the saline control group ($n=8$ /group). ^{***} $P<0.001$ vs. plasmid control group. NF- κ B, nuclear factor- κ B.

A20 attenuated renal injury, improved histological features, decreased cell apoptosis and inhibited NF- κ B activation in

this rat model of renal IRI. Collectively, these results suggest that transfection of the A20 gene may be a simple and highly efficient method of protecting the kidneys from renal IRI.

To evaluate the therapeutic potential of the A20 gene, a liposome-encapsulated plasmid DNA was used for gene delivery as this method is a reliable and simple technique for gene transfection (17,18). Notably, non-viral transfection methods have the advantage of safe application with minimal risk of replication or viral incorporation and a lower chance of immunological rejection. In addition, plasmid DNA can be produced stably and at a low cost, to a high level of purity. As demonstrated by the western blot analysis, *in vivo* delivery of the liposome pcDNA3.1-A20 complex promoted A20 protein expression in kidney tissues 72 h following injection, suggesting that this method of delivery may be an efficient way of upregulating A20 expression *in vivo*. In contrast to A20 protein expression, the A20 mRNA expression levels did not demonstrate a significant change 72 h following transfection. This apparent discrepancy may be due to transient A20 gene expression induced by pcDNA3.1-A20 and differences in mRNA and protein turnover rates.

The present study determined whether A20 overexpression protected rat kidneys from IRI by conducting renal function tests and observing histological alterations. Increased blood levels of Scr and BUN are indicators of renal injury. The results indicated that A20 transfection markedly decreased the Scr and BUN levels following IRI. In response to IRI, the pathological damage was primarily evident in tubular epithelial cells. This damage was localized to the epithelial cells brush border following A20 treatment, and the tubular injury scores were attenuated. These results provided evidence of the beneficial effect of A20 transfection on renal IRI.

The functional and morphological improvement in the A20 group is likely to be attributable to the inhibitory effect of A20 on NF- κ B activation and the resulting suppression of cell apoptosis. Activation of NF- κ B occurs rapidly in response to ischemia/reperfusion in a number of tissues including kidney cells, brain, liver and myocardium (9,19-22). A20, as a key negative regulator of NF- κ B, exhibits anti-inflammatory and anti-apoptotic effects in cultured cells via inhibition of NF- κ B (11,12,23). In the rat model of the present study, the number of apoptotic cells was significantly reduced in animals pretreated with pcDNA3.1-A20. In addition, a large proportion of TUNEL-positive cells were predominantly observed in the distal tubular epithelium in response to renal IRI, coinciding with the morphological features observed. The nuclear NF- κ B binding activity was examined to analyze NF- κ B signaling. Under normal conditions, NF- κ B is retained in the cytoplasm by binding to its specific inhibitor protein, inhibitor of NF- κ B (I κ B). Upon stimulation, I κ B is phosphorylated, degraded and is subsequently separated from NF- κ B. The activated NF- κ B is then translocated into the nucleus to activate transcriptional expression of downstream genes associated with inflammatory responses and apoptosis (21,24). The results demonstrated that A20 transfection resulted in a 53% reduction in NF- κ B DNA-binding activity, further confirming the efficacy of this *in vivo* transfection model and explaining the protective mechanisms of A20 in renal injury.

In conclusion, the results demonstrated that the liposome-encapsulated plasmid DNA vector for gene delivery

upregulated the expression of A20 *in vivo*, conferring protection against IRI which was mediated by NF- κ B. These results offer a potential prophylactic treatment strategy for renal injury and support the use of A20 as a potential therapeutic target in renal IRI. However, further studies are required to assess whether there are side effects in other tissues, including the liver and heart, due to the systemic delivery of A20.

Acknowledgements

The present study was supported by the Natural Science Foundation of Fujian Province (grant no. 2010J0101).

References

1. Agrawal M and Swartz R: Acute renal failure. *Am Fam Physician* 61: 2077-2088, 2000.
2. Giulini SM, Bonardelli S, Portolani N, Giovanetti M, Galvani G, Maffei R, Coniglio A, Tiberio GA, Nodari F, De Lucia M, *et al*: Suprarenal aortic cross-clamping in elective abdominal aortic aneurysm surgery. *Eur J Vasc Endovasc Surg* 20: 286-289, 2000.
3. Basile DP: Rarefaction of peritubular capillaries following ischemic acute renal failure: A potential factor predisposing to progressive nephropathy. *Curr Opin Nephrol Hypertens* 13: 1-7, 2004.
4. Basile DP, Donohoe D, Roethe K and Osborn JL: Renal ischemic injury results in permanent damage to peritubular capillaries and influences long-term function. *Am J Physiol Renal Physiol* 281: F887-F899, 2001.
5. Bonventre JV and Weinberg JM: Recent advances in the pathophysiology of ischemic acute renal failure. *J Am Soc Nephrol* 14: 2199-2210, 2003.
6. Nangaku M: Chronic hypoxia and tubulointerstitial injury: A final common pathway to end-stage renal failure. *J Am Soc Nephrol* 17: 17-25, 2006.
7. Malek M and Nematbakhsh M: Renal ischemia/reperfusion injury; from pathophysiology to treatment. *J Renal Inj Prev* 4: 20-27, 2015.
8. Beyaert R, Heyninck K and Van Huffel S: A20 and A20-binding proteins as cellular inhibitors of nuclear factor-kappa B-dependent gene expression and apoptosis. *Biochem Pharmacol* 60: 1143-1151, 2000.
9. Matsui N, Kasajima K, Hada M, Nagata T, Senga N, Yasui Y, Fukuishi N and Akagi M: Inhibitor of NF-kappaB activation during ischemia reduces hepatic ischemia/reperfusion injury in rats. *J Toxicol Sci* 30: 103-110, 2005.
10. Cao CC, Ding XQ, Ou ZL, Liu CF, Li P, Wang L and Zhu CF: *In vivo* transfection of NF-kappaB decoy oligodeoxynucleotides attenuate renal ischemia/reperfusion injury in rats. *Kidney Int* 65: 834-845, 2004.
11. Cooper JT, Stroka DM, Brostjan C, Palmethofer A, Bach FH and Ferran C: A20 blocks endothelial cell activation through a NF-kappaB-dependent mechanism. *J Biol Chem* 271: 18068-18073, 1996.
12. Longo CR, Arvelo MB, Patel VI, Daniel S, Mahiou J, Grey ST and Ferran C: A20 protects from CD40-CD40 ligand-mediated endothelial cell activation and apoptosis. *Circulation* 108: 1113-1118, 2003.
13. Lutz J, Luong le A, Strobl M, Deng M, Huang H, Anton M, Zakkar M, Enesa K, Chaudhury H, Haskard DO, *et al*: The A20 gene protects kidneys from ischaemia/reperfusion injury by suppressing pro-inflammatory activation. *J Mol Med (Berl)* 86: 1329-1339, 2008.
14. Lee EG, Boone DL, Chai S, Libby SL, Chien M, Lodolce JP and Ma A: Failure to regulate TNF-induced NF-kappaB and cell death responses in A20-deficient mice. *Science* 289: 2350-2354, 2000.
15. Partridge J, Carlsen H, Enesa K, Chaudhury H, Zakkar M, Luong L, Kinderlerer A, Johns M, Blomhoff R, Mason JC, *et al*: Laminar shear stress acts as a switch to regulate divergent functions of NF-kappaB in endothelial cells. *FASEB J* 21: 3553-3561, 2007.
16. Furuichi K, Wada T, Iwata Y, Sakai N, Yoshimoto K, Kobayashi Ki K, Mukaida N, Matsushima K and Yokoyama H: Administration of FR167653, a new anti-inflammatory compound, prevents renal ischaemia/reperfusion injury in mice. *Nephrol Dial Transplant* 17: 399-407, 2002.

17. Nicolau C, Le Pape A, Soriano P, Fargette F and Juhel MF: In vivo expression of rat insulin after intravenous administration of the liposome-entrapped gene for rat insulin I. *Proc Natl Acad Sci USA* 80: 1068-1072, 1983.
18. Levine RM, Pearce TR, Adil M and Kokkoli E: Preparation and characterization of liposome-encapsulated plasmid DNA for gene delivery. *Langmuir* 29: 9208-9215, 2013.
19. Kis A, Yellon DM and Baxter GF: Role of nuclear factor-kappa B activation in acute ischaemia-reperfusion injury in myocardium. *Br J Pharmacol* 138: 894-900, 2003.
20. Donnahoo KK, Shames BD, Harken AH and Meldrum DR: Review article: The role of tumor necrosis factor in renal ischemia-reperfusion injury. *J Urol* 162: 196-203, 1999.
21. Nichols TC: NF-kappaB and reperfusion injury. *Drug News Perspect* 17: 99-104, 2004.
22. Shen WH, Zhang CY and Zhang GY: Antioxidants attenuate reperfusion injury after global brain ischemia through inhibiting nuclear factor-kappa B activity in rats. *Acta Pharmacol Sin* 24: 1125-1130, 2003.
23. Zou XL, Pei DA, Yan JZ, Xu G and Wu P: A20 overexpression inhibits lipopolysaccharide-induced NF-kappaB activation, TRAF6 and CD40 expression in rat peritoneal mesothelial cells. *Int J Mol Sci* 15: 6592-6608, 2014.
24. Mazière C and Mazière JC: Activation of transcription factors and gene expression by oxidized low-density lipoprotein. *Free Radic Biol Med* 46: 127-137, 2009.