

Cassiae semen: A review of its phytochemistry and pharmacology (Review)

XIAOXV DONG, JING FU, XINGBIN YIN, CHUNJING YANG, XIN ZHANG,
WENPING WANG, XUEYING DU, QINGLING WANG and JIAN NI

Department of Chinese Medicine Pharmaceuticals, School of Chinese Materia Medica,
Beijing University of Chinese Medicine, Beijing 100102, P.R. China

Received June 2, 2016; Accepted April 11, 2017

DOI: 10.3892/mmr.2017.6880

Abstract. *Cassiae* semen (*Leguminosae*), a well-known traditional Chinese medicine, has been used for a number of centuries in areas of Southeast Asia, including Korea, Japan and China. The present review aims to provide updated and comprehensive information, on the botany, phytochemistry and pharmacology of *Cassiae* semen. The available information on *Cassiae* semen was collected using several different resources, including classic books on Chinese herbal medicine and a number of scientific databases, including the China Academic Journals full-text database, PubMed, SciFinder, the Web of Science and Science Direct. To date >70 chemical compounds have been isolated from *Cassiae* semen, and the major components have been determined to be anthraquinones, naphthopyrones and volatile oil. The crude extracts and pure compounds of *Cassiae* semen have been used as effective agents in preclinical and clinical practice due to their beneficial activities, including antihyperlipidemic, antidiabetic, neuroprotective, hepatoprotective, antibacterial, antioxidant and hypotensive activities. With the body of reported data, it has been suggested that *Cassiae* semen has convincing medicinal potential. However, the pharmacological mechanisms of the main bioactive compounds and the association between structure and activity require further investigation.

Contents

1. Introduction
2. Ethnobotany

Correspondence to: Professor Jian Ni, Department of Chinese Medicine Pharmaceuticals, School of Chinese Materia Medica, Beijing University of Chinese Medicine, 11 North Third Ring Road, Beijing 100102, P.R. China
E-mail: njtem@263.net

Key words: *Cassiae* semen, phytochemistry, pharmacology

3. Phytochemistry
4. Pharmacology
5. Conclusions

1. Introduction

Cassiae semen, also known as ‘Juemingzi’ in Chinese, is the dry and mature seed of *Cassia obtusifolia* L. or *C. tora* L., which belong to the *Cassia* genus of *Leguminosae* (1). It is cultivated in Korea, Japan and China, and is commonly consumed as a roasted tea (2,3). In traditional Chinese medicine, it has been used in treatments for hyperlipemia, diabetes mellitus, Alzheimer's disease, acute liver injury, inflammation, photophobia, headache, dizziness and hypertension (4-6).

Phytochemical investigations have isolated and identified >70 compounds, including anthraquinones, naphthopyrones, volatile oils and sterols (7,8). Among these, anthraquinones are the primary functional components and possess a wide spectrum of pharmacological properties (9-11), including antihyperlipidemic, neuroprotective, hepatoprotective, antibacterial and antimutagenic activities (12-14). Naphthopyrones, other primary components, exhibit antidiabetic (15,16), antimicrobial (17), antiestrogenic (18), antiallergic (19) and anthelmintic effects (20). At present, the Pharmacopoeia of the People's Republic of China recommends the use of chryso-phenol and aurantio-obtusin as the indicator components, and the quality of *Cassiae* semen is evaluated primarily by assessing the content of these two compounds (1).

The purpose of the present review is to provide comprehensive information on the ethnobotany, phytochemistry and pharmacology of *Cassiae* semen collated from previous studies, in order to facilitate the further study and application of *Cassiae* semen, as well as generate a novel basis for the associations between structure and activity, and their molecular mechanisms of action.

2. Ethnobotany

C. obtusifolia is similar to *C. tora* in terms of botanical morphology. The two are an annual, erect, stout herb, ~1-2 m in length, and their leaves are paripinnate, typically pubescent and are 4-8 cm in length with a conical gland between each of the two lowest pairs of leaflets. Leaflets are formed of 3 leaf

pairs and are glaucous, membranous, glabrous or pubescent, and have obcordate or obovate oblong morphology (2-6 cm long x 1.5-2.5 cm wide); the base is somewhat oblique, usually rounded and there are 8-10 pairs of main nerves. The petioles are 1.5-2 mm in length and their stipules are linear, pilose and caducous. It blossoms from July to September and produces fruit from September to October. Flowers are usually in subsessile pairs in leaf axils, the pedicels are filiform and are 1-1.5 cm in length. Calyces are ovate, glabrous, membranous and comprised of five-parts; there are five petals, which are pale yellow, oblong, obtuse and the upper petal (standard) is two-lobed. The flowers have 10 stamens, while the upper three are reduced to minute staminodes. The pods are slender, puberulous, subtetragonous, obliquely septate and, are ~15 cm in length and 3-4 mm in width.

However, the seeds of *C. obtusifolia* are a dark brown or green-brown, rhombohedral or short cylindrical, and are 3-7 mm in length and 2-4 mm in width. While *C. tora* seeds are a light brown, shiny, short cylindrical, and are 3-5 mm in length and 2-3 mm in width (1,21).

C. obtusifolia is cultivated in multiple provinces of China, including Henan, Hubei, Shanxi, Sichuan, Zhejiang and Anhui, and also other countries, including Korea, India and Japan. It is primarily distributed in moist and sunny places, in hillside shrubs and in the sandy soil of river banks (21).

As a widely used traditional Chinese medicine, there are some adulterants of this plant, including the seeds of *C. occidentalis* (*Leguminosae*), *C. sophera* (*Leguminosae*) and *Sesbania aculeata Pers* (*Papilionaceae*) (22-24). To date, a number of methods have been developed to identify and distinguish these, including experiential identification, morphological identification, ultraviolet spectrophotometry, the thin layer chromatography method, high performance liquid chromatography (HPLC), HPLC-coupled with time-of-flight and ion trap mass spectrometry, and SDS-PAGE (25-28). Among these methods, the HPLC method is regarded as the most popular method for evaluating the quality and authenticity of *Cassiae* semen. Chrysophanol and aurantio-obtusin are used as the indicator compounds to characterize the quality of this plant and the minimum contents are defined as 0.20 and 0.080%, respectively, in the Pharmacopoeia of the People's Republic of China (1).

3. Phytochemistry

A number of compounds, including anthraquinones, naphthopyrones, volatile oils and sterols, have been isolated from *Cassiae* semen. Anthraquinones and naphthopyrones, which exhibit multiple pharmacological activities, are considered the primary active ingredients of *Cassiae* semen. All compounds isolated from *Cassiae* semen are listed in Table I, and their chemical structures are displayed in Figs. 1-3.

Anthraquinones. *Cassiae* semen contains structurally diverse and biologically active anthraquinones. Thus far, ~53 anthraquinones have been isolated and identified. The predominant anthraquinones are emodin-type anthraquinones, which include emodin, chrysophanol, physcion, aloe-emodin, rhein, obtusin, chryso-obtusin, aurantio-obtusin, obtusifolin, questin, 1-desmethylaurantio-obtusin, 1-desmethylobtusin,

1-desmethylchryso-obtusin, chrysophanol-10,10'-bianthrone, 1,2-dihydroxyanthraquinone, 2-hydroxyemodin-1-methylether, alaternin, 1,3-dihydroxy-6-methoxy-7-methyl anthraquinone, 1-hydroxy-3,7-diformyl anthraquinone, chrysarobin, 8-O-methylchrysophanol, 1-O-methylemodin, 1,2-dimethoxy-8-hydroxy-3-methyl-9,10-anthraquinone and 1,2,7-trimethoxyl-6,8-dihydroxy-3-methylanthraquinone (compounds 1-24, respectively; Fig. 1); these have all been isolated from *Cassiae* semen (29-38). There are also many combined anthraquinones (compounds 25-46; Fig. 1), which have been isolated from the seeds of *C. obtusifolia* or *C. tora* (3,7,30,36,39-46).

Naphthopyrones. Naphthopyrones are the other characteristic components in *Cassiae* semen. In 1969, torachryson, rubrofusarin and rubrofusarin-6-O- β -D-gentiobioside were isolated from *C. tora* seeds (compounds 47-49, respectively; Fig. 2) (47,48). Subsequently, toralactone, rubrofusarin triglucoside, cassiaside, nor-rubrofusarin gentiobioside, cassiaside B, cassiaside C, torachryson apiglucoside, torachryson gentiobioside, torachryson tetraglucoside, cassiatoroside, demethylflavasperone gentiobioside and cassialactone gentiobioside were isolated from *C. tora* seeds (compounds 50-61, respectively; Fig. 2) (17,28,49-51). In addition, cassiaside B and cassiaside C were identified in the seeds of *C. obtusifolia* (52). Other naphthopyrones, including torosachryson, isotoralactone, cassialactone, cassiaside B₂, cassiaside C₂, nor-rubrofusarin-6-O- β -D (6'-O-acetyl) glucopyranoside and 1-hydroxyl-2-acetyl-3,8-dimethoxy-6-O-[β -D-apiofuranosyl-(1 \rightarrow 2)- β -D-glucopyranosyl]-naphthalene were also isolated from *C. obtusifolia* seeds (compounds 62-68, respectively; Fig. 2) (19,53-55).

Volatile oils. Li (56) extracted the volatile oils from *Cassiae* semen by steam distillation, and subsequently identified 37 components according to the gas chromatography/mass spectrometry analysis. Among these peaks, the major volatile components were 9-octadecenoic acid (E) (22.15%), n-hexadecanoic acid (12.53%), 9,10-anthracenedione, 1,8-dihydroxy-3-methyl (7.66%), octadecanoic acid (4.56%) and 13-octadecenoic acid methyl ester (Z) (3.84%) (56).

Other compounds. A range of other components have been isolated from *Cassiae* semen, including malvalic acid, sterculic acid, mandelic acid, campesterol, aspidinol and 5,7-dihydroxychromone (compounds 69-74, respectively; Fig. 3) (5,57,58). In addition, the four flavonoid compounds, chrysin, chrysin-7-O- β -D-glucoside, galangin and cyanidenon, were also obtained and identified from *Cassiae* semen (compounds 75-78, respectively; Fig. 3) (58).

4. Pharmacology

Cassiae semen exerts a great variety of pharmacological activities due to its complex bioactive compounds. An overview of the pharmacological studies on *Cassiae* semen is presented in detail in the following sections.

Antihyperlipidemic activity. In traditional Chinese herbal medicine, *Cassiae* semen is used for the prevention and

Table I. Chemical compounds isolated from *Cassiae* semen.

Classification	No.	Chemical component	(Refs.)
Anthraquinones	1	Emodin	(37,38)
	2	Chrysophanol	(34)
	3	Physcion	(34)
	4	Aloe-emodin	(38)
	5	Rhein	(38)
	6	Obtusin	(35)
	7	Chryso-obtusin	(35)
	8	Aurantio-obtusin	(35)
	9	Obtusifolin	(35)
	10	Questin	(36)
	11	1-desmethylaurantio-obtusin	(36)
	12	1-desmethylobtusin	(36)
	13	1-desmethylchryso-obtusin	(36)
	14	Chrysophanol-10,10'-bianthrone	(36)
	15	1,2-dihydroxyanthraquinone	(7)
	16	2-hydroxyemodin-1-methylether	(7)
	17	Alaternin	(29)
	18	1,3-dihydroxy-6-methoxy-7-methyl anthraquinone	(30)
	19	1-hydroxy-3,7-diformyl anthraquinone	(30)
	20	Chrysarobin	(30,31)
	21	8-O-methylchrysophanol	(32)
	22	1-O-methylemodin	(32)
	23	1,2-dimethoxy-8-hydroxy-3-methyl-9,10-anthraquinone	(32)
	24	1,2,7-trimethoxyl-6,8-dihydroxy-3-methylanthraquinone	(33)
	25	Gluco-aurantioobtusin	(39)
	26	Emodin-6-glucoside	(30)
	27	Physcion-8-O- β -D-glucopyranoside	(7,40)
	28	Physcion-8-O- β -gentiobioside	(41)
	29	Emodin-1-O- β -gentiobioside	(41)
	30	Obtusifolin-2-O- β -D-glucoside	(42)
	31	Chrysophanol-1-O- β -gentiobioside	(41)
	32	Alaternin-1-O- β -D-glucopyranoside	(7)
	33	Alaternin-2-O- β -D-glucopyranoside	(29)
	34	Aurantio-obtusin-6-O- β -D-glucopyranoside	(43)
	35	Chryso-obtusin-2-O- β -D-glucopyranoside	(7)
	36	Obtusifolin-2-O- β -D-(6'-O-acetyl) glucopyranoside	(44)
	37	Emodin-8-O- β -D-glucopyranoside	(45)
	38	2-methoxyl-chrysophanol-8-O- β -D-glucopyranoside	(46)
	39	1-demethylaurantio-obtusin-2-O- β -D-glucopyranoside	(43)
	40	1,7,8-trimethoxyl-2-hydroxyl-3-methylanthraquinone-2-O- β -D-glucopyranoside	(33)
	41	1,7-diimethoxyl-2,8-dihydroxyl-3-methylanthraquinone-2-O- β -D-glucopyranoside	(33)
	42	1,2,7-trimethoxyl-6,8-dihydroxy-3-methylanthraquinone-6-O- β -D-glucopyranoside	(33)
	43	2,8-dimethoxyl-1,6-dihydroxy-3-methylanthraquinone-6-O- β -D-glucopyranoside	(33)
	44	1-[(β -D-glucopyranosyl-(1 \rightarrow 3)-O- β -glucopyranosyl-(1 \rightarrow 6)-O- β -D-glucopyranosyl)oxy]-8-hydroxy-3-methyl-9,10-anthraquinone	(42)
	45	1-[(β -D-glucopyranosyl-(1 \rightarrow 6)-O- β -glucopyranosyl-(1 \rightarrow 3)-O- β -D-glucopyranosyl-(1 \rightarrow 6)-O- β -D-glucopyranosyl)oxy]-8-hydroxy-3-methyl-9,10-anthraquinone	(42)
46	4,6,7-trimethoxyl-aloe-emodin-8-O- β -D-glucopyranoside	(46)	
Naphthopyrones	47	Torachryson	(48)
	48	Rubrofusarin	(47)
	49	Rubrofusarin-6-O- β -D-gentiobioside	(47)

Table I. Continued.

Classification	No.	Chemical component	(Refs.)
	50	Toralactone	(49)
	51	Rubrofusarin triglucoside	(17)
	52	Cassiaside	(50)
	53	Nor-rubrofusarin gentiobioside	(17)
	54	Cassiaside B	(50)
	55	Cassiaside C	(50)
	56	Torachryson apiglucoside	(17)
	57	Torachryson gentiobioside	(17)
	58	Torachryson tetraglucoside	(17)
	59	Cassiatoroside	(51)
	60	Demethylflavasperone gentiobioside	(17)
	61	Cassialactone gentiobioside	(28)
	62	Torosachryson	(53)
	63	Isotalactone	(53)
	64	Cassialactone	(53)
	65	Cassiaside B ₂	(19)
	66	Cassiaside C ₂	(19)
	67	Nor-rubrofusarin-6-O-β-D(6'-o-acetyl) glucopyranoside	(54)
	68	1-hydroxyl-2-acetyl-3,8-dimethoxy-6-O-[β-D-apiofuranosyl-(1→2)-β-D-glucopyranosyl]-naphthalene	(55)
Other compounds	69	Malvalic acid	(5,57)
	70	Sterculic acid	(5)
	71	Mandelic acid	(5)
	72	Campesterol	(5)
	73	Aspidinol	(58)
	74	5,7-dihydroxymenone	(58)
	75	Chrysin	(58)
	76	Chrysin-7-O-β-D-glucoside	(58)
	77	Galangin	(58)
	78	Cyanidenon	(58)

treatment of hyperlipidemia. Several Chinese herbal formulations containing *Cassiae* semen is available in the Chinese market for preventing the formation of atherosclerotic plaques (59). In certain Asian countries, including China and Korea, it is also commonly drunk as a roasted tea to reduce body weight (60,61). Previous studies using mice have evaluated the reductions in blood lipid contents induced by different *Cassiae* semen extracts obtained through different methods, including supercritical fluid extraction, systematic solvents (petroleum ether, ethyl acetate, n-butanol, 70% ethanol and water) and ethanol precipitation following water extraction. The results revealed that the n-butanol and ethyl acetate extracts were the most effective (62,63). In addition, the ethanol and aqueous extracts of *Cassiae* semen significantly decreased the serum levels of total cholesterol (TC), triglyceride (TG) and low-density lipoprotein cholesterol (LDL-C), however, they increased the levels of high-density lipoprotein cholesterol (HDL-C) (13,64,65). Similarly, He *et al* (66) reported that treatment with the water extract form of *C. obtusifolia* seeds decreased the blood-lipid level by inhibiting cholesterol synthesis. Cho *et al* (67)

demonstrated that soluble fibers from *C. tora* seeds markedly decreased liver TC and TG levels in rats fed with a high-cholesterol diet. The underlying mechanism may be mediated by increasing fecal bile acid excretion and down-regulating the production of lipogenic enzymes (67). In addition, soluble fibers decreased the serum levels of TC, TG and LDL-C in patients with type II diabetes without serious adverse effects (2). Liu *et al* (68) revealed that the ethanol extract of *Cassiae* semen upregulated the expression levels of peroxisome proliferator-activated receptor (PPAR)-γ, sterol regulatory element-binding protein-1c, hormone-sensitive lipase and triacylglycerol hydrolase, however, tumor necrosis factor receptor superfamily member 6 was downregulated in adipose tissue. The anti-hyperlipidemia activity of *Cassiae* semen is primarily due to its antioxidant components, such as anthraquinones and polysaccharides. There are a variety of bioactive anthraquinone components in *Cassiae* semen, including chrysophanol, physcion, aurantio-obtusin, obtusifolin and emodin, which have been observed to decrease the levels of TC and TG (69,70). Previous studies have demonstrated that anthraquinones isolated from *Cassiae* semen

Table II. Pharmacological activities of *Cassiae* semen.

Pharmacological activities	Actions	Extracts/compounds	Application	(Refs.)
Anti-hyperlipidemia activity	Reduces blood lipid levels	SFE, systematic solvents (petroleum ether, ethyl acetate, n-butanol, 70% ethanol and water)	<i>In vivo</i>	(62,63)
	Decreases the levels of TC, TG and LDL-C; Increases the level of HDL-C	Ethanol and aqueous extracts	<i>In vivo</i>	(13,64,65)
	Inhibits the synthesis of cholesterol	Water extract	<i>In vitro</i>	(66)
	Increases fecal bile acid excretion and downregulates the production of lipogenic enzymes	Soluble fibers	<i>In vivo</i>	(67)
Anti-hyperlipidemia activity	Upregulates the expression levels of PPAR γ , SREBP-1c, HSL and TGH; Downregulates the levels of FAS	Ethanol extract	<i>In vivo</i>	(68)
	Decreases TC and TG levels	Chrysophanol, physcion, aurantio-obtusin, obtusifolin and emodin	<i>In vitro</i>	(69,70)
	Decreases TC, TG and LDL-C levels; Increases HDL-C levels	Anthraquinones	<i>In vivo</i>	(73)
	Binds bile acids and reduces the absorption of cholesterol	Water-soluble polysaccharides	<i>In vitro</i>	(74)
Anti-diabetic activity	Inhibits AGEs activity	Cassiaside, cassiaside C, rubrofusarin-6-O- β -D-gentiobioside	<i>In vitro</i>	(15)
	Inhibits the expression of TGF-1 and ECM proteins	Rubrofusarin-6-O- β -D-gentiobioside, cassiaside	<i>In vitro</i>	(16)
	Inhibits AGEs accumulation and, RAGE and COX-2 expression	Methanol extract	<i>In vitro/vivo</i>	(75,76)
	Downregulates the expression of TGF- β 1, CTGF and smad3, and upregulates the protein expression of smad6	Water extract	<i>In vivo</i>	(77)
Neuroprotective activity	Inhibits AChE activity	Ethanol extract	<i>In vivo</i>	(6)
	Upregulates the expression of pCREB and BDNF	Ethanol extract	<i>In vivo</i>	(78)
	Attenuates secondary calcium dysregulation and cell death	Ethanol extract	<i>In vivo</i>	(79)
	Ameliorates the A β -induced synaptic dysfunction model	Obtusifolin, alaternin	<i>In vivo</i>	(80)
	Inhibits cell damage and protects DA neuronal degeneration	Ethanol extract	<i>In vitro/vivo</i>	(81)
	Improves learning and memory capacity; Inhibits MDA and MAO levels; Enhances the level of SOD	Protein and anthraquinone glucosides	<i>In vivo</i>	(82)

Table II. Continued.

Pharmacological activities	Actions	Extracts/compounds	Application	Reference
Hepatoprotective activity	Hepatoprotective effects	Methanol extract	<i>In vitro</i>	(42,50)
	Increases the serum levels of SOD and decreases the serum levels of AST, ALT and MDA	Aqueous extract	<i>In vivo</i>	(85,86)
	Increases the serum levels of SOD and decreases the serum levels of TG, TC, MDA, AST and ALT	Ethanol extract	<i>In vivo</i>	(10,87)
Antibacterial activity	Anti- <i>Staphylococcus aureus</i> ; Anti- <i>Escherichia coli</i> K12	Naphthalenes, anthraquinones	<i>In vitro</i>	(17)
	Exhibits fungicidal activity against <i>Botrytis cinerea</i> , <i>Erysiphe graminis</i> , <i>Phytophthora infestans</i> , <i>Puccinia recondita</i> , <i>Phacelia grisea</i> , and <i>Rhizoctonia solani</i> .	Anthraquinones	<i>In vitro</i>	(88)
	Anti- <i>Helicobacter pylori</i>	Ethanol and aqueous extracts	<i>In vitro</i>	(89)
	Anti- <i>Clostridium perfringens</i> ; Anti- <i>E. coli</i>	1,2-dihydroxyanthraquinone	<i>In vitro</i>	(90)
	Anti- <i>Fusarium oxysporum</i> ; Anti- <i>B. cinerea</i>	Chloroform extract	<i>In vitro</i>	(91)
Antioxidant activity	Accelerates the oxidation of deoxyribose; inhibits linoleic acid peroxidation	Water extract	<i>In vitro</i>	(92,93)
	Scavenges free oxygen radicals	Water extract	<i>In vitro</i>	(94)
	Inhibits superoxide radicals	Water-soluble polysaccharides	<i>In vitro</i>	(95)
	Decreases MDA serum levels	Water-soluble polysaccharides	<i>In vitro</i>	(96)
	Scavenges hydroxyl and superoxide radicals	Water-soluble polysaccharides	<i>In vitro</i>	(97)
	Scavenges DPPH radicals	Ethyl acetate fraction	<i>In vitro</i>	(98)
	Antioxidant effect	Methanol extract	<i>In vitro</i>	(12)
	Scavenges DPPH radicals	Alaternin, cassiaside and rubrofusarin-6-O- β -D-gentiobioside	<i>In vitro</i>	(100)
	Hypotensive activity	Reduces arterial blood pressure	Water extract	<i>In vivo</i>
Hypotensive activity		Water extract	<i>In vivo</i>	(103)
Decreases the blood pressure		Ethanol extract	<i>In vitro</i>	(104)
Other activities	Estrogenic activity	70% EtOH extract	<i>In vitro</i>	(105)
	Inhibits histamine release from mast cells	Cassiaside C ₂	<i>In vitro</i>	(19)
	Anti-platelet aggregation	Gluco-aurantioobtusin	<i>In vitro</i>	(39)
	Antigenotoxic activity	Water extract	<i>In vitro</i>	(106)
	Antimutagenic activity	Anthraquinone aglycones and naphthopyrone glycosides	<i>In vitro</i>	(14)

Table II. Continued.

Pharmacological activities	Actions	Extracts/compounds	Application	Reference
	Improves myocardial function and attenuates MI/R-induced injury	Water extract	<i>In vivo</i>	(107)

TC, total cholesterol; TG, triglyceride; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol; PPAR γ , peroxisome proliferator-activated receptor; SREBP-1c, sterol regulatory element-binding protein-1c; HSL, hormone-sensitive lipase; TGH, triacylglycerol hydrolase; FAS, tumor necrosis factor receptor superfamily member 6; AGEs, advanced glycation end products; TGF-1, transforming growth factor-1; ECM, extracellular matrix; RAGE, receptor for advanced glycosylation end product; COX-2, cyclooxygenase-2; CTGF, connective tissue growth factor; Smad3/6, mothers against decapentaplegic homolog 3 and 6; AChE, acetylcholinesterase; pCREB, phosphorylated cyclic AMP response element binding protein; BDNF, brain-derived neurotrophic factor; DA, dopaminergic; MDA, malondialdehyde; MAO, monoamine oxidase; SOD, superoxide dismutase; AST, aspartate transaminase; ALT, alanine transaminase; DPPH, 1-diphenyl-2-picrylhydrazyl; MI/R, myocardial ischemia and reperfusion.

were effective substances during hypolipidemic activities (71,72). These results were verified by a previous study, which applied an experimental hyperlipidemic rat model to investigate anthraquinone treatment (80 and 20 mg/kg, per os, for 20 days). The TC, TG and LDL-C levels were significantly reduced in a dose-dependent manner, however, the levels of HDL-C increased. Inhibition of cholesterol synthesis may be one of the underlying mechanisms involved in decreasing blood lipid levels (73). Water-soluble polysaccharides (WSPs) from *Cassiae* semen markedly inhibited the activities of α -amylase and pancreatic lipase, however, protease activity increased. The results demonstrated that WSPs had the ability to bind to bile acids and reduce the absorption of cholesterol, indicating that WSPs may have potential as an effective herbal ingredient in functional food applications (74).

Antidiabetic activity. A number of studies have demonstrated that *Cassiae* semen exhibits anti-diabetic activity. A total of three naphthopyrone glucosides (compounds 49, 52 and 55) isolated from the butanol-soluble extract of *Cassia* semen have been evaluated for their inhibitory activity on advanced glycation end products (AGEs) formation *in vitro*. The results revealed that these compounds possessed more potent inhibitory activity against AGEs compared with the aminoguanidine positive control (15). In addition, rubrofusarin-6-O- β -D-gentiobioside (compound 49) and cassiaside (compound 52) significantly inhibited the expression of transforming growth factor (TGF)-1 and extracellular matrix protein in glomerular mesangial cells cultured under diabetic conditions, suggesting that the active compounds in *Cassiae* semen may be effective in the treatment of renal complications associated with diabetes (16). Similarly, Kim *et al* (75) evaluated the preventive effects of the methanol extract of *Cassia* semen (200 mg/kg/day, for 12 weeks) on the development of diabetic nephropathy in streptozotocin (STZ)-induced diabetic rats. The results indicated that oral treatment with the *Cassia* semen methanol extract inhibited the development of diabetic nephropathy by inhibiting AGEs accumulation, receptor for advanced glycosylation end product and cyclooxygenase-2 expression in the renal cortex of STZ-diabetic rats (75,76). In

addition, Zhu (77) reported that the water extract of *Cassia* semen exhibited protective activity against STZ-induced renal fibrosis in diabetic rats. The underlying mechanisms may be associated with its ability to downregulate the expression of TGF- β 1, connective tissue growth factor and mothers against decapentaplegic homolog 3 (smad3), as well as upregulating the protein expression of smad6 (77).

Neuroprotective activity. The ethanolic extract from the seeds of *C. obtusifolia* has been reported to have a neuroprotective effect in brain disease models. Kim *et al* (6) suggested that *C. obtusifolia* (25, 50 or 100 mg/kg) significantly attenuated scopolamine or transient bilateral common carotid artery occlusion (2VO)-induced memory impairment. These effects are mediated by the enhancement of the cholinergic nervous system via acetylcholinesterase inhibition in a dose-dependent manner [half maximal inhibitory concentration (IC₅₀)=81.6 μ g/ml] (6). In addition, *C. obtusifolia* (10 or 50 mg/kg/day) exhibited a neuroprotective effect in a mouse transient global ischemia model due to its anti-inflammatory properties and the induced upregulated expression of phosphorylated cyclic AMP response element binding protein and brain-derived neurotrophic factor (78).

Drever *et al* (79) demonstrated that treatment with *C. obtusifolia* (0.1-10 μ g/ml) significantly attenuated secondary calcium dysregulation and cell death induced by N-methyl-D-aspartate and 3-nitropropionic acid in mouse hippocampal cultures, and no significant effect on cell death was induced by incubation with naturally-secreted oligomers of amyloid (A) β . Yi *et al* (80) reported for the first time, that *C. obtusifolia* (10 μ g/ml) ameliorated the A β -induced synaptic dysfunction model through anti-inflammatory and protein kinase B (Akt)/glycogen synthase kinase-3 β pathways. The results suggested that the neuroprotective effect may be attributable to obtusifolin (compound 9) and/or alaternin (compound 17) (80). In a further experiment, *C. obtusifolia* (0.1-1 μ g/ml) inhibited cell damage against oxidopamine (6-OHDA)-induced dopaminergic (DA) neural toxicity in PC12 cells through an anti-oxidant and antimitochondrial-mediated apoptosis mechanism. In a mesencephalic DA culture, *C. obtusifolia* (0.1-1 μ g/ml) protected the DA cells against 6-OHDA- and N-methyl-4-phenylpyridinium iodide-induced





Figure 1. Chemical structures of anthraquinones.

toxicities. In addition, *C. obtusifolia* (50 mg/kg/day for 15 days) significantly protected DA neuronal degeneration in a 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced mouse Parkinson's disease (PD) model by inhibiting the movement impairment and the loss of DA neurons, indicating that *C. obtusifolia* may be a useful neuroprotective candidate for PD (81). In addition, protein and anthraquinone glucosides

from *Cassia* semen improved learning and memory capacity, inhibited the malondialdehyde (MDA) and monoamine oxidase levels, and enhanced the level of superoxide dismutase (SOD) in the cerebrum of senile mice (82).

Hepatoprotective activity. It was recorded in the Compendium of Materia Medica that *Cassiae* semen exhibited the functions




26 $R_1=OH, R_2=O-glc$


27 $R_1=O-glc, R_2=OCH_3$

28 $R_1=O-glc-glc, R_2=OCH_3$

29 $R_1=O-glc-glc, R_2=H$




30 $R=O-glc$



31 $R_1=O-glc-glc, R_2=H$


32 $R_1=O-glc, R_2=OH$

33 $R_1=OH, R_2=O-glc$




34 $R_1=OH, R_2=O-glc, R_3=OH$

35 $R_1=O-glc, R_2=OCH_3, R_3=OCH_3$




36




37 $R_1=O-glc, R_2=OH, R_3=H$

38 $R_1=O-glc, R_2=H, R_3=OCH_3$




39 $R=O-glc$




40 $R_1=OCH_3, R_2=H, R_3=O-glc$

41 $R_1=OH, R_2=H, R_3=O-glc$

42 $R_1=OH, R_2=O-glc, R_3=OCH_3$




43 $R=O-glc$



44 $R=O-glc-glc-glc$

45 $R=O-glc-glc-glc-glc$



46 $R=O-glc$

Figure 1. Continued. Chemical structures of anthraquinones.




Figure 2. Chemical structures of naphthopyrones.




Figure 3. Chemical structures of other compounds.

of nourishing the liver and improving vision (83). In Korea, the aqueous extract of *C. tora* L. seeds has been used for the protection of the liver. A weak anti-hepatotoxic activity in CCl_4 -induced mice was observed when the drug was administered orally at a dose of 670 mg/kg (84). The methanol extract of *C. tora* L. seeds exhibited significantly protective effects in primary cultured hepatocytes against CCl_4 and toxicity. A total of two anthraquinone glycosides (compounds 44 and 45) and three naphthoquinone glycosides (compounds 49, 51 and 53) from the methanol extract of *C. tora* seeds were the primary chemical constituents (42,50). The preventative

effects of *Cassiae* semen on acute liver injury in mice induced by CCl_4 have also been investigated (85,86). When compared with the control group, varying concentrations of the aqueous extract of *Cassiae* semen significantly increased the serum levels of SOD, and decreased the serum levels of aspartate transaminase (AST), alanine transaminase (ALT) and MDA. These results indicate that *Cassiae* semen is potentially beneficial in the treatment or prevention of hepatic damage (85,86). In addition, the ethanol extract of *Cassiae* semen has been observed to increase the serum levels of SOD and to decrease the serum levels of TG, TC, MDA, AST and ALT (86). Total

anthraquinones from *Cassiae* semen exhibited a protective effect on alcohol-induced acute liver injury in mice by regulating fat metabolism, improving liver function and increasing the mRNA and protein expression levels of PPAR- γ (10,87).

Antibacterial activity. Antibacterial activity, an important effect of *Cassiae* semen, has been comprehensively investigated. Naphthalenes (compounds 47 and 50) and anthraquinones (compounds 1, 4 and 5) isolated from *C. tora* seeds exhibited significant antibacterial effects on four strains of methicillin-resistant *Staphylococcus aureus* [minimal inhibitory concentration (MIC) was 2-64 $\mu\text{g/ml}$] and a strain of methicillin-sensitive *S. aureus*. In addition, rhein (compound 5) and torachryson (compound 47) from the seeds of *C. tora* exhibited antibacterial activity against *Escherichia coli* K12 with MIC values of 512 and 128 $\mu\text{g/ml}$, respectively (17). Kim *et al.* (88) were the first to demonstrate that emodin (compound 1) from *C. tora* seeds has a median lethal dose (LC₅₀) value of 0.102, 0.163, 0.385 and 0.046 g/l against *Rhizoctonia solani*, *Botrytis cinerea*, *Phytophthora infestans* and *Erysiphe graminis*, respectively, and physcion (compound 3) has an LC₅₀ value of 0.248, 0.263, 0.518, and 0.073 g/l against *R. solani*, *B. cinerea*, *P. infestans* and *E. graminis*, respectively. In addition, the LC₅₀ value of rhein (compound 5) is 0.375, 0.478, and 0.047 g/l against *R. solani*, *B. cinerea* and *P. infestans*, respectively (88).

It has been reported that ethanol and aqueous extracts of *C. obtusifolia* seeds were inhibitory against *Helicobacter pylori* strains (MIC were 100 and 60 $\mu\text{g/ml}$, respectively) (89). In addition, 1,2-dihydroxyanthraquinone (compound 15) isolated from *C. obtusifolia* seeds was revealed to inhibit the growth of *Clostridium perfringens* and *E. coli*, indicating that this drug exhibited potent growth-inhibiting activities towards human intestinal bacteria (90). Li *et al.* (91) demonstrated that the chloroform extract of the seeds of *C. obtusifolia* also exhibited different inhibitory activities against *Fusarium oxysporum* and *B. cinerea* (IC₅₀ values were 0.57 mg/ml and 0.97 mg/ml).

Antioxidant activity. The water extract of *C. tora* seeds accelerated the oxidation of deoxyribose induced by Fe³⁺-EDTA/H₂O₂ and exhibited 94% inhibition of linoleic acid peroxidation at a concentration of 0.2 mg/ml. The underlying mechanisms of this may be mediated by reducing metal ions, scavenging hydroxyl radical and chelating ferrous ion (92,93). Xv and Hu (94) demonstrated that the water extract of *Cassiae* semen exhibited a potent ability to scavenge free oxygen radicals [IC₅₀ values were 2 mg/ml and 2 $\mu\text{g/ml}$ for hydroxyl radicals (OH \cdot) and hydrogen peroxide (H₂O₂), respectively]. WSP from *Cassiae* semen (0.022 mg/ml) effectively inhibited superoxide radicals (O²⁻) induced by pyrogallol autoxidation (95). The inhibitory effects of WSP on serum levels of MDA were used to evaluate its antioxidation capabilities. The results demonstrated that WSP decreased MDA serum levels with an IC₅₀ value of 15.80% (96). In another study, Liu *et al.* (97) optimized the extraction conditions for WSP of *Cassiae* semen (temperature 80°C, extraction time 3.5 h, solid-liquid ratio 1:30) and observed that WSP (94.03 $\mu\text{g/ml}$) had the ability to scavenge hydroxyl and superoxide radicals with scavenging rates of 43.32 and 64.97%, respectively.

In addition, ethyl acetate fraction and n-butanol fraction of *Cassiae* semen were evaluated by DPPH radical scavenging activity. The results revealed that the ethyl acetate fraction had a lower IC₅₀ value of 56.4 g/ml, when compared with the value of 80.6 g/ml for n-butanol fraction. 1-Desmethyllaurantio-obtusin (compound 11) exhibited good scavenging activity on DPPH with an IC₅₀ value of 4.5 \pm 0.7 g/ml, while aurantio-obtusin-6-O- β -D-glucopyranoside (compound 34) and questin (compound 10) exhibited moderate antioxidant activity, and their IC₅₀ values were 103.2 \pm 1.5 g/ml and 185.2 \pm 1.8 g/ml, respectively. When compared with these results, chryso-obtusin (compound 7) and aurantio-obtusin (compound 8) demonstrated weaker antioxidant activity (IC₅₀ >200 $\mu\text{g/ml}$) (98). The methanolic extract of *C. tora* seeds exhibited a high antioxidant activity on lipid peroxidation (99). Similarly, in another study, Yen *et al.* (12) demonstrated that the methanolic extract of *C. tora* seeds exerted a greater antioxidant activity than the other organic solvents (n-hexane and ethyl acetate). Emodin was also revealed to be an antioxidative component (12). In addition, alaternin (compound 17), cassiaside (compound 52) and rubrofusarin-6-O- β -D-gentiobioside (compound 49) isolated from *C. tora* seeds exhibited good scavenging activity against DPPH radicals with IC₅₀ values of 17.59, 32.52 and 18.04 $\mu\text{g/ml}$, respectively (100).

Hypotensive activity. Aqueous and ethanol extracts of *Cassiae* semen have been reported to possess hypotensive effects (101). Koo *et al.* (101) reported that the water extract of *C. tora* seeds (3.75, 7.5, 15, 30, 60 and 250 mg/kg) consistently reduced arterial blood pressure in anesthetized rats. A potential reflex mechanism of this hypotensive action may involve a vagal reflex, which reciprocally inhibits the peripheral vasomotor tone via a reflex reduction in the sympathetic neural outflow to blood vessels (102). In addition, the media portion of the medullary reticular formation has been revealed to be directly involved in the hypotensive effect of *C. tora* seeds (103). Furthermore, the ethanol extract of *Cassiae* semen significantly decreased blood pressure in hypertensive rats by inhibiting receptor-controlled calcium channels on vessels and regulating the secretion of nitric oxide and inducible nitric oxide synthase (104).

Other activities. In addition to the pharmacological effects described above, *Cassiae* semen and its ingredients have other pharmacological effects, including estrogenic, anti-allergic, antigenotoxic, anti-aggregatory, antimutagenic and cardio-protective effects. Some of these effects are discussed briefly below.

The estrogenic activity of *C. obtusifolia* seeds was evaluated by a recombinant yeast screening assay. The results revealed that 70% EtOH extracts of this drug exhibited estrogenic relative potency [half maximal effective concentration (EC₅₀) was 60.2 $\mu\text{g/ml}$] (105). Cassiaside C₂ (compound 66) isolated from *C. obtusifolia* seeds exhibited a potent anti-allergic activity by inhibiting the histamine release from mast cells induced by antigen-antibody reaction (19). Furthermore, gluco-aurantioobtusin (compound 25) from *C. obtusifolia* seeds possessed potent inhibitory activities against arachidonic acid-, ADP- and collagen-induced platelet aggregations (39). Wu and Yen (106) demonstrated that the water extract of *C. tora* seeds exhibited potential

antigenotoxic activities against the dietary mutagens Glu-P-1 and TrpP-1 in the Ames test and the Comet assay. The potential mechanisms may be associated with neutralization of the reactive intermediate of Trp-P-1 and an antioxidant effect of the tested compounds (106). Anthraquinone aglycones (compounds 2, 7 and 8) and naphthopyrone glycosides (compounds 49 and 52) from *C. tora* seeds exhibited significant antimutagenic activity *in vitro*. The mechanism associated with these compounds may be mediated via interactions with a microsomal activating system (14). Fu *et al* (107) reported that the water extract of *Cassiae* semen (10 mg/kg/day, for one week) effectively improved myocardial function, and attenuated myocardial ischemia and reperfusion-induced injury and apoptosis in diabetic animals, which is potentially attributable to the reduced plasma lipid levels and the triggered cell survival Akt and extracellular signal-regulated kinases 1/2 signaling.

5. Conclusions

In traditional Chinese medicine, *Cassiae* semen has long been used to clean the liver, brighten the eye, loosen the bowel to relieve constipation, and for the treatment of inflammation, photophobia, headaches, dizziness, hyperlipemia and Alzheimer's disease. In addition, *Cassiae* semen is commonly used in the composition of other herbs. Although modern experiments have confirmed that this drug alone exhibits multiple pharmacological activities, it is important to investigate the molecular mechanisms of *Cassiae* semen combined with other herbs based on traditional uses.

A number of studies have investigated the effective constituents of *Cassiae* semen from different batches and geographical areas. HPLC-fingerprint chromatography is a common method to compare the differences (108-111). Zhang *et al* (3) developed a sensitive and reliable ultra-high-performance liquid chromatography-electrospray ionization-tandem mass spectrometry (UHPLC-ESI-MS/MS) method to evaluate the quality of *Cassiae* semen through simultaneous determination of 13 components, providing a novel basis for the overall assessment of the quality of this plant. In addition, a novel nonaqueous capillary electrophoresis method was used for the analysis of aurantio-obtusin, emodin and rhein in *Cassiae* semen with satisfactory results (112). Yang *et al* (113) was the first to simultaneously determine 7 anthraquinones in rat plasma by UHPLC-MS/MS following oral administration of *Cassiae* semen extract. These results may support investigations into the bioactivity mechanism and clinical application of this drug (113). Anthraquinones and naphthopyrones are considered to be the major constituents. Therefore, characteristic compounds or a biological index should be established to evaluate the quality and ensure their clinical application is suitable. In the Pharmacopoeia of the People's Republic of China, chrysophanol and aurantio-obtusin are used as the indicator compounds to characterize the quality of *Cassiae* semen with the minimum contents of 0.20 and 0.080%, respectively (1).

A total of 79 compounds including anthraquinones, naphthopyrones and volatile oil have been isolated and identified from *Cassiae* semen (Table I; Figs. 1-3). It has also been suggested that certain efforts should be made to isolate and identify novel compounds from *Cassiae* semen, in order to

strengthen its pharmacological profile to develop it further as a candidate for novel drug developments in the future.

Pharmacological studies have revealed that *Cassiae* semen possesses a variety of biological effects, including anti-hyperlipidemic, anti-diabetic, neuroprotective, hepatoprotective, antimicrobial, anti-oxidant and hypotensive activities (5,90,93,114,115). Extracts and compounds responsible for the pharmacological properties have also been determined, as presented in Table II. Although the pharmacological properties of certain traditional uses of *Cassiae* semen have been validated, these studies were primarily conducted *in vitro* (16,116,117). Therefore, the effects of these compounds require verification *in vivo*. In addition, the association between structure and activity, and the potential synergistic action exerted by the bioactive compounds requires further elucidation. It is anticipated that the comprehensive and current research on the pharmacological activities of extracts, as well as on active molecules isolated from *Cassiae* semen, provided in this review will inspire novel strategies in therapeutics for curing a number of different ailments.

Acknowledgements

The present review was financially supported by the Collaborative Innovation Construction Plan of Beijing University of Chinese Medicine (grant no. 2013-XTCX-03).

References

1. Editorial Committee of Chinese Pharmacopoeia. Chinese Pharmacopoeia, 2015 (eds). Medical Science and Technology Press, Beijing, China, pp145, 2015.
2. Cho SH, Kim TH, Lee NH, Son HS, Cho IJ and Ha TY: Effects of Cassia tora fiber supplement on serum lipids in Korean diabetic patients. *J Med Food* 8: 311-318, 2005.
3. Zhang W, Wang Y, Wang Q, Yang WJ, Gu Y, Wang R, Song XM and Wang XJ: Quality evaluation of Semen Cassiae (*Cassia obtusifolia* L.) by using ultra-high performance liquid chromatography coupled with mass spectrometry. *J Sep Sci* 35: 2054-2062, 2012.
4. Chen JK and Chen TT: Chinese Medical Herbology and Pharmacology. CA: Art of Medicine Press, p803, 2001.
5. Hao YJ, Sang YL and Zhao YQ: The advancement of the studies on the seeds of *Cassia obtusifolia*. *Chinese Tradit Herb Drugs* 32: 858-859, 2001.
6. Kim DH, Yoon BH, Kim YW, Lee S, Shin BY, Jung JW, Kim HJ, Lee YS, Choi JS, Kim SY, *et al*: The seed extract of *Cassia obtusifolia* ameliorates learning and memory impairments induced by scopolamine or transient cerebral hypoperfusion in mice. *J Pharmacol Sci* 105: 82-93, 2007.
7. Kitanaka S, Kimura F and Takido M: Studies on the Constituents of seeds of *Cassia obtusifolia* LINN. The structures of two new anthraquinone glycosides. *Chem Pharm Bull* 33: 1274-1276, 1985.
8. Zhang C, Li GL, Xiao YQ and Pang LZ: Two new glycosides from the seeds of *Cassia obtusifolia*. *Chinese Chem Lett* 20: 1097-1099, 2009.
9. Deng ZY, Zhang JW, Li J, Fan YW, Cao SW, Huang RL, Yin YL, Zhong HY and Li TJ: Effect of polysaccharides of cassiae seeds on the intestinal microflora of piglets. *Asia Pac J Clin Nutr* 16 (Suppl 1): 143-147, 2007.
10. Luo X, Xu X, Huang C, Wu X, Liu J, Lan B and Xu J: Experiment study of total anthraquinone in *Cassiae* Semen on lipid peroxidation and PPAR- γ expression in liver tissues of rats with alcoholic fatty liver. *Zhongguo Zhong Yao Za Zhi* 36: 1654-1658, 2011 (In Chinese).
11. Su H, Wang Z and Tang L: Simultaneous determination of 4 major components in *Cassiae* Semen *obtusifolia* by HPLC. *Zhongguo Zhong Yao Za Zhi* 36: 1327-1329, 2011 (In Chinese).

12. Yen GC, Chen HW and Duh PD: Extraction and identification of an antioxidative component from Jue Ming Zi (*Cassia tora* L.). *J Agr Food Chem* 46: 820-824, 1998.
13. Patil UK, Saraf S and Dixit VK: Hypolipidemic activity of seeds of *Cassia tora* Linn. *J Ethnopharmacol* 90: 249-252, 2004.
14. Choi JS, Lee HJ, Park KY, Ha JO and Kang SS: In vitro antimutagenic effects of anthraquinone aglycones and naphthopyrone glycosides from *Cassia tora*. *Planta Med* 63: 11-14, 1997.
15. Lee GY, Jang DS, Lee YM, Kim JM and Kim JS: Naphthopyrone glucosides from the seeds of *Cassia tora* with inhibitory activity on advanced glycation end products (AGEs) formation. *Arch Pharm Res* 29: 587-590, 2006.
16. Jung DH, Kim YS, Kim NH, Lee J, Jang DS and Kim JS: Extract of *Cassiae* semen and its major compound inhibit S100b-induced TGF- β 1 and fibronectin expression in mouse glomerular mesangial cells. *Eur J Pharmacol* 641: 7-14, 2010.
17. Hatano T, Uebayashi H, Ito H, Shiota S, Tsuchiya T and Yoshida T: Phenolic constituents of *Cassia* seeds and antibacterial effect of some naphthalenes and anthraquinones on methicillin-resistant *Staphylococcus aureus*. *Chem Pharm Bull (Tokyo)* 47: 1121-1127, 1999.
18. Das C, Dash S, Sahoo DC, Mohanty A and Rout D: *Cassia tora*: A phyto-pharmacological overview. *Int J Res Ayurveda Pharm* 2: 1162-1174, 2011.
19. Kitanaka S, Nakayama T, Shibano T, Ohkoshi E and Takido M: Antiallergic agent from natural sources. Structures and inhibitory effect of histamine release of naphthopyrone glycosides from seeds of *Cassia obtusifolia* L. *Chem Pharm Bull (Tokyo)* 46: 1650-1652, 1998.
20. Deore SL, Khadabadi SS, Kamdi KS, Ingle VP, Kawalkar NG, Sawarkar PS, Patil UA and Vyas AJ: In vitro Anthelmintic activity of *Cassia tora*. *Int J Chem Tech Res* 1: 177-179, 2009.
21. Editorial Board of Flora of China. Science Publishing House Press, p126, 1988.
22. Du ZX: Comparative identification of *Cassiae* Semen and *Sesbania aculeata* Pers. *Chinese Pharm J* 30: 204-205, 1995.
23. Zhu SY, Xv HJ and Xv MG: Comparative identification of Semen *Cassiae* and *Cassia sophera*. *Chin Tradit Herb Drugs* 34: 379-380, 2003.
24. Liu J, Zou CC and Sun ZF: Comparative identification of *Cassiae* Semen and *Cassia occidentalis* L. *Heilongjiang Med Pharm* 29: 63-64, 2006.
25. Hu YJ, Wan L, Zhang JX, *et al.*: Identification of *Cassia obtusifolia* L. by TLC. *Lishizhen Med Mater Med Res* 17: 2129, 2006.
26. Sun GF: Identification of *Cassia obtusifolia* L., *Cassia tora*. and *Cassia occidentalis* L. by SDS-Polyacrylamide gelelectrophoresis. *Tianjin Pharm* 8: 69-71, 1996.
27. Wang JB, Zhou X and Hu ZF: Quality evaluation of *Cassiae* Semen by both indicated component determination and HPLC fingerprint. *Chin Tradit Herb Drugs* 39: 917-919, 2008.
28. Luo Y, Zhang L, Wang WH and Li B: Components identification in *Cassiae* Semen by HPLC-IT-TOF MS. *Chin J Pharm Anal* 35: 1408-1516; 2015.
29. Lee HJ, Choi JS, Jung JH and Kang SS: Alaternin glucoside isomer from *Cassia tora*. *Phytochemistry* 49: 1403-1404, 1998.
30. Chen QD, Xv R, Xv ZN, *et al.*: Progress in studies of active constituents of anthraquinones and their biological activities from *Cassiae* Semen. *Chins J Mod Appl Pharm* 20: 120-123, 2003.
31. DiGiovanni J and Boutwell RK: Tumor promoting activity of 1, 8-dihydroxy-3-methyl-9-anthrone (chrysarobin) in female SENCAR mice. *Carcinogenesis* 4: 281-284, 1983.
32. Jia ZB, Chen WW, Jiang JX and DING XL: Study on anthraquinone constituents in the seed of *Cassia tora* L. *Chem Ind Forest Prod* 29: 100-102, 2009.
33. Xv YL: Studies on the chemical constituents from Semen *Cassiae* and the influence of processing. Beijing: Chinese Academy of Medical Sciences 1: 15, 2014
34. Takido M: Studies on the constituents of the seeds of *Cassia obtusifolia* L. I. The Structure of Obtusifolin. *Chem Pharm Bull* 6: 397-400: 1958.
35. Takido M: Studies on the constituents of seeds of *Cassia obtusifolia* L. II. The structure of Obtusin, Chryso-obutsin and Aurantio-obuyusin. *Chem Pharm Bull* 8: 246, 1960.
36. Kitanaka S and Takido M: Studies on the constituents of the seeds of *Cassia obtusifolia*: The structures of three new anthraquinones. *Chem Pharm Bull* 32: 860-864, 1984.
37. Tang LY, Wang ZJ, Fu HM, *et al.*: Study on anthraquinones constituents from Semen *Cassiae*. *J Chinese Med Mater* 32: 717-719, 2009a.
38. Zhang ZX and Liang YF: Isolation and identification of chemical constituents from seeds of *Cassia obtusifolia*. *China Pharm* 23: 1782-1783, 2012.
39. Yun-Choi HS, Kim JH and Takido M: Potential inhibitors of platelet aggregation from plant sources, V. Anthraquinones from seeds of *Cassia obtusifolia* and related compounds. *J Nat Prod* 53: 630-633, 1990.
40. Hao YJ, Sang YL and Zhao YQ: Study on anthraquinone constituents in *Cassiae* Semen. *Chinese Tradit Herb Drugs* 34: 18-19, 2003.
41. Li CH, Wei XY and Li XE: A new anthraquinone glycoside from the seed of *Cassia obtusifolia*. *Chinese Chem Lett* 15: 1448-1450, 2004.
42. Wong SM, Wong MM, Seligmann O and Wagner H: Anthraquinone glycosides from the seeds of *Cassia tora*. *Phytochemistry* 28: 211-214, 1989.
43. Tang LY, Wang ZJ, Fu MH, He Y, Wu HW and Huang LQ: A new anthraquinone glycoside from seeds of *Cassia obtusifolia*. *Chinese Chem Lett* 19: 1083-1085, 2008.
44. Li G, Xiao Y, Li L, Zhang C and Pang Z: Studies on chemical constituents of roasted seeds of *Cassia obtusifolia*. *Zhongguo Zhong Yao Za Zhi* 34: 54-56, 2009 (In Chinese).
45. Tang LY, Wang ZJ, He Y, *et al.*: Glycosides from seeds of *Cassia obtusifolia*. *Chinese J Exp Tradit Med Formulae* 15: 35-37, 2009b.
46. Jia ZB and Ding XL: Anthraquinone constituents from seeds of *Cassia tora* L. *J Chinese Med Mater* 29: 28-29, 2006.
47. Miyuki K, Eisaku M and Shoji S: Chemical Studies on the Oriental Plant Drugs. XXI. The Constituents of *Cassia tora* L. (2). A Glycoside of Rubrofusarin. *Chem Pharm Bull* 17: 458-461, 1969.
48. Shibata S, Morishita E and Kaneda M: Chemical Studies on the Oriental Plant Drugs. XX. The Constituents of *Cassia tora* L.(1). The Structure of Torachryson. *Chem Pharm Bull* 17: 454-457, 1969.
49. Takahashi S and Takido M: Studies on the constituents of the seeds of *Cassia tora* L. II.(On the purgative crude drugs. VII). The structure of the new naphthopyrone derivative, toralactone. *Yakugaku Zasshi* 93: 261, 1973.
50. Wong SM, Wong MM, Seligmann O and Wagner H: New anti-hepatotoxic naphthopyrone glycosides from the seeds of *Cassia tora*. *Planta Med* 55: 276-280, 1989.
51. Choi JS, Jung JH, Lee HJ, Lee HJ, Lee JH and Kang SS: A naphthalene glycoside from *Cassia tora*. *Phytochemistry* 40: 997-999, 1995.
52. Tianaka S and Takido M: Studies on the constituents of the seeds of *Cassia obtusifolia* L. The structures of two naphthopyrone glycosides. *Chem Pharm Bull* 36: 3980-3984, 1988.
53. Kitanaka S and Mieho T: Studies on the constituents of the seeds of *Cassia obtusifolia*: The structures of two new lactones, isotoralactone and cassialactone. *Phytochemistry* 20: 1951-1953, 1981.
54. Li L, Zhang C, Xiao YQ, Li W, Yin X, Chen D, Tian G and Wang Y: Glycosides of roasted seeds of *Cassia obtusifolia*. *Zhongguo Zhong Yao Za Zhi* 35: 1566-1568, 2010.
55. Wang ZJ, Wu QP, Tang LY, Fu MH, He Y, Gong QF and Hung LQ: Two new glycosides from the genus of *Cassia*. *Chinese Chem Lett* 18: 1218-1220, 2007.
56. Li YM: Comparison of volatile components of *Cassiae* Semen and semen seeds tea. *J Med Plants Res* 6: 3865-3869, 2012.
57. Jiao SF and Han HD: Studies on chemical constituents of *Cassiae* Semen. *Chinese J Clin Ration Drug Use* 3: 81-82, 2010.
58. Wu XH: Study on the chemical constituents, quality control and metabolism of *Cassia obtusifolia*. Wuhan: Huazhong University of Science and Technology 1: 11-12, 2010.
59. Guan Y and Zhao S: Yishou jiangzhi (de-blood-lipid) tablets in the treatment of hyperlipidemia. *J Tradit Chin Med* 15: 178-179, 1995.
60. Yang Y, Liu J and Lai XH: Observation on cassia seed tea combined with walking exercise on weight loss in the elderly. *Modern Prev Med* 40: 2468-2474, 2013.
61. Lai XH: Double intervention of cassia seed tea and sports on older women weight loss. *Chinese J Gerontol* 31: 2402-2404, 2011.
62. Zhang R, Feng ML and Wu YP: Experimental study on the active situs of fetid cassia seed to reduce blood lipid and their dose-effect relation. *China Remedies Clin* 5: 183-185, 2005b.
63. Zhang JX, Wan L, Hu YJ, Qu OL and Shi JY: Study on the effective part of reducing blood lipid in Semen *Cassiae*. *Lishizhen Med Mater Med Res* 17: 904-905, 2006.
64. Li CH, Li XE and Guo BJ: The effects of *Cassia obtusifolia* seeds extracts on reducing blood lipid. *J South China Normal Univ* 98: 29-32: 2002.

65. Wang YH, Gao L, Zhou WJ and Ma WJ: Effects of ethanol extraction from Cassiae Semen on Serum IL-6 and TNF- α in hyperlipidemia rats. *Chinese J Exp Tradit Med Formulae* 20: 178-181, 2014.
66. He JY, Liu SQ, Peng YF, *et al*: Study of the Mechanism of Cassia Obtusifolia L in decreasing blood -lipid. *China Pharm* 14: 202-203, 2003.
67. Cho JJ, Lee C and Ha TY: Hypolipidemic effect of soluble fiber isolated from seeds of Cassia tora Linn. In rats fed a high-cholesterol diet. *J Agric Food Chem* 55: 1592-1596, 2007.
68. Liu SM, Sun C and Xie WH: Effect of Semen Cassiae extracts on expression of lipogenesis genes in hyperlipidemia model mice. *Chinese Tradit Herb Drugs* 40: 583-587, 2009.
69. Wei N, Lv HR and Liu MF: Study on the chemical constitutions of reducing blood lipid in Cassiae Semen. *Guangdong Chem Ind* 39: 99-100, 2012.
70. Li BL: The active ingredients of reducing blood lipid in Semen Cassia. *China Prac Med* 7: 172-173, 2012.
71. Guo CY, Horn W and Pin DD: Extraction and identification of an antioxidative component from Jue Ming Zi (Cassia tora L.). *J Agric Food Chem* 46: 820-824, 1998.
72. Li XE and Guo BJ: Effects of protein and anthraquinone glucosides from Cassia seed on serum lipid of hyperlipidemia rats. *Zhongguo Zhong Yao Za Zhi* 27: 374-376, 2002 (In Chinese).
73. Li CH, Li XE, Fang KY and Guo BJ: Effects of anthraquinones from Cassia obtusifolia L. on cholesterol biosynthesis in cells. *J Clin Rehabil Tissue Eng Res* 12: 6593-6596, 2008.
74. Huang YL, Chow CJ and Tsai YH: Composition, characteristics, and in-vitro physiological effects of the water-soluble polysaccharides from Cassia seed. *Food Chem* 134: 1967-1972, 2012.
75. Kim YS, Jung DH, Sohn E, Lee YM, Kim CS and Kim JS: Extract of Cassiae semen attenuates diabetic nephropathy via inhibition of advanced glycation end products accumulation in streptozotocin-induced diabetic rats. *Phytomedicine* 21: 734-739, 2014.
76. Jung DH, Kim YS, Sohn EJ, *et al*: Effects of CS, a BuOH-Soluble Fraction of Cassiae Semen Methanolic Extract, on COX-2 Expression in Renal Cortex of STZ-Induced Diabetic Rats and Cultured Glomerular Mesangial Cells. *Diabetes* 56: pA155-pA155, 2007.
77. Zhu TC: Inhibitory Effects of Cassia Seed on the Renal Fibrosis in Diabetic Rats. *Chin J Exp Tradit Med Formulae* 18: 315-319, 2012.
78. Kim DH, Kim S, Jung WY, Park SJ, Park DH, Kim JM, Cheong JH and Ryu JH: The neuroprotective effects of the seeds of Cassia obtusifolia on transient cerebral global ischemia in mice. *Food Chem Toxicol* 47: 1473-1479, 2009.
79. Drever BD, Anderson WG, Riedel G, Kim DH, Ryu JH, Choi DY and Platt B: The seed extract of Cassia obtusifolia offers neuroprotection to mouse hippocampal cultures. *J Pharmacol Sci* 107: 380-392, 2008.
80. Yi JH, Park HJ, Lee S, Jung JW, Kim BC, Lee YC, Ryu JH and Kim DH: Cassia obtusifolia seed ameliorates amyloid β -induced synaptic dysfunction through anti-inflammatory and Akt/GSK-3 β pathways. *J Ethnopharmacol* 178: 50-57, 2016.
81. Ju MS, Kim HG, Choi JG, Ryu JH, Hur J, Kim YJ and Oh MS: Cassiae semen, a seed of Cassia obtusifolia, has neuroprotective effects in Parkinson's disease models. *Food Chem Toxicol* 48: 2037-2044, 2010.
82. Liu JZ, Lin X, Li XE and Guo BJ: Effect of protein and anthraquinone glucosides from Semen Cassiae on learning and memory capacity and related substances of senile mice induced by D-galactone. *Zhongguo Zhong Yao Za Zhi* 32: 516-519, 2007 (In Chinese).
83. Jiangsu New Medical College: *Encyclopedia of Chinese MateriaMedica*, I. Shanghai Science and Technology Press, Shanghai, China pp94984, 1975.
84. Yun HS and Chang IM: Plants with liver protective activities (I). *Korean J Pharmacogn* 8: 125-129, 1977.
85. Gao Q, Xu H and Chen J: Liver-protective and bowel-lubricating and defecation-promoting effects of crude and processed Semen Cassiae. *Tradit Chinese Drug Res Clin Pharmacol* 18: 194-196, 2007.
86. Lin DJ and Jin Z: Experimental Study on Protective Effect of Semen Cassiae Extract Against Acute Liver Injury. *Lishizhen Med Mater Med Res* 17: 214-215, 2006.
87. Niu YF, Zhao T, Zeng T, *et al*: Study on the protective effect of Cassiae Semen extract against alcohol-induced acute liver injury in mice. *J Toxicol* 24: 58-61, 2010.
88. Kim YM, Lee CH, Kim HG and Lee HS: Anthraquinones isolated from Cassia tora (Leguminosae) seed show an anti-fungal property against phytopathogenic fungi. *J Agr Food Chem* 52: 6096-6100, 2004.
89. Li Y, Xu C, Zhang Q, Liu JY and Tan RX: In vitro anti-Helicobacter pylori action of 30 Chinese herbal medicines used to treat ulcer diseases. *J Ethnopharmacol* 98: 329-333, 2005.
90. Sung BK, Kim MK, Lee WH, Lee DH and Lee HS: Growth responses of Cassia obtusifolia toward human intestinal bacteria. *Fitoterapia* 75: 505-509, 2004.
91. Li XH, Gong CR, Cao H, *et al*: Primary study on inhabiting of the extracts from Cassia obtusifolia seeds against Fusarium oxysporum and Botrytis cinerea. *J Shanxi Agric Univ* 26: 348-350, 2006.
92. Yen GC and Chung DY: Antioxidant Effects of Extracts from Cassia tora L. Prepared under different degrees of roasting on the oxidative damage to biomolecules. *J Agr Food Chem* 47: 1326-1332, 1999.
93. Yen GC and Chuang DY: Antioxidant properties of water extracts from Cassia tora L. In relation to the degree of roasting. *J Agr Food Chem* 48: 2760-2765, 2000.
94. Xv JG and Hu QP: Study on Free Radical Scavenging Capacity by Cassia Seed Water Extract in vitro. *Food Sci* 27: 73-75, 2006.
95. Guo CQ, Yan J, Wu XY, Xu GY, Fan CH and Gou XJ: Study on purification and antioxidation of water-soluble polysaccharide isolated from semen cassia. *Food Sci* 28: 205-208, 2007.
96. Liu J, Deng ZY and Yu HH: Antioxidation Study on Water-soluble Polysaccharide Isolated from Semen Cassiae. *Food Sci* 27: 61-63, 2006.
97. Liu C, Liu Q, Sun J, Jiang Bb, Yan J: Extraction of water-soluble polysaccharide and the antioxidant activity from Cassiae Semen. *J Food Drug Anal* 22: 492-499, 2014.
98. Zeng H, Liu Q, Wang M, Jiang S, Zhang L, He X, Wang J and Chen X: Target-guided separation of antioxidants from Semen cassia via off-line two-dimensional high-speed counter-current chromatography combined with complexation and extrusion elution mode. *J Chromatogr B* 1001: 58-65, 2015.
99. Kim SY, Kim JH, Kim SK, Oh MJ and Jung MY: Antioxidant activities of selected oriental herb extracts. *J Am Oil Chem Soc* 71: 633-640, 1994.
100. Choi JS, Lee HJ and Kang SS: Alaternin, cassiaside and rubrofusarin gentiobioside, radical scavenging principles from the seeds of Cassia tora on 1, 1-diphenyl-2-picrylhydrazyl (DPPH) radical. *Arch Pharm Res* 17: 462-466, 1994.
101. Koo A, Wang JC and Li KM: Extraction of hypotensive principles from seeds of Cassia tora. *Am J Chin Med (Gard City N Y)* 4: 245-248, 1976.
102. Koo A, Chan WS and Li KM: A possible reflex mechanism of hypotensive action of extract from Cassia tora seeds. *Am J Chin Med (Gard City N Y)* 4: 249-255, 1976.
103. Chan SH, Koo A and Li KM: The involvement of medullary reticular formation in the hypotensive effect of extracts from seeds of Cassia tora. *Am J Chin Med (Gard City N Y)* 4: 383-389, 1976.
104. Mao WH, Shang QH, Liu AD, *et al*: Effects of Cassiae Semen Extracts on Vasorelaxation and Its Mechanisms in Rat Aorta. *Chin J Hypertens* 18: 60-63, 2010.
105. Zhang CZ, Wang SX, Zhang Y, Chen JP and Liang XM: In vitro estrogenic activities of Chinese medicinal plants traditionally used for the management of menopausal symptoms. *J Ethnopharmacol* 98: 295-300, 2005.
106. Wu CH and Yen GC: Antigenotoxic properties of Cassia tea (Cassia tora L.): Mechanism of action and the influence of roasting process. *Life Sci* 76: 85-101, 2004.
107. Fu F, Tian F, Zhou H, Lv W, Tie R, Ji L, Li R, Shi Z, Yu L, Liang X, *et al*: Semen cassiae attenuates myocardial ischemia and reperfusion injury in high-fat diet streptozotocin-induced type 2 diabetic rats. *Am J Chinese Med* 42: 95-108, 2014.
108. Zhang P and Chen JW: HPLC Fingerprint of Cassiae Semen. *Chinese Tradit Herb Drugs* 38: 372-375, 2007.
109. Luo W, Liu B, Wang W, *et al*: HPLC fingerprint chromatogram of Cassiae Semen. *J Beijing Univ Tradit Chin Med* 32: 115-117, 2009.
110. Wang WY, Zhao Q, Zhang TJ, *et al*: HPLC Fingerprint and chemical pattern recognition of Cassiae Semen. *Chinese Tradit Herb Drugs* 40: 1638-1641, 2009.
111. Tang YL, Liang TZ, Zhang HX and Xu JY: Study on fingerprint of Cassiae Semen, ultramicro powder and powder particle. *Zhong Yao Cai* 34: 1861-1866, 2011 (In Chinese).

112. Wang N, Wu Y, Wu X, Liang S and Sun H: A novel nonaqueous capillary electrophoresis method for effective separation and simultaneous determination of aurantio-obtusin, emodin and rhein in semen cassiae and cassia seed tea. *Anal Methods* 6: 5133-5139, 2014.
113. Yang C, Wang S, Guo X, Sun J, Liu L and Wu L: Simultaneous determination of seven anthraquinones in rat plasma by Ultra High Performance Liquid Chromatography-tandem Mass Spectrometry and pharmacokinetic study after oral administration of Semen Cassiae extract. *J Ethnopharmacol* 169: 305-313, 2015.
114. Hyun SK, Lee H, Kang SS, Chung HY and Choi JS: Inhibitory activities of Cassia tora and its anthraquinone constituents on angiotensin-converting enzyme. *Phytother Res* 23: 178-184, 2009.
115. Dhanasekaran M, Ignacimuthu S and Agastian P: Potential hepatoprotective activity of ononitol monohydrate isolated from Cassia tora L. on carbon tetrachloride induced hepatotoxicity in Wistar rats. *Phytomedicine* 16: 891-895, 2009.
116. Jung HA, Ali MY, Jung HJ, Jeong HO, Chung HY and Choi JS: Inhibitory activities of major anthraquinones and other constituents from Cassia obtusifolia against β -secretase and cholinesterases. *J Ethnopharmacol* 191: 152-160, 2016.
117. Wen CC, Shyur LF, Jan JT, Liang PH, Kuo CJ, Arulsevan P, Wu JB, Kuo SC and Yang NS: Traditional Chinese medicine herbal extracts of Cibotium barometz, Gentiana scabra, Dioscorea batatas, Cassia tora and Taxillus chinensis inhibit SARS-CoV replication. *J Tradit Complement Med* 1: 41-50, 2011.