Baicalin prevents the apoptosis of endplate chondrocytes by inhibiting the oxidative stress induced by H₂O₂

YUTAO PAN*, DI CHEN*, QINGYOU LU, LIFENG LIU, XIA LI and ZENGCHUN LI

Department of Emergency and Trauma Surgery, Shanghai East Hospital Affiliated to Tongji University, Shanghai 200120, P.R. China

Received July 6, 2016; Accepted May 8, 2017

DOI: 10.3892/mmr.2017.6904

Abstract. Osteoarthritis (OA) is a degenerative disease of articular cartilage. The pathogenesis of OA remains to be fully elucidated, and several studies have found that oxidative stress is important in its pathogenesis. Baicalin is well known and has already been investigated for its role of inhibiting the oxidative stress pathway. Thus, the present study aimed to investigate the role of baicalin on the inhibition of oxidative stress in endplate chondrocytes induced by hydrogen peroxide (H₂O₂). Following treatment of endplate chondrocytes with different doses of H₂O₂ with or without baicalin for different incubation durations, a CCK-8 assay and Annexin V/PI staining were used to measure the cell proliferation and apoptotic rates to identify the optimal experimental conditions. Subsequently, for examining the effects and underlying mechanism of baicalin on oxidative stress, the protein expression levels of cleaved-poly (ADP-ribose) polymerase (PARP), B-cell lymphoma-2-associated X protein (Bax) and pro-caspase-3 were analyzed using western blot analysis, intracellular anti-oxidant activities, including those of malondialdehyde (MDA), superoxide dismutase (SOD) and nitric oxide (NO), were quantified, and the levels of endothelial nitric oxide synthase (eNOS) were examined using reverse transcription-polymerase chain reaction analysis. The results revealed that the oxidative stress of endplate chondrocytes induced by 0.5 mM H₂O₂ for 4 h were the most appropriate conditions for experiments, and pretreatment with 100 μ mol/l baicalin for 1 h effectively reversed the effect of H₂O₂ on the endplate chondrocytes. In addition, Annexin V/PI staining demonstrated that the cell death induced by H₂O₂ was apoptotic, and

baicalin reversed the apoptosis induced by oxidative stress. H₂O₂ activated PARP cleavage, and the expression of Bax and pro-caspase-3; however, baicalin inhibited the expression of these apoptotic signaling indicators. Baicalin also reduced the levels of MDA, and increased the levels of SOD and NO. Baicalin also significantly elevated the mRNA levels of eNOS in endplate chondrocytes. Therefore, the results of the present study showed that baicalin significantly inhibited the oxidative stress in endplate chondrocytes induced by H₂O₂, and decreased cell apoptosis.

Introduction

Osteoarthritis (OA) is a degenerative disease of articular cartilage, which is characterized by local and progressive destruction of articular cartilage, and osteophyte formation in the subcartilage edges, and is often accompanied by synovial inflammation. At present, the pathogenesis of articular cartilage injury in OA remains to be fully elucidated. Studies in previous decades have confirmed that oxidative stress is important in the pathogenesis and progression of OA (1,2). Oxidative stress is characterized by excessive reactive oxygen species (ROS) production and reduced cellular antioxidant activity, leading to membrane lipid peroxidation and mitochondrial dysfunction (3-5). If oxidative insult persists, programmed cell death is initiated, which can cause apoptosis of chondrocytes and extracellular matrix degradation (1,2).

Oxidative stress is primarily caused by the excessive accumulation of ROS, which include hydrogen peroxide (H₂O₂), superoxide anions and hydroxyl radicals. H₂O₂ is a major component of ROS and is used extensively as an inducer in oxidative stress models (5-9). Therefore, treatment strategies with the potential to prevent oxidative stress induced by H₂O₂ are considered valuable for patients with OA. Baicalin is a flavonoid glycoside extracted from a type of traditional Chinese medicine, *Scutellaria baicalensis*, and its chemical structure has been determined (10). Baicalin has been shown to possess antibacterial, anti-inflammatory and anti-apoptotic properties, and is widely used in the treatment of infectious and inflammatory diseases (8,11-14).

The aim of the present study was to investigate the potential protective effects of baicalin treatment on H₂O₂-induced oxidative stress in endplate chondrocytes *in vitro*, particularly its effect on cell apoptosis.

Correspondence to: Dr Zengchun Li or Dr Xia Li, Department of Emergency and Trauma Surgery, Shanghai East Hospital Affiliated to Tongji University, 150 Jimo Road, Shanghai 200120, P.R. China
E-mail: lzc0155@easthospital.cn
E-mail: daxia868@163.com

*Contributed equally

Key words: endplate chondrocyte, baicalin, H₂O₂, oxidative stress, apoptosis

Materials and methods

Isolation and culture of primary rat endplate chondrocytes.

A total of 40 4-week-old male Sprague-Dawley rats weighing 160–200 g were obtained from Shanghai SLAC Laboratory Animal Co., Ltd. (Shanghai, China) and housed at a temperature of 26°C, 40–60% humidity, standard diet and a 14-h light/10-h dark cycle. Rat endplate chondrocytes were obtained following cervical dislocation after anesthesia according to a previous study (15,16). In brief, the region of the lumbar posterior midline was shaved and then covered with sterile drapes. The rats were placed in a prone position on the operating table. Cartilage of the L3–L5 endplates was carefully removed from the vertebrae and placed into a small beaker with high-glucose (4.5 g/l; 25 mM) Dulbecco's modified Eagle's medium (DMEM; Gibco; Thermo Fisher Scientific, Inc., Waltham, MA, USA). Sterile ophthalmic scissors were used to cut the tissues into 1-mm³ sections, and these were transferred to a 50-ml sterile test tube. The tissues were digested with 0.25% trypsin (Hyclone; GE Healthcare Life Sciences; Logan, UT, USA) at 37°C for 30 min, followed by the addition of 0.2% collagenase (Sigma; Merck KGaA; Darmstadt, Germany) at 37°C for 2 h. Following centrifugation (400 x g for 10 min at room temperature), the supernatant was discarded and the tissue suspension was washed three times with DMEM with 10% heat-inactivated fetal bovine serum (FBS; Gibco; Thermo Fisher Scientific, Inc.). The released cells were cultured in DMEM with 10% FBS, 100 U/ml penicillin and 100 µg/ml streptomycin (both from Gibco; Thermo Fisher Scientific, Inc.). The cells were maintained at 37°C in a humidified atmosphere of 5% CO₂, and used at passage 4–6 for all experiments. The present study was approved by the Ethics Committee of Shanghai East Hospital Affiliated to Tongji University (Shanghai, China).

Cell viability of endplate chondrocytes treated with H₂O₂.

The endplate chondrocytes were grown in 96-well plates (BD Biosciences, Franklin Lakes, NJ, USA) at a density of ~1x10⁵/ml (200 µl/well) in DMEM supplemented with 10% FBS, 100 U/ml penicillin and 100 µg/ml streptomycin. After 24 h, the cells were washed with PBS buffer and incubated in serum-free medium at 37°C for 12 h. The cells were then washed again and incubated with medium containing different concentrations (0.1, 0.5 and 1.0 mM) of H₂O₂ (Lingfeng Chemicals Co., Ltd., Shanghai, China) at 37°C for 4 h, respectively. Untreated cells were referred to as the normal control cells. Cell viability was determined using a CCK-8 assay (Dojindo Molecular Technologies, Inc., Kumamoto, Japan). The CCK-8 reagent was diluted 10-fold with DMEM prior to being added (100 µl) to each well. After 2.5 h at 37°C, sample optical density (OD) values were read at 450 nm using a multi-mode microplate reader (FlexStation 3; Molecular Devices LLC, Sunnyvale, CA, USA). The OD₄₅₀ is proportional to the degree of cell viability. Data are shown as the mean of at least three independent experiments.

Annexin V/PI assay of endplate chondrocytes treated with H₂O₂. The apoptosis of endplate chondrocytes induced by H₂O₂ was measured using a commercial Annexin V/PI assay kit (Nanjing KeyGen Biotech Co., Ltd., Nanjing, China)

according to the manufacturer's protocol. Briefly, the endplate chondrocytes were cultured in 6-well plates (BD Biosciences) at a density of ~1.0x10⁵/ml (2 ml/well) and incubated in DMEM supplemented with 10% FBS, 100 U/ml penicillin and 100 µg/ml streptomycin. After 24 h, the cells were washed and incubated in serum-free medium for 12 h. The cells were then washed again and incubated in medium with various concentrations of H₂O₂ (0.1, 0.5 and 1.0 mM) at 37°C for 4 h. Following incubation, the cells were trypsinized and washed with PBS. Following centrifugation at 100 x g for 5 min at room temperature, the cells were resuspended in 500 µl binding buffer. The suspensions were transferred into 1.5-ml tubes, and 5 µl of Annexin V and 5 µl of PI solution were added. The cells were incubated in the dark at room temperature for 20 min, and flow cytometric analysis was performed using a BD FACSaria II flow cytometer (BD Bioscience). Data were analyzed using BD FACSDiva software version 6.1.3 (BD Bioscience), and Flowjo version 7.6.1 (FlowJo LLC, Ashland, OR, USA). The percentage apoptosis was determined as the percentage of Annexin V-positive/PI-negative cells.

Cell viability of endplate chondrocytes treated with baicalin.

The endplate chondrocytes were incubated with 0.5 mM H₂O₂ at 37°C and 5% CO₂ for 4 h, with baicalin added at different treatment time-points (-1, 0, 1 and 2 h). Untreated cells and cells treated with H₂O₂ only were referred as the control groups. The viability of the endplate chondrocytes was determined using a CCK-8 assay, as described above. In addition to determining the optimal incubation duration, the optimal concentration of baicalin on endplate chondrocytes was investigated. The cells were pretreated with baicalin at various concentrations (50, 100 and 200 µmol/l) for 1 h, and were incubated with 0.5 mM H₂O₂ at 37°C for 4 h, respectively. Untreated cells and cells treated with H₂O₂ only were referred as the control groups. As pretreatment of endplate chondrocytes with 100 µmol/l baicalin was optimal for inhibiting the oxidative stress induced by H₂O₂, the cell viability was compared between the H₂O₂-only group and the groups treated with H₂O₂ at different time-points (4, 8, 12 and 24 h). Untreated cells were referred to as the control group. All the above data are shown as the mean of at least three independent experiments.

Annexin V/PI assay of endplate chondrocytes pretreated with baicalin. The cells were pretreated with 100 µmol/l for 1 h, and then incubated in medium with 0.5 mM H₂O₂ at 37°C for 4 h. Based on the previously described method, the inhibitory effect of baicalin on the apoptosis of endplate chondrocytes induced by H₂O₂ was measured using an Annexin V/PI assay kit.

Western blot analysis. The protein expression of endplate chondrocytes was evaluated in the groups of chondrocytes treated with 0.5 mM H₂O₂ only for 4 h, with 0.5 mM H₂O₂ for 4 h+100 µmol/l baicalin pretreatment for 1 h, and in normal control group. For the extraction of proteins, the cells were placed in RIPA lysis buffer (Beyotime Institute of Biotechnology, Haimen, China) for 15 min at 4°C and centrifuged at 1,300 x g for 30 min at 4°C. The protein concentrations were determined using a NanoDrop

instrument, and 40 μ g of protein from each sample were run on a 15% SDS-PAGE gels. The separated proteins were transferred onto PVDF membranes. Following blocking with 5% nonfat dry milk in double-distilled water at room temperature for 1 h, the membranes were washed three times with TBS containing 0.1% Tween-20 (TBS-T) and incubated overnight at 4°C with primary mouse monoclonal anti-PARP (ab203467; 1:500), anti-B-cell lymphoma-2-associated X protein (Bax, ab32503; 1:1,000), anti-pro-caspase-3 (ab13847; 1:500), or anti-GAPDH antibodies (ab8245; 1:500) (all from Abcam, Cambridge, MA, USA). The membranes were then washed three times with TBS-T, followed by 1 h incubation at room temperature in a 1:1,000 dilution of goat anti-mouse/rabbit HRP antibody (715-035-150/415-035-166; Jackson ImmunoResearch Laboratories, Inc., West Grove, PA, USA). Following incubation, the membranes were washed three times with TBS-T. Immune complexes were visualized using enhanced chemiluminescence followed by exposure to a Tanon 5200 instrument (Tanon Science and Technology Co., Ltd., Shanghai, China). Data were normalized to the GAPDH content of the same sample.

Measurement of oxidative activity. The oxidative activities were measured at 6, 12 and 24 h in the 0.5 mM H_2O_2 , 0.5 mM H_2O_2 +100 μ mol/l baicalin pretreatment for 1 h, and normal control groups. The concentrations of malondialdehyde (MDA), superoxide dismutase (SOD) and nitric oxide (NO) were assessed using dedicated kits (Nanjing KeyGen Biotech Co., Ltd.) according to the manufacturer's protocols.

Reverse transcription-quantitative polymerase chain reaction (RT-qPCR) analysis. The mRNA expression levels of endothelial nitric oxide synthase (eNOS) were determined in the groups treated with 0.5 mM H_2O_2 only for 4 h, 0.5 mM H_2O_2 for 4 h+100 μ mol/l baicalin pretreatment for 1 h, and the normal control group. Total cellular RNA was extracted from the endplate chondrocytes using TRIzol (Invitrogen; Thermo Fisher Scientific, Inc.). OLIGO was used as the reverse transcription primer, and the total RNA was extracted from the cells of each group as the template, which was then reverse transcribed into cDNA under the RT enzyme. The reaction system included 500 ng RNA template, 5X buffer 2 μ l, PrimeScript RT enzyme mix 1 0.5 μ l, 10 μ mol/l 50 M OLIGO 0.5 μ l and 100 μ M random 6 mers 0.5 μ l. DEPC water was added up to 10 μ l at 37°C for 15 min. PCR amplification was performed in a 20 μ l reaction volume. The primer sequences were as follows: eNOS forward, 5'-CCAGCTAGCCAAAGTCACCAT-3' and reverse, 5'-GTCTCGGAGCCATACAGGATT-3'; GAPDH forward, 5'-CGGAGTCAACGGATTTGGTCGTAT-3' and reverse, 5'-AGCCTTCTCCATGGTGGTGAAGAC-3'. Triplicate reactions were run with a volume of 20 μ l, containing 2 μ l cDNA, 10 μ l 2X SYBR-Green mix, 6 μ l ddH₂O, 1 μ l PCR forward primer (10 μ M) and 1 μ l PCR reverse primer (10 μ M). Following an initial denaturation at 95°C for 5 min, the PCR conditions were as follows: 35 cycles of denaturation at 95°C for 30 sec, annealing at 55°C for 30 sec, and extension at 72°C for 30 sec. The 2^{- $\Delta\Delta C_q$} (quantification cycle) method (17) was used to calculate the mRNA expression levels of eNOS in each sample. Data were normalized to the GAPDH content of the same sample.

Statistical analysis. All data are expressed as the mean $\pm$ standard deviation. Statistical analysis was performed using Student's t-test and one-way analysis of variance using SPSS version 18.0 software (SPSS, Inc., Chicago, IL, USA). $P < 0.05$ was considered to indicate a statistically significant difference.

Results

H₂O₂ induces apoptotic cell death of endplate chondrocytes. To characterize the effects of H_2O_2 on the induction of cell death of endplate chondrocytes, cell viability and apoptotic rates were detected. As H_2O_2 concentration increased, the cell viability gradually decreased (Fig. 1A; $P < 0.05$), and apoptotic cell death was significantly increased (Fig. 1B-F; $P < 0.05$). On the basis of these experiments on cell viability and the Annexin V/PI staining assay, 0.5 mM H_2O_2 was selected as a model dose for the following experiments.

Baicalin inhibits apoptotic cell death of endplate chondrocytes induced by H₂O₂. As shown in Fig. 2A, cell viability was highest when endplate chondrocytes were pretreated with baicalin for 1 h ($P < 0.05$). Subsequently, the cell viability gradually decreased when the endplate chondrocytes were treated with baicalin at or following the start of incubation with H_2O_2 . Without the coexistence of H_2O_2 , no significant differences in cell viability were observed among the group pretreated with 50 μ mol/l baicalin, the group pretreated with 100 μ mol/l baicalin and the control group ($P > 0.05$). However, when the pretreatment concentration of Baicalin reached 200 μ mol/l, the cell viability of endplate chondrocytes was significantly lower, compared with that in the control ($P < 0.05$, Fig. 2B). Furthermore, in the cells incubated with 0.5 mM H_2O_2 for 4 h with baicalin pretreatment for 1 h, the viability of cells in the group pretreated with 100 μ mol/l baicalin was significantly higher, compared with that in the other two groups ($P < 0.05$), which was considered the optimal dose for application in the subsequent experiments (Fig. 2B). As shown in Fig. 2C, compared with the group treated with H_2O_2 alone, cell death was significantly decreased in the group treated with H_2O_2 +baicalin at 4, 8, 12 and 24 h ($P < 0.05$), which suggested that baicalin may inhibit the cell apoptosis induced by H_2O_2 . The results of the Annexin V/PI staining assay (Fig. 2D-F) also showed that baicalin inhibited the cell apoptosis induced by H_2O_2 .

Baicalin inhibits the activation of the apoptotic cell death pathway induced by H₂O₂. Classic apoptotic cell death is induced through a pathway, which involves the cleavage of PARP and pro-caspase-3, and the activation of Bax. As shown in Fig. 3A and B, the protein expression of cleaved PARP, Bax and pro-caspase-3 proteins were determined following treatment of the endplate chondrocytes with 0.5 mM H_2O_2 for 4 h, which were significantly higher, compared with that in the normal control ($P < 0.05$). By contrast, baicalin pretreatment downregulated the protein expression levels of cleaved PARP, Bax and pro-caspase-3 ($P < 0.05$). The quantified results of western blot analysis are shown in Fig. 3B.

Baicalin decreases the oxidative activity in endplate chondrocytes induced by H₂O₂. As a biomarker of oxidative stress,

Figure 1. Effect of H₂O₂ on endplate chondrocytes. (A) Cell viability of endplate chondrocytes treated with H₂O₂ at different concentrations (0.1, 0.5 and 1.0 mM). Apoptosis of endplate chondrocytes induced by H₂O₂ were examined at different concentrations in the (B) control; (C) 0.1 mM; (D) 0.5 mM; (E) 1.0 mM groups. (F) Apoptotic rates of endplate chondrocytes treated with H₂O₂ at different concentrations (0.1, 0.5 and 1.0 mM). Data are presented as the mean $\pm$ standard error of the mean. * P <0.05 H₂O₂ vs. control groups. H₂O₂, hydrogen peroxide.

the levels of MDA in endplate chondrocytes were increased by H₂O₂ treatment at 6, 12 and 24 h (P <0.05), and this was reversed by baicalin at all time-points (P <0.05; Fig. 4A). As an enzyme, which regulates oxidative stress, the levels

Figure 2. Effects of baicalin on endplate chondrocytes. (A) Effects of baicalin treatment on H₂O₂-induced loss of endplate chondrocytes at different time-points (-1, 0, 1 and 2 h). (B) Effects of 1 h baicalin treatment on endplate chondrocytes at various concentrations (50, 100 and 200 μ M/l) with or without H₂O₂. (C) Effects of 1 h baicalin treatment on endplate chondrocytes at different time-points (4, 8, 12 and 24 h) with H₂O₂. Data are presented as the mean $\pm$ standard error of the mean. * P <0.05 H₂O₂ vs. control group, baicalin+H₂O₂ vs. control groups. Inhibitory effects of baicalin on the apoptosis of endplate chondrocytes induced by H₂O₂ were determined in the (D) control, (E) H₂O₂ (F) baicalin+H₂O₂ groups. The results of the Annexin V/PI staining assay showed that baicalin inhibited the cell apoptosis induced by H₂O₂. H₂O₂, hydrogen peroxide.

of SOD were significantly decreased by exposure to H₂O₂ (P <0.05), however, pretreatment with baicalin for 1 h effectively elevated levels of SOD in endplate chondrocytes at the various time-points (P <0.05; Fig. 4B). The effects of baicalin on oxidative activity were further verified by assessing levels of NO, a free radical signaling mediator. The levels of NO were significantly decreased in the H₂O₂-induced endplate chondrocytes (P <0.05), and this decrease of NO was reversed by baicalin at 6, 12 and 24 h (P <0.05; Fig. 4C). The mRNA

Figure 3. Effect of baicalin on the apoptotic cell death pathway. (A) Western blot analysis of the effects of 1 h baicalin pretreatment on protein levels of apoptosis-related molecules with or without H₂O₂ in endplate chondrocytes. Compared with the control group, H₂O₂ downregulated the protein expression levels of cleaved PARP, Bax and pro-caspase-3, whereas baicalin reversed and upregulated the expression of apoptosis-related proteins. (B) Quantification of the results of western blot analysis. [#], ^sP<0.05 H₂O₂, baicalin+H₂O₂ vs. the control groups. H₂O₂, hydrogen peroxide; PARP, poly (ADP-ribose) polymerase; Bax, B-cell lymphoma-2-associated X protein.

levels of eNOS, an enzyme, which functions in catalyzing the release of NO, were also investigated. The mRNA expression of eNOS was downregulated by H₂O₂, and this was reversed by baicalin (P<0.05; Fig. 5).

Discussion

In the present study, a model of 0.5 mM H₂O₂-induced oxidative stress was successfully induced in endplate chondrocytes, which was characterized by reduced cell viability, increased intracellular ROS and reduced intracellular antioxidant activity. These pathophysiological processes led to apoptosis by activation of the intrinsic apoptotic pathway. As a flavonoid glycoside extracted from a type of traditional Chinese medicine, *Scutellaria baicalensis*, baicalin has been reported to significantly attenuate oxidative injury, and partially prevent apoptosis during the oxidative stress reaction (8,10-14,18). For example, Lin *et al* (8,11) found that baicalin ameliorated H₂O₂-induced cytotoxicity in a human renal proximal tubular epithelial cell line (HK-2 cells), and showed effects against renal ischemia-reperfusion injury through the inhibition of inflammation and apoptosis. Therefore, the present study investigated the potential protective effects of baicalin treatment on H₂O₂-induced oxidative stress in endplate chondrocytes *in vitro*.

Initially, different time-points of induction with baicalin incubation under H₂O₂ stimulation were investigated, which showed that 1 h pretreatment with baicalin exhibited optimal protective effects. Secondly, it was observed that the potential protective effects of baicalin pretreatment against

Figure 4. Effects of 1 h baicalin pretreatment on oxidative activity in endplate chondrocytes induced by H₂O₂ at the different time-points (6, 12 and 24 h). Baicalin suppressed oxidative activity in endplate chondrocytes induced by H₂O₂, (A) including effectively reducing levels of MDA, (B) increasing levels of SOD, and (C) elevating NO activity. Data are presented as the mean ± standard error of the mean. ^{*}, [#]P<0.05 H₂O₂, baicalin+H₂O₂ vs. the control groups. MDA, malondialdehyde; SOD, superoxide dismutase; NO, nitric oxide.

Figure 5. Effects of 1 h baicalin pretreatment on mRNA expression levels of eNOS in endplate chondrocytes induced by H₂O₂ for 4 h. mRNA levels of eNOS were downregulated by H₂O₂, which was reversed by baicalin. Data are presented as the mean ± standard error of the mean. ^{*}, [#]P<0.05 H₂O₂, baicalin+H₂O₂ vs. the control groups. H₂O₂, hydrogen peroxide; eNOS, endothelial nitric oxide synthase.

oxidative stress-related injury in endplate chondrocytes were dose-dependent. However, baicalin significantly decreased cell viability when its concentration reached 200 $\mu\text{mol/l}$. Therefore, to achieve optimal protective effects, 100 $\mu\text{mol/l}$ baicalin was selected as the optimal concentration for inhibiting oxidative stress in endplate chondrocytes induced by H_2O_2 .

In the present study, the protective effects of baicalin on oxidative stress of endplate chondrocytes induced by H_2O_2 were confirmed through the analysis of cell viability. The cell viability of endplate chondrocytes stimulated by H_2O_2 was significantly lower, compared with that in the normal control, as detected by the CCK-8 assay. Notably, a significant increase in cell viability was observed following baicalin pretreatment for 1 h at the various time-points. Baicalin was found to inhibit the activation of the apoptotic cell death pathway triggered by H_2O_2 . Classic apoptotic cell death is induced through a pathway involving the cleavage of PARP and pro-caspase-3, and the activation of Bax (19-21). The abnormally high expression levels of cleaved PARP, Bax and pro-caspase-3 induced by H_2O_2 were significantly reversed by pretreatment with baicalin. In addition, baicalin suppressed oxidative activity in the endplate chondrocytes induced by H_2O_2 , via effectively reducing levels of MDA, increasing levels of SOD, and elevating NO activities (5,9,22). eNOS, an enzyme which activates the expression of NO (23-25), was also shown to be increased following pretreatment with baicalin, which suggested that baicalin also inhibited the apoptosis through upregulating the expression of eNOS.

However, the mechanism of baicalin-mediated protection against oxidative stress remains to be fully elucidated. Chen *et al* (18) reported that baicalin inhibited oxidative-stress-induced apoptosis via modulating the activation of extracellular signal-regulated kinases and inducing the gene expression of heme oxygenase-1 in rat C6 glioma cells. Lin *et al* (8) reported that the effect of baicalin was through the inhibition of endoplasmic reticulum stress and the activation of nuclear factor erythroid 2-related factor 2 signaling, which were important during baicalin-mediated protection. The targets of baicalin also remain to be fully elucidated. A previous study indicated that baicalin activates AMP-activated protein kinase (AMPK) through the Ca^{2+} /calmodulin-dependent protein kinase kinase β (CaMKK β)-dependent pathway in HeLa and A549 cells (26). AMPK is important in cell physiology and also affects the cell response to oxidative stress. However, at present, no direct association between the renal protective effects of baicalin and activation of the Ca^{2+} /CaMKK β -AMPK pathway has been reported. Peroxisome proliferator-activated receptor- γ (PPAR γ) has also been suggested as a target of baicalin (27). Baicalin was found to activate PPAR γ and suppress downstream nuclear factor- κB -mediated inflammation in aged rat kidneys. In addition, studies have suggested other potential targets, including the proteasome (28), macrophages (29) and notch signaling (30). These possibilities were not examined in the present study, and warrant further investigation.

In conclusion, the results of the present study suggested that baicalin pretreatment protected endplate chondrocytes from the oxidative stress-related damage induced by H_2O_2 . The role of baicalin was predominantly based on inhibiting the production of ROS, increasing intracellular antioxidants

and attenuating apoptosis. The results of the present study require verification in future *in vivo* investigations, however, the results provide further insight into the potential benefits of baicalin for patients with oxidative stress-related diseases, including OA.

Acknowledgements

This study was supported by the National Science Foundation for Young Scholars of China (grant no. 81601675 to Dr Yutao Pan).

References

- Henrotin YE, Bruckner P and Pujol JP: The role of reactive oxygen species in homeostasis and degradation of cartilage. *Osteoarthritis Cartilage* 11: 747-755, 2003.
- Henrotin Y, Kurz B and Aigner T: Oxygen and reactive oxygen species in cartilage degradation: Friends or foes? *Osteoarthritis Cartilage* 13: 643-654, 2005.
- Rocha M, Apostolova N, Hernandez-Mijares A, Herance R and Victor VM: Oxidative stress and endothelial dysfunction in cardiovascular disease: Mitochondria-targeted therapeutics. *Curr Med Chem* 17: 3827-3841, 2010.
- Ying W and Xiong ZG: Oxidative stress and NAD⁺ in ischemic brain injury: Current advances and future perspectives. *Curr Med Chem* 17: 2152-2158, 2010.
- Mao CY, Lu HB, Kong N, Li JY, Liu M, Yang CY and Yang P: Levocarnitine protects H9c2 rat cardiomyocytes from H_2O_2 -induced mitochondrial dysfunction and apoptosis. *Int J Med Sci* 11: 1107-1115, 2014.
- Nogueira-Pedro A, Cesário TA, Dias CC, Origassa CS, Eça LP, Paredes-Gamero EJ and Ferreira AT: Hydrogen peroxide (H_2O_2) induces leukemic but not normal hematopoietic cell death in a dose-dependent manner. *Cancer Cell Int* 13: 123, 2013.
- Pomari E, Stefanon B and Colitti M: Effect of plant extracts on H_2O_2 -induced inflammatory gene expression in macrophages. *J Inflamm Res* 7: 103-112, 2014.
- Lin M, Li L, Zhang Y, Zheng L, Xu M, Rong R and Zhu T: Baicalin ameliorates H_2O_2 induced cytotoxicity in HK-2 cells through the inhibition of ER stress and the activation of Nrf2 signaling. *Int J Mol Sci* 15: 12507-12522, 2014.
- Chen S, Tang Y, Qian Y, Chen R, Zhang L, Wo L and Chai H: Allicin prevents H_2O_2 -induced apoptosis of HUVECs by inhibiting an oxidative stress pathway. *BMC Complement Altern Med* 14: 321, 2014.
- Cao Y, Mao X, Sun C, Zheng P, Gao J, Wang X, Min D, Sun H, Xie N and Cai J: Baicalin attenuates global cerebral ischemia/reperfusion injury in gerbils via anti-oxidative and anti-apoptotic pathways. *Brain Res Bull* 85: 396-402, 2011.
- Lin M, Li L, Li L, Pokhrel G, Qi G, Rong R and Zhu T: The protective effect of baicalin against renal ischemia-reperfusion injury through inhibition of inflammation and apoptosis. *BMC Complement Altern Med* 14: 19, 2014.
- Tang YJ, Zhou FW, Luo ZQ, Li XZ, Yan HM, Wang MJ, Huang FR and Yue SJ: Multiple therapeutic effects of adjunctive baicalin therapy in experimental bacterial meningitis. *Inflammation* 33: 180-188, 2010.
- Zhu J, Wang J, Sheng Y, Zou Y, Bo L, Wang F, Lou J, Fan X, Bao R, Wu Y, *et al*: Baicalin improves survival in a murine model of polymicrobial sepsis via suppressing inflammatory response and lymphocyte apoptosis. *PLoS One* 7: e35523, 2012.
- Xiping Z, Guanghua F, Jinxian H, Weihong W, Rujun X, Wei Z, Jing Y, Qijun Y, Meijuan Y, Qing W and Lini F: Baicalin protects thymus of rats with severe acute pancreatitis. *Inflammation* 33: 157-165, 2010.
- Wang YJ, Shi Q, Sun P, Zhou Q, Darowish M, Li TF, Dong YF, Lu WW and Leong JC: Insulin-like growth factor-1 treatment prevents anti-Fas antibody-induced apoptosis in endplate chondrocytes. *Spine (Phila Pa 1976)* 31: 736-741, 2006.
- Yuan FL, Wang HR, Zhao MD, Yuan W, Cao L, Duan PG, Jiang YQ, Li XL and Dong J: Ovarian cancer G protein-coupled receptor 1 is involved in acid-induced apoptosis of endplate chondrocytes in intervertebral discs. *J Bone Miner Res* 29: 67-77, 2014.

17. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
18. Chen YC, Chow JM, Lin CW, Wu CY and Shen SC: Baicalein inhibition of oxidative-stress-induced apoptosis via modulation of ERKs activation and induction of HO-1 gene expression in rat glioma cells C6. *Toxicol Appl Pharmacol* 216: 263-273, 2006.
19. Boulares AH, Yakovlev AG, Ivanova V, Stoica BA, Wang G, Iyer S and Smulson M: Role of poly(ADP-ribose) polymerase (PARP) cleavage in apoptosis. Caspase 3-resistant PARP mutant increases rates of apoptosis in transfected cells. *J Biol Chem* 274: 22932-22940, 1999.
20. Ruemmele FM, Dionne S, Qureshi I, Sarma DS, Levy E and Seidman EG: Butyrate mediates Caco-2 cell apoptosis via up-regulation of pro-apoptotic BAK and inducing caspase-3 mediated cleavage of poly-(ADP-ribose) polymerase (PARP). *Cell Death Differ* 6: 729-735, 1999.
21. Hoetelmans R, van Slooten HJ, Keijzer R, Erkeland S, van de Velde CJ and Dierendonck JH: Bcl-2 and Bax proteins are present in interphase nuclei of mammalian cells. *Cell Death Differ* 7: 384-392, 2000.
22. Liao JK, Shin WS, Lee WY and Clark SL: Oxidized low-density lipoprotein decreases the expression of endothelial nitric oxide synthase. *J Biol Chem* 270: 319-324, 1995.
23. Knowles RG and Moncada S: Nitric oxide synthases in mammals. *Biochem J* 298: 249-258, 1994.
24. Chen CA, Wang TY, Varadharaj S, Reyes LA, Hemann C, Talukder MA, Chen YR, Druhan LJ and Zweier JL: S-glutathionylation uncouples eNOS and regulates its cellular and vascular function. *Nature* 468: 1115-1118, 2010.
25. Wu JR, Hsu JH, Dai ZK, Wu BN, Chen JJ, Liou SF and Yeh JL: Activation of endothelial NO synthase by a xanthine derivative ameliorates hypoxia-induced apoptosis in endothelial progenitor cells. *J Pharm Pharmacol* 68: 810-818, 2016.
26. Ma Y, Yang F, Wang Y, Du Z, Liu D, Guo H, Shen J and Peng H: CaMKK β is involved in AMP-activated protein kinase activation by baicalin in LKB1 deficient cell lines. *PLoS One* 7: e47900, 2012.
27. Lim HA, Lee EK, Kim JM, Park MH, Kim DH, Choi YJ, Ha YM, Yoon JH, Choi JS, Yu BP and Chung HY: PPAR γ activation by baicalin suppresses NF- κ B-mediated inflammation in aged rat kidney. *Biogerontology* 13: 133-145, 2012.
28. Wu YX, Sato E, Kimura W and Miura N: Baicalin and scutellarin are proteasome inhibitors that specifically target chymotrypsin-like catalytic activity. *Phytother Res* 27: 1362-1367, 2013.
29. Liu LL, Gong LK, Wang H, Xiao Y, Wu XF, Zhang YH, Xue X, Qi XM and Ren J: Baicalin inhibits macrophage activation by lipopolysaccharide and protects mice from endotoxin shock. *Biochem Pharmacol* 75: 914-922, 2008.
30. Wang AM, Ku HH, Liang YC, Chen YC, Hwu YM and Yeh TS: The autonomous notch signal pathway is activated by baicalin and baicalein but is suppressed by niclosamide in K562 cells. *J Cell Biochem* 106: 682-692, 2009.