

Roles of miR-4463 in H₂O₂-induced oxidative stress in human umbilical vein endothelial cells

XUEQIN WANG¹, XUEMEI HE², XIAN DENG^{1,3}, YANZHENG HE¹ and XIANGYU ZHOU¹

¹Department of Vascular and Thyroid Surgery; ²Medical Research Center, The Affiliated Hospital of Southwest Medical University, Jiangyang, Luzhou, Sichuan 646000; ³State Key Laboratory of Quality Research in Chinese Medicine, Faculty of Chinese Medicine, Macau University of Science and Technology, Taipa, Macau SAR 519020, P.R. China

Received July 23, 2016; Accepted April 28, 2017

DOI: 10.3892/mmr.2017.7001

Abstract. Oxidative stress is implicated in the pathophysiology of vascular diseases, including atherosclerosis, aneurysm and arteriovenous fistula. A previous study from our lab suggested that microRNA (miR)-4463 may be involved in the pathogenesis of vascular disease; however, the roles of oxidative stress in the molecular mechanisms underlying the actions of miR-4463 in vascular disease have yet to be elucidated. The aim of the present study was to investigate the role of miR-4463 in hydrogen peroxide (H₂O₂)-induced oxidative stress in human umbilical vein endothelial cells (HUVECs). Reverse transcription-quantitative polymerase chain reaction was used to assess the expression levels of miR-4463 in HUVECs treated with various concentrations of H₂O₂. Flow cytometry was used to evaluate the percentage of apoptotic cells, and the protein expression levels of the apoptotic markers cleaved (C)-caspase3, poly (adenosine diphosphate-ribose) polymerase 1 (PARP1), B cell lymphoma-2 (Bcl-2), Bcl-2-associated X protein (Bax) and X-linked inhibitor of apoptosis protein (XIAP) were determined using western blot analysis. The results demonstrated that the apoptotic rate of HUVECs was increased following treatment with H₂O₂ in a concentration-dependent manner, and the expression of miR-4463 was also upregulated in a dose-dependent manner. Following transfection with miR-4463 mimics, the levels of malondialdehyde and reactive oxygen species were increased in HUVECs, with a corresponding increase in the apoptotic rate. Furthermore, western blot analysis revealed that the protein expression levels of C-caspase3, PARP1 and Bax were upregulated, whereas the levels of Bcl-2 and XIAP were

downregulated. In conclusion, the present findings suggested that the upregulation of miR-4463 may enhance H₂O₂-induced oxidative stress and promote apoptosis in HUVECs *in vitro*.

Introduction

Oxidative stress is an imbalance between oxidative and anti-oxidative processes *in vivo* and is involved in the pathogenesis of numerous diseases. Vascular disease is an inflammatory process, and oxidative stress has been implicated in vascular pathology (1,2). A previous study demonstrated that treatment with fucoidan, which is an aggregate name for algal fucose-enriched sulfated polysaccharides extracted from the extracellular matrix of seaweeds, may increase oxidative stress production by activating endothelial nitric oxide synthase and promoting Akt phosphorylation in myocardial disease and stroke (3). A recent study suggested that oxidative stress may be involved in high-fat diet-induced aortic remodeling, which may be exacerbated by Zn²⁺ deficiency, as the anti-oxidative molecule resveratrol was demonstrated to prevent high-fat diet-induced vascular inflammation, oxidative stress and pathological remodeling (4). Therefore, investigating the implications of oxidative stress in the pathogenesis of vascular diseases and exploring the underlying molecular mechanisms are imperative for the development of effective preventive and therapeutic strategies aimed at patients with vascular diseases.

Signaling pathways related to oxidative stress are associated with endothelial cells that line the inner walls of blood vessels, which are continuously exposed to various stresses owing to the mechanical force exerted by blood flow and pressure, and are sensitive to oxidative damage (5,6). A recent study demonstrated that the natural antioxidant allicin protected human umbilical vein endothelial cells (HUVECs) from oxidized low-density lipoprotein-induced injury by preventing apoptosis through the inhibition of caspase-3 and nicotinamide-adenine dinucleotide phosphate oxidase-mediated proapoptotic signaling (7). Another recent study reported that resveratrol (polyphenolic compounds) treatment ameliorated high glucose-induced HUVEC injury through the activation of 5' adenosine monophosphate-activated protein kinase α , leading to an increase in reductive reactions and a corresponding decrease in oxidative stress (8). These studies

Correspondence to: Dr Xiangyu Zhou, Department of Vascular and Thyroid Surgery, The Affiliated Hospital of Southwest Medical University, 25 Taiping Street, Jiangyang, Luzhou, Sichuan 646000, P.R. China
E-mail: xiangyuzhou971@vip.126.com

Key words: microRNA-4463, hydrogen peroxide, human umbilical vein endothelial cells, oxidative stress

suggested that oxidative stress may serve crucial roles during the development of vascular endothelial cell injury. Therefore, the present study aimed to investigate the effects of oxidative stress in HUVECs *in vitro*.

MicroRNAs (miRNAs) are small (~22 nucleotides long) non-coding RNA molecules, which modulate the stability and the translational efficiency of target mRNA transcripts (9). miRNAs are involved in the post-transcriptional regulation of gene expression and are implicated in numerous biological processes, including proliferation, differentiation, senescence and death (10). Previous studies have suggested that miRNAs may be involved in the regulation of oxidative stress during vascular disease pathogenesis, serving important roles in oxidative stress-induced endothelial dysfunction and mitochondrial metabolism dysregulation (1,11). Several miRNAs have been implicated in endothelial development, including miRNA (miR)-20a, miR-23b-3p, miR-150, miR-195 and miR-200b (11,12). Previous studies have also revealed crucial roles for miRNAs in endothelial physiology and pathology, thus suggesting that miRNAs may have potential as novel therapeutic targets for the treatment of patients with vascular diseases (13,14).

miR-4463 has garnered attention as a prognostic and diagnostic biomarker in patients with polycystic ovary syndrome (15). In addition, our previous studies have demonstrated the aberrant expression of miR-4463 in vascular diseases, including arteriosclerosis obliterans (16) and carotid artery stenosis (17), which suggested that miR-4463 may also have potential as a biomarker for the early diagnosis of vascular diseases. However, the roles of miR-4463 in the regulation of oxidative stress in endothelial cells, and the molecular mechanisms underlying its actions, have yet to be elucidated. Therefore, the present study aimed to investigate the putative relationship between miR-4463 and oxidative stress, and explore the molecular mechanisms that may be involved in HUVECs.

Materials and methods

Cell culture and oxidative stress model. The HUVEC cell line was purchased from ScienCell Research Laboratories, Inc. (Carlsbad, CA, USA) and cultured in low-glucose Dulbecco's modified Eagle's medium (DMEM; Hyclone, GE Healthcare Life Sciences, Logan, UT, USA), supplemented with 10% heat-inactivated fetal bovine serum (FBS; Gibco, Thermo Fisher Scientific, Inc., Waltham, MA, USA) and maintained in a humidified 5% CO₂ atmosphere. Cells between passages 4 and 6, in the logarithmic phase of growth, were used in the subsequent experiments. The morphology of the cells was observed using a phase-contrast microscope (cellSens standard software version 1.6, DP72; Olympus Corporation, Tokyo, Japan).

Hydrogen peroxide (H₂O₂) is commonly used in models of oxidative stress-induced apoptosis (18). In the present study, HUVECs were treated with various concentrations of H₂O₂ (200, 500, 700 and 1,000 μ mol/l; Table I) using 0.3% H₂O₂ stock solution, or 0 μ l H₂O₂ (low-glucose DMEM alone containing 10% FBS) for 16 h to induce oxidative stress, as previously described (19-21).

Transfection. The intracellular level of miR-4463 was upregulated or downregulated by the transfection of

Table I. Various concentrations of H₂O₂.

H ₂ O ₂ concentration	0.3% H ₂ O ₂	Low-glucose DMEM with 10% FBS
0 μ mol/l	0.0 μ l	500.0 μ l
200 μ mol/l	11.4 μ l	488.6 μ l
500 μ mol/l	28.5 μ l	471.5 μ l
700 μ mol/l	39.9 μ l	460.1 μ l
1,000 μ mol/l	57.0 μ l	443.0 μ l

H₂O₂, hydrogen peroxide; DMEM, Dulbecco's modified Eagle's medium; FBS, fetal bovine serum.

miRNA mimics and inhibitor respectively, according to the manufacturer's protocol (Guangzhou RiboBio Co., Ltd., Guangzhou, China). The miRNA mimics synthesized by the chemical method induces a high level of expression of mature miRNA in the cell, and it enhances the regulation of endogenous miRNA and gains certain functions in cells. miRNA inhibitors inhibit the effects of mature miRNA, and weaken the role of endogenous gene regulation and cell function. When the cell density reached 50-70%, the transfection efficiency was optimal (Table II). Transfection was performed using 100 μ mol/l miR-4463 mimic or inhibitor for 15 min at room temperature, and the efficiency of transfection was determined by RT-qPCR. Finally, 100 nmol/l NC, miR-4463 mimics and miR-4463 inhibitor was chosen for the next step after 3 days transfection. NC, miR-4463 mimics and miR-4463 inhibitor sequences are listed on the Guangzhou RiboBio Co., Ltd. Website: <http://ribobio.bioon.com.cn>.

RNA exaction and reverse transcription-quantitative polymerase chain reaction (RT-qPCR). Following treatment with H₂O₂, total RNA was extracted from HUVECs (2-3x10⁵ cells) using TRIzol reagent (Invitrogen; Thermo Fisher Scientific, Inc.). Total RNA (500 ng) was reverse transcribed into cDNA using an miScript II RT kit (catalog no. 218160 and 21816; Qiagen GmbH, Hilden, Germany), according to manufacturer's protocol. The specific reaction components of reverse-transcription are mentioned in Table III. The mix was incubated for 60 min at 37°C, incubated for 5 min at 95°C and stored at -20°C. Acquired cDNA was used for RT-qPCR according to the manufacturer's protocol using the miScript SYBR-Green PCR kit (catalog no. 208054; Qiagen GmbH). Mature has-miR-4463 primer (catalog no. MIMAT0018987) and U6 primer (catalog no. MS00044996) were purchased from Qiagen GmbH. The specific reaction components of RT-qPCR are mentioned in Table IV, and the cycling conditions are mentioned in Table V. qPCR was performed using the StepOnePlus version 2.2.3 real-time PCR system (Applied Biosystems; Thermo Fisher Scientific, Inc.). Relative expression levels were quantified using the 2^{- $\Delta\Delta$ C_q} method (22) and normalized to the expression of U6; Ct values >35 were excluded. Experiments were performed in triplicate.

Apoptosis assay. Apoptosis and necrosis were assessed using the fluorescein isothiocyanate (FITC) Annexin V and

Table II. Transfection reagent components.

Final concentration	Final volume	Culture medium	1x riboFECT™ CP buffer	NC, miR-4463 mimics or miR-4463 inhibitor	riboFECT™ CP reagent
100 nmol/l NC	500 μ l	464.5 μ l	30 μ l	2.5 μ l NC	3 μ l
100 nmol/l miR-4463 mimic	500 μ l	464.5 μ l	30 μ l	2.5 μ l miR-4463 mimic	3 μ l
100 nmol/l miR-4463 inhibitor	500 μ l	464.5 μ l	30 μ l	2.5 μ l miR-4463 inhibitor	3 μ l

NC, negative control; miR, microRNA.

Table III. Reverse-transcription reaction components.

Component	Volume/reaction
5X miScript HiSpec buffer	4 μ l
10x nucleics mix	2 μ l
RNase-free water	Variable
miScript reverse transcriptase mix	2 μ l
Template RNA	Variable (500 ng)
Total volume	20 μ l

propidium iodide (PI) detection kit (BD Pharmingen; BD Biosciences, Franklin Lakes, NJ, USA), according to the manufacturer's protocol. Following treatment with H₂O₂, HUVECs (5x10⁵ cells) were washed with 10X Annexin V binding buffer, and stained for 10 min with 5 μ l FITC Annexin V and 5 μ l PI at room temperature in the dark. Stained cells were analyzed by flow cytometry using a BD FACSVerse and BD FACSuite™ software version 1.0.3 (BD Biosciences), and the experiment was repeated three times.

Oxidative stress assays. Reactive oxygen species (ROS) production and lipid peroxidation were measured in order to assess oxidative stress in HUVECs, using ROS and malondialdehyde (MDA) detection kits (Beyotime Institute of Biotechnology, Haimen, China). Experiments were repeated >3 times. To detect ROS production, H₂O₂-treated HUVECs (1x10⁵ cells) were incubated with the fluorescent probe 2',7'-dichlorodihydrofluorescein diacetate (chloromethyl derivative) at 37°C for 30 min, as previously described (23-25). Subsequently, samples were digested by trypsin-EDTA solution (Beyotime Institute of Biotechnology), centrifuged (1,000 x g for 5 min at room temperature) and transferred to black 96-well plates. The fluorescence was measured using an excitation wavelength of 488 nm and an emission wavelength of 525 nm using a microplate reader. MDA is a natural product of lipid oxidation in organisms. The level of lipid oxidation may be measured by detecting the level of MDA, and therefore the determination of MDA is widely used as an indicator of lipid oxidation. To assess MDA contents in HUVECs (1x10⁵ cells), cells were lysed and then centrifuged (1,600 x g for 10 min at 4°C). A total of 100 μ l supernatant and 200 μ l MDA detection working fluid (150 μ l TBA dilute solution, 50 μ l TBA storage solution and 3 μ l antioxidant) were mixed, heated at 100°C for 15 min

Table IV. Reaction mix for miScript SYBR Green polymerase chain reaction.

Component	Volume/reaction (96-well)
2x QuantiTect SYBR-Green	12.5 μ l
PCR Master Mix	
10x miScript universal primer	2.5 μ l
10x miScript primer assay	2.5 μ l
RNase-free water	6.5 μ l
Diluted cDNA	1.0 μ l
Total volume	25.0 μ l



and cooled to room temperature. The reaction mixture was centrifuged (1,000 x g for 10 min at room temperature), and subsequently transferred to 96-well plates and the absorbance was measured at 532 nm using a microplate reader, as previously described (26,27).

Cell proliferation assay. Cellular proliferation was evaluated using a Cell Counting Kit-8 (CCK-8; Beyotime Institute of Biotechnology), as previously described (28,29). Following treatment with H₂O₂ for 16 h, HUVECs (1x10⁴ cells/well) were seeded in 96-well plates (5 wells/experimental group) in 100 μ l serum-free DMEM (Hyclone; GE Healthcare Life Sciences) and 10 μ l CCK-8 solution were added to each well and cells were incubated at 37°C for 35 min. The absorbance of each sample was measured at 450 nm using a microplate reader as previously described (30). The percentage of living cells was calculated as a ratio of the optical density of treated cells over untreated control cells.

Western blot analysis. Following treatment with H₂O₂, total protein was extracted from HUVECs (2x10⁶ cells) using Radioimmunoprecipitation Assay Lysis buffer (Beyotime Institute of Biotechnology) as previously described (2,31). Equal amounts (20 μ g) of protein in each sample (extracted using a bicinchoninic acid assay) were separated by 12% SDS-PAGE and transferred onto polyvinylidene difluoride membranes. Then, membranes were blocked for 2 h in 0.5% nonfat milk at room temperature, and washed three times for 5 mins with PBS containing 0.1% Tween-20. Next, membranes were incubated with the following primary antibodies at 4°C overnight: Anti-poly (adenosine diphosphate-ribose) polymerase

Table V. Cycling conditions of miScript SYBR Green polymerase chain reaction.

Cycling conditions			
Step	Time	Temperature	Additional comments
Initial activation step	15 min	95°C	HotStarTaq [®] DNA polymerase is activated by this heating step.
3-step cycling:			
Denaturation	15 sec	94°C	
Annealing	30 sec	55°C	
Extension	30 sec	70°C	Perform fluorescence data collection.
Cycle number	40 cycles		Cycle number depends on the amount of template cDNA and abundance of the target.

Figure 1. Morphological alterations in HUVECs. Untreated control group (0 $\mu\text{mol/l}$), 200, 500, 700 and 1,000 $\mu\text{mol/l}$ H₂O₂ treatment for 16 h. The morphology of the cells was observed by phase-contrast microscope, using x200 magnification 200.Figure 2. miR-4463 expression levels in HUVECs following treatment with H₂O₂. HUVECs were treated with 200, 500, 700 and 1 000 $\mu\text{mol/l}$ H₂O₂ for 16 h and miR-4463 expression was assessed using reverse transcription-quantitative polymerase chain reaction. Control cells were not exposed to H₂O₂. Data are expressed as the mean \pm standard deviation (n=3). *P<0.05 vs. Control. H₂O₂, hydrogen peroxide; HUVEC, human umbilical vein endothelial cell; miR, microRNA.

and anti-B cell lymphoma-2 (Bcl-2)-associated X protein (Bax; 1:1,000; catalog no. 14796) were purchased from Cell Signaling Technology, Inc. (Danvers, MA, USA); anti-active (cleaved)-caspase-3 (C-caspase-3; 1:500; catalog no. ab32042) was purchased from Abcam (Cambridge, UK); and anti-Bcl-2 (1:1,000; catalog no. AB112-1) and anti-GAPDH (1:2,000; catalog no. AF0006) were purchased from Beyotime Institute of Biotechnology. GAPDH served as the loading control (18,19). Subsequently, membranes were incubated with the following horseradish peroxidase-conjugated secondary antibodies for 1 h at 37°C: Anti-rabbit immunoglobulin (Ig)G (1:2,000; catalog no. A0208) and anti-mouse IgG (1:3,000; catalog no. A0216) purchased from Beyotime Institute of Biotechnology. Protein bands were visualized by enhanced chemiluminescence detection reagent (Bio-Rad Laboratories, Inc., Hercules, CA, USA), and blots were semi-quantified by densitometric analysis using Quantity One version 4.6.2 software (Bio-Rad Laboratories, Inc.). Experiments were performed >3 times.

1 (PARP1; 1:1,000; catalog no. 9532), anti-X-linked inhibitor of apoptosis protein (XIAP; 1:1,000; catalog no. 14334),

Statistical analysis. Statistical analysis was performed using SPSS software version 19.0 (IBM Corp., Armonk, NY, USA).




Figure 3. H_2O_2 induces apoptosis in HUVECs. HUVECs were treated with 200, 500 and 700 $\mu\text{mol/l}$ H_2O_2 for 16 h. Control cells were not exposed to H_2O_2 . (A) Cellular apoptosis was assessed using flow cytometry following staining with FITC Annexin V and propidium iodide. (B) Treatment with H_2O_2 significantly increased the apoptotic rate of HUVECs. Data are expressed as the mean \pm standard deviation ($n=3$). * $P<0.05$ vs. Control. FITC, fluorescein isothiocyanate; H_2O_2 , hydrogen peroxide; HUVEC, human umbilical vein endothelial cell.

Data are expressed as the mean \pm standard deviation. The statistical significance of the differences between groups was assessed using one-way analysis of variance followed by a post hoc Dunnett's two-tailed test for multiple comparisons. $P<0.05$ was considered to indicate a statistically significant difference.

Results

Alterations in HUVEC morphology and miR-4463 expression following treatment with H_2O_2 . When examined under an inverted phase-contrast microscope, HUVECs in the untreated control group exhibited physiological morphology, characterized by spindle- or round-shaped cell architecture, cells that were adherent and formed paving stone-like cultures (32,33).

Conversely, the number of HUVECs in culture appeared to be markedly reduced following H_2O_2 treatment, with undefined boundaries and observable necrotic and disintegrative phenomena among the cells, which concurred with previously published data (34-36) (Fig. 1).

The expression levels of miR-4463 were assessed in HUVECs treated with various concentrations of H_2O_2 . The results demonstrated that miR-4463 expression levels were significantly upregulated in H_2O_2 -treated HUVECs in a dose-dependent manner compared with in untreated control cells (Fig. 2). miR-4463 expression appeared to be the highest following treatment with H_2O_2 at a concentration of 700 $\mu\text{mol/l}$. The high concentration of H_2O_2 (1,000 $\mu\text{mol/l}$) may promote cell death and inhibit the expression of the cells under oxidative stress.




Figure 4. H₂O₂ induces oxidative stress in HUVECs. HUVECs were treated with 200, 500 and 700 μmol/l H₂O₂ for 16 h; Control cells were not exposed to H₂O₂. (A) MDA levels and (B) ROS production were assessed in HUVECs using commercially available kits. (C) A Cell Counting Kit-8 assay was used to evaluate cell viability. Data are expressed as the mean ± standard deviation (n=3). *P<0.05 vs. Control. H₂O₂, hydrogen peroxide; HUVEC, human umbilical vein endothelial cell; MDA, malondialdehyde; ROS, reactive oxygen species.

H₂O₂ induces apoptosis in HUVECs. Flow cytometric analysis revealed that treatment with H₂O₂ was able to induced apoptosis in HUVECs (Fig. 3). Annexin V-positive cells and cellular uptake of PI were evaluated in order to distinguish between living, apoptotic and necrotic cells, as previously described (37). The proapoptotic effects of H₂O₂ treatment appeared to be concentration-dependent (Fig. 3B): No significant difference in the apoptotic rate was detected following treatment with 200 μmol/l H₂O₂; however, HUVEC apoptosis was significantly increased following treatment with 500 and 700 μmol/l H₂O₂ compared with control untreated cells (P<0.05). As H₂O₂ at a concentration of 700 μmol/l appeared to produce the greatest increase in apoptotic rate (43%) and

necrotic rate (8.25%), this concentration was selected for subsequent experiments.

H₂O₂ induces oxidative stress in HUVECs. To confirm the induction of oxidative stress following H₂O₂ treatment in HUVECs, oxidative stress products were detected and cellular viability was evaluated, using commercially available kits. MDA is an end product of lipid oxidation that is involved in the oxidative damage of the cellular membrane; therefore, MDA contents may be measured as an indication of the degree of lipid peroxidation and the extent of membrane damage (38,39). In HUVECs treated with 700 μmol/l H₂O₂, MDA levels were increased by 1.5-fold compared with the untreated control cells (P<0.05; Fig. 4A); whereas no significant differences were indicated in HUVECs treated with 200 or 500 μmol/l H₂O₂. In addition, intracellular ROS production was significantly enhanced in H₂O₂-treated cells, as indicated by the 3.75-fold increase in ROS levels following treatment with 700 μmol/l H₂O₂ compared with the control. Treatment with 500 μmol/l H₂O₂ increased ROS levels by ~2-fold compared with the control, and there was no significant increase after treatment with 200 μmol/l H₂O₂ (Fig. 4B). Notably, no significant difference in cellular proliferation was detected between 200 μmol/l H₂O₂-treated HUVECs and control cells (Fig. 4C). However, cell proliferation and viability was significantly reduced following treatment with 500 or 700 μmol/l H₂O₂ (Fig. 4C). These findings suggested that treatment with H₂O₂ may induce oxidative stress in HUVECs and suppress their viability in a concentration-dependent manner.

miR-4463 promotes H₂O₂-induced increases in oxidative stress. The apoptotic rate of HUVECs pretreated with miR-4463 mimics prior to H₂O₂ exposure was significantly increased compared with negative control-treated H₂O₂-exposed cells (Fig. 5), thus suggesting that miR-4463 may serve a role in promoting apoptosis following H₂O₂ treatment. Detecting NC-treated H₂O₂ as the control group eliminates any background effects of the transfection reagent. The roles of miR-4463 in H₂O₂-induced oxidative stress were also investigated, and the results revealed that intracellular ROS and MDA levels were significantly increased in H₂O₂-treated HUVECs following transfection with miR-4463 mimics; however, the miR-4463 inhibitor did not induce a significant difference. This suggested that miR-4463 overexpression enhanced H₂O₂-induced oxidative stress (Fig. 6A and B). However, the miR-4463 mimics did not have an exact effect on cell proliferation (Fig. 6C). These findings suggested that miR-4463 may enhance oxidative stress on HUVECs but have no apparent effect on the proliferation of HUVECs.

miR-4463 promotes proapoptotic signaling in H₂O₂-treated HUVECs. The protein expression levels of pro- and antiapoptotic markers, including C-caspase-3, Bax, Bcl-2, XIAP and PARP1, were detected in H₂O₂-treated HUVECs by western blot analysis (Fig. 7). The results demonstrated that following H₂O₂ treatment, the expression levels of the antiapoptotic proteins Bcl-2 and XIAP were downregulated, whereas the expression of the proapoptotic C-caspase3, PARP1 and Bax was enhanced. Notably, HUVECs pretreated with miR-4463




Figure 5. miR-4463 enhances H₂O₂-induced HUVEC apoptosis. HUVECs were pretreated with a miR-4463 mimic, a miR-4463 inhibitor or NC miRNA, and then exposed to 700 μmol/l H₂O₂ for 16 h. Control cells did not receive treatment. (A) Cellular apoptosis was assessed using flow cytometry following staining with FITC Annexin V and propidium iodide. (B) miR-4463 overexpression enhanced the proapoptotic effects of H₂O₂. Data are expressed as the mean ± standard deviation (n=3). *P<0.05 vs. 700+NC group. miR, microRNA; H₂O₂, hydrogen peroxide; HUVEC, human umbilical vein endothelial cell; NC, negative control; FITC, fluorescein isothiocyanate.

mimics prior to H₂O₂ exposure exhibited significantly increased C-caspase3, cleaved PARP and Bax protein expression levels, whereas the expression of Bcl-2 and XIAP was significantly suppressed. The present findings suggested that miR-4463 overexpression may increase the proapoptotic effects of H₂O₂ on HUVECs.

Discussion

miRNAs are able to combine with complementary sequences in the 3'-untranslated region of target mRNA transcripts, and thus regulate the transcription of target genes and ultimately inhibit the translation of specific proteins (9). miRNAs have been demonstrated previously to be involved in the regulation

of HUVEC proliferation, migration and apoptosis (12). In addition, miRNAs have been associated with processes involved in the development of oxidative stress (24). However, the roles of miRNAs in the pathophysiology of vascular diseases, as well as their clinical value, have yet to be elucidated. Our previous study reported that miR-4463 has a greater change in the plasma levels of vascular patients (16). Therefore, the present study hypothesized that miR-4463 may have an effect on the pathophysiological processes associated with vascular diseases.

H₂O₂ is one of the main ROS, which has been reported to decrease cellular proliferation and induce apoptosis, in a process involving several miRNAs (28). Exposure of endothelial cells to H₂O₂ has been reported to cause cell dysfunction




Figure 6. miR-4463 enhances H₂O₂-induced oxidative stress in HUVECs. HUVECs were pretreated with miR-4463 mimics, a miR-4463 inhibitor or a NC miRNA, and then exposed to 700 μmol/l H₂O₂ for 16 h; Control cells did not receive treatment. (A) ROS production and (B) MDA levels were assessed using commercially available kits. (C) A Cell Counting Kit-8 assay was used to evaluate cell viability. Data are expressed as the mean ± standard deviation (n=3). *P<0.05 vs. 700+NC group. H₂O₂, hydrogen peroxide; HUVEC, human umbilical vein endothelial cell; MDA, malondialdehyde; miR, microRNA; NC, negative control; ROS, reactive oxygen species.

and promote apoptosis, thus inducing endothelial inflammation. For example, Yu *et al* (40) previously demonstrated that miR-200c expression was upregulated following 100 μmol/l H₂O₂ treatment in BV-2 mouse microglial cells *in vitro*, whereas Zhang *et al* (38) reported that miR-92a overexpression in HUVECs enhanced capillary tube formation under conditions of oxidative stress. The present study investigated the expression of miR-4463 in H₂O₂-treated HUVECs, and RT-qPCR analysis revealed that miR-4463 expression was upregulated following treatment with H₂O₂ in a concentration-dependent manner. Notably, in HUVECs treated with 1,000 μmol/l H₂O₂, miR-4463 expression was comparable to baseline levels. Yu *et al* (40) also reported similar changes in miR-200c expression levels following spinal cord injury in murine BV-2 cells; however, the reason remains unclear and further studies are required to investigate whether high H₂O₂ concentration (1,000 μmol/l) may increase cell necrosis. Regardless of this discrepancy, the present results suggested that miR-4463 may serve a role in oxidative processes in HUVECs, and its expression may be regulated by H₂O₂ in a concentration-dependent manner.

Oxidative stress has been implicated in cellular apoptosis. In the present study, HUVEC apoptosis was investigated using

flow cytometry following cell exposure to H₂O₂. The present results demonstrated that H₂O₂ increased the apoptotic rate in HUVECs in a concentration-dependent manner. Once generated, ROS readily react with unsaturated fatty acids and cholesterol molecules present in cell membranes, thus leading to cellular apoptosis through mitochondrial pathways involving nuclear factor-κB, p53 and stress-activated protein kinase (41). In the present study, oxidative stress was investigated in HUVECs by assessing ROS production and MDA concentration; the results of which revealed that ROS and MDA levels were significantly increased following H₂O₂ treatment, whereas cell viability was significantly reduced, thus indicating that H₂O₂ was able to induced apoptosis in HUVECs, which is in accordance with previous studies (27,37).

To investigate the roles of miR-4463 in oxidative pathways and cellular apoptosis, HUVECs were transfected with miR-4463 mimics or inhibitors prior to H₂O₂ exposure. The present results demonstrated that the apoptotic rate and the levels of oxidation products were significantly increased in HUVECs overexpressing miR-4463. However, miR-4463 inhibition did not produce a significant effect. These results suggested that miR-4463 upregulation may enhance H₂O₂-induced HUVEC apoptosis; however, further studies are




Figure 7. miR-4463 promotes proapoptotic signaling in H_2O_2 -treated HUVECs. (A) HUVECs were pretreated with miR-4463 mimics, a miR-4463 inhibitor or a NC miRNA, and then exposed to 700 $\mu\text{mol/l}$ H_2O_2 for 16 h; Control cells did not receive treatment. (B) Western blot analysis was used to assess the protein expression levels of the proapoptotic factors PARP1, Bax and C-caspase3, and the antiapoptotic factors XIAP and Bcl-2 (n=3). *P<0.05 vs. 700+NC. Bax, Bcl-2-associated X protein; Bcl-2, B-cell lymphoma-2; C, cleaved; H_2O_2 , hydrogen peroxide; HUVEC, human umbilical vein endothelial cell; miR, microRNA; NC, negative control; PARP1, poly (adenosine diphosphate-ribose) polymerase I; XIAP, X-linked inhibitor of apoptosis protein.

required to confirm the exact roles of miR-4463 in HUVECs under oxidative stress and to determine the possible underlying molecular mechanisms that are involved.

Oxidative stress is a harmful mechanism for self-protection, therefore, excessive activation may enhance this effect. The results of the present study revealed that H_2O_2 treatment altered miR-4463 expression in HUVECs, and miR-4463 overexpression enhanced H_2O_2 -induced oxidative stress. Therefore, the molecular mechanisms underlying the involvement of miR-4463 in apoptotic pathways were investigated in HUVECs, by assessing the expression levels of apoptosis-associated marker proteins. Following miR-4463 overexpression in H_2O_2 -treated HUVECs, the proapoptotic

proteins C-caspase3, PARP1 and Bax were significantly upregulated, whereas the protein expression of the anti-apoptotic effectors Bcl-2 and XIAP was significantly reduced. These findings suggested that miR-4463 may be implicated in the regulation of apoptosis-related protein expression in HUVECs treated with H_2O_2 . However, further studies are required to fully elucidate the roles of miR-4463 and its target genes in apoptotic pathways.

In conclusion, the present results suggested that miR-4463 may be implicated in the pathogenesis of cardiovascular disease through the regulation of oxidative processes. Future studies will address the roles of specific target genes during the regulation of HUVEC oxidative stress and apoptosis, and

the effects of miRNAs in the pathophysiological mechanisms involved in oxidative processes will be explored. The results of the present study suggested that miR-4463 may have potential for the development of novel strategies for the diagnosis, prognosis and treatment of cardiovascular diseases.

References

- Ono K: MicroRNA-133a in the development of arteriosclerosis obliterans. *J Atheroscler Thromb* 22: 342-343, 2015.
- Rajput C, Tauseef M, Farazuddin M, Yazbeck P, Amin MR, Avin Br V, Sharma T and Mehta D: MicroRNA-150 suppression of angiopoietin-2 generation and signaling is crucial for resolving vascular injury. *Arterioscler Thromb Vasc Biol* 36: 380-388, 2016.
- Li X, Li J, Li Z, Sang Y, Niu Y, Zhang Q, Ding H and Yin S: Fucoidan from *Undaria pinnatifida* prevents vascular dysfunction through PI3K/Akt/eNOS-dependent mechanisms in the l-NAME-induced hypertensive rat model. *Food Funct* 7: 2398-2408, 2016.
- Guo R, Li W, Liu B, Li S, Zhang B and Xu Y: Resveratrol protects vascular smooth muscle cells against high glucose-induced oxidative stress and cell proliferation in vitro. *Med Sci Monit Basic Res* 20: 82-92, 2014.
- Csordas A, Kreutmayer S, Ploner C, Braun PR, Karlas A, Backovic A, Wick G and Bernhard D: Cigarette smoke extract induces prolonged endoplasmic reticulum stress and autophagic cell death in human umbilical vein endothelial cells. *Cardiovasc Res* 92: 141-148, 2011.
- Jeon BK, Kwon K, Kang JL and Choi YH: Csk-induced phosphorylation of Src at tyrosine 530 is essential for H₂O₂-mediated suppression of ERK1/2 in human umbilical vein endothelial cells. *Sci Rep* 5: 12725, 2015.
- Chen X, Pang S, Lin J, Xia J and Wang Y: Allicin prevents oxidized low-density lipoprotein-induced endothelial cell injury by inhibiting apoptosis and oxidative stress pathway. *BMC Complement Altern Med* 16: 133, 2016.
- Zhou DY, Su Y, Gao P, Yang QH, Wang Z and Xu Q: Resveratrol ameliorates high glucose-induced oxidative stress injury in human umbilical vein endothelial cells by activating AMPK. *Life Sci* 136: 94-99, 2015.
- Zheng D, Yu Y, Li M, Wang G, Chen R, Fan GC, Martin C, Xiong S and Peng T: Inhibition of microRNA 195 prevents apoptosis and multiple-organ injury in mouse models of sepsis. *J Infect Dis* 213: 1661-1670, 2016.
- Hecker M, Thamilarasan M, Koczan D, Schröder I, Flechtner K, Freisleben S, Füllen G, Thiesen HJ and Zettl UK: MicroRNA expression changes during interferon-beta treatment in the peripheral blood of multiple sclerosis patients. *Int J Mol Sci* 14: 16087-16110, 2013.
- Zhao S, Li T, Li J, Lu Q, Han C, Wang N, Qiu Q, Cao H, Xu X, Chen H and Zheng Z: miR-23b-3p induces the cellular metabolic memory of high glucose in diabetic retinopathy through a SIRT1-dependent signalling pathway. *Diabetologia* 59: 644-654, 2016.
- Li X, Yao N, Zhang J and Liu Z: MicroRNA-125b is involved in atherosclerosis obliterans in vitro by targeting podocalyxin. *Mol Med Rep* 12: 561-568, 2015.
- Correia AC, Moonen JR, Brinker MG and Krenning G: FGF2 inhibits endothelial-mesenchymal transition through microRNA-20a-mediated repression of canonical TGF- β signaling. *J Cell Sci* 129: 569-579, 2016.
- Savoia C, Sada L, Zezza L, Pucci L, Lauri FM, Befani A, Alonzo A and Volpe M: Vascular inflammation and endothelial dysfunction in experimental hypertension. *Int J Hypertens* 2011: 281240, 2011.
- Ding CF, Chen WQ, Zhu YT, Bo YL, Hu HM and Zheng RH: Circulating microRNAs in patients with polycystic ovary syndrome. *Hum Fertil (Camb)* 18: 22-29, 2015.
- He XM, Zheng YQ, Liu SZ, Liu Y, He YZ and Zhou XY: Altered plasma microRNAs as novel biomarkers for arteriosclerosis obliterans. *J Atheroscler Thromb* 23: 196-206, 2016.
- Lu G, Wong MS, Xiong MQ, Leung CK, Su XW, Zhou JY, Poon WS, Zheng VZY, Chan WY and Wong GKC: Circulating MicroRNAs in delayed cerebral infarction after aneurysmal subarachnoid hemorrhage. *J Am Heart Assoc* 6: pii: e005363, 2017.
- Liu X and Sun J: Endothelial cells dysfunction induced by silica nanoparticles through oxidative stress via JNK/P53 and NF-kappaB pathways. *Biomaterials* 31: 8198-8209, 2010.
- Lesniewska MA, Dereziński P, Klupczyńska A, Kokot ZJ, Ostrowski T, Zeidler J and Muszalska I: HPLC and HPLC/MS/MS studies on stress, accelerated and intermediate degradation tests of antivirally active tricyclic analog of acyclovir. *J AOAC Int* 98: 1240-1247, 2015.
- Lennicke C, Rahn J, Lichtenfels R, Wessjohann LA and Seliger B: Hydrogen peroxide - production, fate and role in redox signaling of tumor cells. *Cell Commun Signal* 13: 39, 2015.
- Zhang H, Sheng C, Yin Y, Wen S, Yang G, Cheng Z and Zhu Q: PABPC1 interacts with AGO2 and is responsible for the microRNA mediated gene silencing in high grade hepatocellular carcinoma. *Cancer Lett* 367: 49-57, 2015.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
- Nagababu P, Barui AK, Thulasiram B, Devi CS, Satyanarayana S, Patra CR and Sreedhar B: Antiangiogenic activity of mononuclear copper(II) polypyridyl complexes for the treatment of cancers. *J Med Chem* 58: 5226-5241, 2015.
- Ma Y, Li W, Yin Y and Li W: AST IV inhibits H₂O₂-induced human umbilical vein endothelial cell apoptosis by suppressing Nox4 expression through the TGF- β 1/Smad2 pathway. *Int J Mol Med* 35: 1667-1674, 2015.
- Yan S, Zhang X, Zheng H, Hu D, Zhang Y, Guan Q, Liu L, Ding Q and Li Y: Clematicinenoside inhibits VCAM-1 and ICAM-1 expression in TNF- α -treated endothelial cells via NADPH oxidase-dependent I κ B kinase/NF- κ B pathway. *Free Radic Biol Med* 78: 190-201, 2015.
- Liu S, Luo R, Xiang Q, Xu X, Qiu L and Pang J: Design and synthesis of novel xyloketal derivatives and their protective activities against H₂O₂-induced HUVEC injury. *Mar Drugs* 13: 948-973, 2015.
- Hou X, Tong Q, Wang W, Xiong W, Shi C and Fang J: Dihydromyricetin protects endothelial cells from hydrogen peroxide-induced oxidative stress damage by regulating mitochondrial pathways. *Life Sci* 130: 38-46, 2015.
- Wu W, Zhang D, Pan D, Zuo G, Ren X and Chen S: Downregulation of vascular endothelial growth factor receptor-2 under oxidative stress conditions is mediated by b-transduction repeat-containing protein via glycogen synthase kinase-3 β signaling. *Int J Mol Med* 37: 911-920, 2016.
- Chen AY, Lü JM, Yao Q and Chen C: Entacapone is an antioxidant more potent than vitamin C and vitamin E for scavenging of hypochlorous acid and peroxynitrite and the inhibition of oxidative stress-induced cell death. *Med Sci Monit* 22: 687-696, 2016.
- Zhang HB, Wen JK, Zhang J, Miao SB, Ma GY, Wang YY, Zheng B and Han M: Flavonoids from *Inula britannica* reduces oxidative stress through inhibiting expression and phosphorylation of p47(phox) in VSMCs. *Pharm Biol* 49: 815-820, 2011.
- Fetahu IS, Tennakoon S, Lines KE, Gröschel C, Aggarwal A, Mesteri I, Baumgartner-Parzer S, Mader RM, Thakker RV and Kállay E: miR-135b- and miR-146b-dependent silencing of calcium-sensing receptor expression in colorectal tumors. *Int J Cancer* 138: 137-145, 2016.
- Hu W, Wang M, Yin H, Yao C, He Q, Yin L, Zhang C, Li W, Chang G and Wang S: MicroRNA-1298 is regulated by DNA methylation and affects vascular smooth muscle cell function by targeting connexin 43. *Cardiovasc Res* 107: 534-545, 2015.
- Liu J, Li W, Wang S, Wu Y, Li Z, Wang W, Liu R, Ou J, Zhang C and Wang S: miR-142-3p attenuates the migration of CD4⁺ T cells through regulating actin cytoskeleton via RAC1 and ROCK2 in arteriosclerosis obliterans. *PLoS One* 9: e95514, 2014.
- Mostafavi N, Haghjooy-Javanmard S, Presidend N, Manssori NS and Kelishadi R: Persistence of endothelial cell damage late after Kawasaki disease in patients without coronary artery complications. *Adv Biomed Res* 4: 25, 2015.
- Du P, Suhaeri M, Subbiah R, Van SY, Park J, Kim SH, Park K and Lee K: Elasticity modulation of fibroblast-derived matrix for endothelial cell vascular morphogenesis and mesenchymal stem cell differentiation. *Tissue Eng Part A* 22: 415-426, 2016.
- Song Z, Liu Y, Hao B, Yu S, Zhang H, Liu D, Zhou B, Wu L, Wang M, Xiong Z, *et al*: Ginsenoside Rb1 prevents H₂O₂-induced HUVEC senescence by stimulating sirtuin-1 pathway. *PLoS One* 9: e112699, 2014.

37. Yin YL, Li P, Yang J, Liu DW, Sun RL, Pan GP, Wan GM and Wan GR: Protective effect of ultrafiltered XinMaiJia extract against H₂O₂-induced injury in human umbilical vein endothelial cells through NHE1 downregulation. *Genet Mol Res* 13: 8436-8449, 2014.
38. Zhang L, Zhou M, Qin G, Weintraub NL and Tang Y: miR-92a regulates viability and angiogenesis of endothelial cells under oxidative stress. *Biochem Biophys Res Commun* 446: 952-958, 2014.
39. Kojima H, Otani A, Oishi A, Makiyama Y, Nakagawa S and Yoshimura N: Granulocyte colony-stimulating factor attenuates oxidative stress-induced apoptosis in vascular endothelial cells and exhibits functional and morphologic protective effect in oxygen-induced retinopathy. *Blood* 117: 1091-1100, 2011.
40. Yu DS, Lv G, Mei XF, Cao Y, Wang YF, Wang YS and Bi YL: miR-200c regulates ROS-induced apoptosis in murine BV-2 cells by targeting FAP-1. *Spinal cord*: Dec 2, 2014 (Epub ahead of print).
41. Kong BS, Cho YH and Lee EJ: G protein-coupled estrogen receptor-1 is involved in the protective effect of protocatechuic aldehyde against endothelial dysfunction. *PLoS One* 9: e113242, 2014.