

CORRIGENDUM

DOI: 10.3892/mmr.2017.7234

Death receptor 5 expression is inversely correlated with prostate cancer progression

ANGELES HERNANDEZ-CUETO, DANIEL HERNANDEZ-CUETO, GABRIELA ANTONIO-ANDRES, MARISELA MENDOZA-MARIN, CARLOS JIMENEZ-GUTIERREZ, ANA LILIA SANDOVAL-MEJIA, ROSARIO MORA-CAMPOS, CESAR GONZALEZ-BONILLA, MARIO I. VEGA, BENJAMIN BONAVIDA and SARA HUERTA-YEPEZ

MolMedRep 10: 2279-2286, 2014; DOI: 10.3892/mmr.2014.2504

During the preparation of the figures in the above article, the authors inadvertently duplicated in Fig. 1B, a and b (high and low magnification images) the images that had already appeared as Figs. 5A, a and c (high and low magnification images), respectively, of the following paper: Huerta-Yepez S, Baritaki S, Baay-Guzman G, Hernandez-Luna MA, Hernandez-Cueto A, Vega MI and Bonavida B: Contribution of either YY1 or BclXL-induced inhibition by the NO-donor DETANONOate in the reversal of drug resistance, both *in vitro* and *in vivo*. Nitric Oxide 29: 17-24, 2013.

The revised version of Fig. 1 containing the corrected data for Fig. 1B, a and b (high and low magnification images; the YY1 data) is shown opposite protein expression. All those authors whom the corresponding author was able to contact have agreed to this Corrigendum. The authors regret this error, and apologize for any confusion that it may have caused.




Figure 1. YY1 and DR5 expression in PIN and PCa in tissue microarray constructions. (A) Immunohistochemical staining for YY1 and DR5 in LG PIN and HG PIN samples. (a) LG PIN tissue showed weak nuclear and cytoplasmic epithelial staining of glandular cells. (b) HG PIN tissue showed frequently higher staining. Scoring was derived from the nuclear expression. (c) LG PIN tissue showed high cytoplasmic membrane staining of glandular cells. (d) HG PIN tissue showed frequently high staining. Scoring was derived from the cytoplasmic membrane staining expression. (B) Immunohistochemical staining for YY1 and DR5 in LG PCa and HG PCa samples. (a) LG PCa tissue showed moderate nuclear epithelial staining of glandular cells. (b) HG PCa tissue showed frequently strong nuclear epithelial staining of glandular cells. (c) LG PCa tissue showed weak cytoplasmic membrane staining of glandular cells. (d) HG PCa tissue showed the weakest to absent staining. Scoring was derived from the cytoplasmic staining expression. Magnification, x100 with x400 inserts. YY1, Yin Yang 1; DR5, death receptor 5; LG PCa, low grade prostate carcinoma; HG PCa, high grade prostate carcinoma; LG PIN, low grade prostatic intraepithelial neoplasia; HG PIN, high grade prostatic intraepithelial neoplasia.