

Detection of circulating tumor cells by reverse transcription-quantitative polymerase chain reaction and magnetic activated cell sorting in the peripheral blood of patients with hepatocellular carcinoma

LIMIN GUO^{1*}, DONGHAI XU^{2*}, YAN LU¹, JIRUN PENG³ and LI JIANG¹

Departments of ¹Hepatobiliary Surgery and ²Radiology, Beijing Ditan Hospital, Capital Medical University, Beijing 100015;

³Department of Surgery, Beijing Shijitan Hospital, Capital Medical University, Beijing 100038, P.R. China

Received February 28, 2016; Accepted February 23, 2017

DOI: 10.3892/mmr.2017.7372

Abstract. Hepatocellular carcinoma (HCC) is one of the most lethal malignancies worldwide. Circulating tumor cells (CTCs) are considered a major cause of recurrence and metastasis in cancer; however, the detection of CTCs is challenging owing to their very low numbers in peripheral blood (around 10 CTCs per 1,000,000 erythrocytes). Cancer-testis antigens (CTAs) are specific tumor markers for CTCs. The present study aimed to evaluate the sensitivity and specificity of reverse transcription-quantitative polymerase chain reaction (RT-qPCR) for the detection of nine CTAs as well as placenta-specific antigen 1 (PLAC1) in peripheral blood mononuclear cell (PBMC) samples collected from 51 patients with HCC. The effectiveness of magnetic-activated cell sorting (MACS) for tumor-cell enrichment, through the depletion of CD45⁺ leukocytes in PBMC samples, was also assessed. Immunocytochemistry along with hematoxylin and eosin staining demonstrated that RT-qPCR achieved an overall positive detection rate for CTAs and PLAC1 of 70.6%; the highest rates were observed for melanoma-associated antigen A3 (MAGEA3), synovial sarcoma X breakpoint 1, MAGEA1, NY-ESO-1, L antigen 1 and PLAC1. MACS-detected intact CTCs in PBMCs were confirmed by H&E staining and morphological assessment; 12 out of 19 (63.2%) patients were identified as positive for CTAs. Screening for these five CTAs and PLAC1 by RT-qPCR may offer a potentially valuable prognostic tool with good

sensitivity and specificity in patients with HCC that may be enhanced by MACS.

Introduction

Hepatocellular carcinoma (HCC) is one of the most lethal malignancies worldwide, and is the third most common cause of cancer-associated mortality. Previous epidemiological studies revealed that ~626,000 new diagnoses of HCC are made annually (1). Despite improved therapeutic modalities, the prognosis for patients with HCC remains poor, primarily owing to difficulties in early diagnosis and the high rates of recurrence and metastasis following curative resection; 54-62% of patients develop recurrence within 5 years of surgery, and the overall 5-year survival rate of patients with HCC is <20% (2).

Circulating tumor cells (CTCs) are tumor cells that are found in the peripheral blood of patients with malignant disease, and are increasingly being recognized as a major cause of recurrence and metastasis (3). Therefore, early detection of CTCs may facilitate prognostic evaluations and contribute to effective therapeutic regimens in patients with HCC. However, the number of CTCs in peripheral blood is extremely low, with only a few cells/ml, compared with the billions of leukocytes and erythrocytes that may be found (4). Advances in molecular technologies and the discovery of novel tumor markers are making it possible to detect these rare cells with higher sensitivity and specificity. Although reverse transcription-polymerase chain reaction (RT-PCR) and nested-primer RT-PCR may be very sensitive, they sometimes yield false positive results (5). By contrast, quantitative RT-PCR (RT-qPCR) may measure very low expression levels of tumor marker genes with precise and reproducible results, and therefore may be a powerful tool to detect CTCs in peripheral blood mononuclear cell (PBMC) samples (6).

Albumin and α -fetoprotein (AFP) are the most commonly used tumor markers for the detection of CTCs in patients with HCC. However, they are non-specific and may occasionally be observed in the peripheral blood of patients with nonmalignant liver diseases, including cirrhosis or hepatitis. Previous reports have revealed that the positive detection

Correspondence to: Professor Li Jiang, Department of Hepatobiliary Surgery, Beijing Ditan Hospital, Capital Medical University, 8 Jingshun East Street, Chaoyang, Beijing 100015, P.R. China
E-mail: jiangli1903@163.com

*Contributed equally

Key words: hepatocellular carcinoma, circulating tumor cells, reverse transcription-quantitative polymerase chain reaction, immunomagnetic beads, tumor cell enrichment

rates for these markers using RT-PCR or nested RT-PCR range between 23.4 and 72.1% (7,8). Cancer-testis antigens (CTAs) have recently been identified as specific tumor markers for CTCs (9,10). CTAs belong to a superfamily of proteins that are solely expressed in tumors and in the testes, but not in normal tissues (11,12). Members of this superfamily include: Fetal and adult testis-expressed 1 (FATE1), melanoma-associated antigen C2 (MAGEC2), L antigen 1 (LAGE1), MAGEA1, MAGEA3, NY-ESO-1, sperm protein associated with the nucleus on the X chromosome (SPANX) A1, SPANXC and synovial sarcoma X breakpoint 1 (SSX1). Placenta-specific antigen 1 (PLAC1) exhibits a similar expression pattern as CTAs, and is expressed in a variety of tumors and in the placenta, but rarely in normal tissues (13).

In the present study, the effectiveness of RT-qPCR for detecting CTCs in patients with HCC was assessed by evaluating the mRNA expression levels of nine CTAs and PLAC1 in PBMC samples from 51 preoperative patients with HCC. In addition, immunomagnetic beads were used to enrich the PBMC samples for tumor cells to identify the presence of CTCs morphologically. By using RT-qPCR, CTCs were detected in ~70% of patients with HCC, indicating that this technique may be enhanced by magnetic-activated cell sorting (MACS).

Materials and methods

Ethical approval and informed consent. The experiments of the present study were approved by the ethics committee of Beijing Ditan Hospital (Capital Medical University, Beijing, China) and written informed consent was obtained from patients prior to enrollment in the study.

Cell lines and cell culture. The QGY7703 human HCC cell line was used in the present study as it expresses the nine target CTAs and PLAC1 genes at moderate levels, as determined by our preliminary experiment (data not shown). The QGY7703 cell line was purchased from the Institute of Biochemistry and Cell Biology (Shanghai, China). Cells were cultured in RPMI-1640 medium (Gibco; Thermo Fisher Scientific, Inc., Waltham, MA, USA) supplemented with 10% fetal calf serum (FBS; Hyclone; GE Healthcare Life Sciences, Logan, UT, USA), 2 mmol glutamine and antibiotics (penicillin 100 U/ml and streptomycin 100 µg/ml), at 37°C in a humidified atmosphere containing 5% CO₂.

Patient characteristics and peripheral blood sample preparation. Peripheral blood (10–15 ml) was collected from 51 preoperative patients with HCC, 10 patients with liver cirrhosis and 20 healthy donors. Clinicopathological characteristics of the patients and the controls are summarized in Table I. Samples were mixed with heparin to prevent coagulation, and were diluted with 10 ml RPMI-1640 medium supplemented with 10% FBS, 2 mmol glutamine and antibiotics. The samples were placed on a Ficoll-Hypaque layer (Thermo Fisher Scientific, Inc.) and PBMCs were collected by density gradient centrifugation at 670 x g and room temperature for 20 min.

Total RNA extraction and cDNA synthesis. Total RNA was isolated from 1x10⁷ cultured QGY7703 cells and PBMC samples using an RNeasy kit (Qiagen GmbH, Hilden,

Germany) and treated with RNase-free DNase I (Qiagen GmbH), following the manufacturer's protocol. First strand cDNA was reverse transcribed from total RNA (2 µg) using random hexamers [d(N)₆] (10 pmol/µl; 2 µl) and Superscript II Reverse Transcriptase (200 U/µl; 1 µl) at 50°C for 50 min (Thermo Fisher Scientific, Inc.), according to manufacturer's protocol.

Preparation of recombinant plasmids as copy number standards for RT-qPCR. Human FATE1, MAGEC2, LAGE1, MAGEA1, MAGEA3, NY-ESO-1, SPANXA1, SPANXC and SSX1 cDNAs were amplified from human testis cDNA by qPCR; PLAC1 cDNA was amplified from placenta cDNA by PCR. Human testis (from patients with advanced prostate cancer following castration therapy) and placenta (from patients following normal delivery) samples were obtained from Beijing Ditan Hospital. Amplification reactions were carried out using cDNA (1 µl), forward and reverse primers (10 pM each), 2X PCR Master Mix (12.5 µl; Invitrogen; Thermo Fisher Scientific, Inc.), and double-distilled H₂O to a final volume of 25 µl. PCR products were purified using the Wizard® SV Gel and PCR Clean-Up System (Promega Corporation, Madison, WI, USA) and cloned into a pGEM-T Easy Vector system (Promega Corporation), following the manufacturer's protocol. Recombinant plasmids were confirmed by DNA sequencing. Briefly, plasmids were linearized by digestion with an EcoRI/NcoI restriction endonuclease at 37°C for 4 h (Takara Bio, Inc., Otsu, Japan) before being purified by 1.5% agarose gel electrophoresis and the Wizard® SV Gel and PCR Clean-Up System (Promega Corporation), and quantified using a spectrophotometer at a wavelength of 260 nm. The plasmid copy numbers (copies/ml) were calculated using the following formula: [plasmid concentration (g/ml) x 6.023x10²³]/(number of base pairs in the plasmid x 660). The number of base pairs in the plasmid includes the plasmid vector and the gene insert. These were used as the copy number standards for subsequent RT-qPCR analyses.

RT-qPCR and construction of standard curves. RT-qPCR primers were designed to specifically amplify the target cDNA fragments containing at least one intron, while avoiding contamination from genomic DNA. Primer sequences and TaqMan probes are listed in Table II; the housekeeping gene GAPDH was used as an internal control. Five 10-fold dilutions of the resulting constructs, ranging from 1x10⁴ to 1x10⁸ copies µl⁻¹, were used to construct the standard curves for GAPDH, LAGE, NY-ESO-1, MAGEA1, MAGEA3, HCA587, FATE, SPANXA1, SPANXC, PLAC1 and SSX-1. Absolute quantification of mRNA using RT-qPCR was performed, which employed standard curves for quantification (14).

TaqMan probes were labeled at the 5' end with the 6-carboxyfluorescein fluorophore, and at the 3' end with the fluorescence quencher tetramethylrhodamine. qPCR reactions were prepared as follows: 1 µl cDNA (~100 ng total RNA) or plasmid DNA (containing 10⁴–10⁸ copies of the target sequence) was mixed with 7.5 pM of each forward and reverse primers, 6.25 pM TaqMan probe and 2X qPCR buffer to a final volume of 25 µl. A positive control containing 10 copies of plasmid DNA was prepared. To test for potential contaminants, two reactions were prepared that contained either 1 µl

Table I. Clinical parameters of the 51 patients with hepatocellular carcinoma and the controls.

Clinicopathological characteristics	Hepatocellular carcinoma, n=51	Cirrhosis donors, n=10	Healthy donors, n=20
Sex			
Male	35	7	13
Female	16	3	7
Age (years)			
<40	6	1	3
≥40	45	9	17
AFP (ng/ml)			
<400	34	-	-
≥400	17	-	-
Tumor size (cm)			
<3	40	NA	NA
≥3	11	NA	NA
Invasion of main portal vein			
Yes	3	NA	NA
No	48	NA	NA
TNM stage			
I and II	33	NA	NA
III and IV	18	NA	NA
Hepatitis B virus			
Positive	41	8	0
Negative	10	2	20
Grade			
Well differentiated	6	NA	NA
Moderately differentiated	38	NA	NA
Poorly differentiated	7	NA	NA
Modus operandi			
Excision	46	NA	NA
Orthotopic liver transplantation	5	NA	NA

Levels of AFP were not measured in patients with cirrhosis or healthy donors. AFP, α -fetoprotein; TNM, tumor-node-metastasis; NA, not applicable.

water or 1 μ l reverse transcriptase without RNA. qPCR was performed at 50°C for 2 min and 95°C for 10 min, followed by 40 cycles of 92°C for 15 sec and 60°C for 1 min, using an ABI PRISM 7700 Sequence Detection System (Applied Biosystems; Thermo Fisher Scientific, Inc.). Each experiment was performed in triplicate.

Sensitivity of RT-qPCR and standard RT-PCR for the detection of MAGEA1 and SSX1 cDNA in different ratios of HCC cells to normal PBMC. HCC to normal PBMC ratios of 10^{-2} , 10^{-3} , 10^{-4} , 10^{-5} and 10^{-6} were prepared by mixing 10^5 , 10^4 , 10^3 , 10^2 and 10^1 QGY7703 cells, respectively, with 10^7 normal PBMCs from healthy individuals. A control with only normal PBMCs was additionally prepared. The cells were recovered by Ficoll-Hypaque density gradient centrifugation (670 x g for 20 min at room temperature) and total RNA was extracted as aforementioned. The expression levels of MAGEA1 and SSX1 cDNA in each of the six samples were measured by RT-qPCR

and standard RT-PCR. Amplification reactions for standard RT-PCR were performed using cDNA (1 μ l), forward and reverse primers (10 pM each), 2X PCR Master Mix (12.5 μ l; Thermo Fisher Scientific, Inc.) and double-distilled H₂O to a final volume of 25 μ l. The following optimal thermal cycling conditions were used: 94°C for 10 min, followed by 35 cycles of 94°C for 25 sec, 54°C for 25 sec and 72°C for 25 sec; followed by 72°C for 8 min. The housekeeping gene GAPDH was used as a control. The optimal thermal cycling conditions for GAPDH were the same for SSX1 and MAGEA1 except that the annealing temperature was 57°C for GAPDH. The same forward and reverse primers were used in RT-qPCR and standard RT-PCR.

Enrichment of HCC cells in PBMC samples by depletion of CD45⁺ cells by MACS. PBMC samples were acquired from 19 HCC patients that were identified to have positive expression levels of the tested CTAs by RT-qPCR. PBMC samples from 10 patients with liver cirrhosis were additionally used,

Table II. List of primer sequences and TaqMan probes.

Gene	Primer sequence (5'-3')	Expected size (bp)	Probe (5'-3')
FATE1	F: GCCAAGGGGAAAACCAAGAG R: TCGGGTGGCAGTCATATTCC	152	CAAAAAGCTGCTGGCTCTGCTTCAGC
MAGEC2	F: AGGGACCTCCCACCATAGAGA R: CCTCCTCTGCTGTGCTGACTT	130	CCGGCCTGTACTGCGCTGCC
LAGE1	F: CCGCCTGCTTCAGTTGC R: CTGCAGCAGTCAGTCGGATA	167	ACGATGCCTTTCTCGTCGCCCCAT
MAGEA1	F: AGAGGAGCACCAAGGAGAAGA R: GGCAGCAGGCAGGAGTGT	74	CTGCCTGTGGGTCTTCATTGCCCA
MAGEA3	F: GGAGCACTGAAGGAGAAGATCTG R: TGA CTCTGGTCAGGGCAACA	142	TGGGTCTCCATTGCCCAGCTCC
NY-ESO-1	F: TGCTTGAGTTCTACCTCGCCA R: TATGTTGCCGGACACAGTGAA	137	AG GATGCCCCACCGCTTCCC
SPANXA1	F: GCCACTGACATTGAAGAACCAA R: GGGTCTCCGGCATCATCTC	115	ATACAATGGACAAACAATCCAGTGCCGG
SPANXC	F: CCGCCTGCTTCAGTTGC R: CTGCAGCAGTCAGTCGGATA	167	TCTGAGTCCTCGACCATACTAGTGGTTCGC
SSX1	F: GTCACCAACTGCTGCCAAC R: TTGCTTCTCTTCTCTGATGCTTT	153	TGCCATGAACGGAGACGACACCTTT
PLAC1	F: CAAGTTCCTTCTTCCCTTTGGAA R: CGTCCAGTGAGGATTTCTTCTTC	120	CAGCTGCCTTCACCAGTGAGCACAA
GAPDH	F: GAAGGTGAAGGTCGGAGTC R: GAAGATGGTGATGGGATTTC	226	CAAGCTTCCCGTTCTCAGCC

Bp, base pairs; F, forward; FATE1, fetal and adult testis-expressed 1; LAGE1, L antigen family 1; MAGE, melanoma-associated antigen; PLAC1, placenta-specific antigen 1; R, reverse; SPANX, sperm protein associated with the nucleus on the X chromosome; SSX1, synovial sarcoma X breakpoint 1.

along with the 10 healthy donors as control. Cells ($\sim 1 \times 10^7$) were washed with PBS and incubated with anti-CD45 antibody-coated immunomagnetic microbeads (Miltenyi Biotec GmbH; Bergisch Gladbach, Germany) at 4°C for 15 min. The cells were subsequently separated on an autoMACS Column (Miltenyi Biotec GmbH) using the sensitive depletion mode for gentle separation to ensure cellular integrity. The CD45-negative fraction, which is effectively enriched in HCC tumor cells due to the depletion of PBMCs, was eluted and cells were collected onto slides using a cytospin device; slides were air dried before being stored at -20°C until required.

Immunocytochemistry of cytokeratin to detect HCC cells in enriched PBMC samples. Slides were fixed in 4% paraformaldehyde for 20 min at room temperature and examined for the presence of cytokeratin-positive cells by immunocytochemistry. Following permeabilization with 0.3% Triton-X-100 for 10 min at room temperature, and treatment with antigen retrieval solution for 30 min at 80°C (Beyotime Institute of Biotechnology, Haimen, China), the slides were incubated in 3% H₂O₂ for 20 min and rinsed with distilled water and PBS. Blocking was performed using 5% normal goat serum (Shanghai Yanjin Biotechnology Co., Ltd., Shanghai, China) at room temperature for 15 min. This was followed by incubation with the mouse anti-cytokeratin, clone CK3-6H5 primary

antibody (1:50; cat. no. 130-098-802; Miltenyi Biotec GmbH) overnight at 4°C. The next day, the slides were rinsed and incubated with the biotinylated horse anti-mouse immunoglobulin G (1:500; cat. no. orb153695; Biorbyt, Ltd., Cambridge, UK) at room temperature for 30 min. The signal was visualized with peroxidase-conjugated streptavidin and 3,3'-diaminobenzidine (DAB). The slides were subsequently examined and photographed using an Olympus BX53 fluorescence microscope (Olympus Corporation, Tokyo, Japan). DAB staining was analyzed by Image-Pro Plus 6.0 software (Media Cybernetics, Inc., Rockville, MD, USA). Unenriched QGY7703 cells were used as the positive control; healthy PBMCs were used as the negative control.

Hematoxylin and eosin (H&E) staining of PBMC samples enriched in HCC tumor cells. Slides were fixed in 4% paraformaldehyde and stained with H&E according to routine protocols at room temperature ($\sim 25^\circ\text{C}$). Briefly, slides were washed with distilled water and stained with hematoxylin solution for 5-10 min, followed by soaking the stained slides in distilled water for 10 min to remove excess dye. Subsequently, after soaking in 95% ethanol for 5 sec, slides were stained with eosin staining solution for 2 min, which was followed by dehydration with graded alcohol and clearing in xylene. Finally, slides were sealed with neutral gum. The mounted slides were




Figure 1. Standard plots (cycle threshold vs. plasmid copy number) for 10 circulating tumor cell markers, including nine cancer-testis antigens and PLAC1 cDNAs. The similar slopes confirm the reproducibility and specificity of reverse transcription-quantitative polymerase chain reaction for detecting circulating tumor cells in the peripheral blood from patients with hepatocellular carcinoma. GAPDH was used as a reference marker. FATE1, fetal and adult testis-expressed 1; LAGE1, L antigen family 1; MAGE, melanoma-associated antigen; PLAC1, placenta-specific antigen 1; SPANX, sperm protein associated with the nucleus on the X chromosome; SSX1, synovial sarcoma X breakpoint 1.




Figure 2. Expression of MAGEA1 and SSX1 mRNAs in different ratios of QGY7703 hepatocellular carcinoma cells and control peripheral blood mononuclear cells were assessed by RT-qPCR (left-hand panels) and standard RT-PCR (right-hand panels) for mRNA expression levels of (A) MAGEA1 and (B) SSX1. (C) GAPDH was used as a reference marker. M, 100 bp marker gene; MAGEA1, melanoma-associated antigen A1; NTC, negative control for PCR; SSX1, synovial sarcoma X breakpoint 1; RT-qPCR, reverse transcription-quantitative polymerase chain reaction; ΔRn , normalized reporter value.




Figure 3. Visualization of HCC cells in peripheral blood samples enriched for HCC cells by magnetic-activated cell sorting. (A) Immunocytochemistry staining of a cytokeratin-positive tumor cell (green arrow) and leukocytes (black arrow) in peripheral blood from a patient with HCC. Magnification, x200. H&E staining at magnifications of (B) x200 and (C) x1,000 demonstrating tumor cells characterized by large, dark blue and heterogeneously stained nuclei with scant cytoplasm around the nuclei (green arrow). The sizes of the nuclei are approximately five times greater than those of leukocytes (black arrow). (D) H&E staining demonstrating that HCC tumor cells (green arrows), characterized by large, heterogeneously stained nuclei, are clearly different from other leukocytes (black arrow). Magnification, x1,000. H&E, hematoxylin and eosin; HCC, hepatocellular carcinoma.

examined and photographed using an Olympus BX53 fluorescence microscope (Olympus Corporation). Cell morphology was assessed by a chief pathologist who was blinded to the study. Unenriched QGY7703 cells were used as the positive control and healthy PBMCs were used as the negative control. Experiments were repeated three times.

Results

Standard curves for RT-qPCR. Standard curves for the nine CTAs, PLAC1 and GAPDH cDNAs were constructed using gene-specific recombinant plasmids as copy number standards (Fig. 1). The reducible amplification curves of the 11 standard curves exhibited similar slopes (between -3.20 and -3.49), confirming that the RT-qPCR protocol had acceptable reproducibility and specificity for the detection of the nine CTAs and PLAC1. However, the y-axis intercepts (cycle thresholds) ranged between 40.71 and 48.47, indicating variations in amplification sensitivities and efficiencies between the different cDNAs.

Sensitivity of the RT-qPCR assay. Using defined ratios of QGY7703 cells and normal PBMCs, the present study was able to detect as few as 10 CTCs among 10^7 PBMCs using RT-qPCR to measure the expression levels of SSX1 and MAGEA1 cDNA in HCC cells (Fig. 2). In contrast, standard RT-PCR was only able to detect as low as 10^3 CTCs in 10^7 PBMCs. This result demonstrated that RT-qPCR was ~100X more sensitive than RT-PCR for the detection of CTCs through CTA cDNA expression analysis in PBMCs.

Expression rates of CTA and PLAC1 cDNAs in peripheral blood samples from patients with HCC. The overall positive detection rate for the 10 tested cDNAs (nine CTAs and PLAC1) in PBMC samples from 51 patients with HCC was 70.6% (36/51 patients detected positive for ≥ 1 target cDNA). However, most of the patients only exhibited between 1 and 3 of these 10 markers. By contrast, none of the tested cDNAs were detected in PBMC samples from the 10 patients with cirrhosis and the 20 healthy patients. The individual positive detection rates were as follows: MAGEA3, 43.1% (22/51); SSX1, 39.2% (20/51); MAGEA1, 25.5% (13/51); NY-ESO-1, 19.6% (10/51); PLAC1, 19.6% (10/51); LAGE1, 17.6% (9/51); MAGEC2, 7.8% (4/51); SPANXC, 7.8% (4/51); SPANXA1, 2.0% (1/51); and FATE1, 0% (0/51). Six of the genes (MAGEA3, SSX1, MAGEA1, NY-ESO-1, PLAC1 and LAGE1) demonstrated high positive rates in the PBMCs of patients with HCC and were selected for further analysis. The expression levels of these six cDNAs, relative to GAPDH, were as follows: MAGEA3, $1/10^3$ - $1/10^7$; SSX1, $1/10^{3.5}$ - $1/10^{6.5}$; MAGEA1, $1/10^{2.5}$ - $1/10^{6.5}$; NY-ESO-1, $1/10^{3.5}$ - $1/10^6$; PLAC1, $1/10^4$ - $1/10^{5.5}$; and LAGE1, $1/10^2$ - $1/10^{5.5}$.

HCC cell enrichment in PBMC and detection by immunocytochemistry and H&E staining. A total of 19 PBMC samples that tested positive for CTA and PLAC1 cDNAs by RT-qPCR were randomly selected for tumor cell enrichment through the depletion of leukocytes by MACS. H&E staining and immunocytochemistry with an anti-cytokeratin antibody were used to detect HCC cells. A total of 12/19 (63.2%) samples exhibited intact tumor cells (Fig. 3), of which HCC cells were identified using both techniques (10/12 samples) or were only detected using one of the two techniques (2/12 samples). By contrast, HCC cells were not observed by either technique in PBMC samples from patients with cirrhosis or normal individuals.

Discussion

CTCs contribute to recurrence and metastasis and are often an indication of advanced stage cancer and poor prognosis (15). Recent reports (10,12) have demonstrated that by examining CTA expression levels, CTCs may be detected in the blood with greater specificity compared with traditional techniques, such as AFP evaluation, RT-PCR or nested-primer RT-PCR. To determine whether RT-qPCR was able to detect CTCs in patients with HCC with greater sensitivity and specificity than standard RT-PCR, the present study evaluated the expression levels of 10 CTC marker cDNAs (nine CTAs and PLAC1) by RT-qPCR in blood samples collected from 51 preoperative patients with HCC. In addition, anti-CD45 antibody coated immunomagnetic beads were used to enrich PBMC samples for tumor cells to establish the potential value of this technique for prognostic evaluation of patients with HCC.

Results from the present study revealed that RT-qPCR of CTA cDNAs was able to detect 10 tumor cells per 10^7 PBMCs, demonstrating that this technique was ~100X more sensitive than standard RT-PCR for the detection of CTCs. Of the 10 tested tumor markers, MAGEA3, SSX1, MAGEA1, NY-ESO-1, PLAC1 and LAGE1 cDNAs had positive detection rates $>10\%$; the overall positive detection rate was 70.6%. However, the expression levels of these six cDNAs varied among the 51 patients with HCC that were examined; most patients

tested positive for only 1-3 of the 10 CTC marker cDNAs. In contrast, all patients with cirrhosis and healthy individuals that were examined tested negative for the 10 cDNAs.

Enrichment of tumor cells by MACS, combined with H&E staining and morphological assessment, additionally emerged as a potential method for detecting CTCs in patients with HCC; positive identification of intact tumor cells was achieved in 63.2% (12/19) of patients that had been identified as positive by RT-qPCR. This result suggested that the two techniques may be combined as a valuable prognostic tool; however, comparable studies on alternative methods for identifying intact tumor cells in HCC are limited.

In conclusion, the present study demonstrated that the use of RT-qPCR to evaluate the expression rates of CTAs and PLAC1 in PBMC samples may offer a highly sensitive and specific technique to detect CTCs in the peripheral blood of patients with HCC. Furthermore, the accuracy of this method may be enhanced by MACS tumor cell enrichment.

Acknowledgements

The present study was supported by the Navigating Plan of Research Fund of Beijing Ditan Hospital, Capital Medical University (grant no. DTQH201405) and the Beijing Municipal Administration of Hospitals Incubating Program (grant no. PX2016043).

References

1. Parkin DM, Bray F, Ferlay J and Pisani P: Global cancer statistics, 2002. *CA Cancer J Clin* 55: 74-108, 2005.
2. Parkin DM: Global cancer statistics in the year 2000. *Lancet Oncol* 2: 533-543, 2001.
3. Elshimali YI and Grody WW: The clinical significance of circulating tumor cells in the peripheral blood. *Diagn Mol Pathol* 15: 187-194, 2006.
4. Hosokawa M, Hayata T, Fukuda Y, Arakaki A, Yoshino T, Tanaka T and Matsunaga T: Size-selective microcavity array for rapid and efficient detection of circulating tumor cells. *Anal Chem* 82: 6629-6635, 2010.
5. Samija I, Lukac J, Kusić Z, Situm M and Samija M: Detection of disseminated melanoma cells by reverse-transcription-polymerase chain reaction. *Coll Antropol* 31: 1187-1194, 2007.
6. Schefe JH, Lehmann KE, Buschmann IR, Unger T and Funke-Kaiser H: Quantitative real-time RT-PCR data analysis: Current concepts and the novel 'gene expression's CT difference' formula. *J Mol Med (Berl)* 84: 901-910, 2006.
7. Morimoto O, Nagano H, Miyamoto A, Fujiwara Y, Kondo M, Yamamoto T, Ota H, Nakamura M, Wada H, Damdinsuren B, *et al*: Association between recurrence of hepatocellular carcinoma and alpha-fetoprotein messenger RNA levels in peripheral blood. *Surg Today* 35: 1033-1041, 2005.
8. Yao F, Guo JM, Xu CF, Lou YL, Xiao BX, Zhou WH, Chen J, Hu YR, Liu Z and Hong GF: Detecting AFP mRNA in peripheral blood of the patients with hepatocellular carcinoma, liver cirrhosis and hepatitis. *Clin Chim Acta* 361: 119-127, 2005.
9. Lu Y, Wu LQ, Lü ZH, Wang XJ and Yang JY: Expression of SSX-1 and NY-ESO-1 mRNA in tumor tissues and its corresponding peripheral blood expression in patients with hepatocellular carcinoma. *Chin Med J (Engl)* 120: 1042-1046, 2007.
10. Mou DC, Cai SL, Peng JR, Wang Y, Chen HS, Pang XW, Leng XS and Chen WF: Evaluation of MAGE-1 and MAGE-3 as tumour-specific markers to detect blood dissemination of hepatocellular carcinoma cells. *Br J Cancer* 86: 110-116, 2002.
11. Zendman AJ, Ruiter DJ and Van Muijen GN: Cancer/testis-associated genes: Identification, expression profile, and putative function. *J Cell Physiol* 194: 272-288, 2003.
12. Peng JR, Chen HS, Mou DC, Cao J, Cong X, Qin LL, Wei L, Leng XS, Wang Y and Chen WF: Expression of cancer/testis (CT) antigens in Chinese hepatocellular carcinoma and its correlation with clinical parameters. *Cancer Lett* 219: 223-232, 2005.
13. Koslowski M, Sahin U, Mitnacht-Kraus R, Seitz G, Huber C and Türeci O: A placenta-specific gene ectopically activated in many human cancers is essentially involved in malignant cell processes. *Cancer Res* 67: 9528-9534, 2007.
14. Bustin SA: Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays. *J Mol Endocrinol* 25: 169-193, 2000.
15. Walzer N and Kulik LM: Hepatocellular carcinoma: Latest developments. *Curr Opin Gastroenterol* 24: 312-319, 2008.