

Adiponectin receptor agonist AdipoRon suppresses adipogenesis in C3H10T1/2 cells through the adenosine monophosphate-activated protein kinase signaling pathway

SHU-JUAN WANG, WEN-YI LU and KAI-YAN LIU

Beijing Key Laboratory of Hematopoietic Stem Cell Transplantation, Peking University People's Hospital and Institute of Hematology, Beijing 100044, P.R. China

Received January 3, 2017; Accepted July 24, 2017

DOI: 10.3892/mmr.2017.7450

Abstract. The aim of the present study was to investigate the effects of AdipoRon, an adiponectin receptor agonist, on adipogenesis in C3H10T1/2 cells and to explore the underlying mechanisms. C3H10T1/2 cells were treated with increasing doses of AdipoRon for 8 days, and Oil Red O staining was used to assess lipid accumulation. The protein and mRNA expression levels of adipogenic transcription factors and adipocyte-specific genes were examined by western blotting and reverse transcription quantitative polymerase chain reaction, respectively. AdipoRon treatment inhibited lipid accumulation in C3H10T1/2 cells in a dose-dependent manner and significantly suppressed the expression of adipogenic transcription factors, including peroxisome proliferator-activated receptor γ , CAAT/enhancer binding protein (C/EBP)- β and C/EBP α . In addition, cells treated with AdipoRon exhibited a significant decrease in the expression of adipocyte-specific genes, including fatty acid binding protein 4, fatty acid synthase, leptin, adiponectin, and stearoyl-CoA desaturase-1. Notably, AdipoRon significantly increased the phosphorylation of adenosine monophosphate-activated protein kinase (AMPK) and acetyl-CoA carboxylase (ACC). The results indicated that AdipoRon exerted an inhibitory effect on adipogenesis in C3H10T1/2 cells by downregulating the expression of adipogenic transcription factors and adipocyte-specific genes and by promoting the phosphorylation of AMPK and ACC, which suggested that AdipoRon may be a potential drug to prevent and treat diseases caused by abnormal adipogenesis, such as obesity.

Introduction

Obesity is a common public health problem and is a significant risk factor for a number of metabolic disorders, including type 2 diabetes, hypertension and cardiovascular disease (1). The development of obesity is characterized by an increase in the number and size of mature adipocytes that are produced by differentiation and mitogenesis (2). Adipogenesis is controlled by many transcription factors, including peroxisome proliferator-activated receptor γ (PPAR γ) and CCAAT/enhancer binding proteins (C/EBPs) (3-5). At the initial stage of adipogenesis, C/EBP β is activated, followed by the sequential activation of C/EBP α and PPAR γ , two major late-adipogenic transcription factors. Activation of C/EBP α and PPAR γ subsequently leads to the expression of a number of genes involved in adipocyte differentiation, including genes encoding lipid metabolizing enzymes [such as fatty acid binding protein (FABP)-4 and fatty acid synthase (FAS)] and adipokines (such as adiponectin and leptin) (6).

Adenosine monophosphate-activated protein kinase (AMPK) is an important cellular energy sensor, and its activation is closely related to the balance between lipid accumulation and carbohydrate metabolism (7). AMPK phosphorylation inhibits metabolic enzymes involved in fatty acid synthesis, leading to suppressed adipogenesis (8,9).

Adiponectin is an adipokine that is mainly produced by adipocytes. Unlike other adipokines, plasma adiponectin levels are reduced in obese subjects (10). Adiponectin binds to the adiponectin receptors, AdipoR1 and AdipoR2, and reduces obesity-related insulin resistance (11). AdipoRon, a newly discovered, orally active small molecule, is an adiponectin receptor agonist that binds to and activates both AdipoR1 and AdipoR2, and has been previously reported to improve obesity-related insulin resistance and type 2 diabetes by activating AMPK and PPAR α pathways (12). However, the effects of adiponectin on obesity itself and on adipogenesis are controversial. Adiponectin may promote adipogenesis induced by weight gain (11,13). By contrast, elevated adiponectin levels were recently reported to reduce lipid content and lipid metabolism in 3T3-L1 cells (14). However, the effects of AdipoRon on adipogenesis have not yet been investigated. The present study investigated the effects of AdipoRon on

Correspondence to: Professor Kai-Yan Liu, Beijing Key Laboratory of Hematopoietic Stem Cell Transplantation, Peking University People's Hospital and Institute of Hematology, 11 Xi-Zhi-Men South Street, Beijing 100044, P.R. China
E-mail: liukaiyan@medmail.com.cn

Key words: adiponectin, AdipoRon, antiadipogenesis, adenosine monophosphate-activated protein kinase, C3H10T1/2 cells

adipogenesis and explored the underlying mechanism in C3H10T1/2 cells.

Materials and methods

Reagents. AdipoRon, 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranoside (AICAR), dexamethasone (Dex), 3-isobutyl-1-methylxanthine (IBMX), indomethacin, insulin and Oil Red O were purchased from Sigma-Aldrich (Merck KGaA, Darmstadt, Germany). Minimum essential Eagle's medium with Earle's Balanced Salts (MEM-EBSS) was purchased from HyClone (GE Healthcare Life Sciences, Logan, UT, USA). Fetal bovine serum (FBS) was purchased from Gibco (Thermo Fisher Scientific, Inc., Waltham, MA, USA). Cell Counting Kit-8 (CCK-8) was purchased from Dojindo Molecular Technologies, Inc. (Kumamoto, Japan). Monoclonal antibodies, including rabbit anti-PPAR γ (cat no. 2443), rabbit anti-FABP4 (cat no. 3544), rabbit anti-C/EBP β (cat no. 3087), rabbit anti-C/EBP α (cat no. 2295), rabbit anti-AMPK α (cat no. 5831), rabbit anti-phosphorylated-acetyl-CoA carboxylase (p-ACC; cat no. 11818), rabbit anti-adiponectin (cat no. 2789) and rabbit anti-GAPDH (cat no. 5174) were purchased from Cell Signaling Technology (Danvers, MA, USA). Rabbit polyclonal anti-p-AMPK α 1/2 antibody (cat no. sc-33524) and goat polyclonal anti-AdipoR1 antibody (cat no. sc-46748) and anti-AdipoR2 antibody (cat no. sc-46751) were purchased from Santa Cruz Biotechnology, Inc. (Dallas, TX, USA).

Cell culture and adipocyte differentiation. The C3H10T1/2 mouse embryonic mesenchymal stem cell line was purchased from the Chinese Academy of Medical Sciences (Beijing, China). Cells were cultured in MEM-EBSS supplemented with 10% FBS at 37°C in a 5% CO₂ atmosphere until adipocyte differentiation. Two days after reaching confluence (day 0), the C3H10T1/2 cells were cultured in differentiation medium (DM; MEM/EBSS containing 10% FBS, 1 μ M Dex, 0.5 mM IBMX, 50 mM indomethacin and 1 μ g/ml insulin) for 2 days to induce differentiation. Following incubation, MEM/EBSS supplemented with 10% FBS and 1 μ g/ml insulin was added for 2 days as a differentiation medium and was changed every 2 days.

Cell viability assay. Cell viability was detected by using a CCK-8 kit, according to the manufacturer's instructions. Briefly, C3H10T1/2 cells were seeded in 96-well plates at a density of 5x10⁴ cells/ml at 37°C in a humidified 5% CO₂ atmosphere and treated with increasing doses of AdipoRon (1, 5, 10, 20 and 40 μ M) for 24, 48 or 72 h. Subsequently, 10 μ l of kit reagent was added to each well and incubated at 37°C for 2 h. The plates were scanned with a microplate reader (Benchmark; Bio-Rad Laboratories, Inc., Hercules, CA, USA) at 450 nm.

Oil Red O (ORO) staining. Cells were cultured in 24-well plates with a seeding density of 5x10⁴ cells/ml and differentiated as described above. On day 8, the cells were stained with ORO as previously described (2). Briefly, the cells were fixed with 10% formalin (Solarbio, Beijing, China) at 4°C for 30 min and stained with ORO at room temperature for 30 min following washes with phosphate-buffered saline. Oil droplets

in the cells were observed under an inverted microscope (Olympus-CKX41; Olympus Corporation, Tokyo, Japan). The ORO-stained plates were then washed and treated with 100% isopropanol (Beijing Solarbio Science & Technology Co., Ltd., Beijing, China), and lipid accumulation was detected by measuring the absorbance at 490 nm using a Benchmark microplate reader (Bio-Rad Laboratories, Inc.).

Western blot analysis. Cells were cultured in 6-well plates with a seeding density of 5x10⁴ cells/ml and differentiated as described above. Cells were lysed in radioimmunoprecipitation assay buffer (Beyotime Institute of Biotechnology, Shanghai, China) supplemented with a protease inhibitor cocktail and phosphatase inhibitors (Roche Molecular Diagnostics, Pleasanton, CA, USA) for 20 min and centrifuged at 12,000 x g for 20 min at 4°C. Protein concentration was determined using the Bicinchoninic Acid Protein Assay reagent kit (Beijing Solarbio Science & Technology Co., Ltd.). A total of 30 μ g of protein lysates from each sample were separated on 10 or 12% SDS-PAGE and transferred to a polyvinylidene difluoride membrane (EMD Millipore, Billerica, MA, USA). The membrane was incubated with 5% bovine serum albumin in Tris-buffered saline with 0.1% Tween-20 (TBST) (both from Beijing Solarbio Science & Technology Co., Ltd.) for 1 h at room temperature, followed by hybridization with primary antibodies against PPAR γ (1:1,000), C/EBP β (1:1,000), C/EBP α (1:1,000), FABP4 (1:2,000), adiponectin (1:500), AMPK α (1:1,000), p-ACC (1:500), GAPDH (1:1,000), p-AMPK α 1/2 (1:200), AdipoR1 (1:200) and AdipoR2 (1:200) overnight at 4°C. Following three washes with TBST, the membrane was incubated with a horseradish peroxidase-conjugated goat anti-rabbit IgG secondary antibody (cat no. sc2004; 1:10,000; Santa Cruz Biotechnology, Inc.) or a horseradish peroxidase-conjugated rabbit anti-goat IgG secondary antibody (cat no. ZB-2306; 1:10,000; Beijing Zhongshan Jinqiao Biotechnology Co., Ltd., Beijing, China) for 1 h at room temperature. Following three washes with TBST, the bands were visualized with an Enhanced Chemiluminescence Substrate (Thermo Fisher Scientific, Inc.). Densitometry of the western blot bands was performed using ImageJ software version 1.31 (National Institutes of Health, Bethesda, ML, USA).

RNA preparation and reverse transcription-quantitative polymerase chain reaction (RT-qPCR). Cells were cultured in 6-well plates with a seeding density of 5x10⁴ cells/ml and differentiated as described above. RNA was extracted from C3H10T1/2 cells on day 8 using TRIzol reagent (Invitrogen; Thermo Fisher Scientific, Inc.). A High Capacity cDNA Reverse Transcription kit (Applied Biosystems; Thermo Fisher Scientific, Inc.) was used to synthesize the cDNA, according to the manufacturer's protocol. mRNA expression levels were determined using an ABI 7500 Fast real-time PCR system (Applied Biosystems; Thermo Fisher Scientific, Inc.) and a SYBR Green PCR Master Mix (Applied Biosystems; Thermo Fisher Scientific, Inc.). Primers were obtained from Genotech (Shanghai, China) and the sequences are provided in Table I. GAPDH was used as the internal control. qPCR was performed using the following previous described cycling conditions (15): Initial denaturation at 95°C for 10 min, followed by 40 cycles

at 95°C for 15 sec and 60°C for 1 min. Each experiment was performed in triplicate. Gene transcript levels were calculated using the $2^{-\Delta\Delta Cq}$ method (16).

Statistical analysis. All experiments were performed at least three times. The results are expressed as the mean \pm standard deviations (SD). The data were analyzed using Student's t-test or one-way analysis of variance followed by least significance difference test. All analyses were performed with GraphPad Prism version 5.01 (GraphPad Software, Inc., La Jolla, CA, USA). $P < 0.05$ was considered to indicate a statistically significant difference.

Results

AdipoRon suppresses lipid accumulation in C3H10T1/2 cells without cytotoxicity. Following treatment for 24, 48 or 72 h, AdipoRon exhibited no significant effects on C3H10T1/2 cell viability at concentrations $\leq 40 \mu\text{M}$ (Fig. 1A). Therefore, 5-20 μM AdipoRon was used in all subsequent experiments. The antiadipogenic effects of AdipoRon treatment were not due to cytotoxicity. C3H10T1/2 cells were induced into adipocytes for 8 days in the presence or absence of various doses of AdipoRon (0-20 μM) to investigate the effects of AdipoRon on adipocyte differentiation. Based on the results of ORO staining, lipid accumulation in C3H10T1/2 cells was dose-dependently decreased by AdipoRon (Fig. 1B and C). Protein expression of the adiponectin receptors, AdipoR1 and AdipoR2, were detected in C3H10T1/2 cells (Fig. 1D).

Effects of AdipoRon on adipogenic transcription factors expression in C3H10T1/2 cells. Adipogenesis is controlled by numerous transcription factors; the present study examined the effects of various concentrations of AdipoRon (0-20 μM) on the mRNA expression levels C/EBP β , C/EBP α and PPAR γ , to investigate whether AdipoRon inhibits adipogenesis by down-regulating the expression of these adipogenic transcription factors. AdipoRon treatment significantly reduced the expression levels of the C/EBP β , C/EBP α and PPAR γ mRNAs in a dose-dependent manner (Fig. 2A-C). In addition, treatment with 20 μM AdipoRon was used to examine the expression levels of the C/EBP β , C/EBP α and PPAR γ proteins. The expression levels of the C/EBP β , C/EBP α and PPAR γ proteins were decreased by 20 μM AdipoRon compared with those of the controls (Fig. 2D).

Effects of AdipoRon on the expression levels of adipogenesis-related genes in C3H10T1/2 cells. As AdipoRon treatment significantly downregulated the expression of adipogenic transcription factors, the effects of various concentrations of AdipoRon on the expression levels of adipogenesis-related genes were also examined. AdipoRon exposure dose-dependently suppressed the mRNA expression of FAS, leptin, stearoyl-CoA desaturase (SCD)-1, adiponectin and FABP4 (Fig. 3A-E). Consistent with the changes in mRNA expression levels, 20 μM AdipoRon decreased the expression levels of the adiponectin and FABP4 proteins (Fig. 3F).

Effects of AdipoRon on AMPK phosphorylation in C3H10T1/2 cells. AdipoRon treatment significantly increased the levels

Table I. Primers used for reverse transcription quantitative polymerase chain reaction.

Gene	Primer sequence (5'→3')
C/EBP α	F: CGGGAACGCAACAACATCGC
C/EBP α	R: CGGTCATTGTCACTGGTCAACTC
C/EBP β	F: GTTTCGGGAGTTGATGCAATC
C/EBP β	R: AACAAACCCCGCAGGAACAT
PPAR γ	F: CATTTCGCATTCCTTTGAC
PPAR γ	R: CGCACTTTGGTATTCTTGGAG
FAS	F: CAAGTGTCCACCAACAAGCG
FAS	R: GGAGCGCAGGATAGACTCAC
FABP4	F: TGTGCGAAACTGAATTTCTCTGC
FABP4	R: GAGATCGGTCTGAGCCAGC
Adiponectin	F: GCCGCTTATGTGTATCGCTCAG
Adiponectin	R: GCCAGTGTCTGCCGTCATAATG
Leptin	F: TCGCCTTTCTCCTGATGACG
Leptin	R: GCAATCACACGGATGGCTTC
SCD-1	F: CGCTGGCACATCAACTTCAC
SCD-1	R: AGGAACTCAGAAGCCCAAAGC
GAPDH	F: TCAATGACAACCTTTGTCAAGCTCA
GAPDH	R: GTGGGTGGTCCAGGGTTTCTTACT

C/EBP, CAAT/enhancer binding protein; FABP, fatty acid binding protein; FAS, fatty acid synthase; PPAR, peroxisome proliferator-activated receptor; SCD-1, stearoyl-CoA desaturase 1.

of p-AMPK and p-ACC, which is the downstream target of AMPK (Fig. 4A-C). The AMPK activator AICAR was used as a positive control to further explore the association between the activation of AMPK signaling and the antiadipogenic effects mediated by AdipoRon. As shown in Fig. 4D and E, both AICAR (1 mM) and AdipoRon (10 and 20 μM) were able to activate the AMPK signaling pathway. Both AICAR and AdipoRon treatments were also able to significantly reduce lipid accumulation and the expression of adipogenesis-related genes, PPAR γ and adiponectin, in C3H10T1/2 cells (Fig. 4F-H).

Discussion

Obesity and its associated complications have become common problems worldwide. AdipoRon, an adiponectin receptor agonist, is a promising new orally administered drug for the treatment of obesity-related diabetes (12). However, its effects on obesity itself and adipogenesis have not yet been investigated. Results from the present study demonstrated that AdipoRon treatment inhibited the differentiation of C3H10T1/2 cells into adipocytes by downregulating the expression of adipogenic transcription factors and regulating the AMPK signaling pathway.

Abnormal adipogenesis and lipid accumulation have previously been associated with the development of obesity (2). A potential treatment strategy for obesity is to inhibit adipogenesis. The presents study results indicated that AdipoRon dose-dependently inhibited lipid accumulation in




Figure 1. Effects of AdipoRon on viability and adipocyte differentiation of C3H10T1/2 cells. (A) The effects of AdipoRon on the viability of C3H10T1/2 cells were examined at (A-a) 24, (A-b) 48 and (A-c) 72 h. (B) Images of the ORO staining at day 8 of adipocyte differentiation; (B-a) non-induced; (B-b) DM-induced adipocytes; DM-induced and treated with (B-c) 5 μ M, (B-d) 10 μ M or (B-e) 20 μ M AdipoRon; magnification, x200. (C) Lipid accumulation was determined by measuring the absorbance at 490 nm. (D) The protein expression levels of AdipoR1 and AdipoR2 in C3H10T1/2 cells. Results are presented as the mean \pm standard deviation; n=3; *P<0.05 vs. DM-induced control). AdipoR, adiponectin receptor; DM, differentiation medium; ORO, Oil Red O.




Figure 2. Effects of AdipoRon on the expression levels of adipogenic transcription factors in C3H10T1/2 cells. C3H10T1/2 cells were incubated with AdipoRon during adipogenesis. Reverse transcription-quantitative polymerase chain reaction was used to analyze mRNA expression, and western blotting was used to analyze protein expression at day 8 post-treatment. Effects of AdipoRon (0-20 μ M) on the mRNA expression levels of the (A) C/EBP β , (B) C/EBP α and (C) PPAR γ . (D) Effects of AdipoRon (20 μ M) on the protein expression levels of the C/EBP β , C/EBP α and PPAR γ . Results are presented as the mean \pm standard deviation; n=3; *P<0.05 vs. 0 μ M AdipoRon. C/EBP, CAAT/enhancer binding protein; PPAR, peroxisome proliferator-activated receptor.




Figure 3. Effects of AdipoRon on the expression levels of adipocyte-specific markers in C3H10T1/2 cells. C3H10T1/2 cells were incubated with AdipoRon during adipogenesis. Reverse transcription-quantitative polymerase chain reaction was used to analyze mRNA expression levels, and western blotting was used to analyze protein expression levels at day 8 post-treatment. The effects of different concentrations of AdipoRon on the mRNA expression levels of (A) FAS, (B) FABP4, (C) adiponectin, (D) leptin and (E) SCD-1. (F) The effects of 20 μ M AdipoRon on the protein expression levels of FABP4 and adiponectin. Results are presented as the mean \pm standard deviation; n=3; *P<0.05 vs. 0 μ M AdipoRon. FABP, fatty acid binding protein; FAS, fatty acid synthase; SCD-1, stearoyl-CoA desaturase 1.

C3H10T1/2 cells without cytotoxicity. Adipogenesis is coordinated by many transcription factors, such as C/EBPs and PPARs; AdipoRon exposure downregulated the expression levels of the C/EBP β mRNA, which subsequently suppressed the expression of the C/EBP α and PPAR γ mRNAs. PPAR γ and C/EBP α are major regulators of adipogenesis by activating the transcription of terminal adipocyte differentiation marker genes, including FABP4, FAS, leptin, SCD-1 and adiponectin (3-5); the expression levels of these mRNAs were demonstrated to be significantly decreased by AdipoRon in the present study. Therefore, AdipoRon treatment suppressed the expression of transcription factors during adipogenesis, which then inhibited adipocyte differentiation of C3H10T1/2 cells.

Obesity occurs when total energy intake exceeds total energy expenditure. AMPK is a metabolic energy sensor that senses nutritional stress for the purpose of regulating glucose and lipid metabolism and may be a potential target for the treatment of obesity (17). AMPK has been reported to

serve an important role in inhibiting adipogenesis in 3T3-L1 cells that have been induced by a number of different natural compounds (2,6,8,18-20). Activated AMPK inhibits the differentiation of adipocytes and the expression of pro-adipogenesis transcription factors PPAR γ and C/EBP α (21). AMPK phosphorylation induces the phosphorylation of its substrate, ACC, which functions to strictly regulate the enzymes that are involved in the fatty acid synthesis pathway that produces malonyl-CoA. Notably, p-ACC lacks the ability to synthesize fatty acids (22). AdipoRon was previously reported to activate AMPK signaling in muscle and liver of obese mice (12). To elucidate whether AdipoRon inhibited adipogenesis through the AMPK signaling pathway, the present study measured the levels of p-AMPK and p-ACC; AdipoRon treatment significantly increased the phosphorylation of AMPK and ACC. In addition, the AMPK activator AICAR was used as a positive control to further elucidate the association between the anti-adipogenesis ability of AdipoRon and the activation of the AMPK signaling pathway. AICAR was




Figure 4. Effects of AdipoRon on the phosphorylation of proteins in the AMPK signaling pathway. (A) C3H10T1/2 cells were treated with or without DM in the presence or absence of AdipoRon for 15 min. Protein expression levels of p-AMPK, AMPK and p-ACC were determined. (B and C) Densitometric analyses for (B) AMPK phosphorylation and (C) ACC phosphorylation from part (A). (D) C3H10T1/2 cells were treated with DM in the presence or absence of AdipoRon (10 or 20 μ M) or AICAR (1 mM) for 15 min. The expression levels of the p-AMPK and AMPK proteins were determined. (E) Densitometric analyses for AMPK phosphorylation from part (D). (F) Images of ORO staining at day 8 of adipocyte differentiation; (F-a) DM-induced adipocytes; (F-b) DM-induced adipocytes treated with 20 μ M AdipoRon; (F-c) DM-induced adipocytes treated with 1 mM AICAR; magnification, \times 200. (G) Lipid accumulation was determined by measuring the absorbance at 490 nm. (H) Effects of AdipoRon (20 μ M) or AICAR (1 mM) on the expression levels of the PPAR γ and adiponectin proteins. Results are presented as the mean \pm standard deviation; $n=3$; * $P<0.05$ vs. DM-induced control. ACC, acetyl-CoA carboxylase; AICAR, 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranoside; AMPK, adenosine monophosphate-activated protein kinase; DM, differentiation medium; ORO, Oil Red O; p, phosphorylated; PPAR, peroxisome proliferator-activated receptor.

previously reported to inhibit adipogenesis in 3T3-L1 adipocytes and restore metabolic alterations in a diet-induced mouse model of obesity (23-25). Results from the present study revealed that both AICAR and AdipoRon were able to activate the AMPK signaling pathway and markedly reduce lipid accumulation in C3H10T1/2 cells, which indicated that the anti-adipogenic effects of AdipoRon may be related to the activation of the AMPK signaling pathway. As C3H10T1/2 is a mouse embryonic mesenchymal stem cell line, the anti-adipogenesis effects of AdipoRon require further validation in human-sourced mesenchymal stem cell line.

In conclusion, AdipoRon, an adiponectin receptor agonist, suppressed adipocyte differentiation of C3H10T1/2 cells, probably by activating the AMPK signaling pathway and downregulating the expression of key regulators of adipogenesis. Therefore, AdipoRon may be a drug with the potential to prevent and treat obesity.

Acknowledgements

This study was supported by grants from the National Natural Science Foundation of China (grant nos. 81670098 and

81270572) and the National Basic Research 973 Program (grant no. 2013CB733701).

References

- Despres JP and Lemieux I: Abdominal obesity and metabolic syndrome. *Nature* 444: 881-887, 2006.
- Ko SC, Lee M, Lee JH, Lee SH, Lim Y and Jeon YJ: Dieckol, a phlorotannin isolated from a brown seaweed, *Ecklonia cava*, inhibits adipogenesis through AMP-activated protein kinase (AMPK) activation in 3T3-L1 preadipocytes. *Environ Toxicol Pharmacol* 36: 1253-1260, 2013.
- White UA and Stephens JM: Transcriptional factors that promote formation of white adipose tissue. *Mol Cell Endocrinol* 318: 10-14, 2010.
- Rosen ED and Spiegelman BM: PPARgamma: A nuclear regulator of metabolism, differentiation, and cell growth. *J Biol Chem* 276: 37731-37734, 2001.
- Wu Z, Rosen ED, Brun R, Hauser S, Adelmant G, Troy AE, McKeon C, Darlington GJ and Spiegelman BM: Cross-regulation of C/EBP alpha and PPAR gamma controls the transcriptional pathway of adipogenesis and insulin sensitivity. *Mol Cell* 3: 151-158, 1999.
- Lee JH, Kim T, Lee JJ, Lee KJ, Kim HK, Yun B, Jeon J, Kim SK and Ma JY: The herbal medicine KBH-1 inhibits fat accumulation in 3T3-L1 adipocytes and reduces high fat diet-induced obesity through regulation of the AMPK pathway. *PLoS One* 10: e0142041, 2015.
- Lin JT, Chen HM, Chiu CH and Liang YJ: AMP-activated protein kinase activators in diabetic ulcers: From animal studies to phase II drugs under investigation. *Expert Opin Investig Drugs* 23: 1253-1265, 2014.
- Kim SK and Kong CS: Anti-adipogenic effect of dioxinohydroeckol via AMPK activation in 3T3-L1 adipocytes. *Chem Biol Interact* 186: 24-29, 2010.
- Ceddia RB: The role of AMP-activated protein kinase in regulating white adipose tissue metabolism. *Mol Cell Endocrinol* 366: 194-203, 2013.
- Hotta K, Funahashi T, Arita Y, Takahashi M, Matsuda M, Okamoto Y, Iwahashi H, Kuriyama H, Ouchi N, Maeda K, *et al*: Plasma concentrations of a novel, adipose-specific protein, adiponectin, in type 2 diabetic patients. *Arterioscler Thromb Vasc Biol* 20: 1595-1599, 2000.
- Yamauchi T, Kamon J, Waki H, Terauchi Y, Kubota N, Hara K, Mori Y, Ide T, Murakami K, Tsuboyama-Kasaoka N, *et al*: The fat-derived hormone adiponectin reverses insulin resistance associated with both lipoatrophy and obesity. *Nat Med* 7: 941-946, 2001.
- Okada-Iwabu M, Yamauchi T, Iwabu M, Honma T, Hamagami K, Matsuda K, Yamaguchi M, Tanabe H, Kimura-Someya T, Shirouzu M, *et al*: A small-molecule AdipoR agonist for type 2 diabetes and short life in obesity. *Nature* 503: 493-499, 2013.
- Holland WL and Scherer PE: Cell Biology. Ronning after the adiponectin receptors. *Science* 342: 1460-1461, 2013.
- Lazra Y, Falach A, Frenkel L, Rozenberg K, Sampson S and Rosenzweig T: Autocrine/paracrine function of globular adiponectin: Inhibition of lipid metabolism and inflammatory response in 3T3-L1 adipocytes. *J Cell Biochem* 116: 754-766, 2015.
- Poudel B, Nepali S, Xin M, Ki HH, Kim YH, Kim DK and Lee YM: Flavonoids from *Triticum aestivum* inhibit adipogenesis in 3T3-L1 cells by upregulating the insig pathway. *Mol Med Rep* 12: 3139-3145, 2015.
- Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
- Cantó C, Gerhart-Hines Z, Feige JN, Lagouge M, Noriega L, Milne JC, Elliott PJ, Puigserver P and Auwerx J: AMPK regulates energy expenditure by modulating NAD⁺ metabolism and SIRT1 activity. *Nature* 458: 1056-1060, 2009.
- Liou CJ, Lai XY, Chen YL, Wang CL, Wei CH and Huang WC: Ginkgolide C suppresses adipogenesis in 3T3-L1 adipocytes via the AMPK signaling pathway. *Evid Based Complement Alternat Med* 2015: 298635, 2015.
- Lee YK, Lee WS, Hwang JT, Kwon DY, Surh YJ and Park OJ: Curcumin exerts antidifferentiation effect through AMPKalpha-PPAR-gamma in 3T3-L1 adipocytes and antiproliferatory effect through AMPKalpha-COX-2 in cancer cells. *J Agric Food Chem* 57: 305-310, 2009.
- Ahn J, Lee H, Kim S, Park J and Ha T: The anti-obesity effect of quercetin is mediated by the AMPK and MAPK signaling pathways. *Biochem Biophys Res Commun* 373: 545-549, 2008.
- Chen S, Li Z, Li W, Shan Z and Zhu W: Resveratrol inhibits cell differentiation in 3T3-L1 adipocytes via activation of AMPK. *Can J Physiol Pharmacol* 89: 793-799, 2011.
- Daval M, Fougelle F and Ferré P: Functions of AMP-activated protein kinase in adipose tissue. *J Physiol* 574: 55-62, 2006.
- Giri S, Rattan R, Haq E, Khan M, Yasmin R, Won JS, Key L, Singh AK and Singh I: AICAR inhibits adipocyte differentiation in 3T3L1 and restores metabolic alterations in diet-induced obesity mice model. *Nutr Metab (Lond)* 3: 31, 2006.
- Habinowski SA and Witters LA: The effects of AICAR on adipocyte differentiation of 3T3-L1 cells. *Biochem Biophys Res Commun* 286: 852-856, 2001.
- Lee H, Kang R, Bae S and Yoon Y: AICAR, an activator of AMPK, inhibits adipogenesis via the WNT/beta-catenin pathway in 3T3-L1 adipocytes. *Int J Mol Med* 28: 65-71, 2011.