

$\alpha 7$ nicotinic acetylcholine receptor agonist attenuates the cerebral injury in a rat model of cardiopulmonary bypass by activating the Akt/GSK3 β pathway

YINGJIE SUN^{1*}, DANDAN SONG^{1*}, MEI WANG¹, KEYAN CHEN² and TIEZHENG ZHANG¹

¹Department of Anesthesiology, General Hospital of Shenyang Military Area Command, Shenyang, Liaoning 110016;

²Department of Laboratory Animal Science, China Medical University, Shenyang, Liaoning 110122, P.R. China

Received September 29, 2016; Accepted July 18, 2017

DOI: 10.3892/mmr.2017.7600

Abstract. $\alpha 7$ nicotinic acetylcholine receptor ($\alpha 7$ nAChR) agonist treatment may provide a promising therapeutic effect for cerebral injuries. However, it is unclear whether the activation of $\alpha 7$ nAChR agonist may reduce cerebral injuries induced by cardiopulmonary bypass (CPB). A total of 96 male Sprague-Dawley rats were randomly divided into four groups (n=24/group): i) Sham operation group; ii) CPB group; iii) CPB + $\alpha 7$ nAChR agonist group; and iv) CPB + $\alpha 7$ nAChR agonist + $\alpha 7$ nAChR antagonist group. Following treatment, 24 rats from each group were sacrificed and the serum and hippocampal tissues were collected. The serum expression levels of S100 β , interleukin 6 and tumor necrosis factor α were evaluated by ELISA, hippocampal tissues were analyzed by histopathological examination using hematoxylin & eosin and terminal deoxynucleotidyl-transferase-mediated dUTP nick-end labeling (TUNEL) staining and Caspase 3 expression in the hippocampal tissues was evaluated by immunohistochemistry. In addition, Caspase 3, Akt and glycogen synthase kinase 3 β (GSK3 β), as well as phosphorylated (p)-Akt and (p)-GSK3 β were examined by western blot assay. The present study demonstrated that $\alpha 7$ nAChR agonist treatment was able to alleviate pathological damage and inhibit hippocampal cell apoptosis and inflammatory response. $\alpha 7$ nAChR agonist

treatment also increased the expression levels of p-Akt and p-GSK3 β , which indicated an upregulation in Akt/GSK3 β signaling. These data suggested that $\alpha 7$ nAChR agonist may provide a promising new therapeutic approach for cerebral injury caused by CPB.

Introduction

Cardiopulmonary bypass (CPB) is considered indispensable during heart operations, but the potential adverse effects on sensitive organs, such as the brain or the kidneys, cannot be ignored (1). In particular, many of the patients who undergo CPB surgery suffer from adverse cerebral outcomes, which may include stroke, postoperative cognitive dysfunction and transient ischemic attacks (2). The underlying molecular mechanism of cerebral injuries induced by CPB is unknown; however, the pathological changes may in part be due to micro-emboli and impaired cerebral perfusion, as well as cerebral ischemia and inflammatory damage (3,4).

It has been previously reported that CPB may initiate systemic inflammatory reaction syndrome (SIRS) owing to the blood comprehensive contact with non-biological materials (5); CPB may also activate cerebral inflammation in the presence of blood-brain barrier injury or disruption (3,6). Therefore, inflammatory responses serve important roles in the progression of cerebral injuries induced by CPB, and reducing inflammation would be of great benefit for CPB surgery of (5,7). For example, ulinastatin treatment exhibited neuroprotective effects on an animal model of CPB, possibly through beneficial effects on anti-inflammatory systems (8).

The cholinergic anti-inflammatory pathway (CAP) is an endogenous neural feedback regulation mechanism and can regulate peripheral inflammatory responses (9). Therefore, the physiological regulation of CAP has been used to treat infectious or inflammatory animal models (9-14). Stimulation of the efferent vagus nerve releases the important neurotransmitter acetylcholine, which acts through the $\alpha 7$ nicotinic acetylcholine receptor ($\alpha 7$ nAChR) expressed in the macrophages and the brain. Notably, it has been revealed that activation of $\alpha 7$ nAChR may effectively decrease the expression of proinflammatory cytokines and inhibit the inflammation process (10,15-18). In addition, the $\alpha 7$ nAChR agonist PHA568487 has been used to

Correspondence to: Professor Tiezheng Zhang, Department of Anesthesiology, General Hospital of Shenyang Military Area Command, 83 Wenhua Road, Shenyang, Liaoning 110016, P.R. China
E-mail: tzzhang@hotmail.com

Dr Keyan Chen, Department of Laboratory Animal Science, China Medical University, 77 Puhe Road, Shenyang, Liaoning 110122, P.R. China
E-mail: kyachen@cmu.edu.cn

*Contributed equally

Key words: $\alpha 7$ nicotinic acetylcholine receptor, cardiopulmonary bypass, apoptosis, Caspase 3, Akt/glycogen synthase kinase 3 β pathway

treat neuroinflammation following tibia fracture and endotoxemia in mice (15), as well as ischemic stroke injury (16) and brain injury in a subarachnoid hemorrhage model rats (19). Therefore, the $\alpha 7$ nAChR agonist may provide promising therapeutic effects for cerebral injuries. However, it is still unclear whether activation of the $\alpha 7$ nAChR agonist is able to reduce cerebral injuries induced by CPB.

The present study evaluated the therapeutic effects and the molecular mechanisms of the $\alpha 7$ nAChR agonist on CPB-induced brain injury in a rat model. The results indicated that the $\alpha 7$ nAChR agonist may effectively inhibit the inflammatory response and reduce apoptosis by activating the Akt/GSK3 β signaling pathway.

Materials and methods

Animals and ethical approval. A total of 96 adult male Sprague-Dawley rats (age, 8-9 weeks; weight, 350-450 g) were obtained from Shenyang Military Region General Hospital Laboratory Animal Center [Shenyang, China; license no. SCXK (Liao) 2012-00022012-0002]. Animals were housed at a constant temperature (22 \pm 1 $^{\circ}$ C), with 50% relative humidity and a 12-h light/dark cycle. The rats had access to food and autoclaved water *ad libitum*. All animal procedures were approved by the Animal Experiments Ethics Committee of the General Hospital of Shenyang Military Region (Shenyang, China).

CPB animal model establishment. CPB surgery was performed as previously reported (7), with minor modifications. Briefly, rats received an intraperitoneal (i.p.) injection of 10% chloral hydrate (300 mg/kg; Shanghai Ziyuan Pharmaceutical Co., Ltd., Shanghai, China) for anesthesia. Photopic oral intubation was performed using a 16 G intravenous (i.v.) catheter, and animals were mechanically ventilated with a small animal ventilator (settings: Frequency, 60 beats/min; tidal volume, 3 ml/kg; inspiratory to expiratory ratio, 1:1.5) connected to a monitor to observe the heart rate, oxygen saturation and rectal temperature of the rats. During surgery, anesthesia was maintained with i.v. injection of pipecuronium bromide (0.1 mg/kg; Hangzhou Minsheng Pharmaceutical Co., Ltd., Hangzhou, China).

The puncture site was sterilized with iodophor (Shandong Lierkang Disinfection Technology Co., Ltd., Dezhou, China), followed by exposure and puncture of the vein. Right femoral vein catheterization (24G) was performed to open the fluid path, which was transfused with 6% hydroxyethyl starch (Guangdong Jiabao Pharmaceutical Co., Ltd., Qingyuan, China) and connected to a microinfusion pump. The left femoral artery was catheterized (22G) and used to monitor blood pressure. Coccygeal artery catheterization (22G) and right internal jugular vein catheterization (18G) were performed to drain blood for CPB. The drainage tube, a homemade blood storage device, a constant peristaltic pump (Baoding Longer Precision Pump Co., Ltd., Baoding, China), silicone tubing (internal diameter, 4 mm) and a rat membrane oxygenator (Guangdong Kewei Medical Instrument Co. Ltd., Dongguan, China) were installed between the two puncture sites to establish the CPB circuit. Heparin sodium (300 IU/kg; Shenyang Haitong Pharmaceutical Co., Ltd., Shenyang, China)

was injected into the left femoral vein once the activated clotting time reached 480 sec.

CPB was performed with the membrane oxygenator to supply oxygen. The low-flow CPB velocity was 35 ml/kg/min, which was later increased to 100-120 ml/kg/min at full-flow bypass. To prevent air embolism, 1-2 ml of blood was retained in the blood storage device. Mean arterial pressure was maintained at >60 mmHg, partial CO₂ pressure at 35-45 mmHg, base excess at -3-3 mmol/l mmHg, pH at 7.35-7.45 and hematocrit at >0.25. Rats were treated with 2-20 μ g/100 g epinephrine hydrochloride (Wuhan Grand Pharmaceutical Group Co., Ltd., Wuhan, China) and fluids during surgery to maintain a stable circulation.

Groups and treatments. Rats were randomly divided into four groups (n=24/group): i) The Sham group (S group), in which intubation and mechanical ventilation were performed in the right femoral artery only and the right internal jugular vein was catheterized without bypass; ii) the CPB surgery group (C group), which received the CPB surgery aforementioned; iii) the $\alpha 7$ nAChR agonist group (P group), which received an i.p. injection of the $\alpha 7$ nAChR agonist PHA568487 (0.8 mg/kg; Tocris Bioscience; Bio-Techne, Minneapolis, MN, USA) 30 min prior to CPB establishment; and iv) the PHA568487 + $\alpha 7$ nAChR antagonist group (M group), which were also pretreated with PHA568487 (0.8 mg/kg) for 30 min, followed by i.p. injection of the $\alpha 7$ nAChR antagonist methyllycaconitine (MLA; 6 mg/kg; Sigma-Aldrich; Merck KGaA, Darmstadt, Germany). CPB surgery was performed 60 min after MLA injection.

Specimen collection and processing. Arterial and venous blood samples were collected prior to CPB (T0), upon completion of CPB surgery (T1), 2 h post-CPB (T2) and 6 h post-CPB (T3); subsequently, rats were sacrificed with 2% pentobarbital sodium (40 mg/kg by i.p. injection; Merck Sharp & Dohme, Shanghai, China). The systemic circulation system of the rats was infused with saline (250-400 ml), and the whole brain was collected on the ice and divided into two halves along the median sagittal line. The hippocampus was isolated from each of the two halves, the right half was fixed in 4% paraformaldehyde (PFA) at room temperature for 24 h, and the left side was stored at -80 $^{\circ}$ C for western blot analysis. Sera were separated by centrifugation at 1,000 x g for 10 min at 4 $^{\circ}$ C, and stored at -80 $^{\circ}$ C.

Histopathological assessment. Fixed hippocampal tissues were gradually dehydrated with ethanol and embedded in paraffin. Paraffin blocks were subsequently sectioned (5 μ m) and stained with a Hematoxylin & Eosin (H&E) staining kit (Nanjing Jiancheng Bioengineering Institute, Nanjing, China). Double-blind evaluation of hippocampal injury was performed by two expert pathologists. Images of the histopathological examination were captured by a light microscope (Olympus Corporation, Tokyo, Japan) at x400 magnification.

Tissue apoptosis assay. A terminal deoxynucleotidyl-transferase-mediated dUTP nick-end labeling (TUNEL) Assay kit (Shanghai Fusheng Industrial Co., Ltd., Shanghai, China) was used, according to the manufacturer's protocol, to determine the effects of $\alpha 7$ nAChR agonist treatment on apoptosis in the

fixed and mounted hippocampal sections. DAPI was used as a nuclear stain, with sections stained with 100 ng/ml DAPI for 5 min. Apoptotic rates were examined and images captured using a light microscope (Olympus Corporation, Japan) at a magnification of x400, and the densitometric scanning was finally analyzed by using the MetaMorph BX41 Image Analysis System (Olympus Corporation, Japan). A total of 5 images were captured randomly for each section at x400 magnification and integral optical density was calculated using Microscopic Image Analyzer (MetaMorph BX41 Image Analysis System). Percentages of TUNEL-positive cells above untreated controls were calculated as follows: %apoptosis = (number of TUNEL-positive cells / number of total cells) x 100.

Immunohistochemistry. To further determine the effects of $\alpha 7$ nAChR on apoptosis in the hippocampus, expression levels of the cellular apoptosis maker Caspase 3 was examined by immunohistochemical analysis. Briefly, dimethylbenzene was used to remove the paraffin from the hippocampal sections, followed by immersion in distilled water. Subsequently, antigen retrieval was conducted by placing the slides in a microwave in 10 mmol/l citrate buffer, pH 6.0, for 15 min. The slides were washed with 0.01 mmol/l PBS (HyClone; GE Healthcare Life Sciences, Logan, UT, USA) every 5 min for 3 times, followed by incubating in TBS + 0.3% H_2O_2 + 0.1% saponin at room temperature for 15 min to block the endogenous peroxidase. The slides were blocked with goat serum (Sigma-Aldrich; Merck KGaA) in TBS + 0.1% saponin for 20 min at room temperature, followed by incubating with polyclonal rabbit anti-Caspase 3 (1:300; ab13847; Abcam, Cambridge, UK) overnight at 4°C. The slides were incubated with biotin-conjugated secondary antibody (1:2,000; ab6720; Abcam) for 30 min, and 3,3'-diaminobenzidine stain (8 min at room temperature) was used to visualize Caspase 3 expression in the hippocampus. Images of Caspase 3 expression were captured with a light microscope (Olympus, Japan) at a magnification of 400x. A total of 5 images were captured randomly for each section at x400 magnification and integral optical density was calculated using Microscopic Image Analyzer (MetaMorph BX41 Image Analysis System).

ELISA determination of S100 β , tumor necrosis factor (TNF)- α and interleukin (IL)-6 levels in rat serum. Serum expression levels (in 100 μ l) of S100 β , TNF- α and IL-6 were determined by ELISA kits (S100 β , JM-E10007507; TNF- α , JM-E10009363; IL-6, JM-E10004387; TSZ Biosciences, San Francisco, CA, USA), according to the manufacturer's protocol. Optical density was measured at 450 nm using a Spectra Max M5 Microplate Reader (Molecular Devices, LLC, Sunnyvale, CA, USA).

Western blot analysis. The frozen hippocampal tissues (100 mg) were ground with a glass homogenizer and subsequently homogenized with Radioimmunoprecipitation Assay Buffer (1 ml; Beijing Solarbio Science & Technology Co., Ltd., Beijing, China) using an IKA T10 homogenizer (IKA-Werke GmbH & Co. KG, Staufen, Germany), followed by centrifugation at 12,000 x g for 15 min at 4°C. The supernatant was collected and protein quantification was performed by

bicinchoninic acid assay, and equal amounts of protein lysate (40 μ g) were separated by 12% SDS-PAGE. Proteins were transferred to nitrocellulose membranes in transfer buffer [12 mM Tris base, 96 mM glycine (pH 8.3) and 15% methanol]. Membranes were blocked for 2 h in TBS + 0.5% Tween-20 (TBST) with 5% bovine serum albumin (Sigma-Aldrich; Merck KGaA) at room temperature and subsequently probed with polyclonal rabbit anti-Akt (1:500; ab8805; Abcam), polyclonal rabbit anti-GSK3 β (1:1,000; ab115774; Abcam), monoclonal rabbit anti-p-Akt (1:500; 13038; Cell Signaling Technology, Inc. Danvers, MA, USA), polyclonal rabbit anti-p-GSK3 β ser9 (1:500; ab131097; Abcam), polyclonal rabbit anti-Caspase 3 (1:300; ab13847; Abcam) or monoclonal rabbit anti- β -actin antibody (1:100; 8457; CST, USA) overnight at 4°C. Membranes were washed with TBST buffer three times, followed by incubating with monoclonal goat anti-rabbit IgG horseradish peroxidase-conjugated secondary antibody (1:4,000; HSI01; Beijing TransGen Biotech Co., Ltd., Beijing, China) for 1 h at room temperature. ECL chemiluminescence was used to detect protein expression levels, which were visualized by scanning densitometry (170-8070 Molecular Imager ChemiDoc XRS System; Bio-Rad Laboratories, Inc. Hercules, CA, USA) using ImageJ Software (version 1.37; National Institutes of Health, Bethesda, MD, USA).

Statistical analysis. Quantitative data were expressed as the mean \pm standard deviation. Statistical analyses were performed with GraphPad Prism software, (version 6.00; GraphPad Software, Inc., La Jolla, CA, USA). Multiple comparisons were analyzed with one-way analysis of variance, followed by an appropriate multiple comparison test (Tukey's procedure). $P < 0.05$ was considered to indicate a statistically significant difference.

Results

$\alpha 7$ nAChR agonist alleviates pathological injury caused by CPB. To determine the protective effects of $\alpha 7$ nAChR agonist on the morphological alterations of the hippocampus, sections were evaluated at 6 h post-CPB, the T3 time point, by H&E staining. There was no detectable morphological damage to the hippocampal tissues in the S group (Fig. 1A), whereas clear cellular degeneration and abnormal cell arrangements were observed in the samples of CPB-injured rats (Fig. 1B), which indicated that the rat model of cerebral injury caused by CPB was successfully established. Following pretreatment with the $\alpha 7$ nAChR agonist, only a slight morphological change was observed in the P group as compared to those in the C group (Fig. 1C), which suggested that the $\alpha 7$ nAChR agonist may have alleviated the pathological injury of the CPB-injured rats; however, the typical vacuolated degenerations in hippocampal neurons were observed in those co-treated with $\alpha 7$ nAChR antagonist (Fig. 1D), indicating the protective effects of $\alpha 7$ nAChR agonist may be inhibited by MLA treatment. These results suggested that activation of $\alpha 7$ nAChR may alleviate CPB-induced pathological injury.

$\alpha 7$ nAChR agonist inhibits CPB-induced apoptosis of hippocampal neurons. To determine the effects of $\alpha 7$ nAChR agonist treatment on apoptosis in hippocampal neurons, the T3




Figure 1. $\alpha 7$ nAChR agonist pretreatment alleviates pathological injury in the hippocampus. Hippocampal tissues were examined at T3 by hematoxylin & eosin staining to evaluate the protective effects of $\alpha 7$ nAChR agonist. The hippocampal samples collected from the (A) Sham control group exhibited no evidence of morphological damage, whereas (B) hippocampal tissues of the rats with CPB model group exhibited signs of obvious cellular degeneration and abnormal cell arrangements. (C) Rats pretreated with $\alpha 7$ nAChR agonist prior to CPB surgery exhibited only slight morphological alterations compared with the CPB group; however, more obvious pathological changes were observed in rats (D) co-treated with the $\alpha 7$ nAChR antagonist. Magnification, x400; scale bar, 20 μ m. $\alpha 7$ nAChR, $\alpha 7$ nicotinic acetylcholine receptor; CPB, cardiopulmonary bypass; T3, 6 h post-CPB.

sections were also evaluated by TUNEL staining. Compared with the control neurons in the S group, the neurons in the C group exhibited typical signs of apoptosis (Fig. 2A); neuronal apoptosis appeared to be lower in the P and M groups when compared with the C group (Fig. 2A). To further determine the effects of the $\alpha 7$ nAChR agonist on hippocampal neuron apoptosis, the integrated OD average of apoptosis positive area was quantified in captured images from all experimental groups. Compared with the S group, apoptosis was significantly increased in CPB-injured rats in groups C, P and M, which suggested that hippocampal cell apoptosis may be induced following CPB surgery. Notably, a lower rate of neuronal apoptosis was observed in rats pretreated with the $\alpha 7$ nAChR agonist compared with the C group ($P < 0.05$; Fig. 2B); however, apoptosis was significantly increased in rats co-treated with the $\alpha 7$ nAChR antagonist compared with the P group (Fig. 2B). These results indicated that CPB-induced apoptosis of hippocampal neurons may be effectively reduced by pretreatment with the $\alpha 7$ nAChR agonist.

In the light of the inhibition of $\alpha 7$ nAChR agonist on apoptosis of hippocampal neurons, the protein expression levels of Caspase 3, a key downstream inducer of apoptosis (20), was evaluated by western blot assay. In the T0 and T1 tissue specimen, no significant differences were detected in Caspase 3 expression between any of the groups, which suggested that apoptosis was not induced at this period in time. Conversely, tissues at T2 and T3 exhibited increased Caspase 3 expression in the CPB-injured rats in groups C, P and M compared with expression in the S group rats ($P < 0.05$; Fig. 3A and B), which implied that apoptosis was activated 3-6 h post-CPB surgery. Caspase 3 expression was significantly decreased in P group rats following pretreatment with $\alpha 7$ nAChR agonist compared

with the C group ($P < 0.05$), whereas this effect was reversed in M group rats co-treated with the $\alpha 7$ nAChR antagonist ($P < 0.05$; Fig. 3A and B).

To confirm the location of Caspase 3 expression in the hippocampus, immunohistochemical analysis was used to determine the expression at T2, as the Caspase 3 expression reached a peak in the CPB-injured rats at T2 according to the western blotting data aforementioned. Caspase 3 expression was detected in the neurons of hippocampus (Fig. 3C), and Caspase 3 expression was significantly inhibited in the P group compared with the C group ($P < 0.05$; Fig. 3D), which was consistent with western blotting results. Therefore, these results indicated that the $\alpha 7$ nAChR agonist may effectively inhibit apoptosis in hippocampal neurons, which may partly be accomplished by suppressing the expression of Caspase 3.

$\alpha 7$ nAChR agonist pretreatment reduces serum levels of S100 β , TNF- α and IL-6 in CPB-injured rats. Serum expression levels of S100 β , TNF- α and IL-6 were measured to evaluate the inflammatory response in rats with CPB injury. Compared with the control S group, rats in the CPB groups C, P and M exhibited significantly increased levels of S100 β , TNF α and IL6 at experimental time points T1-T3 ($P < 0.05$; Fig. 4A-C, respectively), which was considered as an indicator of serious cerebral injury. The levels of S100 β , TNF α and IL6 were significantly decreased in the P group following pretreatment with $\alpha 7$ nAChR agonist compared with the expression levels in CPB model rats in the C group at the T1-T3 experimental time points ($P < 0.05$); however, rats in the M group exhibited an increase in serum expression levels compared with the P group ($P < 0.05$; Fig. 4A-C). The apparent improvement of




Figure 2. $\alpha 7$ nAChR agonist pretreatment inhibits neuronal apoptosis in the hippocampus. Hippocampal tissues at T3 were examined by TUNEL assay to evaluate the effects of $\alpha 7$ nAChR agonist on apoptosis. (A) Hippocampal neurons exhibited typical apoptosis, whereas a lower neuronal apoptosis can be observed after pretreatment of $\alpha 7$ nAChR agonist. Magnification, x400; scale bar, 20 μ m; red arrows indicated positive expressions. (B) Quantitative results of TUNEL assay from part A. Data are presented as the mean \pm standard deviation; n=24/group; *P<0.05 vs. S group; **P<0.05 vs. C group; ΔP<0.05 vs. P group. $\alpha 7$ nAChR, $\alpha 7$ nicotinic acetylcholine receptor; C group, CPB surgery only; CPB, cardiopulmonary bypass; M group, CBP + $\alpha 7$ nAChR agonist PHA568487 + $\alpha 7$ nAChR antagonist methyllycaconitine; P group, CBP + $\alpha 7$ nAChR agonist PHA568487; S group, Sham operation; T3, 6 h post-CPB.




Figure 3. $\alpha 7$ nAChR agonist pretreatment inhibits Caspase 3 protein expression in the hippocampus. (A) Western blot analysis for Caspase 3 in hippocampus at different time points in the different experiments groups. (B) Densitometric analysis Caspase 3 expression presented in (A); decreased expressions of Caspase 3 were observed in rats in the P group following pretreatment of $\alpha 7$ nAChR agonist compared with those in the C group at T2 and T3. (C) Immunohistochemistry for Caspase 3 in hippocampus at T2. Magnification, x400; scale bar, 20 μ m; red arrows indicate positive expressions. (D) Integrated OD average analysis indicated the decreased expression levels of Caspase 3 in P group rats pretreated with the $\alpha 7$ nAChR agonist compared with expression in the C group model rats. Data are presented as the mean \pm standard deviation; n=24/group; *P<0.05 vs. S group; **P<0.05 vs. C group; ΔP<0.05 vs. P group. $\alpha 7$ nAChR, $\alpha 7$ nicotinic acetylcholine receptor; C group, CPB surgery only; CPB, cardiopulmonary bypass; M group, CBP + $\alpha 7$ nAChR agonist PHA568487 + $\alpha 7$ nAChR antagonist methyllycaconitine; P group, CBP + $\alpha 7$ nAChR agonist PHA568487; S group, Sham operation; T0, prior to CPB; T1, upon completion of CPB; T2, 3 h post-CPB; T3, 6 h post-CPB.




Figure 4. $\alpha 7$ nAChR agonist pretreatment reduces the serum expression levels of S100 β , TNF- α and IL-6 in CPB-injured rats. The serum levels of (A) S100 β , (B) TNF- α and (C) IL-6 were measured to evaluate the anti-inflammation effects of $\alpha 7$ nAChR agonist on CPB-injured rats. Serum levels of S100 β , TNF α and IL6 were significantly decreased in P group rats following pretreatment with the $\alpha 7$ nAChR agonist compared with levels in the C group model rats. Data are presented as the mean \pm standard deviation; n=24/group; *P<0.05 vs. S group; #P<0.05 vs. C group; Δ P<0.05 vs. P group. $\alpha 7$ nAChR, $\alpha 7$ nicotinic acetylcholine receptor; C group, CPB surgery only; CPB, cardiopulmonary bypass; M group, CPB + $\alpha 7$ nAChR agonist PHA568487 + $\alpha 7$ nAChR antagonist methyllycaconitine; IL, interleukin; P group, CPB + $\alpha 7$ nAChR agonist PHA568487; S group, Sham operation; T0, prior to CPB; T1, upon completion of CPB; T2, 3 h post-CPB; T3, 6 h post-CPB; TNF- α , tumor necrosis factor α .




Figure 5. $\alpha 7$ nAChR agonist promotes expression of p-Akt and p-GSK3 β . (A) Western blot analysis for Akt and p-Akt in the hippocampus at T3. (B) p-Akt expression levels were significantly increased in the P group following pretreatment with the $\alpha 7$ nAChR agonist compared with expression in the C group model rats. (C) Western blot analysis for GSK3 β and p-GSK3 β in the hippocampus at T3. (D) p-GSK3 β expression levels were significantly increased in the P group following pretreatment with the $\alpha 7$ nAChR agonist compared in the C group model rats. Data are presented as the mean \pm standard deviation; n=24/group; *P<0.05 vs. S group; #P<0.05 vs. C group; Δ P<0.05 vs. P group. $\alpha 7$ nAChR, $\alpha 7$ nicotinic acetylcholine receptor; C group, CPB surgery only; CPB, cardiopulmonary bypass; GSK3 β , glycogen synthase kinase 3 β ; M group, CPB + $\alpha 7$ nAChR agonist PHA568487 + $\alpha 7$ nAChR antagonist methyllycaconitine; p, phosphorylated; P group, CPB + $\alpha 7$ nAChR agonist PHA568487; S group, Sham operation; T3, 6 h post-CPB.

inflammation suggested that $\alpha 7$ nAChR agonist pretreatment may have a beneficial effect on anti-inflammatory systems of the rat model of CPB.

$\alpha 7$ nAChR agonist pretreatment promotes phosphorylation of Akt and GSK3 β . To further explore the underlying mechanisms by which $\alpha 7$ nAChR agonist alleviated the cerebral injuries caused by CPB, Akt/GSK3 β pathway activation was examined

to determine the protective effects of $\alpha 7$ nAChR agonist on the CPB-injured rats, as the Akt/GSK3 β pathway was previously identified as a significant cell survival pathway (21). CPB rats in the C group exhibited a significant increase in the expression of p-Akt and p-GSK3 β compared with expression levels in the S group (P<0.05; Fig. 5); whereas the expression levels of p-Akt and p-GSK3 β were significantly increased in the P group following pretreatment with the $\alpha 7$ nAChR agonist,

compared with the S group and C group, which suggested that the $\alpha 7$ nAChR agonist may promote the phosphorylation of Akt and GSK3 β . Rats in the M group that were co-treated with the $\alpha 7$ nAChR antagonist exhibited a significant decrease in p-Akt and p-GSK3 β expression levels compared with the P group (Fig. 5). By contrast, no significant differences in the expression levels of total Akt and total GSK3 β were identified between the groups (Fig. 5), which implied that the $\alpha 7$ nAChR agonist did not affect the expressions of Akt and GSK3 β . Therefore, these results indicated that the $\alpha 7$ nAChR agonist may effectively upregulate the activation of the Akt/GSK3 β signaling pathway in the CPB-injured rats.

Discussion

Cerebral injury is a serious complication following the use of CPB in the cardiac surgery (2,22,23). This pathological lesion may be due to several aspects, including impaired cerebral perfusion and oxygenation, cerebral microemboli and SIRS (3,4). Among these factors, SIRS is one of great significance for CPB; therefore, minimizing SIRS is widely considered as a prerequisite strategy for inhibiting the inflammatory response (5). It is generally accepted that proinflammatory cytokines such as TNF- α may further increase the permeability of the blood-brain barrier and subsequently promote the invasion of inflammatory cytokines and immune cells (24). Results from the present study demonstrated that expression levels of the proinflammatory cytokines, including the TNF- α and IL-6, were significantly increased in CPB-injured rats, which was consistent with previous reports (7,25). Therefore, reducing proinflammatory cytokines levels may alleviate neuronal injury and improve functional recovery.

As a physiological regulation of the innate immune system, CAP has been widely used to inhibit the expression of proinflammatory cytokines for treating infectious and inflammatory diseases (16). According to that report, activation of the main regulatory target, $\alpha 7$ nAChR, may aid in the reduction of proinflammatory cytokines. Therefore, the present study hypothesized that the $\alpha 7$ nAChR agonist may effectively inhibit the serum levels of TNF- α and IL-6 in the CPB-injured rats, which suggested that the $\alpha 7$ nAChR agonist may provide a promising strategy for reducing SIRS post-CPB.

S100 β is regarded as a reliable serum maker of cerebral injury following the breakdown of the blood-brain barrier (26-28). In the present study, an increased serum level of S100 β was observed in the CPB model rats compared with normal rats, whereas the $\alpha 7$ nAChR agonist was able to decrease the serum level of S100 β , which demonstrated that the CPB model was successfully established and that the neuroprotective effects may be achieved by pretreatment with the $\alpha 7$ nAChR agonist.

Several previous reports suggested that the hippocampus is sensitive to ischemia and reperfusion injury caused by CPB (1,29). In the present study, clear pathological damage and an increase in cell apoptosis and Caspase 3 expression levels in the hippocampus were observed in the CPB-injured rats, which confirmed that pathological changes occur in the hippocampus following CPB surgery. Notably, these pathological injuries were effectively inhibited in rats pretreated

with the $\alpha 7$ nAChR agonist, which demonstrated the protective effects of the $\alpha 7$ nAChR agonist on CPB rats.

Additional studies have demonstrated that the Akt/GSK3 β pathway serves a central role in cell survival in a number of neurological diseases (30-32). In particular, activation of the Akt/GSK3 β pathway may attenuate apoptosis, which is closely related to the regulation of Caspase 3 expression (33-35). Based on the present results that demonstrated the inhibitory effects of the $\alpha 7$ nAChR agonist on apoptosis and Caspase 3 expression, activation of the Akt/GSK3 β pathway was further examined for the protective effects of $\alpha 7$ nAChR agonist on CPB. The results indicated that p-Akt and p-GSK3 β expressions were upregulated following $\alpha 7$ nAChR agonist pretreatment, which suggested that the $\alpha 7$ nAChR agonist may be able to inhibit hippocampal cell apoptosis by activating the Akt/GSK3 β pathway.

To further determine the protective effects of the $\alpha 7$ nAChR agonist on CPB, the $\alpha 7$ nAChR antagonist was concurrently administered in the present study. By contrast to pretreatment with the $\alpha 7$ nAChR agonist alone, co-treatment with the $\alpha 7$ nAChR antagonist resulted in significant increases in the serum levels of S100 β , TNF- α and IL-6, as well as the pathological damage, increased apoptosis and increased Caspase 3 expression, and a significant decrease in the expression levels of p-Akt and p-GSK3 β . These results further demonstrated the neuroprotective effects of $\alpha 7$ nAChR agonist on CPB-injured rats.

In conclusion, the present study demonstrated that the $\alpha 7$ nAChR agonist may reduce pathological damage and apoptosis in the hippocampus by upregulating Akt/GSK3 β signaling. The $\alpha 7$ nAChR agonist may provide a promising therapeutic approach for cerebral injury caused by CPB.

Acknowledgements

This study was supported by The Natural Science Foundation of China (grant nos. 81471121 and 3120175) and The Teaching Project of China Medical University (grant no. XZR20160036).

References

1. Salameh A and Dhein S: Strategies for pharmacological organoprotection during extracorporeal circulation targeting ischemia-reperfusion injury. *Front Pharmacol* 6: 296, 2015.
2. Roach GW, Kanchuger M, Mangano CM, Newman M, Nussmeier N, Wolman R, Aggarwal A, Marschall K, Graham SH and Ley C: Adverse cerebral outcomes after coronary bypass surgery. Multicenter study of perioperative ischemia research group and the ischemia research and education foundation investigators. *N Engl J Med* 335: 1857-1863, 1996.
3. van Harten AE, Scheeren TW and Absalom AR: A review of postoperative cognitive dysfunction and neuroinflammation associated with cardiac surgery and anaesthesia. *Anaesthesia* 67: 280-293, 2012.
4. Cao HJ, Sun YJ, Zhang TZ, Zhou J and Diao YG: Penhexylidine hydrochloride attenuates the cerebral injury in a rat model of cardiopulmonary bypass. *Can J Physiol Pharmacol* 91: 521-527, 2013.
5. Evora PR, Bottura C, Arcêncio L, Albuquerque AA, Evora PM and Rodrigues AJ: Key Points for curbing cardiopulmonary bypass inflammation. *Acta Cir Bras* 31 (Suppl 1): S45-S52, 2016.
6. Ouk T, Amr G, Azzaoui R, Delassus L, Fossaert E, Tailleux A, Bordet R and Modine T: Lipid-lowering drugs prevent neurovascular and cognitive consequences of cardiopulmonary bypass. *Vascul Pharmacol* 80: 59-66, 2016.

7. Zhou J, Zhou N, Wu XN, Cao HJ, Sun YJ, Zhang TZ, Chen KY and Yu DM: Role of the Toll-like receptor 3 signaling pathway in the neuroprotective effect of sevoflurane pre-conditioning during cardiopulmonary bypass in rats. *Mol Med Rep* 12: 7859-7868, 2015.
8. Wang X, Xue Q, Yan F, Li L, Liu J, Li S and Hu S: Ulinastatin as a neuroprotective and anti-inflammatory agent in infant piglets model undergoing surgery on hypothermic low-flow cardiopulmonary bypass. *Paediatr Anaesth* 23: 209-216, 2013.
9. Borovikova LV, Ivanova S, Zhang M, Yang H, Botchkina GI, Watkins LR, Wang H, Abumrad N, Eaton JW and Tracey KJ: Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin. *Nature* 405: 458-462, 2000.
10. Ulloa L: The vagus nerve and the nicotinic anti-inflammatory pathway. *Nat Rev Drug Discov* 4: 673-684, 2005.
11. Liu JS, Wei XD, Lu ZB, Xie P, Zhou HL, Chen YY, Ma JM and Yu LZ: Liang-Ge-San, a classic traditional Chinese medicine formula, protects against lipopolysaccharide-induced inflammation through cholinergic anti-inflammatory pathway. *Oncotarget* 7: 21222-21234, 2016.
12. Cheng Z, Li-Sha G, Jing-Lin Z, Wen-Wu Z, Xue-Si C, Xing-Xing C and Yue-Chun L: Protective role of the cholinergic anti-inflammatory pathway in a mouse model of viral myocarditis. *PLoS One* 9: e112719, 2014.
13. Koopman FA, Vosters JL, Roescher N, Broekstra N, Tak PP and Vervoordeldonk MJ: Cholinergic anti-inflammatory pathway in the non-obese diabetic mouse model. *Oral Dis* 21: 858-865, 2015.
14. Jiang Y, Li L, Liu B, Zhang Y, Chen Q and Li C: Vagus nerve stimulation attenuates cerebral ischemia and reperfusion injury via endogenous cholinergic pathway in rat. *PLoS One* 9: e102342, 2014.
15. Terrando N, Yang T, Ryu JK, Newton PT, Monaco C, Feldmann M, Ma D, Akassoglou K and Maze M: Stimulation of the $\alpha 7$ nicotinic acetylcholine receptor protects against neuroinflammation after tibia fracture and endotoxemia in mice. *Mol Med* 20: 667-675, 2015.
16. Han Z, Shen F, He Y, Degos V, Camus M, Maze M, Young WL and Su H: Activation of $\alpha 7$ nicotinic acetylcholine receptor reduces ischemic stroke injury through reduction of pro-inflammatory macrophages and oxidative stress. *PLoS One* 9: e105711, 2014.
17. Su X, Matthey MA and Malik AB: Requisite role of the cholinergic $\alpha 7$ nicotinic acetylcholine receptor pathway in suppressing Gram-negative sepsis-induced acute lung inflammatory injury. *J Immunol* 184: 401-410, 2010.
18. Terrando N, Eriksson LI, Ryu JK, Yang T, Monaco C, Feldmann M, Jonsson Fagerlund M, Charo IF, Akassoglou K and Maze M: Resolving postoperative neuroinflammation and cognitive decline. *Ann Neurol* 70: 986-995, 2011.
19. Duris K, Manaenko A, Suzuki H, Rolland WB, Krafft PR and Zhang JH: $\alpha 7$ nicotinic acetylcholine receptor agonist PNU-282987 attenuates early brain injury in a perforation model of subarachnoid hemorrhage in rats. *Stroke* 42: 3530-3536, 2011.
20. McComb S, Mulligan R and Sad S: Caspase-3 is transiently activated without cell death during early antigen driven expansion of CD8(+) T cells in vivo. *PLoS One* 5: e15328, 2010.
21. Wang G, Fang H, Zhen Y, Xu G, Tian J, Zhang Y, Zhang D, Zhang G, Xu J, Zhang Z, *et al*: Sulforaphane prevents neuronal apoptosis and memory impairment in diabetic rats. *Cell Physiol Biochem* 39: 901-907, 2016.
22. Salazar JD, Wityk RJ, Grega MA, Borowicz LM, Doty JR, Petrofski JA and Baumgartner WA: Stroke after cardiac surgery: Short- and long-term outcomes. *Ann Thorac Surg* 72: 1195-1202, 2001.
23. Vedel AG, Holmgaard F, Rasmussen LS, Paulson OB, Thomsen C, Danielsen ER, Langkilde A, Goetze JP, Lange T, Ravn HB and Nilsson JC: Perfusion Pressure Cerebral Infarct (PPCI) trial-the importance of mean arterial pressure during cardiopulmonary bypass to prevent cerebral complications after cardiac surgery: Study protocol for a randomised controlled trial. *Trials* 17: 247, 2016.
24. Han Z, Li L, Wang L, Degos V, Maze M and Su H: Alpha-7 nicotinic acetylcholine receptor agonist treatment reduces neuroinflammation, oxidative stress, and brain injury in mice with ischemic stroke and bone fracture. *J Neurochem* 131: 498-508, 2014.
25. Li YP, Huang J, Huang SG, Xu YG, Xu YY, Liao JY, Feng X, Zhang XG, Wang JH and Wang J: The compromised inflammatory response to bacterial components after pediatric cardiac surgery is associated with cardiopulmonary bypass-suppressed Toll-like receptor signal transduction pathways. *J Crit Care* 29: 312.e7-e13, 2014.
26. Yuan SM: S100 and S100 β : Biomarkers of cerebral damage in cardiac surgery with or without the use of cardiopulmonary bypass. *Rev Bras Cir Cardiovasc* 29: 630-641, 2014.
27. Einav S, Shoshan Y, Ovadia H, Matot I, Hersch M and Itshayek E: Early postoperative serum S100 beta levels predict ongoing brain damage after meningioma surgery: A prospective observational study. *Crit Care* 10: R141, 2006.
28. Zhang B, Yu JY, Liu LQ, Peng L, Chi F, Wu CH, Jong A, Wang SF, Cao H and Huang SH: Alpha7 nicotinic acetylcholine receptor is required for blood-brain barrier injury-related CNS disorders caused by *Cryptococcus neoformans* and HIV-1 associated comorbidity factors. *Bmc Infect Dis* 15: 352, 2015.
29. Chugani HT: Biological basis of emotions: Brain systems and brain development. *Pediatrics* 102 (5 Suppl E): S1225-S1229, 1998.
30. Zhu YM, Wang CC, Chen L, Qian LB, Ma LL, Yu J, Zhu MH, Wen CY, Yu LN and Yan M: Both PI3K/Akt and ERK1/2 pathways participate in the protection by dexmedetomidine against transient focal cerebral ischemia/reperfusion injury in rats. *Brain Res* 1494: 1-8, 2013.
31. Zhang HY, Zhang X, Wang ZG, Shi HX, Wu FZ, Lin BB, Xu XL, Wang XJ, Fu XB, Li ZY, *et al*: Exogenous basic fibroblast growth factor inhibits ER stress-induced apoptosis and improves recovery from spinal cord injury. *CNS Neurosci Ther* 19: 20-29, 2013.
32. Krafft PR, Altay O, Rolland WB, Duris K, Lekic T, Tang J and Zhang JH: $\alpha 7$ nicotinic acetylcholine receptor agonism confers neuroprotection through GSK-3 β inhibition in a mouse model of intracerebral hemorrhage. *Stroke* 43: 844-850, 2012.
33. Dong M, Hu N, Hua Y, Xu X, Kandadi MR, Guo R, Jiang S, Nair S, Hu D and Ren J: Chronic Akt activation attenuated lipopolysaccharide-induced cardiac dysfunction via Akt/GSK3 β -dependent inhibition of apoptosis and ER stress. *Biochim Biophys Acta* 1832: 848-863, 2013.
34. Pan JJ, Chang QS, Wang X, Son YO, Liu J, Zhang Z, Bi YY and Shi X: Activation of Akt/GSK3 β and Akt/Bcl-2 signaling pathways in nickel-transformed BEAS-2B cells. *Int J Oncol* 39: 1285-1294, 2011.
35. Hong Y, Shao A, Wang J, Chen S, Wu H, McBride DW, Wu Q, Sun X and Zhang J: Neuroprotective effect of hydrogen-rich saline against neurologic damage and apoptosis in early brain injury following subarachnoid hemorrhage: Possible role of the Akt/GSK3 β signaling pathway. *PLoS One* 9: e96212, 2014.