

# Identification of key genes in glioma CpG island methylator phenotype via network analysis of gene expression data

LIJUAN BO<sup>1\*</sup>, BO WEI<sup>2\*</sup>, ZHANFENG WANG<sup>2</sup>, DALIANG KONG<sup>3</sup>, ZHENG GAO<sup>2</sup> and ZHUANG MIAO<sup>2</sup>

Departments of <sup>1</sup>Infections, <sup>2</sup>Neurosurgery and <sup>3</sup>Orthopaedics, China-Japan Union Hospital of Jilin University, Changchun, Jilin 130033, P.R. China

Received March 6, 2017; Accepted August 16, 2017

DOI: 10.3892/mmr.2017.7834

**Abstract.** Gene expression data were analysed using bioinformatic tools to demonstrate molecular mechanisms underlying the glioma CpG island methylator phenotype (CIMP). A gene expression data set (accession no. GSE30336) was downloaded from Gene Expression Omnibus, including 36 CIMP<sup>+</sup> and 16 CIMP<sup>-</sup> glioma samples. Differential analysis was performed for CIMP<sup>+</sup> vs. CIMP<sup>-</sup> samples using the limma package in R. Functional enrichment analysis was subsequently conducted for differentially expressed genes (DEGs) using Database for Annotation, Visualization and Integration Discovery. Protein-protein interaction (PPI) networks were constructed for upregulated and downregulated genes with information from STRING. MicroRNAs (miRNAs) targeting DEGs were also predicted using WebGestalt. A total of 439 DEGs were identified, including 214 upregulated and 198 downregulated genes. The upregulated genes were involved in extracellular

matrix organisation, defence and immune response, collagen fibril organisation and regulation of cell motion and the downregulated genes in cell adhesion, sensory organ development, regulation of system process, neuron differentiation and membrane organisation. A PPI network containing 134 nodes and 314 edges was constructed from the upregulated genes, whereas a PPI network consisting of 85 nodes and 80 edges was obtained from the downregulated genes. miRNAs regulating upregulated and downregulated genes were predicted, including miRNA-124a and miRNA-34a. Numerous key genes associated with glioma CIMP were identified in the present study. These findings may advance the understanding of glioma and facilitate the development of appropriate therapies.

## Introduction

Malignant glioma is the most common central nervous system tumour in adults and is associated with significant morbidity and mortality (1). Gliomas are highly invasive and poorly respond to conventional treatments; therefore, further studies to support the development of therapy for them are warranted (2).

Alterations in methylation serve a critical role in the pathogenesis of numerous human malignancies, including gliomas (3). CpG island methylator phenotype (CIMP) has emerged as a distinct molecular subclass of tumours (4). It features extensive, coordinated hypermethylation at specific loci (5,6). Several key genes regulated by methylation have been previously identified. O<sup>6</sup>-methylguanine-DNA methyltransferase (MGMT), which is responsible for DNA repair, is associated with chemotherapy resistance (7). Previous studies indicated that epigenetic silencing of MGMT via promoter methylation serves an important role in the regulation of MGMT expression in gliomas (8). Bruna *et al* (9) demonstrated that the methylation of platelet-derived growth factor (PDGF)-B can dictate transforming growth factor- $\beta$  as an oncogenic factor to promote cell proliferation in human glioma. In addition, Wiencke *et al* (10) reported that methylation of the phosphatase and tensin homolog promoter defines low-grade gliomas and secondary glioblastoma. Mueller *et al* (11) also suggested that epigenetic dysregulation of runt-related transcription factor 3 and testin is involved in glioblastoma tumorigenesis. Abnormal DNA methylation of CD133 (12) and tumor protein 53 (13) is also observed in glioma. Additionally, Turcan *et al* (14) indicated that isocitrate dehydrogenase 1 mutation is sufficient to

---

*Correspondence to:* Dr Zhuang Miao, Department of Neurosurgery, China-Japan Union Hospital of Jilin University, 126 Xiantai Street, Changchun, Jilin 130033, P.R. China  
E-mail: miaozhuang99@163.com

\*Contributed equally

**Abbreviations:** CIMP, CpG island methylator phenotype; DEGs, differentially expressed genes; PPI, protein-protein interaction; MGMT, O<sup>6</sup>-methylguanine-DNA methyltransferase; PDGF, platelet-derived growth factor; GO, Gene Ontology; COL3A1, collagen type III  $\alpha$ 1; COL5A2, collagen type V  $\alpha$ 2; TIMP1, TIMP metalloproteinase inhibitor 1; COL5A1, collagen type V  $\alpha$ 1; VIM, vimentin; GRIA2, glutamate receptor ionotropic AMPA2; BMP2, bone morphogenetic protein 2; PRKX, protein kinase X-linked; MYC, v-myc avian myelocytomatosis viral oncogene homolog; TJP2, tight junction protein 2; PDGFRA, platelet-derived growth factor receptor  $\alpha$  polypeptide; IQGAP1, IQ motif-containing GTPase activating protein 1; ALCAM, activated leukocyte cell adhesion molecule

**Key words:** glioma, CpG island methylator phenotype, differentially expressed gene, functional enrichment analysis, protein-protein interaction network, microRNA

Table I. GO biological process terms enriched in the differentially expressed genes.

A, Upregulated genes			
GO term	Count	(%)	P-value
GO:0030198 extracellular matrix organization	13	5.676855895	2.62x10 <sup>-8</sup>
GO:0030199 collagen fibril organization	8	3.493449782	1.25x10 <sup>-7</sup>
GO:0002504 antigen processing and presentation of peptide or polysaccharide antigen via major histocompatibility complex class II	8	3.493449782	3.25x10 <sup>-7</sup>
GO:0009611 response to wounding	25	10.91703057	4.95x10 <sup>-7</sup>
GO:0006955 immune response	28	12.22707424	1.58x10 <sup>-6</sup>
GO:0043062 extracellular structure organization	13	5.676855895	3.55x10 <sup>-6</sup>
GO:0006952 defense response	25	10.91703057	6.64x10 <sup>-6</sup>
GO:0016064 immunoglobulin mediated immune response	8	3.493449782	1.05x10 <sup>-5</sup>
GO:0019724 B cell mediated immunity	8	3.493449782	1.34x10 <sup>-5</sup>
GO:0006954 inflammatory response	17	7.423580786	1.59x10 <sup>-5</sup>
B, Downregulated genes			
GO term	Count	(%)	P-value
GO:0007423 sensory organ development	11	5.882352941	2.00x10 <sup>-4</sup>
GO:0007155 cell adhesion	20	10.69518717	2.07x10 <sup>-4</sup>
GO:0022610 biological adhesion	20	10.69518717	2.10x10 <sup>-4</sup>
GO:0030182 neuron differentiation	15	8.021390374	2.97x10 <sup>-4</sup>
GO:0048666 neuron development	13	6.951871658	3.27x10 <sup>-4</sup>
GO:0044057 regulation of system process	12	6.417112299	5.55x10 <sup>-4</sup>
GO:0048592 eye morphogenesis	6	3.20855615	9.10x10 <sup>-4</sup>
GO:0051966 regulation of synaptic transmission, glutamatergic	4	2.139037433	1.07x10 <sup>-3</sup>
GO:0031175 neuron projection development	10	5.347593583	1.93x10 <sup>-3</sup>
GO:0015672 monovalent inorganic cation transport	11	5.882352941	2.48x10 <sup>-3</sup>

GO, Gene Ontology.

establish the glioma hypermethylator phenotype. However, the identification of glioma-CIMP (G-CIMP) tumours based on gene expression data has rarely been reported (15). In the present study, gene expression profiles of CIMP-positive (CIMP<sup>+</sup>) samples were compared with those of CIMP-negative (CIMP<sup>-</sup>) samples to identify differentially expressed genes (DEGs), which were further subjected to functional enrichment analysis and network analyses. The findings of the present study may extend the understanding of the molecular mechanisms of CIMP<sup>+</sup> glioma.

## Materials and methods

**Gene expression data.** A gene expression data set (accession no. GSE30336) was downloaded from Gene Expression Omnibus (14), including 36 CIMP<sup>+</sup> glioma and 16 CIMP<sup>-</sup> samples. Gene expression levels were measured using the GPL571 (HG-U133A\_2) Affymetrix Human Genome U133A 2.0 Array (Affymetrix; Thermo Fisher Scientific Inc., Waltham, MA, USA). Probe annotations were also acquired.

**Pretreatment and differential analysis.** Raw data were pre-treated with the Robust Multichip Average method using the Affy package of R ([www.bioconductor.org/packages/release/bioc/html/affy.html](http://www.bioconductor.org/packages/release/bioc/html/affy.html)). Differential analysis was performed for CIMP<sup>+</sup> vs. CIMP<sup>-</sup> using the limma package (16) of R. |Log (fold change)| >1.0 and P<0.05 were set as cut-offs for significant differential expression.

**Functional enrichment analysis.** The Gene Ontology (GO; [www.geneontology.org/](http://www.geneontology.org/)) database is a bioinformatics resource that can provide functional categorization and annotations for gene products via the use of structured, controlled vocabularies (17). The Kyoto Encyclopaedia of Genes and Genome (KEGG; [www.genome.jp/kegg](http://www.genome.jp/kegg)) is a database for systematic analysis of the functions of genes or proteins in several specific metabolic and regulatory pathways (18). Functional enrichment analyses of the GO and KEGG databases were conducted using the Database for Annotation, Visualization and Integration Discovery ([david.abcc.ncifcrf.gov/](http://david.abcc.ncifcrf.gov/)) (19). The statistical method for this was based on hypergeometric

distribution.  $P < 0.05$  was considered to indicate significant functions and pathways.

**Construction of protein-protein interaction (PPI) network.** Proteins work together to complete certain biological functions. Therefore, revealing PPI is useful in elucidating underlying molecular mechanisms. In the present study, PPI networks were constructed for upregulated and downregulated genes using information from STRING (20). Interactions with the required level of confidence (i.e., score  $> 0.4$ ) were retained in the network. The two networks were visualised using Cytoscape (21).

Proteins in the network were presented as 'nodes', and each pairwise protein interaction was represented by an undirected link and the 'degree' of a node corresponded to the number of interactions by the protein. 'Degree' was calculated for each node.

**Prediction of miRNAs and construction of the whole regulatory network.** Web-based Gene Set Enrichment Analysis Toolkit (WebGestalt; [www.webgestalt.org/option.php](http://www.webgestalt.org/option.php)) is a comprehensive and powerful analysis toolkit, which can be used for enrichment analysis and microRNA (miRNA)-target prediction by identifying miRNA-binding site motifs. In the present study, miRNAs regulating DEGs were predicted using WebGestalt (22). Count  $\geq 2$  was set as the cut-off for predicted miRNAs and the top 10 miRNAs were selected. Following miRNA-target gene network pairs using WebGestalt, PPI networks and miRNA-target gene interactions were integrated. Subsequently, the whole regulatory network was visualised using Cytoscape (21).

## Results

**DEGs.** A total of 41,335 genes were detected and 439 DEGs between CIMP<sup>+</sup> and CIMP<sup>-</sup> samples were identified, including 241 upregulated and 198 downregulated genes in CIMP<sup>+</sup> samples.

**Functional enrichment analysis.** The GO biological pathway terms enriched for the 241 upregulated genes in CIMP<sup>+</sup> samples could be divided into 13 clusters. They were associated with extracellular matrix organisation, defence response, immune response, collagen fibril organisation, and regulation of cell motion. The top 10 terms are listed in Table I.

The GO biological pathway terms enriched for the 198 downregulated genes were divided into 12 clusters. They were associated with cell adhesion, sensory organ development, system process regulation, neuron differentiation and membrane organisation. The top 10 terms are listed in Table I.

KEGG pathway enrichment analysis revealed 16 significant pathways associated with upregulated genes (Table II), including focal adhesion (hsa04510), asthma (hsa05310), ECM-receptor interaction (hsa04512), intestinal immune network for immunoglobulin A production (hsa04672) and allograft rejection (hsa05330). No significant pathway was identified for the downregulated genes.

**PPI networks of the DEGs.** A PPI network containing 134 nodes and 314 edges was constructed for the upregulated genes (Fig. 1), whereas a PPI network consisting of 85 nodes and 80 edges was obtained for the downregulated genes (Fig. 2).

Table II. Kyoto Encyclopaedia of Genes and Genome pathways enriched in the upregulated genes.

Term	Count	P-value
hsa04510:Focal adhesion	15	2.96x10 <sup>-6</sup>
hsa05310:Asthma	7	5.16x10 <sup>-6</sup>
hsa04512:Extracellular matrix-receptor interaction	10	6.52x10 <sup>-6</sup>
hsa04672:Intestinal immune network for immunoglobulin A production	8	1.09x10 <sup>-5</sup>
hsa05330:Allograft rejection	7	1.94x10 <sup>-5</sup>
hsa05322:Systemic lupus erythematosus	10	2.52x10 <sup>-5</sup>
hsa05332:Graft-vs.-host disease	7	3.12x10 <sup>-5</sup>
hsa04514:Cell adhesion molecules	11	4.31x10 <sup>-5</sup>
hsa04940:Type I diabetes mellitus	7	4.83x10 <sup>-5</sup>
hsa05416:Viral myocarditis	8	1.27x10 <sup>-4</sup>
hsa05320:Autoimmune thyroid disease	7	1.48x10 <sup>-4</sup>

Table III. Top 10 nodes with a high degree in the up and down-regulated protein-protein interaction network.

Gene	Degree
Upregulated	
COL3A1	22
COL5A2	18
TIMP1	16
COL5A1	16
VIM	15
ANXA2	13
S100A6	12
ANXA1	12
COL4A1	11
CXCL10	11
Downregulated	
GRIA2	6
BMP2	5
PRKX	5
MYC	5
TJP2	5
PDGFRA	5
DCX	4
SH3GL2	4
RTN1	4
ID1	4

PPI, protein-protein interaction.

The top ten nodes with a high degree in the up and down-regulated PPI networks are listed in Table III. The top five nodes in the network of upregulated genes were collagen type III  $\alpha 1$  (COL3A1), collagen type V  $\alpha 2$  (COL5A2), TIMP metalloproteinase inhibitor 1 (TIMP1), collagen type V  $\alpha 1$




Figure 1. PPI network for upregulated genes. The circular nodes represent the names of genes and lines represent the interactions between the genes. PPI, protein-protein interaction.




Figure 2. PPI network for downregulated genes. The circular nodes represent the name of genes and lines represent the interactions between the genes. PPI, protein-protein interaction.

(COL5A1) and vimentin (VIM). In the network of down-regulated genes, the top six nodes were glutamate receptor

ionotropic AMPA2 (GRIA2), bone morphogenetic protein 2 (BMP2), protein kinase X-linked (PRKX), v-myc avian




Figure 3. The distribution of upregulated genes in biological processes categories, as assessed by WebGestalt.




Figure 4. The distribution of downregulated genes in biological processes categories, as assessed by WebGestalt.

myelocytomatosis viral oncogene homolog (MYC), tight junction protein 2 (TJP2) and platelet-derived growth factor receptor  $\alpha$  polypeptide (PDGFRA).

*miRNA prediction and regulatory network analysis.* The distribution of upregulated and downregulated genes in biological processes was analysed using WebGestalt (Figs. 3 and 4,

Table IV. Predicted miRs targeting the differentially expressed genes.

A, Upregulated genes		
miR	Gene	Statistics
hsa_TTGCCT, miR-130a, miR-301, miR-130b	2	C=52; O=2; E=6.41; R=0.31; raw P=1.0000; adj P=1.0000
hsa_TTTGCAC, miR-19a, miR-19b	2	C=71; O=2; E=8.76; R=0.23; raw P=1.0000; adj P=1.0000
hsa_TGGTGCT, miR-29a, miR-29b, miR-29c	3	C=59; O=3; E=7.28; R=0.41; raw P=1.0000; adj P=1.0000
hsa_TGCCTTA, miR-124a	3	C=84; O=3; E=10.36; R=0.29; raw P=1.0000; adj P=1.0000
hsa_GTGCCTT, miR-506	5	C=105; O=5; E=12.95; R=0.39; raw P=1.0000; adj P=1.0000
hsa_ACATTC, miR-1, miR-206	3	C=61; O=3; E=7.52; R=0.40; raw P=1.0000; adj P=1.0000
hsa_ACACTCC, miR-122a	2	C=13; O=2; E=1.60; R=1.25; raw P=0.4895; adj P=1.0000
hsa_ATGTTTC, miR-494	2	C=28; O=2; E=3.45; R=0.58; raw P=1.0000; adj P=1.0000
hsa_GGGACCA, miR-133a, miR-133b	2	C=37; O=2; E=4.56; R=0.44; raw P=1.0000; adj P=1.0000
hsa_ATTCTTT, miR-186	2	C=45; O=2; E=5.55; R=0.36; raw P=1.0000; adj P=1.0000
B, Downregulated genes		
miR	Gene	Statistics
hsa_ACTGCCT, miR-34b	4	C=41; O=4; E=4.11; R=0.97; raw P=1.0000; adj P=1.0000
hsa_CACTGCC, miR-34a, miR-34c, miR-449	3	C=47; O=3; E=4.71; R=0.64; raw P=1.0000; adj P=1.0000
hsa_AAACCAC, miR-140	2	C=25; O=2; E=2.50; R=0.80; raw P=1.0000; adj P=1.0000
hsa_TAGCTTT, miR-9	2	C=31; O=2; E=3.11; R=0.64; raw P=1.0000; adj P=1.0000
hsa_TGCACTT, miR-519c, miR-519b, miR-519A	3	C=54; O=3; E=5.41; R=0.55; raw P=1.0000; adj P=1.0000
hsa_GTTAAAG, miR-302b	2	C=9; O=2; E=0.90; R=2.22; raw P=0.2255; adj P=1.0000
hsa_AACTGGA, miR-145	2	C=33; O=2; E=3.31; R=0.61; raw P=1.0000; adj P=1.0000
hsa_ACCAAAG, miR-9	2	C=67; O=2; E=6.71; R=0.30; raw P=1.0000; adj P=1.0000
hsa_TGCTGCT, miR-15a, miR-16, miR-15b, miR-195, miR-424, miR-497	2	C=86; O=2; E=8.61; R=0.23; raw P=1.0000; adj P=1.0000
hsa_CTGAGCC, miR-24	3	C=35; O=3; E=3.51; R=0.86; raw P=1.0000; adj P=1.0000

miR, microRNA; C, number of reference genes in the category; O, number of genes in the gene set and also in the category; E, expected number in the category; R, ratio of enrichment; raw P, P-value from hypergeometric test; adj P, adjusted P-value.

respectively). The regulatory miRNAs of DEGs were also predicted (Table IV). Among these predicted miRNAs, miRNA-506 and miR-34b targeted the most DEGs in the up- and downregulated regulatory networks. In the upregulated regulatory network, miRNA-506 (miR-506) regulated five upregulated genes: VIM, aryl hydrocarbon receptor, proteolipid protein 2, IQ motif-containing GTPase activating protein 1 (IQGAP1) and syndecan 4. In the downregulated regulatory network, miR-34b regulated four downregulated genes: Sex determining region Y-box 4, PDGFRA, activated leukocyte cell adhesion molecule (ALCAM) and MYC. All predicted miRNAs with their target DEG pairs are presented in the regulatory network (Fig. 5).

## Discussion

In the present study, a total of 439 DEGs were identified, including 241 upregulated and 198 downregulated genes. Functional enrichment analysis predicted that upregulated genes were associated with extracellular matrix organisation, defence response, immune response, collagen fibril organisation

and regulation of cell motion, whereas downregulated genes were associated with cell adhesion, sensory organ development, regulation of system process, neuron differentiation and membrane organisation. These findings are consistent with previous reports (23-26). Ulrich *et al* (27) pointed out that the mechanical rigidity of the extracellular matrix regulates the structure, motility and proliferation of glioma cells. Cell motion and cell adhesion were closely associated with the invasion of glioma cells.

In the present study, a PPI network containing 134 nodes and 314 edges was constructed for upregulated genes, whereas a PPI network consisting of 85 nodes and 80 edges was also obtained for downregulated genes. The top five nodes in the network of upregulated genes were COL3A1, COL5A2, TIMP1, COL5A1 and VIM. TIMP1, as an inhibitor of matrix metalloproteinases, can promote cell proliferation and may have anti-apoptotic function (28). Groft *et al* (29) reported the differential expression and localisation of TIMP-1 and TIMP-4 in human gliomas and suggested that they may contribute to the pathophysiology of human malignant gliomas. In addition, Aaberg-Jessen *et al* (30) demonstrated



dataset used. Additionally, experimental or data verification for the DEGs identified in glioma CIMP was not conducted, and in future, samples should be divided into different CIMPs based on methylation analysis to conduct the experimental validation.

In conclusion, several key genes were identified in glioma CIMP, some of which (TIMP1, VIM, BMP2, c-MYC and PDGFRA) may be viewed as potential markers or therapeutic targets for gliomas. In addition, relevant miRNAs, such as miR-124a and miR-34a that regulate genes involved in gliomas were also detected. These findings may provide helpful guidance to reveal molecular mechanisms underlying glioma CIMP.

## References

- Kim TY, Zhong S, Fields CR, Kim JH and Robertson KD: Epigenomic profiling reveals novel and frequent targets of aberrant DNA methylation-mediated silencing in malignant glioma. *Cancer Res* 66: 7490-7501, 2006.
- Pan D, Wei X, Liu M, Feng S, Tian X, Feng X and Zhang X: Adenovirus mediated transfer of p53, GM-CSF and B7-1 suppresses growth and enhances immunogenicity of glioma cells. *Neurol Res* 32: 502-509, 2010.
- Christensen BC, Smith AA, Zheng S, Koestler DC, Houseman EA, Marsit CJ, Wiemels JL, Nelson HH, Karagas MR, Wrensch MR, *et al*: DNA methylation, isocitrate dehydrogenase mutation, and survival in glioma. *J Natl Cancer Inst* 103: 143-153, 2011.
- Noushmehr H, Weisenberger DJ, Diefes K, Phillips HS, Pujara K, Berman BP, Pan F, Pelloski CE, Sulman EP, Bhat KP, *et al*: Identification of a CpG island methylator phenotype that defines a distinct subgroup of glioma. *Cancer Cell* 17: 510-522, 2010.
- Fang F, Turcan S, Rimner A, Kaufman A, Giri D, Morris LG, Shen R, Seshan V, Mo Q, Heguy A, *et al*: Breast cancer methylomes establish an epigenomic foundation for metastasis. *Sci Transl Med* 3: 75ra25, 2011.
- Cheng YW, Pincas H, Bacolod MD, Schemmann G, Giardina SF, Huang J, Barral S, Idrees K, Khan SA, Zeng Z, *et al*: CpG island methylator phenotype associates with low-degree chromosomal abnormalities in colorectal cancer. *Clin Cancer Res* 14: 6005-6013, 2008.
- Weller M, Stupp R, Reifenberger G, Brandes AA, Van Den Bent MJ, Wick W and Hegi ME: MGMT promoter methylation in malignant gliomas: Ready for personalized medicine? *Nat Rev Neurol* 6: 39-51, 2010.
- Hegi ME, Liu L, Herman JG, Stupp R, Wick W, Weller M, Mehta MP and Gilbert MR: Correlation of O<sup>6</sup>-methylguanine methyltransferase (MGMT) promoter methylation with clinical outcomes in glioblastoma and clinical strategies to modulate MGMT activity. *J Clin Oncol* 26: 4189-4199, 2008.
- Bruna A, Darken RS, Rojo F, Ocaña A, Peñuelas S, Arias A, Paris R, Tortosa A, Mora J, Baselga J and Seoane J: High TGFbeta-Smad activity confers poor prognosis in glioma patients and promotes cell proliferation depending on the methylation of the PDGF-B gene. *Cancer Cell* 11: 147-160, 2007.
- Wiencke JK, Zheng S, Jelluma N, Tihan T, Vandenberg S, Tamgüney T, Baumber R, Parsons R, Lamborn KR, Berger MS, *et al*: Methylation of the PTEN promoter defines low-grade gliomas and secondary glioblastoma. *Neuro Oncol* 9: 271-279, 2007.
- Mueller W, Nutt CL, Ehrlich M, Riemenschneider MJ, Von Deimling A, Van Den Boom D and Louis DN: Down-regulation of RUNX3 and TES by hypermethylation in glioblastoma. *Oncogene* 26: 583-593, 2007.
- Yi JM, Tsai HC, Glöckner SC, Lin S, Ohm JE, Easwaran H, James CD, Costello JF, Riggins G, Eberhart CG, *et al*: Abnormal DNA methylation of CD133 in colorectal and glioblastoma tumors. *Cancer Res* 68: 8094-8103, 2008.
- Amatya VJ, Naumann U, Weller M and Ohgaki H: TP53 promoter methylation in human gliomas. *Acta Neuropathol* 110: 178-184, 2005.
- Turcan S, Rohle D, Goenka A, Walsh LA, Fang F, Yilmaz E, Campos C, Fabius AW, Lu C, Ward PS, *et al*: IDH1 mutation is sufficient to establish the glioma hypermethylator phenotype. *Nature* 483: 479-483, 2012.
- Baysan M, Bozdogan S, Cam MC, Kotliarova S, Ahn S, Walling J, Killian JK, Stevenson H, Meltzer P and Fine HA: G-cimp status prediction of glioblastoma samples using mRNA expression data. *PLoS One* 7: e47839, 2012.
- Diboun I, Wernisch L, Orengo CA and Koltzenburg M: Microarray analysis after RNA amplification can detect pronounced differences in gene expression using limma. *BMC Genomics* 7: 252, 2006.
- Gene Ontology Consortium, Blake JA, Dolan M, Drabkin H, Hill DP, Li N, Sitnikov D, Bridges S, Burgess S, Buza T, *et al*: Gene Ontology annotations and resources. *Nucleic Acids Res* 41 (Database issue): D530-D535, 2013.
- Kanehisa M and Goto S: KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res* 28: 27-30, 2000.
- Huang DW, Sherman BT, Tan Q, Collins JR, Alvord WG, Roayaei J, Stephens R, Baseler MW, Lane HC and Lempicki RA: The DAVID gene functional classification tool: A novel biological module-centric algorithm to functionally analyze large gene lists. *Genome Biol* 8: R183, 2007.
- Szklarczyk D, Franceschini A, Kuhn M, Simonovic M, Roth A, Minguez P, Doerks T, Stark M, Muller J, Bork P, *et al*: The STRING database in 2011: Functional interaction networks of proteins, globally integrated and scored. *Nucleic Acids Res* 39 (Database issue): D561-D568, 2011.
- Smoot ME, Ono K, Ruscheinski J, Wang PL and Ideker T: Cytoscape 2.8: New features for data integration and network visualization. *Bioinformatics* 27: 431-432, 2011.
- Zhang B, Kirov S and Snoddy J: WebGestalt: An integrated system for exploring gene sets in various biological contexts. *Nucleic Acids Res* 33 (Web server issue): W741-W748, 2005.
- Yao ZQ and Lu YC: Research on molecular mechanism of human glioma and its clinical application. *Chin J Cancer Biother* 15: 90-94, 2008.
- Huse JT and Holland EC: Targeting brain cancer: Advances in the molecular pathology of malignant glioma and medulloblastoma. *Nat Rev Cancer* 10: 319-331, 2010.
- Clubb BH and Shivers RR: Extracellular matrix regulates microfilament and vinculin organization in C6-glioma cells. *Acta Neuropathol* 91: 31-40, 1996.
- Bauke AC, Sasse S, Matzat T and Klämbt C: A transcriptional network controlling glial development in the *Drosophila* visual system. *Development* 142: 2184-2193, 2015.
- Ulrich TA, de Juan Pardo EM and Kumar S: The mechanical rigidity of the extracellular matrix regulates the structure, motility, and proliferation of glioma cells. *Cancer Res* 69: 4167-4174, 2009.
- Lee SJ, Yoo HJ, Bae YS, Kim HJ and Lee ST: TIMP-1 inhibits apoptosis in breast carcinoma cells via a pathway involving pertussis toxin-sensitive G protein and c-Src. *Biochem Biophys Res Commun* 312: 1196-1201, 2003.
- Groft LL, Muzik H, Rewcastle NB, Johnston RN, Knäuper V, Lafleur MA, Forsyth PA and Edwards DR: Differential expression and localization of TIMP-1 and TIMP-4 in human gliomas. *Br J Cancer* 85: 55-63, 2001.
- Aaberg-Jessen C, Christensen K, Offenberg H, Bartels A, Dreehsen T, Hansen S, Schröder HD, Brünner N and Kristensen BW: Low expression of tissue inhibitor of metalloproteinases-1 (TIMP-1) in glioblastoma predicts longer patient survival. *J Neurooncol* 95: 117-128, 2009.
- Crocker M, Ashley S, Giddings I, Petrik V, Hardcastle A, Aherne W, Pearson A, Bell BA, Zacharoulis S and Papadopoulos MC: Serum angiogenic profile of patients with glioblastoma identifies distinct tumor subtypes and shows that TIMP-1 is a prognostic factor. *Neuro Oncol* 13: 99-108, 2011.
- Yamasaki T, Seki N, Yamada Y, Yoshino H, Hidaka H, Chiyomaru T, Nohata N, Kinoshita T, Nakagawa M and Enokida H: Tumor suppressive microRNA-138 contributes to cell migration and invasion through its targeting of vimentin in renal cell carcinoma. *Int J Oncol* 41: 805-817, 2012.
- Satelli A and Li S: Vimentin in cancer and its potential as a molecular target for cancer therapy. *Cell Mol Life Sci* 68: 3033-3046, 2011.
- Lee J, Son MJ, Woolard K, Donin NM, Li A, Cheng CH, Kotliarova S, Kotliarov Y, Walling J, Ahn S, *et al*: Epigenetic-mediated dysfunction of the bone morphogenetic protein pathway inhibits differentiation of glioblastoma-initiating cells. *Cancer Cell* 13: 69-80, 2008.
- Liu C, Tian G, Tu Y, Fu J, Lan C and Wu N: Expression pattern and clinical prognostic relevance of bone morphogenetic protein-2 in human gliomas. *Jpn J Clin Oncol* 39: 625-631, 2009.

36. Wang J, Wang H, Li Z, Wu Q, Lathia JD, McLendon RE, Hjelmeland AB and Rich JN: c-Myc is required for maintenance of glioma cancer stem cells. *PLoS One* 3: e3769, 2008.
37. Jensen NA, Pedersen KM, Lihme F, Rask L, Nielsen JV, Rasmussen TE and Mitchelmore C: Astroglial c-Myc overexpression predisposes mice to primary malignant gliomas. *J Biol Chem* 278: 8300-8308, 2003.
38. Puputti M, Tynninen O, Sihto H, Blom T, Mäenpää H, Isola J, Paetau A, Joensuu H and Nupponen NN: Amplification of KIT, PDGFRA, VEGFR2, and EGFR in gliomas. *Mol Cancer Res* 4: 927-934, 2006.
39. Giannini C, Sarkaria JN, Saito A, Uhm JH, Galanis E, Carlson BL, Schroeder MA and James CD: Patient tumor EGFR and PDGFRA gene amplifications retained in an invasive intracranial xenograft model of glioblastoma multiforme. *Neuro Oncol* 7: 164-176, 2005.
40. Silber J, Lim DA, Petritsch C, Persson AI, Maunakea AK, Yu M, Vandenberg SR, Ginzinger DG, James CD, Costello JF, *et al*: miR-124 and miR-137 inhibit proliferation of glioblastoma multiforme cells and induce differentiation of brain tumor stem cells. *BMC Med* 6: 14, 2008.
41. Fowler A, Thomson D, Giles K, Maleki S, Mreich E, Wheeler H, Leedman P, Biggs M, Cook R, Little N, *et al*: miR-124a is frequently down-regulated in glioblastoma and is involved in migration and invasion. *Eur J Cancer* 47: 953-963, 2011.
42. Kuroda S, Fukata M, Nakagawa M, Fujii K, Nakamura T, Ookubo T, Izawa I, Nagase T, Nomura N, Tani H, *et al*: Role of IQGAP1, a target of the small GTPases Cdc42 and Rac1, in regulation of E-cadherin-mediated cell-cell adhesion. *Science* 281: 832-835, 1998.
43. Noritake J, Watanabe T, Sato K, Wang S and Kaibuchi K: IQGAP1: A key regulator of adhesion and migration. *J Cell Sci* 118: 2085-2092, 2005.
44. Li Y, Guessous F, Zhang Y, Dipierro C, Kefas B, Johnson E, Marcinkiewicz L, Jiang J, Yang Y, Schmittgen TD, *et al*: MicroRNA-34a inhibits glioblastoma growth by targeting multiple oncogenes. *Cancer Res* 69: 7569-7576, 2009.
45. Heinrich MC, Corless CL, Duensing A, McGreevey L, Chen CJ, Joseph N, Singer S, Griffith DJ, Haley A, Town A, *et al*: PDGFRA activating mutations in gastrointestinal stromal tumors. *Science* 299: 708-710, 2003.
46. Silber J, Jacobsen A, Ozawa T, Harinath G, Holland EC, Sander C and Huse JT: Abstract B15: Repression of PDGFRA-targeting miR-34a promotes tumorigenesis in proneural malignant gliomas. *Cancer Res* 72 (Suppl 2): B15, 2012.
47. Swart GW: Activated leukocyte cell adhesion molecule (CD166/ALCAM): Developmental and mechanistic aspects of cell clustering and cell migration. *Eur J Cell Biol* 81: 313-321, 2002.