

# Isorhamnetin prevents H<sub>2</sub>O<sub>2</sub>-induced oxidative stress in human retinal pigment epithelial cells

JING WANG<sup>1,2</sup>, HUI-MIN GONG<sup>3</sup>, HUI-HUI ZOU<sup>2</sup>, LING LIANG<sup>2</sup> and XIN-YI WU<sup>1</sup>

<sup>1</sup>Department of Ophthalmology, Qi Lu Hospital of Shandong University, Jinan, Shandong 250012; <sup>2</sup>Department of Ophthalmology, Dezhou People's Hospital, Dezhou, Shandong 253014; <sup>3</sup>Ophthalmic Center, Qingdao Municipal Hospital, Qingdao, Shandong 266000, P.R. China

Received April 15, 2016; Accepted March 24, 2017

DOI: 10.3892/mmr.2017.7916

**Abstract.** Isorhamnetin, a 3-O-methylated metabolite of quercetin, exhibits antioxidant effects. However, to the best of our knowledge, no study to date has focused on the effects of isorhamnetin on retinal pigment epithelium (RPE) cells, and its underlying molecular mechanisms. Therefore, the present study aimed to examine the potential protective effect of isorhamnetin against oxidative stress in human RPE cells. The results demonstrated that pretreatment of RPE cells with isorhamnetin significantly protected cell viability against oxidative stress. In addition, isorhamnetin pretreatment inhibited hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>)-induced reactive oxygen species (ROS) production and caspase-3 activation in RPE cells. Furthermore, isorhamnetin pretreatment significantly increased the phosphorylation of phosphoinositide 3-kinase (PI3K) and AKT serine/threonine kinase 1 (Akt) in RPE cells exposed to H<sub>2</sub>O<sub>2</sub>, compared with cells treated with H<sub>2</sub>O<sub>2</sub> alone. Taken together, the present results demonstrated that isorhamnetin protected human RPE cells from oxidative stress-induced cell death, and this effect was associated with activation of the PI3K/Akt signaling pathway. Thus, isorhamnetin may be considered as a potential antioxidant useful for the prevention of age-related macular degeneration.

## Introduction

Age-related macular degeneration (AMD) is a leading cause of visual impairment in the elderly in developed countries (1). Epidemiological studies have demonstrated that the prevalence of AMD in the aged population is increasing steadily every year (2). The retinal pigment epithelium (RPE), which forms a blood-retinal barrier between the neural retina and

choriocapillaris, serves a pathophysiological role in the process of AMD (3). Oxidative stress is crucial in the development of AMD (4,5). Previous studies have reported that RPE cells are susceptible to oxidative damage, and oxidative stress from hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) leads to RPE cell death by causing preferential damage to mitochondrial DNA (6,7). Thus, strategies for protecting RPE cells against oxidative damage may be helpful for preventing the progression of AMD.

Isorhamnetin is one of the major active components isolated from herbal medicinal plants, such as *Hippophae rhamnoides* and *Ginkgo biloba* (8). Previous studies have indicated that isorhamnetin exhibits anti-inflammatory and antitumor effects, and that it protects ventricular myocytes from ischemia and reperfusion injury (9-11). In addition, isorhamnetin has been reported to exhibit antioxidant properties (12-14). Choi (15) reported that isorhamnetin reverses H<sub>2</sub>O<sub>2</sub>-induced growth inhibition and exhibits scavenging activity against intracellular reactive oxygen species (ROS) in mouse-derived C2C12 myoblasts (15). However, the effects of isorhamnetin on RPE cells and the underlying molecular mechanism remain unclear. Therefore, the aim of the present study was to examine the effect of isorhamnetin against oxidative stress in human RPE cells.

## Materials and methods

**Reagents and antibodies.** Isorhamnetin (purity >98%) was purchased from Shanghai Tongtian Biotechnology Co., Ltd. (Shanghai, China). Antibodies against phosphoinositide 3-kinase (PI3K; 1:2,000; cat no. sc-365290), phosphorylated (p)-PI3K (1:2,000; cat no. sc-293115), AKT serine/threonine kinase 1 (Akt; 1:2,000; cat no. sc-5298), p-Akt (1:2,000; cat no. sc-52940), GAPDH (1:1,000; cat no. sc-365062) and horseradish peroxidase-conjugated secondary antibodies (1:3,000; cat no. sc-2370) were from Santa Cruz Biotechnology, Inc. (Dallas, TX, USA). All other chemicals and reagents were purchased from Sigma-Aldrich (Merck KGaA, Darmstadt, Germany).

**Cell culture.** The human RPE cell line ARPE-19 was purchased from the American Type Culture Collection (Manassas, VA, USA) and cultured in DMEM/Nutrient Mixture F-12 (Gibco; Thermo Fisher Scientific, Inc., Waltham, MA, USA),

**Correspondence to:** Dr Xin-Yi Wu, Department of Ophthalmology, Qi Lu Hospital of Shandong University, 107 Culture West Road, Jinan, Shandong 250012, P.R. China  
E-mail: wuxyoph@163.com

**Key words:** age-related macular degeneration, isorhamnetin, retinal pigment epithelium cells, oxidative stress

supplemented with 10% fetal bovine serum (Sigma-Aldrich; Merck KGaA) and 1% penicillin/streptomycin, in a 37°C incubator under a humidified atmosphere containing 5% CO<sub>2</sub>.

**Cell viability assay.** Cell viability was evaluated using an MTT assay. Briefly, ARPE-19 cells (4x10<sup>4</sup> cells/well) were pretreated with various concentrations of isorhamnetin (25, 50 and 100 µM) for 24 h prior to exposure to 250 µM H<sub>2</sub>O<sub>2</sub> for 24 h. The medium was then replaced with fresh medium containing 0.5 mg/ml MTT for 4 h at 37°C. The MTT formazan crystals were dissolved in dimethyl sulfoxide, and the absorbance was measured at 450 nm with a microplate reader (Omega Bio-Tek, Inc., Norcross, GA, USA). The relative cell viability was defined as the absorbance of treated cells divided by that of the control untreated cells.

**Measurement of intracellular ROS generation.** Intracellular ROS production was measured using the 2',7'-dichlorodihydrofluorescein diacetate (H<sub>2</sub>DCFDA) probe (Sigma-Aldrich; Merck KGaA). Briefly, following treatment, ARPE-19 cells were washed twice with PBS, and then incubated with 10 mM H<sub>2</sub>DCFDA for 30 min at 37°C in the dark. Then, the cells were harvested and fluorescence of the resulting dichlorofluorescein (DCF) was measured using an F-2500 spectrophotometer (Hitachi, Ltd., Tokyo, Japan) at 488 nm excitation and 525 nm emission. The results were expressed as the fluorescence intensity of DCF of each sample relative to the control.

**Caspase-3 activity assay.** Caspase-3 activity was detected using a caspase-3 cellular activity assay kit (Merck KGaA), as per the manufacturer's instructions. Briefly, following treatment, ARPE-19 cells were harvested and then suspended in the cell lysis buffer. Cell lysates were incubated in the presence or absence of 5 µl Asp-Glu-Val-Asp-p-nitroanilide at 37°C for 1 h. Following washing, the fluorescence released by active caspase-3 was measured using a microplate reader at 405 nm wavelength.

**Western blot analysis.** ARPE-19 cells were harvested, washed twice with PBS, and lysed in radioimmunoprecipitation assay buffer containing 1 mM phenylmethylsulfonyl fluoride (both from Shenneng Bocai Biotechnology Co., Ltd., Shanghai, China) on ice for 10 min. Cell lysates were centrifuged at 13,000 x g for 15 min at 4°C. The supernatants were collected, and the protein content of each lysate was measured by BCA protein assay (Pierce; Thermo Fisher Scientific, Inc.). Protein (30 µg/lane) was separated by electrophoresis on a 10% SDS-polyacrylamide gel and transferred electrophoretically to a nitrocellulose membrane (Bio-Rad Laboratories, Inc., Hercules, CA, USA). The membranes were blocked in 5% non-fat dry milk and TBST buffer (5 mM Tris-HCl, pH 7.4, 136 mM NaCl and 0.1% Tween-20) for 1 h at room temperature, and incubated with specific antibodies (p-PI3K, PI3K, p-Akt, Akt, and GAPDH) overnight at 4°C. Then, the membranes were incubated with horseradish peroxidase-conjugated secondary antibodies (Santa Cruz Biotechnology, Inc.) for another 1 h at room temperature. The blots were visualized with an enhanced chemiluminescence system (GE Healthcare Life Sciences, Chalfont, UK) and grey intensity analysis




Figure 1. Effect of isorhamnetin treatment on ARPE-19 cell viability. ARPE-19 cells were pretreated with various concentrations of isorhamnetin (0, 25, 50 and 100 µM) for 24 h, and cell viability was determined by MTT assay. Data are expressed as mean ± standard deviation of three independent experiments.

was performed using Image-Pro Plus 6.0 software (Media Cybernetics, Inc., Rockville, MD, USA).

**Statistical analysis.** The SPSS software package (version 17.0; SPSS Inc., Chicago, IL, USA) was used for statistical analysis. Data are expressed as mean ± standard deviation of three independent experiments performed in triplicate. Significant differences were determined using a Student's t-test or one-way analysis of variance followed by Tukey's post hoc test.  $P < 0.05$  was considered to indicate a statistically significant difference.

## Results

**Isorhamnetin is not cytotoxic in ARPE-19 cells.** First, the potential cytotoxicity of isorhamnetin was examined on human ARPE-19 cells using the MTT assay. As demonstrated in Fig. 1, isorhamnetin did not significantly affect the viability of ARPE-19 cells in doses up to 100 µM. Therefore, 25–100 µM of isorhamnetin was used for treatments in the following experiments.

**Isorhamnetin treatment protects human RPE cells against oxidative stress.** The effect of isorhamnetin on the cell viability of H<sub>2</sub>O<sub>2</sub>-treated ARPE-19 cells was examined by MTT assay. The results demonstrated that H<sub>2</sub>O<sub>2</sub> treatment resulted in a significant decrease in cell viability compared with untreated cells, whereas pretreatment with isorhamnetin significantly reversed this decrease in a dose-dependent manner (Fig. 2).

**Isorhamnetin treatment inhibits H<sub>2</sub>O<sub>2</sub>-induced ROS production in human RPE cells.** Increased ROS levels result in oxidative stress and are hypothesized to be critical in the pathogenesis of AMD. Therefore, the effect of isorhamnetin on ROS generation was examined in ARPE-19 cells exposed to H<sub>2</sub>O<sub>2</sub>. As indicated in Fig. 3, treatment with H<sub>2</sub>O<sub>2</sub> for 24 h dramatically increased the production of ROS compared with control untreated cells. However, pretreatment with isorhamnetin significantly suppressed the H<sub>2</sub>O<sub>2</sub>-induced ROS production in a dose-dependent manner (Fig. 3).

**Isorhamnetin treatment inhibits H<sub>2</sub>O<sub>2</sub>-induced caspase-3 activity in human RPE cells.** Caspase-3 is a main executor and




Figure 2. Isorhamnetin treatment protects human retinal pigment epithelium cells against oxidative stress. ARPE-19 cells were pretreated with various concentrations of isorhamnetin (0, 25, 50 and 100 μM) for 24 h, prior to exposure to 250 μM H<sub>2</sub>O<sub>2</sub> for 24 h. Cell viability was determined by MTT assay. Data are expressed as mean ± standard deviation of three independent experiments. \*P<0.05 vs. control untreated group; #P<0.05 vs. H<sub>2</sub>O<sub>2</sub> group. H<sub>2</sub>O<sub>2</sub>, hydrogen peroxide.




Figure 3. Isorhamnetin treatment inhibits H<sub>2</sub>O<sub>2</sub>-induced ROS production in human retinal pigment epithelium cells. ARPE-19 cells were pretreated with various concentrations of isorhamnetin (0, 25, 50 and 100 μM) for 24 h prior to exposure to 250 μM H<sub>2</sub>O<sub>2</sub> for 24 h. Intracellular ROS production was measured using a 2',7'-dichlorodihydrofluorescein diacetate probe. Data are expressed as mean ± standard deviation of three independent experiments. \*P<0.05 vs. control untreated group; #P<0.05 vs. H<sub>2</sub>O<sub>2</sub> group. H<sub>2</sub>O<sub>2</sub>, hydrogen peroxide.




Figure 4. Isorhamnetin treatment inhibits H<sub>2</sub>O<sub>2</sub>-induced caspase-3 activation in retinal pigment epithelium cells. ARPE-19 cells were pretreated with various concentrations of isorhamnetin (0, 25, 50 and 100 μM) for 24 h prior to exposure to 250 μM H<sub>2</sub>O<sub>2</sub> for 24 h. Caspase-3 activity was measured using a caspase-3 cellular activity assay kit. Data are expressed as mean ± standard deviation of three independent experiments. \*P<0.05 vs. control untreated group; #P<0.05 vs. H<sub>2</sub>O<sub>2</sub> group. H<sub>2</sub>O<sub>2</sub>, hydrogen peroxide.

an established marker of cell apoptosis, so caspase-3 activity was measured as a surrogate for cell death initiation (16). As




Figure 5. Isorhamnetin treatment results in PI3K/Akt pathway activation in retinal pigment epithelium cells. ARPE-19 cells were pretreated with various concentrations of isorhamnetin (0, 25, 50 and 100 μM) for 24 h prior to exposure to 250 μM H<sub>2</sub>O<sub>2</sub> for 24 h. Protein expression levels of p-PI3K, total PI3K, p-Akt, total Akt and GAPDH (as a loading control) were examined by western blotting. (A) Representative images. (B) Quantitative analysis of p-PI3K/PI3K ratio. (C) Quantitative analysis of p-Akt/Akt ratio. Data are expressed as mean ± standard deviation of three independent experiments. \*P<0.05 vs. control untreated group; #P<0.05 vs. H<sub>2</sub>O<sub>2</sub> group. PI3K, phosphoinositide 3-kinase; Akt, AKT serine/threonine kinase 1; H<sub>2</sub>O<sub>2</sub>, hydrogen peroxide; p, phosphorylated.

demonstrated in Fig. 4, H<sub>2</sub>O<sub>2</sub> treatment significantly increased caspase-3 activity, compared with untreated control. This H<sub>2</sub>O<sub>2</sub>-induced increase in caspase-3 activity, however, was dramatically attenuated by isorhamnetin pretreatment in a dose-dependent manner (Fig. 4).

*Isorhamnetin treatment activates the PI3K/Akt pathway in human RPE cells.* Activation of the PI3K/Akt signaling pathway is important in protecting RPE cells from oxidative stress (17). Therefore, the hypothesis that the PI3K/Akt pathway may be involved in the protective effects of isorhamnetin on



H<sub>2</sub>O<sub>2</sub>-induced cell death was tested in ARPE-19 cells. As demonstrated in Fig. 5, phosphorylation of PI3K and Akt were significantly decreased by H<sub>2</sub>O<sub>2</sub> treatment, compared with the control group, as evident by the decreased p-PI3K/PI3K and p-Akt/Akt ratios. However, pretreatment with isorhamnetin for 24 h markedly increased phosphorylation of both PI3K and Akt in ARPE-19 cells exposed to H<sub>2</sub>O<sub>2</sub>, compared with cells treated with H<sub>2</sub>O<sub>2</sub> alone (Fig. 5).

## Discussion

The present study focused on the effect of isorhamnetin on oxidative stress in human RPE cells. The results provide the first evidence that pretreatment of RPE cells with isorhamnetin significantly protected cell viability against oxidative stress. Furthermore, isorhamnetin inhibited H<sub>2</sub>O<sub>2</sub>-induced ROS production and caspase-3 activation in ARPE-19 cells. This protective effect of isorhamnetin may be mediated through PI3K/Akt signaling in ARPE-19 cells.

H<sub>2</sub>O<sub>2</sub> is a well-established model of oxidative stress in RPE cells because of its rapid membrane permeability (18). A vast array of studies have demonstrated that oxidative stress induced by chemical oxidants, such as H<sub>2</sub>O<sub>2</sub>, leads to RPE damage and cell death (5,19,20). For these reasons, H<sub>2</sub>O<sub>2</sub> was selected in the present study as an inducer of oxidative stress in human RPE cells to investigate the effect of isorhamnetin. The results demonstrated that H<sub>2</sub>O<sub>2</sub> treatment resulted in a significant decrease in viability of ARPE-19 cells. However, pretreatment with isorhamnetin significantly reversed this decrease in a dose-dependent manner, implying that isorhamnetin exhibits a protective effect against oxidative stress in RPE cells.

Oxidative damage is important in the pathogenesis of AMD. ROS levels are increased in retinas of patients with AMD, resulting in oxidization of key cellular components in local RPE cells and severe RPE damage (21). It has been reported that isorhamnetin pretreatment blocks tertiary-butyl hydroperoxide-induced ROS production in hepatocytes (12). Sun *et al* (22) reported that isorhamnetin significantly decreases ROS generation in H9c2 cardiomyocytes exposed to H<sub>2</sub>O<sub>2</sub>. In accordance with these previous reports, the present results demonstrated that H<sub>2</sub>O<sub>2</sub> treatment significantly increased intracellular ROS production in human RPE cells compared with untreated cells, and that this effect was significantly inhibited by isorhamnetin pretreatment. These results suggest that isorhamnetin protected ARPE-19 cells from H<sub>2</sub>O<sub>2</sub>-induced cell damage through its antioxidant activity.

Caspase-3 activation is an initiative process of cell apoptosis (23). It has been reported that exposure of RPE cells to increased ROS results in mitochondrial DNA damage and induces caspase-3 activation (24). In the present study, pretreatment with isorhamnetin was demonstrated to significantly suppress H<sub>2</sub>O<sub>2</sub>-induced caspase-3 activation, compared with cells treated with H<sub>2</sub>O<sub>2</sub> alone. These data suggest that isorhamnetin protected ARPE-19 cells from H<sub>2</sub>O<sub>2</sub>-induced cell damage via its anti-apoptotic effect.

There is substantial evidence that the PI3K/Akt signaling pathway is involved in regulating the antioxidant function in RPE cells (25-27). Akt becomes phosphorylated in a PI3K-dependent manner in response to oxidants, such as H<sub>2</sub>O<sub>2</sub>, in human RPE cells, and treatment with a PI3K inhibitor

(LY294002) blocks H<sub>2</sub>O<sub>2</sub>-mediated Akt phosphorylation and significantly enhances caspase-associated RPE cell death (28); this result is different to that observed in the present study which, may be due to the length of H<sub>2</sub>O<sub>2</sub> exposure. In the study by Yang *et al* (28), when RPE cells were exposed to H<sub>2</sub>O<sub>2</sub> for 15 min, p-Akt levels increased. However, in the present study, cells were exposed to H<sub>2</sub>O<sub>2</sub> for 24 h, and the phosphorylation of PI3K and Akt significantly decreased. A recent study reported that isorhamnetin attenuates atherosclerosis by suppressing apoptosis in THP-1-derived macrophages via PI3K/Akt activation (29). In the present study, it was demonstrated that pretreatment with isorhamnetin significantly enhanced phosphorylation of PI3K and Akt in RPE cells exposed to H<sub>2</sub>O<sub>2</sub>, compared to cells exposed to H<sub>2</sub>O<sub>2</sub> alone. These data suggest that isorhamnetin may have protected human RPE cells from oxidative stress-induced cell death, through activation of the PI3K/Akt signaling pathway.

In summary, the present report demonstrated that isorhamnetin protected human RPE cells from oxidative stress-induced cell death, which may be associated with the activation of PI3K/Akt signaling. Thus, isorhamnetin may be considered as a potential antioxidant agent towards the prevention of AMD.

## References

1. Congdon NG, Friedman DS and Lietman T: Important causes of visual impairment in the world today. *JAMA* 290: 2057-2060, 2003.
2. Fletcher EC and Scholl HP: Ophthalmic disease in the ageing society. *Ophthalmology and the Ageing Society*: 1-9, 2013.
3. Curcio CA, Zanzottera EC, Messinger JD, Ach T, Smith T and Freund KB: Retinal pigment epithelium (RPE) transdifferentiation and death in age-related macular degeneration (AMD), seen in the Project MACULA grading system. *Invest Ophthalmol Vis Sci* 56: 893-893, 2015.
4. Beatty S, Koh HH, Phil M, Henson D and Boulton M: The role of oxidative stress in the pathogenesis of age-related macular degeneration. *Surv Ophthalmol* 45: 115-134, 2000.
5. Jarrett SG and Boulton ME: Consequences of oxidative stress in age-related macular degeneration. *Mol Aspects Med* 33: 399-417, 2012.
6. Liang FQ and Godley BF: Oxidative stress-induced mitochondrial DNA damage in human retinal pigment epithelial cells: A possible mechanism for RPE aging and age-related macular degeneration. *Exp Eye Res* 76: 397-403, 2003.
7. Drobek-Słowik M, Karczewicz D and Safranow K: The potential role of oxidative stress in the pathogenesis of the age-related macular degeneration (AMD). *Postepy Hig Med Dosw (Online)* 61: 28-37, 2007 (In Polish).
8. Park JC, Young HS, Yu YB and Lee JH: Isorhamnetin sulphate from the leaves and stems of oenanthе javanica in korea. *Planta Med* 61: 377-378, 1995.
9. Chen TL, Zhu GL, Wang JA, Zhang GD, Liu HF, Chen JR, Wang Y and He XL: Protective effects of isorhamnetin on apoptosis and inflammation in TNF- $\alpha$ -induced HUVECs injury. *Int J Clin Exp Pathol* 8: 2311-2320, 2015.
10. Saud SM, Young MR, Jones-Hall YL, Ileva L, Evbuomwan MO, Wise J, Colburn NH, Kim YS and Bobe G: Chemopreventive activity of plant flavonoid isorhamnetin in colorectal cancer is mediated by oncogenic Src and  $\beta$ -catenin. *Cancer Res* 73: 5473-5484, 2013.
11. Zhang N, Pei F, Wei H, Zhang T, Yang C, Ma G and Yang C: Isorhamnetin protects rat ventricular myocytes from ischemia and reperfusion injury. *Exp Toxicol Pathol* 63: 33-38, 2011.
12. Yang JH, Shin BY, Han JY, Kim MG, Wi JE, Kim YW, Cho IJ, Kim SC, Shin SM and Ki SH: Isorhamnetin protects against oxidative stress by activating Nrf2 and inducing the expression of its target genes. *Toxicol Appl Pharmacol* 274: 293-301, 2014.
13. Kong CS, Kim JA, Qian ZJ, Kim YA, Lee JI, Kim SK, Nam TJ and Seo Y: Protective effect of isorhamnetin 3-O- $\beta$ -d-glucopyranoside from *Salicornia herbacea* against oxidation-induced cell damage. *Food Chem Toxicol* 47: 1914-1920, 2009.

14. Bouhlef I, Skandrani I, Nefatti A, Valenti K, Ghedira K, Mariotte AM, Hininger-Favier I, Laporte F, Dijoux-Franca MG and Chekir-Ghedira L: Antigenotoxic and antioxidant activities of isorhamnetin 3-O neohesperidoside from *Acacia salicina*. *Drug Chem Toxicol* 32: 258-267, 2009.
15. Choi YH: The cytoprotective effect of isorhamnetin against oxidative stress is mediated by the upregulation of the Nrf2-dependent HO-1 expression in C2C12 myoblasts through scavenging reactive oxygen species and ERK inactivation. *Gen Physiol Biophys* 35: 145-154, 2016.
16. Jänicke RU, Sprengart ML, Wati MR and Porter AG: Caspase-3 is required for DNA fragmentation and morphological changes associated with apoptosis. *J Biol Chem* 273: 9357-9360, 1998.
17. Faghiri Z and Bazan NG: PI3K/Akt and mTOR/p70S6K pathways mediate neuroprotection D1-induced retinal pigment epithelial cell survival during oxidative stress-induced apoptosis. *Exp Eye Res* 90: 718-725, 2010.
18. Kim MH, Chung J, Yang JW, Chung SM, Kwag NH and Yoo JS: Hydrogen peroxide-induced cell death in a human retinal pigment epithelial cell line, ARPE-19. *Korean J Ophthalmol* 17: 19-28, 2003.
19. Hanneken A, Lin FF, Johnson J and Maher P: Flavonoids protect human retinal pigment epithelial cells from oxidative-stress-induced death. *Invest Ophthalmol Vis Sci* 47: 3164-3177, 2006.
20. Sreekumar PG, Kannan R, Yaung J, Spee CK, Ryan SJ and Hinton DR: Protection from oxidative stress by methionine sulfoxide reductases in RPE cells. *Biochem Biophys Res Commun* 334: 245-253, 2005.
21. Cai J, Nelson KC, Wu M, Sternberg P Jr and Jones DP: Oxidative damage and protection of the RPE. *Prog Retin Eye Res* 19: 205-221, 2000.
22. Sun B, Sun GB, Xiao J, Chen RC, Wang X, Wu Y, Cao L, Yang ZH and Sun XB: Isorhamnetin inhibits H<sub>2</sub>O<sub>2</sub>-induced activation of the intrinsic apoptotic pathway in H9c2 cardiomyocytes through scavenging reactive oxygen species and ERK inactivation. *J Cell Biochem* 113: 473-485, 2012.
23. Porter AG and Jänicke RU: Emerging roles of caspase-3 in apoptosis. *Cell Death Differ* 6: 99-104, 1999.
24. Kook D, Wolf AH, Alice LY, Neubauer AS, Priglinger SG, Kampik A and Welge-Lüssen UC: The protective effect of quercetin against oxidative stress in the human RPE in vitro. *Invest Ophthalmol Vis Sci* 49: 1712-1720, 2008.
25. Wang R, Peng L, Zhao J, Zhang L, Guo C, Zheng W and Chen H: Gardenamide A protects RGC-5 cells from H<sub>2</sub>O<sub>2</sub>-induced oxidative stress insults by activating PI3K/Akt/eNOS signaling pathway. *Int J Mol Sci* 16: 22350-22367, 2015.
26. Zha X, Wu G, Zhao X, Zhou L, Zhang H, Li J, Ma L and Zhang Y: PRDX6 protects ARPE-19 cells from oxidative damage via PI3K/AKT signaling. *Cell Physiol Biochem* 36: 2217-2228, 2015.
27. Li Z, Dong X, Liu H, Chen X, Shi H, Fan Y, Hou D and Zhang X: Astaxanthin protects ARPE-19 cells from oxidative stress via upregulation of Nrf2-regulated phase II enzymes through activation of PI3K/Akt. *Mol Vis* 19: 1656-1666, 2013.
28. Yang P, Peairs JJ, Tano R and Jaffe GJ: Oxidant-mediated Akt activation in human RPE cells. *Invest Ophthalmol Vis Sci* 47: 4598-4606, 2006.
29. Luo Y, Sun G, Dong X, Wang M, Qin M, Yu Y and Sun X: Isorhamnetin attenuates atherosclerosis by inhibiting macrophage apoptosis via PI3K/AKT activation and HO-1 induction. *PLoS One* 10: e0120259, 2015.