

High expression of *FLT3* is a risk factor in leukemia

JIE CHENG^{1,2}, LIJUN QU^{1,2}, JIAN WANG^{1,2}, LEMEI CHENG^{1,2} and YI WANG¹

¹Department of Hematology and Oncology, Children's Hospital of Soochow University, Suzhou, Jiangsu 215025;

²Department of Hematology and Oncology, Anhui Provincial Children's Hospital, Hefei, Anhui 230041, P.R. China

Received January 31, 2017; Accepted August 23, 2017

DOI: 10.3892/mmr.2017.8232

Abstract. Several studies have shown that internal tandem duplication (ITD) of FMS-like tyrosine kinase 3 (*FLT3*) can result in the failure of leukemia treatment and contribute to a poor prognosis. However, the role of the overexpression of *FLT3* in leukemia remains to be fully elucidated. By mining public database, the present study first identified that the expression of *FLT3* in leukemia was markedly higher, compared with that in other types of tumor and cell lines, indicating that *FLT3* is important in leukemia. In leukemia, *FLT3* was found to be significantly upregulated in acute myeloid leukemia and acute lymphoblastic leukemia, and a high expression of *FLT3* contributed to reduced survival rates. By analyzing Gene Expression Omnibus and The Cancer Genome Atlas data, it was found that genetic alterations and modification of DNA methylation increased the expression of *FLT3* in leukemia. *FLT3*-ITD and *FLT3* tyrosine kinase domain point mutations increased the expression of *FLT3* in four independent datasets. In addition, the status of *FLT3* gene methylation was negatively correlated with the expression of *FLT3*, and haploinsufficiency of DNA methyltransferase 1 increased the expression of *Flt3* in mouse leukemia cells. By analyzing the enrichment of differentially-expressed genes in chemical and genetic perturbation datasets, it was found that genes, which were upregulated in the *FLT3* high expression group had myeloid lymphoid leukemia- and nucleophosmin 1-like signatures, indicating that the overexpression of *FLT3* may use the same mechanism to promote leukemia. Collectively, the results of the present study showed that the overexpression of *FLT3* is a potential risk factor in leukemia.

Introduction

The FMS-like tyrosine kinase 3 (*FLT3*) gene, encoding a membrane-bound receptor tyrosine kinase, is crucial

in normal hematopoiesis (1,2). It has been reported that *FLT3* has two mutation types in leukemia, the most common form of *FLT3* mutation is an internal tandem duplication (ITD) within the juxtamembrane domain, which occurs in 15-35% of patients with acute myeloid leukemia (AML) (3-13) and 5-10% of patients with myelodysplasia (MDS) (14,15). Another mutation type is the missense point mutation on the tyrosine kinase domain (TKD), which occurs in 5-10% of patients with AML, 2-5% of patients with MDS and 1-3% of patients with acute lymphoblastic leukemia (ALL) (9,16,17). *FLT3*-ITD can promote ligand-independent dimerization, autophosphorylation and constitutive activation of the receptor, which lead to the aberrant activation of multiple signaling pathways, including phosphatidylinositol 3-kinase/AKT, mitogen-activated protein kinase/extracellular signal-regulated kinase (MAPK/ERK) and signal transducer and activator of transcription 5 (STAT5) (18-20). *FLT3*-TKD also promotes constitutive phosphorylation of the receptor and ligand-independent cell growth (16,17,21).

FLT3 mutations are high risk factors in leukemia, and contribute to increased risk of treatment failure and poor prognosis (7,8). Mutations of *FLT3*-ITD have been reported to confer resistance to multiple tyrosine kinase inhibitors (22). The mutant allelic burden and ITD length are significantly associated with reduced overall survival and disease-free survival rates (23,24). The detection of ITD mutations at diagnosis is now a routine clinical practice to provide guidance for the optimal treatment of patients with AML.

Several previous studies have shown that the expression of *FLT3* is activated in acute promyelocytic leukemia (APL) and adult B lymphoblastic leukemia (25,26), and the upregulation of *FLT3* is a passive event in Hoxa9/Meis1-induced AML (27) indicating that the overexpression of *FLT3* may have a tumor-promotion effect. In addition, the overexpression of *FLT3* has been reported to activate the AKT and MAPK pathways, but not the STAT5 pathway (28), which suggests that the overexpression of *FLT3* has overlapping downstream pathways with *FLT3*-ITD.

In the present study, it was found that *FLT3* was upregulated in leukemia, and that the high expression of *FLT3* indicated a poor prognosis. By analyzing differentially-expressed genes (DEGs), certain leukemic oncogenes were identified, and the high expression of *FLT3* was found to have myeloid lymphoid leukemia (MLL)- and nucleophosmin 1-like (NPM1)-like signatures.

Correspondence to: Professor Yi Wang, Department of Hematology and Oncology, Children's Hospital of Soochow University, 92 Zhong-Nan Street, Suzhou, Jiangsu 215025, P.R. China
E-mail: yi.wang1510@gmail.com

Key words: FMS-like tyrosine kinase 3, acute myeloid leukemia, prognostic factor, DNA methylation, differentially-expressed gene

Materials and methods

Expression profile analysis. *FLT3* expression data were collected from the Gene Expression Omnibus (GEO; <http://www.ncbi.nlm.nih.gov/geo/>), cBioportal (29,30), Oncomine (<http://www.oncomine.org>), Cancer Cell Line Encyclopedia (CCLE; <https://portals.broadinstitute.org/ccle/home>) (31) and the Human Protein Atlas (HPA; <http://www.proteinatlas.org>) (32). The expression levels of *FLT3* in various normal human tissues and cells were analyzed using three independent databases, which included the GEO GDS3834 dataset, and the HPA and Genotype-Tissue Expression (GTEx; <https://www.gtexportal.org/home/>) databases. The *FLT3* expression data were directly downloaded from the GEO and HPA databases; the tissue with the highest expression of *FLT3* was normalized to 1.

To compare the expression of *FLT3* in different types of cancer, data were directly downloaded from cBioportal, Oncomine and CCLE, and cancer cell lines with the same tissue origin were classified as the same group.

The expression levels of *FLT3* in different types of leukemia, including AML, B-cell ALL, B-cell childhood ALL, chronic lymphocytic leukemia, chronic myelogenous leukemia, myelodysplastic syndrome, pro-B ALL, and T-cell ALL, were analyzed using Oncomine. The peripheral blood mononuclear cells were considered a normal control.

The GDS4306 dataset was used to evaluate the effect of a DNA methyltransferase 1 (*DNMT1*) haploinsufficiency on the expression of *FLT3*. To compare the correlation between *FLT3* methylation and expression, data were downloaded from cBioportal to perform linear regression. The *FLT3* mutated samples and wild-type samples were analyzed separately.

Overall survival analysis. The GSE12417 GEO dataset, which contains data on survival rates and survival status, was selected to draw the overall survival curve, and the median expression of *FLT3* was used as the cut-off, according to a previous report (33), to classify patients into a high expression group and low expression group. The top 50% of patients were classified as the *FLT3* high expression group and the lowest 50% patients were defined as the *FLT3* low expression group, according to the expression of *FLT3* from high to low. The statistical difference between two curves was calculated using a log-rank test.

Analysis of genetic alterations. In order to summarize the genetic alterations of *FLT3* in different types of cancer, The Cancer Genome Atlas (TCGA) data were downloaded through cBioportal. A total of 30 types of cancer, including AML, skin cutaneous melanoma, colorectal adenocarcinoma, esophageal carcinoma, lung adenocarcinoma, stomach adenocarcinoma, lung squamous cell cancer, lymphoid neoplasm diffuse large B-cell lymphoma, bladder urothelial carcinoma, uterine corpus endometrial carcinoma, sarcoma, cholangiocarcinoma, prostate adenocarcinoma, breast invasive carcinoma, glioblastoma multiforme, liver hepatocellular carcinoma, ovarian serous cystadenocarcinoma, cervical squamous cell carcinoma and endocervical adenocarcinoma, head and neck squamous cell carcinoma, kidney renal clear cell carcinoma, uterine carcinosarcoma, brain lower grade glioma, pancreatic

adenocarcinoma, pheochromocytoma and paraganglioma, adrenocortical carcinoma, thyroid carcinoma, testicular germ cell cancer, kidney renal papillary cell carcinoma, thymoma, kidney chromophobe and mesothelioma, were selected to analyze *FLT3* mutations and copy number alterations.

Screening of DEGs. Three GEO datasets of AML (GSE10358, GSE14468 and GSE34860) were selected to analyze the DEGs between the *FLT3* high expression group and *FLT3* low expression group. Initially, the raw data in the CEL file were downloaded from the GEO database (34), and the robust multiarray average algorithm in the affy R-3.3.1 package was used to perform background correction, normalization and expression calculation (35-37). The Limma package in R (38) was used to identify the DEGs at the probe level between these two groups. $P < 0.05$ and \log_2 fold change (FC) > 0.585 were used as the cut-off criteria. Finally, these probes were annotated into gene names.

Chemical and genetic perturbations enrichment of DEGs. In order to identify the association between the overexpression of *FLT3* and other risk factors for leukemia, the enrichment of DEGs in the chemical and genetic perturbations gene set were determined using Gene Set Enrichment Analysis (<http://software.broadinstitute.org/gsea>) (39,40). Input of the upregulated and downregulated genes was performed on the website separately, and chemical and genetic perturbations was selected to calculate the enrichment. $P < 0.05$ and FDR q -value < 0.05 were used as the cut-off criteria.

Statistical analysis. Student's t-test was used to calculate statistically significant differences between quantitative variables. The log-rank test was used to compare the overall survival curve. $P < 0.05$ was considered to indicate a statistically significant difference. GraphPad Prism 5.01 was used for statistical analysis (GraphPad Software, Inc., La Jolla, CA, USA).

Results

***FLT3* is upregulated in leukemia and a high expression of *FLT3* is a prognostic biomarker.** It has been reported that the *FLT3*-ITD mutation is correlated with prognosis and drug response in leukemia. To investigate the role of the expression of *FLT3* in leukemia, the present study firstly evaluated the expression of *FLT3* in different types of cancer and normal tissues. In normal tissues, *FLT3* was expressed at high levels in bone marrow, lymph nodes, the thymus and spleen in three independent databases (Fig. 1A), which indicated that *FLT3* was involved in hematopoiesis. In cancer, the expression of *FLT3* was significantly higher in leukemia, compared with other types of tumor (Fig. 1B and C), and *FLT3* was specifically upregulated in AML and ALL (Fig. 1D), suggesting that *FLT3* contributed to the progression of AML and ALL. In order to analyze the association between the expression of *FLT3* and survival rates in AML, the GSE12417 dataset, in which patients with AML exhibited a normal karyotype, was selected for analysis. Kaplan-Meier survival analysis showed that patients with AML with high expression levels of *FLT3* ($n=121$) had reduced survival rates, compared with those with low expression levels of *FLT3* ($n=121$; $P < 0.05$; Fig. 1E). These




Figure 1. Tissue expression profile identifying the overexpression of *FLT3* in leukemia. (A) RNA expression of *FLT3* in various normal human tissues and cells, analyzed using three independent databases (HPA, GTEx and GDS3834). Red color indicates relative expression of *FLT3*, black indicates expression data are not available. (B) TCGA and Oncomine databases were used to evaluate expression levels of *FLT3* in different types of tumor. (C) Expression of *FLT3* in different cancer cell lines was analyzed using CCLE and Oncomine databases. Cell lines of the same tissue origin were classified into the same group. (D) *FLT3* was significantly upregulated in several types of leukemia. The X-axis shows the fold change between different types of leukemia and peripheral blood mononuclear cells. The Y-axis shows the log 10 transformed P-value. Different colors of points indicate different types of leukemia. (E) Results of Kaplan-Meier survival analysis showed that patients with AML and a high expression of *FLT3* (n=121) had shorter overall survival rates, compared with those with a low expression of *FLT3* (n=121). *FLT3*, FMS-like tyrosine kinase; HPA, Human Protein Atlas; GTEx, Genotype-Tissue Expression; CCLE, Cancer Cell Line Encyclopedia; AC, adrenocortical carcinoma; AML, acute myeloid leukemia; BUC, bladder urothelial carcinoma; BLGG, brain lower grade glioma; BIC, breast invasive carcinoma, CSCCEA, cervical squamous cell carcinoma and endocervical adenocarcinoma; CC, cholangiocarcinoma; CA, colorectal adenocarcinoma; EC esophageal carcinoma; GM, glioblastoma multiforme; HCC hepatocellular carcinoma; HNSCC, head and neck squamous cell carcinoma; KC, kidney chromophobe; KRCCC, kidney renal clear cell carcinoma; KRCCC, kidney renal papillary cell carcinoma; LA, lung adenocarcinoma; LSCC, lung squamous cell carcinoma; LNDLBL, lymphoid neoplasm diffuse large B-cell lymphoma; OSC, ovarian serous cystadenocarcinoma; Pan-ad, pancreatic adenocarcinoma; PP, pheochromocytoma and paraganglioma; Pro-ad, prostate adenocarcinoma; SCM, skin cutaneous melanoma; SA, stomach adenocarcinoma; TGCC, testicular germ cell cancer; TC, thyroid carcinoma; UC, uterine carcinosarcoma; UCEC, uterine corpus endometrial carcinoma; B-ALL, B-cell acute lymphoblastic leukemia; BA, breast adenocarcinoma; DLBL, diffuse large B-cell lymphoma; EA, endometrial adenocarcinoma; FL, follicular lymphoma GB, glioblastoma; MB, medulloblastoma; OA, ovarian adenocarcinoma; PM, pleural mesothelioma; Pro-ca, prostate carcinoma; RCC, renal cell carcinoma.




Figure 2. Genetic and epigenetic alterations increase the expression of *FLT3* in AML. (A) cBioportal database was used to analyze mutations and copy number alterations of *FLT3*. Genetic alterations of *FLT3* in different types of cancer showed the most mutations of *FLT3* in leukemia. (B) Four databases (GSE10358, GSE14468, GSE34860 and TCGA) were utilized to examine the effect of *FLT3* mutations on the expression of *FLT3*. (C) cBioportal result showed the expression of *FLT3* was negatively correlated with *FLT3* methylation in leukemia. (D) Gene Expression Omnibus dataset (GDS4306) showed that *DNMT1* haploinsufficiency (haplo) increased expression of *Flt3* in mouse leukemia cells. *FLT3*, FMS-like tyrosine kinase; AML, acute myeloid leukemia; DLBC, diffuse large B-cell; GBM, glioblastoma multiforme; ccRCC, clear cell renal cell carcinoma; UC, uterine carcinosarcoma; PCPG, pheochromocytoma and paraganglioma; ACC, adenoid cystic carcinoma; pRCC, papillary renal cell carcinoma; chRCC, chromophobe renal cell carcinoma; WT, wild-type; ITD, internal tandem duplication; TKD, tyrosine kinase domain. *DNMT1*, DNA methyltransferase 1; * $P<0.05$, ** $P<0.01$ and *** $P<0.001$; ns, not significant.

collective data indicated that *FLT3* was overexpressed in leukemia and was a prognostic biomarker.

Expression of FLT3 is regulated by genetic and epigenetic alterations. In order to determine the mechanism increasing the expression of *FLT3* in leukemia, the present study evaluated genetic and epigenetic alterations of *FLT3* in AML. By mining TCGA data, it was found that *FLT3* was significantly mutated in AML, compared with other types of cancer, which

was similar to the results obtained on the expression of *FLT3*. The results revealed ~28% of patients with AML had somatic mutations in *FLT3* (Fig. 2A). Therefore, it was hypothesized that *FLT3* mutations may increase the expression of *FLT3* in AML. To confirm this, the expression of *FLT3* was compared between wild-type and mutated groups in four independent databases, (GSE10358, GSE14468, GSE34860 and TCGA). It was found that ITD and TKD mutations significantly increased the expression of *FLT3* (Fig. 2B). In addition, methylation data




Figure 3. DEGs between the *FLT3* high expression group and *FLT3* low expression group. Three databases (GSE10358, GSE14468 and GSE34860) were used to analyze DEGs. (A) Genes with $P < 0.05$ and $FC > 1.5$ are indicated in red and green colors in the volcano plot. Red indicates genes upregulated in the *FLT3* high expression group, and green indicates genes downregulated in the *FLT3* high expression group. The X-axis is the \log_2 -transformed fold change, and the Y-axis is the \log_{10} -transformed P-value. (B) Pie charts shows number of DEGs in the three databases. A total of seven genes were upregulated in the *FLT3* high expression group (left) and 11 genes were downregulated (right). (C) Gene Set Enrichment Analysis was used to assess the association between DEGs and chemical and genetic perturbations. The seven upregulated genes were mainly associated with the MLL signature and NPM1 signature (above). The majority of the downregulated genes were also reduced in HSCs (below). DEGs, differentially-expressed genes; FC, fold change; HSCs, hematopoietic stem cells; MLL, myeloid lymphoid leukemia; NPM1, nucleophosmin 1-like; AML, acute myeloid leukemia; GBM, glioblastoma multiforme.

from TCGA was used to compare the association between the expression of *FLT3* and methylation status. In AML, the expression of *FLT3* was negatively correlated with its methylation (Fig. 2C), indicating that the hypomethylation of *FLT3* may be a potential mechanism resulting in the upregulation of *FLT3*. Of note, mining of the GEO dataset (GDS4306) revealed that the haploinsufficiency of *DNMT1* significantly increased the expression of *Flt3* in mouse leukemia cells (Fig. 2D), suggesting that DNMT1 modified the methylation of *FLT3*. Collectively, these data showed that genetic and epigenetic alterations may be potential mechanisms by which the expression of *FLT3* is increased in AML.

High expression of *FLT3* has MLL- and NPM1-like signatures.
In order to identify the DEGs associated with a high expression

of *FLT3*, three independent AML databases (GSE10358, GSE14468 and GSE34860) were examined (Fig. 3A). The expression of seven genes were significantly upregulated in the *FLT3* high expression group, comprising *ATP8B4*, *CCNA1*, *HOXA10*, *IQCJ*, *PBX3*, *SEL1L3* and *SLC2A5*. *CCNA1*, *HOXA10*, *PBX3* and *SLC2A5* have been reported to function as oncogenes in leukemia (41-44). A total of 11 genes were down-regulated, comprising *AHSP*, *ALAS2*, *EPB42*, *FCAR*, *GBP1*, *HBG1*, *LEF1*, *LTF*, *SLC6A8*, *TUBB2A* and *ZFP36L1*, as shown in Fig. 3B. The deletion of RNA-binding protein *ZFP36L1* has been reported to lead to perturbed thymic development and T lymphoblastic leukemia (45). Subsequently, the present study analyzed the enrichment of these DEGs in the chemical and genetic perturbations dataset. Of note, the seven overexpressed genes were enriched in MLL and NPM1 signatures,

indicating that the majority were also upregulated in MLL- or NPM1-mutated AML (Fig. 3C). The 11 downregulated genes were reduced in hematopoietic stem cells (Fig. 3C). Taken together, these results showed that a high expression of *FLT3* was associated with MLL- and NPM1-like signatures.

Discussion

In the present study, it was found that the expression of *FLT3* was high in normal hematopoietic tissues (Fig. 1A) and, in leukemia, *FLT3* was specifically upregulated in AML and ALL (Fig. 1D), indicating that a high expression of *FLT3* may contribute to the progression of leukemia. It has been reported that the overexpression of *FLT3* can induce autophosphorylation (46), and activate the AKT and MAPK pathways in AML (47). ITD mutations clustered in the juxtamembrane domain of *FLT3* are the most frequent forms in AML, and *FLT3*-ITD mutations are associated with a poor prognosis (47-49). In the present study, it was also found that a high expression of *FLT3* was a prognostic factor for poor prognosis in AML using the GEO database (Fig. 1E).

FLT3-related pathways were activated in AML by ITD mutations or the overexpression of *FLT3*. Several interacting proteins are also reported to interact with *FLT3* to negatively or positively regulate *FLT3* pathways. Spleen tyrosine kinase (SYK) and the mucin 1-C-terminal subunit (MUC1-C) oncoprotein are reported to directly bind to and activate *FLT3*-related pathways (50,51), whereas suppressor of cytokine signaling 2 (SOCS2) and src-like adaptor protein 2 (SLAP2) interact with *FLT3* protein to inhibit its signaling (52,53). In addition, the transcription of *FLT3* can be regulated in AML. Certain AML-related transcription factors, including CCAAT/enhancer binding protein α and the proto-oncogene MYB, can bind to the *FLT3* promoter to activate the transcription of *FLT3* (54). In the present study, another two mechanisms were found to increase the expression of *FLT3* in AML. ITD and TKD mutations, and the methylation of *FLT3* increased its expression in AML (Fig. 2B and C). In addition, DNMT1 was identified as a potential regulator of *FLT3* (Fig. 2D).

By analyzing DEGs in the *FLT3* high expression group, a total of 18 genes were identified using three independent AML datasets, seven of which were upregulated and 11 of which were downregulated (Fig. 3B). These upregulated genes were enriched in MLL and NPM1 signatures (Fig. 3C), and *CCNA1*, *HOXA10*, *PBX3* and *SLC2A5* were reported to function as oncogenes in leukemia. *CCNA1* was overexpressed in ALL, and patients with high levels of *CCNA1* exhibit poor event-free survival rates (55). *CCNA1* transgenic mice have also been shown to exhibit abnormal myelopoiesis and progressed to overt AML (40). Similarly, the overexpression of *HOXA10* can cooperate with active SHP2 to induce AML (41), and *HOXA10* was usually fused with the *NUP98*, collaborating with overexpressed *FLT3* receptor tyrosine kinase to induce aggressive AML (56). *PBX3* is an important cofactor of *HOXA9* in leukemogenesis (57) and the coexpression of *PBX3* and *MEIS1* (*PBX3/MEIS1*) can cause AML *in vivo* (58). *SLC2A5* is also overexpressed in AML and can increase the fructose utilization of leukemic cells, with a high expression of *SLC2A5* being associated with poor outcomes (44). Among these 11 downregulated genes, *ZFP36L1* has a tumor suppressor role

in leukemia. *ZFP36L1* is an RNA-binding protein and leads to mRNA degradation; deletion of *ZFP36L1* in mice can induce T cell ALL (45).

Collectively the findings obtained in the present study showed that *FLT3* is overexpressed in leukemia and is a prognostic factor for poor prognosis in AML. Patients with a high expression of *FLT3* simultaneously express high levels of leukemic oncogenes. Therefore, a high expression of *FLT3* is a risk factor in leukemia.

Acknowledgements

This study was supported by a grant from the National Natural Science Foundation of China (grant no. 8130044).

References

- Volpe G, Clarke M, Garcia P, Walton DS, Vegiopoulos A, Del Pozzo W, O'Neill LP, Frampton J and Dumon S: Regulation of the *Flt3* gene in haematopoietic stem and early progenitor cells. *PLoS One* 10: e0138257, 2015.
- Ebihara Y, Wada M, Ueda T, Xu MJ, Manabe A, Tanaka R, Ito M, Mugishima H, Asano S, Nakahata T and Tsuji K: Reconstitution of human haematopoiesis in non-obese diabetic/severe combined immunodeficient mice by clonal cells expanded from single CD34⁺CD38⁻ cells expressing *Flk2/Flt3*. *Br J Haematol* 119: 525-534, 2002.
- Nakao M, Yokota S, Iwai T, Kaneko H, Horiike S, Kashima K, Sonoda Y, Fujimoto T and Misawa S: Internal tandem duplication of the *flt3* gene found in acute myeloid leukemia. *Leukemia* 10: 1911-1918, 1996.
- Kiyoi H, Towatari M, Yokota S, Hamaguchi M, Ohno R, Saito H and Naoe T: Internal tandem duplication of the *FLT3* gene is a novel modality of elongation mutation which causes constitutive activation of the product. *Leukemia* 12: 1333-1337, 1998.
- Kiyoi H, Naoe T, Yokota S, Nakao M, Minami S, Kuriyama K, Takeshita A, Saito K, Hasegawa S, Shimodaira S, *et al*: Internal tandem duplication of *FLT3* associated with leukocytosis in acute promyelocytic leukemia. *Leukemia Study Group of the Ministry of Health and Welfare (Kohseisho)*. *Leukemia* 11: 1447-1452, 1997.
- Stirewalt DL, Kopecky KJ, Meshinchi S, Appelbaum FR, Slovak ML, Willman CL and Radich JP: *FLT3*, *RAS*, and *TP53* mutations in elderly patients with acute myeloid leukemia. *Blood* 97: 3589-3595, 2001.
- Meshinchi S, Woods WG, Stirewalt DL, Sweetser DA, Buckley JD, Tjoa TK, Bernstein ID and Radich JP: Prevalence and prognostic significance of *Flt3* internal tandem duplication in pediatric acute myeloid leukemia. *Blood* 97: 89-94, 2001.
- Schnittger S, Schoch C, Dugas M, Kern W, Staib P, Wuchter C, Löffler H, Sauerland CM, Serve H, Büchner T, *et al*: Analysis of *FLT3* length mutations in 1003 patients with acute myeloid leukemia: Correlation to cytogenetics, FAB subtype, and prognosis in the AMLCG study and usefulness as a marker for the detection of minimal residual disease. *Blood* 100: 59-66, 2002.
- Thiede C, Steudel C, Mohr B, Schaich M, Schäkel U, Platzbecker U, Wermke M, Bornhäuser M, Ritter M, Neubauer A, *et al*: Analysis of *FLT3*-activating mutations in 979 patients with acute myelogenous leukemia: Association with FAB subtypes and identification of subgroups with poor prognosis. *Blood* 99: 4326-4335, 2002.
- Abu-Duhier FM, Goodeve AC, Wilson GA, Gari MA, Peake IR, Rees DC, Vandenbergh EA, Winship PR and Reilly JT: *FLT3* internal tandem duplication mutations in adult acute myeloid leukaemia define a high-risk group. *Br J Haematol* 111: 190-195, 2000.
- Kottaridis PD, Gale RE, Frew ME, Harrison G, Langabeer SE, Belton AA, Walker H, Wheatley K, Bowen DT, Burnett AK, *et al*: The presence of a *FLT3* internal tandem duplication in patients with acute myeloid leukemia (AML) adds important prognostic information to cytogenetic risk group and response to the first cycle of chemotherapy: Analysis of 854 patients from the United Kingdom Medical Research Council AML 10 and 12 trials. *Blood* 98: 1752-1759, 2001.

12. Xu F, Taki T, Yang HW, Hanada R, Hongo T, Ohnishi H, Kobayashi M, Bessho F, Yanagisawa M and Hayashi Y: Tandem duplication of the FLT3 gene is found in acute lymphoblastic leukaemia as well as acute myeloid leukaemia but not in myelodysplastic syndrome or juvenile chronic myelogenous leukaemia in children. *Br J Haematol* 105: 155-162, 1999.
13. Kiyoi H, Naoe T, Nakano Y, Yokota S, Minami S, Miyawaki S, Asou N, Kuriyama K, Jinnai I, Shimazaki C, *et al*: Prognostic implication of FLT3 and N-RAS gene mutations in acute myeloid leukemia. *Blood* 93: 3074-3080, 1999.
14. Horiike S, Yokota S, Nakao M, Iwai T, Sasai Y, Kaneko H, Taniwaki M, Kashima K, Fujii H, Abe T and Misawa S: Tandem duplications of the FLT3 receptor gene are associated with leukemic transformation of myelodysplasia. *Leukemia* 11: 1442-1446, 1997.
15. Yokota S, Kiyoi H, Nakao M, Iwai T, Misawa S, Okuda T, Sonoda Y, Abe T, Kahsima K, Matsuo Y and Naoe T: Internal tandem duplication of the FLT3 gene is preferentially seen in acute myeloid leukemia and myelodysplastic syndrome among various hematological malignancies. A study on a large series of patients and cell lines. *Leukemia* 11: 1605-1609, 1997.
16. Yamamoto Y, Kiyoi H, Nakano Y, Suzuki R, Kodaera Y, Miyawaki S, Asou N, Kuriyama K, Yagasaki F, Shimazaki C, *et al*: Activating mutation of D835 within the activation loop of FLT3 in human hematologic malignancies. *Blood* 97: 2434-2439, 2001.
17. Abu-Duhier FM, Goodeve AC, Wilson GA, Care RS, Peake IR and Reilly JT: Identification of novel FLT-3 Asp835 mutations in adult acute myeloid leukaemia. *Br J Haematol* 113: 983-988, 2001.
18. Hayakawa F, Towatari M, Kiyoi H, Tanimoto M, Kitamura T, Saito H and Naoe T: Tandem-duplicated Flt3 constitutively activates STAT5 and MAP kinase and introduces autonomous cell growth in IL-3-dependent cell lines. *Oncogene* 19: 624-631, 2000.
19. Kiyoi H, Ohno R, Ueda R, Saito H and Naoe T: Mechanism of constitutive activation of FLT3 with internal tandem duplication in the juxtamembrane domain. *Oncogene* 21: 2555-2563, 2002.
20. Mizuki M, Fenski R, Halfter H, Matsumura I, Schmidt R, Müller C, Grüning W, Kratz-Albers K, Serve S, Steur C, *et al*: FLT3 mutations from patients with acute myeloid leukemia induce transformation of 32D cells mediated by the Ras and STAT5 pathways. *Blood* 96: 3907-3914, 2000.
21. Spiekermann K, Bagrintseva K, Schoch C, Haferlach T, Hiddemann W and Schnittger S: A new and recurrent activating length mutation in exon 20 of the FLT3 gene in acute myeloid leukemia. *Blood* 100: 3423-3425, 2002.
22. Williams AB, Nguyen B, Li L, Brown P, Levis M, Leahy D and Small D: Mutations of FLT3/ITD confer resistance to multiple tyrosine kinase inhibitors. *Leukemia* 27: 48-55, 2013.
23. Kim Y, Lee GD, Park J, Yoon JH, Kim HJ, Min WS and Kim M: Quantitative fragment analysis of FLT3-ITD efficiently identifying poor prognostic group with high mutant allele burden or long ITD length. *Blood Cancer J* 5: e336, 2015.
24. Stirewalt DL, Kopecky KJ, Meshinchi S, Engel JH, Pogossova-Agadjanyan EL, Linsley J, Slovak ML, Willman CL and Radich JP: Size of FLT3 internal tandem duplication has prognostic significance in patients with acute myeloid leukemia. *Blood* 107: 3724-3726, 2006.
25. Griffith M, Griffith OL, Krysiak K, Skidmore ZL, Christopher MJ, Klcio JM, Ramu A, Lamprecht TL, Wagner AH, Campbell KM, *et al*: Comprehensive genomic analysis reveals FLT3 activation and a therapeutic strategy for a patient with relapsed adult B-lymphoblastic leukemia. *Exp Hematol* 44: 603-613, 2016.
26. Lilakos K, Viniou NA, Mavrogianni D, Vassilakopoulos TP, Dimopoulou MN, Plata E, Angelopoulou MK, Variami E, Stavrogianni N, Liapi D, *et al*: FLT3 overexpression in acute promyelocytic leukemia patients without detectable FLT3-ITD or codon 835-836 mutations: A pilot study. *Anticancer Res* 26: 1201-1207, 2006.
27. Staffas A, Arabanian LS, Wei SY, Jansson A, Ståhlman S, Johansson P, Fogelstrand L, Cammenga J, Kuchenbauer F and Palmqvist L: Upregulation of Flt3 is a passive event in Hoxa9/Meis1-induced acute myeloid leukemia in mice. *Oncogene* 36: 1516-1524, 2017.
28. Spiekermann K, Bagrintseva K, Schwab R, Schmiele K and Hiddemann W: Overexpression and constitutive activation of FLT3 induces STAT5 activation in primary acute myeloid leukemia blast cells. *Clin Cancer Res* 9: 2140-2150, 2003.
29. Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A, Sinha R, Larsson E, *et al*: Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* 6: pii, 2013.
30. Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A, Byrne CJ, Heuer ML, Larsson E, *et al*: The cBio cancer genomics portal: An open platform for exploring multidimensional cancer genomics data. *Cancer Discov* 2: 401-404, 2012.
31. Barretina J, Caponigro G, Stransky N, Vekatesan K, Margolin AA, Kim S, Wilson CJ, Lehár J, Kryukov GV, Sonkin D, *et al*: The cancer cell line encyclopedia enables predictive modelling of anticancer drug sensitivity. *Nature* 483: 603-607, 2012.
32. Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson Å, Kampf C, Sjöstedt E, Asplund A, *et al*: Proteomics. Tissue-based map of the human proteome. *Science* 347: 1260419, 2015.
33. Wu BH, Chen H, Cai CM, Fang JZ, Wu CC, Huang LY, Wang L and Han ZG: Epigenetic silencing of JMJD5 promotes the proliferation of hepatocellular carcinoma cells by down-regulating the transcription of CDKN1A. *Oncotarget* 7: 6847-6863, 2016.
34. Edgar R, Domrachev M and Lash AE: Gene expression omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res* 30: 207-210, 2002.
35. Gautier L, Cope L, Bolstad BM and Irizarry RA: affy-analysis of Affymetrix GeneChip data at the probe level. *Bioinformatics* 20: 307-315, 2004.
36. Troyanskaya O, Cantor M, Sherlock G, Brown P, Hastie T, Tibshirani R, Botstein D and Altman RB: Missing value estimation methods for DNA microarrays. *Bioinformatics* 17: 520-525, 2001.
37. Fujita A, Sato JR, Rodrigues Lde O, Ferreira CE and Sogayar MC: Evaluating different methods of microarray data normalization. *BMC Bioinformatics* 7: 469, 2006.
38. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W and Smyth GK: limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* 43: e47, 2015.
39. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES and Mesirov JP: Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci USA* 102: 15545-15550, 2005.
40. Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, Puigserver P, Carlsson E, Ridderstråle M, Laurila E, *et al*: PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet* 34: 267-273, 2003.
41. Liao C, Wang XY, Wei HQ, Li SQ, Merghoub T, Pandolfi PP and Wolgemuth DJ: Altered myelopoiesis and the development of acute myeloid leukemia in transgenic mice overexpressing cyclin A1. *Proc Natl Acad Sci USA* 98: 6853-6858, 2001.
42. Wang H, Lindsey S, Konieczna I, Bei L, Horvath E, Huang W, Saberwal G and Eklund EA: Constitutively active SHP2 cooperates with HoxA10 overexpression to induce acute myeloid leukemia. *J Biol Chem* 284: 2549-2567, 2009.
43. Garcia-Cuellar MP, Steger J, Füller E, Hetzner K and Slany RK: Pbx3 and Meis1 cooperate through multiple mechanisms to support Hox-induced murine leukemia. *Haematologica* 100: 905-913, 2015.
44. Chen WL, Wang YY, Zhao A, Xia L, Xie G, Su M, Zhao L, Liu J, Qu C, Wei R, *et al*: Enhanced fructose utilization mediated by SLC2A5 is a unique metabolic feature of acute myeloid leukemia with therapeutic potential. *Cancer Cell* 30: 779-791, 2016.
45. Hodson DJ, Janas ML, Galloway A, Bell SE, Andrews S, Li CM, Pannell R, Siebel CW, MacDonald HR, De Keersmaecker K, *et al*: Deletion of the RNA-binding proteins ZFP36L1 and ZFP36L2 leads to perturbed thymic development and T lymphoblastic leukemia. *Nat Immunol* 11: 717-724, 2010.
46. Ozeki K, Kiyoi H, Hirose Y, Iwai M, Ninomiya M, Kodaera Y, Miyawaki S, Kuriyama K, Shimazaki C, Akiyama H, *et al*: Biologic and clinical significance of the FLT3 transcript level in acute myeloid leukemia. *Blood* 103: 1901-1908, 2004.
47. Döhner H, Estey EH, Amadori S, Appelbaum FR, Büchner T, Burnett AK, Dombret H, Fenaux P, Grimwade D, Larson RA, *et al*: Diagnosis and management of acute myeloid leukemia in adults: Recommendations from an international expert panel, on behalf of the European Leukemia Net. *Blood* 115: 453-474, 2010.
48. Estey EH: Acute myeloid leukemia: 2012 update on diagnosis, risk stratification, and management. *Am J Hematol* 87: 89-99, 2012.
49. Patel JP, Gönen M, Figueroa ME, Fernandez H, Sun Z, Racevskis J, Van Vlierberghe P, Dolgalev I, Thomas S, Aminova O, *et al*: Prognostic relevance of integrated genetic profiling in acute myeloid leukemia. *N Engl J Med* 366: 1079-1089, 2012.

50. Puissant A, Fenouille N, Alexe G, Pikman Y, Bassil CF, Mehta S, Du J, Kazi JU, Luciano F, Rönstrand L, *et al*: SYK is a critical regulator of FLT3 in acute myeloid leukemia. *Cancer Cell* 25: 226-242, 2014.
51. Liu S, Yin L, Stroopinsky D, Rajabi H, Puissant A, Stegmaier K, Avigan D, Kharbanda S, Kufe D and Stone R: MUC1-C oncoprotein promotes FLT3 receptor activation in acute myeloid leukemia cells. *Blood* 123: 734-742, 2014.
52. Moharram SA, Chougule RA, Su X, Li T, Sun J, Zhao H, Rönstrand L and Kazi JU: Src-like adaptor protein 2 (SLAP2) binds to and inhibits FLT3 signaling. *Oncotarget* 7: 57770-57782, 2016.
53. Kazi JU and Rönstrand L: Suppressor of cytokine signaling 2 (SOCS2) associates with FLT3 and negatively regulates downstream signaling. *Mol Oncol* 7: 693-703, 2013.
54. Volpe G, Walton DS, Del Pozzo W, Garcia P, Dassé E, O'Neill LP, Griffiths M, Frampton J and Dumon S: C/EBP α and MYB regulate FLT3 expression in AML. *Leukemia* 27: 1487-1496, 2013.
55. Holm C, Ora I, Brunhoff C, Anagnostaki L, Landberg G and Persson JL: Cyclin A1 expression and associations with disease characteristics in childhood acute lymphoblastic leukemia. *Leuk Res* 30: 254-261, 2006.
56. Palmqvist L, Argiropoulos B, Pineault N, Abramovich C, Sly LM, Krystal G, Wan A and Humphries RK: The Flt3 receptor tyrosine kinase collaborates with NUP98-HOX fusions in acute myeloid leukemia. *Blood* 108: 1030-1036, 2006.
57. Li Z, Zhang Z, Li Y, Arnovitz S, Chen P, Huang H, Jiang X, Hong GM, Kunjamma RB, Ren H, *et al*: PBX3 is an important cofactor of HOXA9 in leukemogenesis. *Blood* 121: 1422-1431, 2013.
58. Li Z, Chen P, Su R, Hu C, Li Y, Elkahoul AG, Zuo Z, Gurbuxani S, Arnovitz S, Weng H, *et al*: PBX3 and MEIS1 cooperate in hematopoietic cells to drive acute myeloid leukemias characterized by a core transcriptome of the MLL-rearranged disease. *Cancer Res* 76: 619-629, 2016.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.