Bioinformatics analysis of the molecular mechanisms underlying traumatic spinal cord injury

SHU-JIE ZHAO, WEI ZHOU, JIAN CHEN, YONG-JUN LUO and GUO-YONG YIN

Department of Orthopedics, The First Affiliated Hospital of Nanjing Medical University, Nanjing, Jiangsu 210029, P.R. China

Received October 17, 2017; Accepted February 7, 2018

DOI: 10.3892/mmr.2018.8918

Abstract. Spinal cord injury (SCI) is a cause of disability. The present study aimed to investigate the molecular mechanisms involved in traumatic SCI. Transcriptome data under accession no. GSE5296, including 96 chips, were downloaded from the Gene Expression Omnibus database. The raw data were normalized and differentially expressed genes (DEGs) were identified. Furthermore, Kyoto Encyclopedia of Genes and Genomes pathway and Gene Ontology enrichment analysis of up- and downregulated DEGs was performed. Additionally, a protein-protein interaction network was constructed and the expression patterns of different genes were determined. Compared with sham samples, there were 374, 707, 1,322, 1,475, 1,724 and 1,342 DEGs identified at 0.5, 4, 24 and 72 h, and 7 and 28 days post-injury, respectively. At 24 and 72 h, and 7 days following injury, the upregulated DEGs were markedly enriched in 'inflammatory response' and 'immune process'. Downregulated DEGs were predominantly enriched in neuronal function-associated pathways and 'steroid biosynthesis' process. Protein-protein interaction network analysis demonstrated similar results. Trend charts further demonstrated that the inflammatory and neuronal functions were altered in a temporal and site-specific manner. The present study provided an insight into the molecular mechanisms underlying traumatic SCI, which may benefit future SCI research and aid in therapy development.

Introduction

Spinal cord injury (SCI) entails severe physical and social consequences for patients and their families, and may be of a traumatic or non-traumatic etiology (1,2). Traumatic SCI,

which is primarily caused by traffic accidents, occurs when an external impact acutely damages the spinal cord, and may be temporally divided into four stages: i) Acute at <48 h following injury; ii) subacute at 48 h-14 days following injury; iii) intermediate at 14 days-6 months following injury; and iv) chronic at >6 months following injury (1). Although numerous studies have focused on neuroprotective and neuroregenerative therapies, no major breakthroughs have been achieved (3-5). Therefore, it is necessary to elucidate the underlying mechanism of SCI.

Pathophysiologically, the initial mechanical injury damages neurons and initiates a complex secondary injury cascade, leading to progressive cell death, ischemia and inflammation (6). It has been demonstrated that the transcriptome may reflect the pathophysiological state of the cell (7). In a number of recent bioinformatics studies, transcriptome analysis at different time points post-SCI was performed and various molecular events were characterized (8-12). The immune response, inflammatory-associated functions, vasculature development and neurological functions were demonstrated to serve roles in the development of SCI (6-12). However, certain molecular alterations that occur in a temporal and spatial manner remain to be elucidated, particularly those that occur during the secondary injury process.

In the present study, transcriptome data under accession no. GSE5296 was used to identify SCI-specific molecular programs. Temporally, three different time points were evaluated, including 24 and 72 h, and 7 days post-injury for pathway and functional enrichment analysis. Furthermore, a protein-protein interaction (PPI) network was constructed. Spatially, differentially expressed genes (DEGs) in different locations, including the trauma site (M), and immediately adjacent rostral (R) and caudal (C) regions were determined at the aforementioned time points. The present study revealed molecular mechanisms that may be associated with SCI and provided an insight into potential therapeutic targets for treatment of SCI.

Materials and methods

Transcriptome data. The transcriptome data under accession number GSE5296 based on the GPL1261 platform (Affymetrix Mouse Genome 430 2.0 Array; Affymetrix; Thermo Fisher Scientific, Inc., Waltham, MA, USA) was obtained from the National Center for Biotechnology Information Gene

Correspondence to: Dr Guo-Yong Yin, Department of Orthopedics, The First Affiliated Hospital of Nanjing Medical University, 300 Guangzhou Road, Nanjing, Jiangsu 210029, P.R. China

E-mail: guoyong_yin@sina.com

Key words: traumatic spinal cord injury, transcriptome, Kyoto Encyclopedia of Genes and Genomes pathway, protein-protein interaction network

Expression Omnibus (GEO) database (www.ncbi.nlm.nih.gov/geo). In the original dataset (GSE5296), C57BL6 mice were subjected to moderate contusion injury at the T8 spinal segment. Sections of the spinal cord (4 mm in length) were analyzed from the site of the trauma and at the immediately adjacent R and C regions, at 0.5, 4, 24 and 72 h, and 7 and 28 days following injury. A total of 96 chips were available in this dataset data, including 18 SCI samples for each M, R and C region ($n=3/\text{time-point}$), 12 sham-injury samples for each M, R and C region ($n=2/\text{time-point}$), and two samples in each region obtained from naive mice (data not shown).

Data preprocessing. The Robust Multichip Average algorithm in the Oligo package, version 1.42.0 (<http://www.bioconductor.org>) was used to preprocess the raw transcriptome data included in the GSE5296 dataset (13). Data were subjected to background correction, normalization, probe summary and $\log_2$ transformation. If there were several probes annotated to same gene symbol, the average value was used to represent the expression level of this gene. There were 45,037 probes in the raw data and 21,812 genes remained following data processing.

Identification and analysis of DEGs. In the present study, data were divided into the following paired groups: i) Post-SCI group vs. sham group in the M region at different time-points (0.5, 4, 24 and 72 h, and 7 and 28 days); ii) post-SCI group vs. post-sham group in the R region at different time-points (24 and 72 h, and 7 days); and iii) post-SCI group vs. sham group in the C region at different time-points (24 and 72 h, and 7 days). Fold change $\log_2\text{FC}$ and P-values from a Student's t-test were used to identify the DEGs. An average fold-change >2.0 and $P<0.05$ were used as cutoff criteria. Subsequently, Venny 2.1 (bioinfogp.cnb.csic.es/tools/venny/index.html) was used to compare lists of DEGs and to construct Venn diagrams.

Pathway and functional enrichment analysis. To identify pathways and biological processes enriched by DEGs, the Database for Annotation, Visualization and Integrated Discovery (DAVID 6.8; <http://david.abcc.ncifcrf.gov>) was used to perform Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway and Gene Ontology (GO) functional analysis (14). GO terms were identified under categories of biological process. The threshold was set to $P<0.05$.

PPI network construction. A PPI network was constructed using the Search Tool for the Retrieval of Interacting Genes/Proteins (String 10.5; www.string-db.org). The network was visualized using the Cytoscape software platform (Cytoscape 3.6) based on functional analysis information, including fold-change in gene/protein expression, PPIs and GO/KEGG pathway enrichment (15). A default confidence cutoff of 400 was used in the present study. Experimentally determined interactions were presented as solid lines between genes/proteins, and the dashed lines represent database predicted interactions (16).

Trend charts of neuronal function- and inflammatory response-associated genes. In the present study, neuronal function and synaptic transmission-associated genes were defined as cholinergic receptor nicotinic $\alpha 7$ subunit

(*Chrna7*), synapsin II (*Syn2*), potassium voltage-gated channel subfamily C member 1 (*Kcnc1*), ATPase plasma membrane Ca^{2+} transporting 2 (*Atp2b2*), unc-13 homolog A (*Unc13a*), regulating synaptic membrane exocytosis 1 (*Rims1*), calcium/calmodulin dependent protein kinase II γ (*Camk2g*), calcium/calmodulin dependent protein kinase II α (*Camk2a*), thyrotropin releasing hormone receptor (*Trhr*) and glutamate metabotropic receptor 1 (*Grm1*) (3,4,9). Furthermore, based on literature review, inflammation-associated genes were identified, including interleukin (*IL*)1 β , *IL6*, *IL7*, *IL4*, *IL10*, CD44 molecule (Indian blood group) (*Cd44*), cytochrome b-245 β -chain (*Cybb*), intercellular adhesion molecule 1 (*Icam1*), cytochrome b-245 α -chain (*Cyba*), HCK proto-oncogene, Src family tyrosine kinase (*Hck*), caspase 1 (*Casp1*), transforming growth factor $\beta 1$ (*Tgfb1*), Rac family small GTPase 2 (*Rac2*), integrin subunit $\beta 2$ (*Itgb2*) and C-X-C motif chemokine receptor 4 (*Cxcr4*) (3,4,9). The fold-change of expression of each gene (post-SCI data vs. sham data in each region) at three different time-points (24 and 72 h, and 7 days) was determined.

Results

Data preprocessing and DEG screening. Box plots presenting post-SCI and sham surgery data in the different regions at different time points (M region: 0.5, 4, 24 and 72 h, and 7 and 28 days; R and C regions: 24 and 72 h, and 7 days) prior to and following data normalization are presented in Fig. 1. The results demonstrated that the gene expression values in each sample were similar following normalization. Following data pre-processing, DEGs between the post-SCI groups and sham groups in the M region at six time points were analyzed. There were 226, 495, 566, 816, 1,286 and 1,023 upregulated DEGs at 0.5, 4, 24, and 72 h and 7 and 28 days, respectively. Additionally, a total of 148, 212, 756, 659, 438 and 319 downregulated DEGs were identified at the six time points, respectively (Fig. 2). There was an increased number of upregulated DEGs compared with downregulated DEGs at the five time points (0.5, 4 and 72 h, and 7 and 28 days) and the number of upregulated DEGs reached a peak on day 7. By contrast, the number of downregulated DEGs peaked at 24 h and subsequently decreased over time. The above results indicated that gene expression alterations occurred primarily at 24 and 72 h, and 7 days following injury.

KEGG pathway and GO enrichment analysis of up- and downregulated DEGs. A number of studies have investigated alterations in gene expression between 0.5 and 6 h following SCI in mice (17,18). The present study focused on DEGs at 24 and 72 h, and 7 days following injury. The top five enriched KEGG pathways of up- and downregulated DEGs at each time point (24 and 72 h, and 7 days) are presented in Fig. 3A and B, respectively. At 24 h, upregulated DEGs were enriched in pathways including 'extracellular matrix-receptor interaction' ($P=2.08\times 10^{-6}$), 'focal adhesion' ($P=3.33\times 10^{-6}$) and 'regulation of the actin cytoskeleton' ($P=7.98\times 10^{-6}$). At 72 h and 7 days, upregulated DEGs were primarily involved in inflammation- and immunity-associated pathways. 'Leukocyte transendothelial migration' ($P=1.05\times 10^{-9}$) and 'natural killer cell-mediated cytotoxicity' ($P=1.02\times 10^{-5}$) were enriched at 72 h. 'Lysosome' ($P=5.57\times 10^{-10}$), 'Toll-like

Figure 1. Data normalization of differentially expressed genes. Box plots of gene expression in the post-SCI and sham groups from the site of trauma (A) prior to and (B) following normalization. Box plots of gene expression in the post-SCI and sham groups at the immediately adjacent R and C regions (C) prior to and (D) following normalization. R, rostral; C, caudal; M, trauma site; SCI, spinal cord injury.

Figure 2. Differentially expressed genes between post-spinal cord injury and sham samples at each time-point following spinal cord injury.

receptor signaling pathway' ($P=5.93 \times 10^{-8}$) and 'leukocyte transendothelial migration' ($P=2.48 \times 10^{-7}$) were enriched at 7 days following injury. Notably, downregulated DEGs were enriched in the neuronal function and synaptic transmission-associated pathways, including 'neuroactive ligand-receptor interactions' ($P=1.43 \times 10^{-5}$) at 24 h, 'synaptic vesicle cycle' ($P=2.32 \times 10^{-7}$) and

'calcium signaling pathway' ($P=3.45 \times 10^{-7}$) at 72 h, and 'neuroactive ligand-receptor interaction' ($P=4.95 \times 10^{-5}$) and 'glutamatergic synapse' ($P=2.23 \times 10^{-4}$) at 7 days following injury. Other downregulated DEGs were most significantly enriched in 'steroid biosynthesis' ($P=1.08 \times 10^{-5}$).

The GO terms (biological process) of up- and downregulated DEGs are summarized in Table I. The results demonstrated that upregulated DEGs were most enriched in 'response to stress' ($P=2.30 \times 10^{-27}$) at 24 h. Additionally, 'immune system process' was the most enriched function at 72 h and 7 days ($P=3.59 \times 10^{-27}$ and $P=7.77 \times 10^{-68}$, respectively). By contrast, downregulated DEGs were most enriched in 'single-organism cellular process' ($P=9.30 \times 10^{-8}$) at 24 h following injury. Furthermore, downregulated DEGs were enriched in neuronal function- and synaptic transmission-associated biological process terms at 72 h and 7 days following injury. At 72 h 'synaptic signaling' ($P=1.21 \times 10^{-30}$) and 'chemical synaptic transmission' ($P=7.68 \times 10^{-25}$) were most enriched. At 7 days following injury, 'synaptic signaling' ($P=5.83 \times 10^{-9}$) and 'anterograde trans-synaptic signaling' ($P=2.51 \times 10^{-8}$) were most enriched. 'Sterol biosynthetic process' ($P=4.96 \times 10^{-10}$) was the most significantly enriched biological process at 7 days following injury. Collectively, in the present study, upregulated

Figure 3. Kyoto Encyclopedia of Genes and Genomes pathways of (A) upregulated and (B) downregulated differentially expressed genes at three time-points (24 and 72 h, and 7 days) following spinal cord injury.

DEGs were predominantly associated with immune and inflammatory functions, while downregulated DEGs were involved in neuronal function, synaptic transmission and steroid biosynthesis.

PPI network construction and functional module analysis. Venn diagram analysis of DEGs at 24 and 72 h, and 7 days is presented in Fig. 4A. There were 138 common upregulated and 20 overlapping downregulated DEGs at these three time points. Subsequently, a more comprehensive bioinformatics analysis was performed using Cytoscape software, a tool for predicting PPI networks. The results revealed that immune and inflammatory functions were enriched in co-upregulated DEGs (Fig. 4B). Additionally, neuronal functions and 'steroid hormone biosynthesis' were enriched in the co-downregulated DEGs (Fig. 4C).

Trends in the expression levels of neuronal function- and inflammatory response-associated genes. The results of pathway, functional enrichment and PPI network analyses demonstrated that inflammatory- and neuronal-associated

functions serve roles in post-SCI mice in the M region at three different time points (24 and 72 h, and 7 days). Alterations in these functions were further investigated via temporal and spatial analysis of the expression of numerous genes. Neuronal function- and synaptic transmission-associated genes, including *Chrna7*, *Atp2b2*, *Rims1*, *Camk2g*, and *Trhr*, were downregulated at 72 h compared with the expression at 24 h post-SCI in the M region (Fig. 5A). Furthermore, neuronal function associated genes, *Chrna7*, *Syn2* and *Unc13a*, were significantly upregulated on day 7 compared with the expression at 72 h post-SCI. Similar alterations were observed in R and C regions at three different time points post-SCI, although the overall fold-changes with time were minimal (Fig. 5A). Genes involved in inflammatory processes exhibited different and complex alterations (Fig. 5B). The expression levels of inflammatory-associated genes (*IL1b* and *IL6*) decreased significantly at 72 h following injury in the M region compared with levels at the 24 h time interval. However, similar alterations were not observed in the R and C regions. By contrast, the expression levels of other genes associated with inflammatory processes (*Cd44*, *Cybb*, *Cyba*, *Hck*, *Caspl*, *Itgb2* and *Cxcr4*)

Table I. GO terms enriched by differentially expressed genes at three time-zpoints following spinal cord injury.

A, Injury vs. sham (24 h)				
Category	Term	Biological process	No. genes	P-value
Upregulated				
	GO:0006950	Response to stress	144	2.30x10 ⁻²⁷
	GO:0070887	Cellular response to chemical stimulus	107	9.95x10 ⁻²²
	GO:0050896	Response to stimulus	233	3.47x10 ⁻²¹
	GO:0010033	Response to organic substance	110	9.68x10 ⁻²¹
	GO:0006952	Defense response	73	3.99x10 ⁻²⁰
Downregulated				
	GO:0044763	Single-organism cellular process	329	9.30x10 ⁻⁸
	GO:0044699	Single-organism process	363	1.70x10 ⁻⁷
	GO:0048512	Circadian behavior	9	3.21x10 ⁻⁷
	GO:0007622	Rhythmic behavior	9	6.19x10 ⁻⁷
	GO:0007275	Multicellular organism development	154	1.11x10 ⁻⁶
B, Injury vs. sham (72 h)				
Category	Term	Biological process	No. genes	P-value
Upregulated				
	GO:0002376	Immune system process	148	3.59x10 ⁻²⁷
	GO:0006952	Defense response	90	4.21x10 ⁻¹⁸
	GO:0006955	Immune response	87	5.70x10 ⁻¹⁸
	GO:0006954	Inflammatory response	52	9.35x10 ⁻¹⁸
	GO:0002684	Positive regulation of immune system process	64	6.12x10 ⁻¹⁶
Downregulated				
	GO:0099536	Synaptic signaling	56	1.21x10 ⁻³⁰
	GO:0098916	Anterograde trans-synaptic signaling	54	3.26x10 ⁻²⁹
	GO:0099537	Trans-synaptic signaling	54	3.65x10 ⁻²⁹
	GO:0007268	Chemical synaptic transmission	57	7.68x10 ⁻²⁵
	GO:0007267	Cell-cell signaling	74	2.96x10 ⁻²⁰
C, Injury vs. sham (day 7)				
Category	Term	Biological process	No. genes	P-value
Upregulated				
	GO:0002376	Immune system process	286	7.77x10 ⁻⁶⁸
	GO:0006952	Defense response	188	7.30x10 ⁻⁵²
	GO:0006950	Response to stress	319	6.25x10 ⁻⁴⁵
	GO:0006955	Immune response	166	2.74x10 ⁻⁴¹
	GO:0002682	Regulation of immune system process	164	1.22x10 ⁻³⁹
Downregulated				
	GO:0016126	Sterol biosynthetic process	9	4.96x10 ⁻¹⁰
	GO:0099536	Synaptic signaling	19	5.83x10 ⁻⁹
	GO:1902653	Secondary alcohol biosynthetic process	7	2.09x10 ⁻⁸
	GO:0098916	Anterograde trans-synaptic signaling	18	2.51x10 ⁻⁸
	GO:0099537	Trans-synaptic signaling	18	2.59x10 ⁻⁸
GO, Gene Ontology.

Experimental modeling of SCI in animals has been widely used to investigate the complex secondary injury cascade (1,19). The GSE5296 database consists of transcriptome data obtained from spinal cord sections from injury sites, and immediately

In the present study, at the M site, there were 374, 707, 1,322, 1,475, 1,724 and 1,342 DEGs identified at 0.5, 4, 24 and 72 h, and 7 and 28 days following injury, respectively. The results of the present study indicated that the majority of alterations in molecular events occurred at 7 days following injury and decreased over time. In addition, Wang *et al* (8)

Figure 5. Trend charts of gene expression in R, M and C regions. The trend charts of expression of genes associated with (A) neuronal function, including *Chrna7*, *Syn2*, *Kcnc1*, *Atp2b2*, *Unc13a*, *Rims1*, *Camk2g*, *Camk2a*, *Trhr* and *Grm1* and inflammatory response, including (B) *IL1b*, *IL6*, *IL7*, *IL4* and *IL10*, and (C) *Cd44*, *Icam1*, *Cybb*, *Cyba*, *Hck*, *Casp1*, *Tgfb1*, *Rac2*, *Itgb2* and *Cxcr4*. Significant differences between expression levels at different time points were identified (* $P < 0.05$ and ** $P < 0.01$, 24 h vs. 72 h; # $P < 0.05$ and ## $P < 0.01$, 72 h and 7 days). R, rostral; C, caudal; M, trauma site. IL, interleukin; *Cd44*, CD44 molecule (Indian blood group); *Cybb*, cytochrome b-245 β -chain; *Icam1*, intercellular adhesion molecule 1; *Cyba*, cytochrome b-245 α -chain; *Hck*, HCK proto-oncogene, Src family tyrosine kinase; *Casp1*, caspase 1; *Tgfb1*, transforming growth factor β 1; *Rac2*, Rac family small GTPase 2; *Itgb2*, integrin subunit β 2; *Cxcr4*, C-X-C motif chemokine receptor 4; *Chrna7*, cholinergic receptor nicotinic α 7 subunit; *Syn2*, synapsin II; *Kcnc1*, potassium voltage-gated channel subfamily C member 1; *Atp2b2*, ATPase plasma membrane Ca^{2+} transporting 2; *Unc13a*, unc-13 homolog A; *Rims1*, regulating synaptic membrane exocytosis 1; *Camk2g*, calcium/calmodulin dependent protein kinase II γ ; *Camk2a*, calcium/calmodulin dependent protein kinase II α ; *Trhr*, thyrotropin releasing hormone receptor; *Grm1*, glutamate metabotropic receptor 1.

demonstrated that the number of DEGs decreased in a time-dependent manner (1,942, 396, 188 and 193 DEGs were identified at 3, 7, 14 days and 1 month, respectively). Although the R and C regions exhibited decreased fold changes in gene expression compared with the M region, further studies of these two areas may improve the understanding of the overall process of SCI.

Numerous molecular events were detected by analyzing up- and downregulated DEGs. As demonstrated in a previous study, downregulated DEGs were primarily enriched in neuronal functions (9). The spectra of expression changes in neuronal function- and synaptic transmission-associated genes (*Chrna7*, *Syn2*, *Kcnc1*, *Atp2b2*, *Unc13a*, *Rims1*, *Camk2g*, *Camk2a*, *Trhr* and *Grm1*) from 24 h to 7 days post-SCI reflects the regulation between the degeneration and survival of injured tissues. Insight into time-dependent alterations in structural and functional neuronal biomarkers may be useful for developing protective or regenerative therapeutic interventions.

As a regulator of degeneration and regeneration of the spinal cord, inflammation is a hallmark of the secondary SCI process (20). In the present study, KEGG pathway and GO enrichment analyses, and the PPI network, revealed that the upregulated DEGs were primarily associated with 'immune system' process and the inflammatory response, particularly at 72 h and 7 days post-SCI. As previously demonstrated, there is an association between the severity of SCI and the intensity of the acute inflammatory response, which includes proinflammatory cytokines and immune cells (20,21). A significant increase in the expression of proinflammatory (*IL-1b*, *IL-6* and *IL-7*) and anti-inflammatory cytokines (*IL-4* and *IL-10*) in the present study reflects both the regulation between degeneration and survival of injured tissues. A protective strategy is to target the process of inflammation and limit the infiltration of immune cells into the injury site (22-24). Notably, another group of inflammation-associated genes, including *Cd44*, *Cybb*, *Icam1*, *Cyba*, *Hck*, *Casp1*, *Tgfb1*, *Rac2*, *Itgb2* and *Cxcr4*

exhibited a different temporal pattern compared with proinflammatory genes (*IL-1 β* , *IL-6* and *IL-7*), indicating a complex inflammatory immune microenvironment at the damaged site that requires further analysis. These findings the importance of monitoring inflammation over time following SCI (20).

In the present study, downregulated DEGs at day 7 were primarily enriched in the 'steroid biosynthesis' process. Steroids may be functionally divided into cholesterol, corticosteroids, sex steroids, neuroactive steroids and vitamin D (25,26). A number of studies have investigated the association between steroid metabolism and SCI (27-34). Estrogen may attenuate inflammation and promote neural survival and regeneration following SCI (27,28). Statins, known as cholesterol-controlling drugs, may significantly enhance neuronal and oligodendrocyte survival, in addition to decreasing the levels of proinflammatory cytokines (29). Previous studies additionally suggested that individuals with SCI are at an increased risk of vitamin D deficiency (30-32). Furthermore, neuroactive steroids are naturally occurring steroids that impact behavior, alter the excitability of neurons and interact with specific neurotransmitter receptors (33,34). Therefore, targeting steroid biosynthesis as a therapeutic approach for neuroprotective and neuroregenerative purposes merits further investigation.

However, one limitation of the present study was that the raw data did not include gene expression data from samples at 3 or 6 months following injury, limiting the information regarding molecular processes during the progression of SCI. KEGG pathway and GO enrichment analyses of data for the very acute phase (0.5 and 4 h post-SCI) or at 1 month post-SCI were not included in the present study. Additionally, future comprehensive analysis of transcriptome data from the adjacent R and C regions at each time point post-SCI may reveal the complex alterations that occur during the pathophysiological process.

In conclusion, the present study revealed that inflammatory response, immune processes, neuronal-associated functions and 'steroid biosynthesis' serve roles in the progression of SCI. Furthermore, the M region exhibited increased fold-changes in the expression of genes associated with inflammatory responses and neuronal function compared with the R and C regions at different time-points post-SCI. However, *in vivo* and *in vitro* studies are required to determine the specific roles of these molecular events in SCI.

Acknowledgements

We would like to thank Dr Yu-qing Jiang (Department of Orthopedics, the Affiliated Hospital of Nanjing Medical University, Changzhou No. 2 People's Hospital) for assistance with the present study.

Funding

The present study was supported by grants from the National Natural Science Foundation of China (grant nos. 8177235, 81520108018 and 81472080), the Jiangsu Committee of Science and Technology-Social Development Plan (grant no. BE2017755) and the Nanjing Committee of Science and Technology (grant no. 201505005).

Availability of data and materials

All data generated or analysed during this study are included in this published article.

Authors' contributions

GYG designed the present study. SJZ, WZ, JC and YJL performed the data analysis and statistical analysis. SJZ and WZ wrote and revised the manuscript. GYG supervised the present study. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Ahuja CS, Wilson JR, Nori S, Kotter MRN, Druschel C, Curt A and Fehlings MG: Traumatic spinal cord injury. *Nat Rev Dis Primers* 3: 17018, 2017.
- Jia X, Kowalski RG, Sciubba DM and Geocadin RG: Critical care of traumatic spinal cord injury. *J Intensive Care Med* 28: 12-23, 2013.
- Siddiqui AM, Khazaei M and Fehlings MG: Translating mechanisms of neuroprotection, regeneration and repair to treatment of spinal cord injury. *Prog Brain Res* 218: 15-54, 2015.
- Ahuja CS and Fehlings M: Concise review: Bridging the gap: Novel neuroregenerative and neuroprotective strategies in spinal cord injury. *Stem Cells Transl Med* 5: 914-924, 2016.
- Kim YH, Ha KY and Kim SI: Spinal cord injury and related clinical trials. *Clin Orthop Surg* 9: 1-9, 2017.
- Bareyre FM and Schwab ME: Inflammation, degeneration and regeneration in the injured spinal cord: Insights from DNA microarrays. *Trends Neurosci* 26: 555-563, 2003.
- Ydens E, Palmers I, Hendrix S and Somers V: The next generation of biomarker research in spinal cord injury. *Mol Neurobiol* 54: 1482-1499, 2017.
- Wang W, Liu R, Xu Z, Niu X, Mao Z, Meng Q and Cao X: Further insight into molecular mechanism underlying thoracic spinal cord injury using bioinformatics methods. *Mol Med Rep* 12: 7851-7858, 2015.
- Chen G, Fang X and Yu M: Regulation of gene expression in rats with spinal cord injury based on microarray data. *Mol Med Rep* 12: 2465-2472, 2015.
- Wen T, Hou J, Wang F, Zhang Y, Zhang T and Sun T: Comparative analysis of molecular mechanism of spinal cord injury with time based on bioinformatics data. *Spinal Cord* 54: 431-438, 2016.
- Zhu Z, Shen Q, Zhu L and Wei X: Identification of pivotal genes and pathways for spinal cord injury via bioinformatics analysis. *Mol Med Rep* 16: 3929-3937, 2017.
- Duan H, Ge W, Zhang A, Xi Y, Chen Z, Luo D, Cheng Y, Fan KS, Horvath S, Sofroniew MV, *et al*: Transcriptome analyses reveal molecular mechanisms underlying functional recovery after spinal cord injury. *Proc Natl Acad Sci USA* 112: 13360-13365, 2015.
- Carvalho BS and Irizarry RA: A framework for oligonucleotide microarray preprocessing. *Bioinformatics* 26: 2363-2367, 2010.
- Huang da W, Sherman BT and Lempicki RA: Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 4: 44-57, 2009.

15. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B and Ideker T: Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Res* 13: 2498-2504, 2003.
16. Sun N, Sun W, Li S, Yang J, Yang L, Quan G, Gao X, Wang Z, Cheng X, Li Z, *et al*: Proteomics analysis of cellular proteins co-immunoprecipitated with nucleoprotein of influenza A virus (H7N9). *Int J Mol Sci* 16: 25982-25998, 2015.
17. Di Giovanni S, Knoblich SM, Brandoli C, Aden SA, Hoffman EP and Faden AI: Gene profiling in spinal cord injury shows role of cell cycle in neuronal death. *Ann Neurol* 53: 454-468, 2003.
18. Carmel JB, Galante A, Soteropoulos P, Tolias P, Recce M, Young W and Hart RP: Gene expression profiling of acute spinal cord injury reveals spreading inflammatory signals and neuron loss. *Physiol Genomics* 7: 201-213, 2001.
19. Kjell J and Olson L: Rat models of spinal cord injury: From pathology to potential therapies. *Dis Model Mech* 9: 1125-1137, 2016.
20. Saghaizadeh A and Rezaei N: The role of timing in the treatment of spinal cord injury. *Biomed Pharmacother* 92: 128-139, 2017.
21. Kinoshita T, Nakamura T, Umemoto Y, Kojima D, Moriki T, Mitsui T, Goto M, Ishida Y and Tajima F: Increase in interleukin-6 immediately after wheelchair basketball games in persons with spinal cord injury: Preliminary report. *Spinal Cord* 51: 508-510, 2013.
22. Hu JZ, Huang JH, Xiao ZM, Li JH, Li XM and Lu HB: Tetramethylpyrazine accelerates the function recovery of traumatic spinal cord in rat model by attenuating inflammation. *J Neurol Sci* 324: 94-99, 2013.
23. David BT, Ratnayake A, Amarante MA, Reddy NP, Dong W, Sampath S, Heary RF and Elkabes S: A toll-like receptor 9 antagonist reduces pain hypersensitivity and the inflammatory response in spinal cord injury. *Neurobiol Dis* 54: 194-205, 2013.
24. Jin W, Wang J, Zhu T, Yuan B, Ni H, Jiang J, Wang H and Liang W: Anti-inflammatory effects of curcumin in experimental spinal cord injury in rats. *Inflamm Res* 63: 381-387, 2014.
25. Schroepfer GJ Jr: Sterol biosynthesis. *Ann Rev Biochem* 50: 585-621, 1981.
26. Ghayee HK and Auchus RJ: Basic concepts and recent developments in human steroid hormone biosynthesis. *Rev Endocr Metab Disord* 8: 289-300, 2007.
27. Elkabes S and Nicot AB: Sex steroids and neuroprotection in spinal cord injury: A review of preclinical investigations. *Exp Neurol* 259: 28-37, 2014.
28. Brotfain E, Gruenbaum SE, Boyko M, Kutz R, Zlotnik A and Klein M: Neuroprotection by estrogen and progesterone in traumatic brain injury and spinal cord injury. *Curr Neuropharmacol* 14: 641-653, 2016.
29. Eftekharpour E, Nagakannan P, Iqbal MA and Chen QM: Mevalonate cascade and small Rho GTPase in spinal cord injury. *Curr Mol Pharmacol* 10: 141-151, 2017.
30. Flueck JL and Perret C: Vitamin D deficiency in individuals with a spinal cord injury: A literature review. *Spinal Cord* 55: 428-434, 2017.
31. Bauman WA, Morrison NG and Spungen AM: Vitamin D replacement therapy in persons with spinal cord injury. *J Spinal Cord Med* 28: 203-207, 2005.
32. Oleson CV, Patel PH and Wuermser LA: Influence of season, ethnicity and chronicity on vitamin D deficiency in traumatic spinal cord injury. *J Spinal Cord Med* 33: 202-213, 2010.
33. Giatti S, Garcia-Segura LM and Melcangi RC: New steps forward in the neuroactive steroid field. *J Steroid Biochem Mol Biol* 153: 127-134, 2015.
34. Tuem KB and Atey TM: Neuroactive steroids: Receptor interactions and responses. *Front Neurol* 8: 442, 2017.