

Identification and functional analysis of differentially expressed genes associated with cerebral ischemia/reperfusion injury through bioinformatics methods

XIAOLI SHAO^{1*}, WANGXIAO BAO^{2*}, XIAOQIN HONG¹, HUIHUA JIANG¹ and ZHI YU¹

¹Department of Neurology, Chun'an First People's Hospital, Zhejiang People's Hospital Chun'an Branch, Hangzhou, Zhejiang 311700; ²Department of Neurology, First Affiliated Hospital of Zhejiang University School of Medicine, Hangzhou, Zhejiang 310003, P.R. China

Received October 11, 2017; Accepted January 18, 2018

DOI: 10.3892/mmr.2018.9135

Abstract. Cerebral ischemia/reperfusion (I/R) injury results in detrimental complications. However, little is known about the underlying molecular mechanisms involved in the reperfusion stage. The aim of the present study was to identify a gene expression profile associated with cerebral ischemia/reperfusion injury. The GSE23160 dataset, which comprised data from sham control samples and post-I/R injury brain tissues that were obtained using a middle cerebral artery occlusion (MCAO) model at 2, 8 and 24 h post-reperfusion, was downloaded from the Gene Expression Omnibus database. The differentially expressed genes (DEGs) in the MCAO samples compared with controls were screened using the GEO2R web tool. Gene Ontology (GO) analysis and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis for DEGs was performed using the online tool DAVID. Furthermore, a protein-protein interaction (PPI) network was constructed using the STRING database and Cytoscape software. In total, 32 DEGs at 2 h post-reperfusion, 39 DEGs at 8 h post-reperfusion and 91 DEGs at 24 h post-reperfusion were identified, while 15 DEGs were common among all three groups. GO analysis revealed that the DEGs at all three time-points were enriched in 'chemotaxis' and 'inflammatory response' terms, while KEGG pathway analysis demonstrated that DEGs were significantly enriched in the 'chemokine signaling pathway'. Furthermore, following PPI network construction, Cxcl1 was identified as the only hub gene that was common among all three time-points. In conclusion,

the present study has demonstrated a global view of the potential molecular differences following cerebral I/R injury and may contribute to an improved understanding of the reperfusion stage, which may ultimately aid in the development of future clinical strategies.

Introduction

Globally, stroke is the second most frequent cause of mortality and the primary cause of serious long-term disability worldwide (1). Of all strokes, 87% are ischemic (2). Various mechanisms underlying ischemic stroke are driven by cell-cell interactions within brain, including excitotoxicity, calcium dysregulation, oxidative and nitrosative Stress, cortical spreading depolarizations, inflammation, necrosis, necroptosis and autophagy (3). In addition to a narrow therapeutic time window (4), ischemic stroke remains difficult to manage.

Although reperfusion has been proven to be beneficial for ischemic stroke (5), reperfusion may result in detrimental secondary damage, which is termed ischemia/reperfusion (I/R) injury. Early reperfusion of ischemic brain tissue has been associated with various negative consequences, including blood-brain barrier breakdown, which may result in cerebral edema and/or brain hemorrhage, neurovascular damage and neuronal death (6). Angiogenesis and vasculogenesis have also been detected following reperfusion (7). In addition, inflammation is induced by reperfusion injury and contributes negatively to long-term disease prognosis (8). The inflammatory response may result in subsequent oxidative injury, excitotoxicity and neuronal cell death (9). Chemokines, produced by resident microglial cells and other immune cells in the brain, contribute to the recruitment of circulating leukocytes and exaggerate the inflammatory response. Chemokines have been demonstrated to have both deleterious and beneficial roles in ischemia/reperfusion injury (10).

Microarray analysis has been previously employed to identify molecular variations in cerebral I/R injury (11,12). However, gene expression profiles at different reperfusion periods have not been investigated extensively. Therefore, the present study employed a microarray dataset from the Gene Expression Omnibus (GEO) database and screened for differentially expressed genes (DEGs) between control samples and

Correspondence to: Dr Xiaoli Shao, Department of Neurology, Chun'an First People's Hospital, Zhejiang People's Hospital Chun'an Branch, 1869 Huanhu North Road, Hangzhou, Zhejiang 311700, P.R. China
E-mail: shaoxiaoli188@163.com

*Contributed equally

Key words: differentially expressed genes, gene ontology, cerebral ischemia/reperfusion injury, inflammatory response, chemokines

cerebral I/R samples at 2, 8 and 24 h post-reperfusion, and subsequently analyzed the functions and interactions of these DEGs. The results of the current study may aid in improving the understanding of the molecular mechanisms underlying cerebral I/R injury.

Materials and methods

Microarray data. Microarray gene expression profiles from GSE23160 (12) were obtained from the GEO database (<http://www.ncbi.nlm.nih.gov/geo/>), which is based on the platform of GPL6885 using Illumina MousRef-8 v2.0 Expression BeadChip (Illumina, Inc., San Diego, CA, USA). All of the samples were taken from male C57BL/6J mice (8–10 weeks). Following 2 h suture-induced middle cerebral artery occlusion (MCAO), the animals underwent reperfusion for 2, 8 or 24 h. Tissue extractions at 2, 8 and 24 h post-reperfusion and sham controls (n=4 per group) were included in this dataset.

Identification of DEGs. GEO2R (<http://www.ncbi.nlm.nih.gov/geo/geo2r/>), an R-based web application (13), was employed to analyze DEGs between MCAO samples and sham samples. $P < 0.05$ and $|\log_{2}FC| \geq 1.2$ were set as the threshold criteria to identify genes that were differentially expressed in MCAO models. Subsequently, the DEGs at 2, 8 and 24 h post-reperfusion were screened for subsequent analyses. A Venn diagram was produced to indicate the intersection among DEGs in the various MCAO groups using FunRich software (version 2.1.1; www.funrich.org) (14).

Functional enrichment analysis of DEGs. To identify the biological processes, cellular components, molecular functions and biological pathways that the DEGs were significantly enriched in, Gene Ontology (GO) enrichment (15) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analyses (16) were performed using the online tool Database for Annotation, Visualization and Integrated Discovery (DAVID; <http://david.abcc.ncifcrf.gov/>). $P < 0.05$ was considered to indicate a significantly enriched term or pathway, which was calculated using a hypergeometric test. Heat map illustration of DEGs was performed with heat map illustrator software (version 1.0.3.7; <http://hemi.biocuckoo.org>) (17).

Construction of the PPI network. To further investigate the underlying molecular mechanisms of cerebral I/R injury, protein-protein interaction (PPI) networks for the DEGs were constructed using the Search Tool for the Retrieval of Interacting Genes (STRING) database (<http://www.string-db.org/>) (18). A combined score of > 0.4 was selected to construct the PPI networks. The obtained PPI networks at 24 h post-reperfusion were subsequently visualized using Cytoscape software (version 3.5.1) (19). Finally, the topological properties of the networks at 2, 8 and 24 h post-reperfusion were analyzed and the degree of each node was calculated; genes with a degree > 10 were defined as hub genes.

Results

Identification of DEGs. As demonstrated in Fig. 1, 32 DEGs at 2 h post-reperfusion, 39 DEGs at 8 h post-reperfusion




Figure 1. Venn diagram presenting the DEGs between cerebral I/R and sham control samples at 2, 8 and 24 h post-reperfusion. A total of 15 DEGs were common among all three post-reperfusion time-points. DEGs, differentially expressed genes; I/R, ischemia/reperfusion.

and 91 DEGs at 24 h post-reperfusion were identified in the MCAO samples compared with the controls. Among them, 15 DEGs were common to all three injury samples, including C-C motif chemokine ligand (CCL)7, suppressor of cytokine signaling 3, CCL4, activating transcription factor 3, lipocalin 2, hemoglobin α adult chain 1, gap junction protein $\beta 2$, CD14 antigen, CCL3, heat shock protein 1A, S100 calcium-binding protein A8 (calgranulin A), C-X-C motif chemokine ligand 1 (CXCL1), epithelial membrane protein 1, tissue inhibitor of metalloproteinase 1 and zinc finger protein 36, all of which were upregulated in the MCAO samples (Table I). Heat map and PPI network analysis at 2 and 8 h post-reperfusion are not presented due to the small number of identified DEGs.

Bioinformatics analyses of DEGs. To further the understanding of the screened DEGs and determine their potential roles following I/R injury, GO functional and KEGG pathway enrichment analyses were performed.

A total of 51 GO enriched terms for biological processes at 2, 8 and 24 h post-reperfusion were obtained. The 10 most enriched GO terms according to the P-value for 2, 8 and 24 h post-reperfusion groups are presented in Table II. Furthermore, ‘chemotaxis’ (GO:0006935), ‘inflammatory response’ (GO:0006954), ‘immune response’ (GO:0006955), ‘G-protein coupled receptor signaling pathway’ (GO:0007186), ‘response to toxic substance’ (GO:0009636), ‘neutrophil chemotaxis’ (GO:0030593), ‘positive regulation of tumor necrosis factor production’ (GO:0032760), ‘positive regulation of GTPase activity’ (GO:0043547), ‘lymphocyte chemotaxis’ (GO:0048247), ‘positive regulation of inflammatory response’ (GO:0050729), ‘cell chemotaxis’ (GO:0060326), ‘chemokine-mediated signaling pathway’ (GO:0070098), ‘positive regulation of ERK1 and ERK2 cascade’ (GO:0070374), ‘cellular response to interferon-gamma’ (GO:0071346), ‘cellular response to interleukin-1’ (GO:0071347), ‘monocyte chemotaxis’ (GO:0002548) and ‘cellular response to tumor necrosis factor’ (GO:0071356) were significantly enriched at all three post-reperfusion time-points (2, 8 and 24 h).

Additionally, DEGs were enriched in various GO cellular component terms; at 2 and 8 h post-reperfusion, DEGs were enriched in ‘extracellular region’ (GO:0005576), while

Table I. DEGs between cerebral I/R and sham control samples that were common among 2, 8 and 24 h post-reperfusion time-points.

Gene symbol	Gene name	Post-reperfusion time-point					
		2 h		8 h		24 h	
		P-value	Log FC	P-value	Log FC	P-value	Log FC
ATF3	Activating transcription factor 3	5.01x10 ⁻⁶	2.026964	3.56x10 ⁻⁵	1.692171	7.49x10 ⁻³	1.591308
CCL3	C-C motif chemokine ligand 3	5.13x10 ⁻¹⁰	3.259239	7.05x10 ⁻⁶	3.291652	4.81x10 ⁻⁴	2.300334
CCL4	C-C motif chemokine ligand 4	1.32x10 ⁻¹⁰	4.429954	1.29x10 ⁻⁶	4.47245	5.83x10 ⁻⁵	3.679271
CCL7	C-C motif chemokine ligand 7	2.18x10 ⁻⁵	1.220192	7.63x10 ⁻⁶	1.426973	4.00x10 ⁻³	1.707457
CD14	CD14 antigen	1.38x10 ⁻⁷	1.757282	5.92x10 ⁻⁷	2.578573	2.56x10 ⁻³	2.353608
CXCL1	C-X-C motif chemokine ligand 1	3.29x10 ⁻⁶	1.806643	4.81x10 ⁻⁸	2.878735	1.43x10 ⁻³	2.596958
EMP1	Epithelial membrane protein 1	1.68x10 ⁻⁶	1.688657	6.78x10 ⁻⁷	1.383669	1.89x10 ⁻³	2.161470
GJB2	Gap junction protein β 2	3.16x10 ⁻⁸	1.999569	1.15x10 ⁻⁷	1.717101	9.13x10 ⁻⁵	1.822626
HBA-A1	Hemoglobin α , adult chain 1	5.38x10 ⁻⁷	2.436023	2.15x10 ⁻⁸	3.757753	4.48x10 ⁻⁴	3.807134
HSPA1A	Heat shock protein 1A	1.69x10 ⁻⁶	3.054788	3.80x10 ⁻⁵	2.545466	1.59x10 ⁻²	1.299110
LCN2	Lipocalin 2	4.06x10 ⁻⁹	1.312245	4.47x10 ⁻¹⁵	3.250233	1.72x10 ⁻⁵	3.408973
S100A8	S100 calcium-binding protein A8 (calgranulin A)	2.29x10 ⁻⁶	1.240885	1.66x10 ⁻⁶	1.655962	1.90x10 ⁻³	3.780381
SOCS3	Suppressor of cytokine signaling 3	3.54x10 ⁻⁸	2.009771	1.74x10 ⁻⁸	2.004998	5.48x10 ⁻⁴	2.361302
TIMP1	Tissue inhibitor of metalloproteinase 1	1.02x10 ⁻⁸	2.376190	3.61x10 ⁻¹⁸	3.403190	1.52x10 ⁻⁵	3.874533
ZFP36	Zinc finger protein 36	2.01x10 ⁻⁵	1.595428	6.26x10 ⁻⁷	1.404682	1.17x10 ⁻²	1.330419

Table II. Top 10 enriched GO biological process terms for DEGs between cerebral ischemia/reperfusion and sham control samples.

A, Top 10 enriched GO biological process terms for DEGs at 2 h post-reperfusion

GO ID	GO term	Count	%	P-value	Genes
GO:0030593	Neutrophil chemotaxis	5	15.63	5.65x10 ⁻⁶	CXCL1, CCL3, S100A8, CCL4, CCL7
GO:0032570	Response to progesterone	4	12.50	1.80x10 ⁻⁵	FOS, OXT, FOSB, GJB2
GO:0006954	Inflammatory response	7	21.88	2.24x10 ⁻⁵	CXCL1, CCL3, S100A8, CCL4, CCL7, CD14, IL1A
GO:0045944	Positive regulation of transcription from RNA polymerase II promoter	10	31.25	2.99x10 ⁻⁵	FOS, CCL3, EGR2, ATF3, EGR4, FOSB, NPAS4, JUNB, IL1A, CYR61
GO:0071356	Cellular response to tumor necrosis factor	5	15.63	3.59x10 ⁻⁵	LCN2, ZFP36, CCL3, CCL4, CCL7
GO:2000503	Positive regulation of natural killer cell chemotaxis	3	9.38	4.25x10 ⁻⁵	CCL3, CCL4, CCL7
GO:0006935	Chemotaxis	5	15.63	4.73x10 ⁻⁵	CCL3, S100A8, CCL4, CCL7, CYR61
GO:0051591	Response to cAMP	4	12.50	8.99x10 ⁻⁵	FOS, OXT, FOSB, JUNB
GO:0070098	Chemokine-mediated signaling pathway	4	12.50	1.13x10 ⁻⁴	CXCL1, CCL3, CCL4, CCL7
GO:0050729	Positive regulation of inflammatory response	4	12.50	1.69x10 ⁻⁴	CCL3, S100A8, CCL4, CCL7

B, Top 10 enriched GO biological process terms for DEGs at 8 h post-reperfusion

GO ID	GO term	Count	%	P-value	Genes
GO:0030593	Neutrophil chemotaxis	10	25.64	8.69x10 ⁻¹⁵	CXCL1, CCL12, CCL3, S100A8, LGALS3, CCL9, CCL4, CCL7, FCGR3, CCL17

Table II. Continued.

B, Top 10 enriched GO biological process terms for DEGs at 8 h post-reperfusion

GO ID	GO term	Count	%	P-value	Genes
GO:0002548	Monocyte chemotaxis	8	20.51	1.19×10^{-12}	CCL12, CCL3, FLT1, LGALS3, CCL9, CCL4, CCL7, CCL17
GO:0070098	Chemokine-mediated signaling pathway	7	17.95	1.09×10^{-9}	CXCL1, CCL12, CCL3, CCL9, CCL4, CCL7, CCL17
GO:0071356	Cellular response to tumor necrosis factor	8	20.51	1.83×10^{-9}	LCN2, ZFP36, CCL12, CCL3, CCL9, CCL4, CCL7, CCL17
GO:0050729	Positive regulation of inflammatory response	7	17.95	2.52×10^{-9}	CCL12, CCL3, S100A8, CCL9, TLR2, CCL4, CCL7
GO:0006935	Chemotaxis	8	20.51	3.01×10^{-9}	CCL12, CCL3, FLT1, S100A8, CCL9, CCL4, CCL7, CCL17
GO:0071346	Cellular response to interferon-gamma	7	17.95	4.02×10^{-9}	CCL12, CCL3, CCL9, CCL4, GBP2, CCL7, CCL17
GO:0048247	Lymphocyte chemotaxis	6	15.38	5.34×10^{-9}	CCL12, CCL3, CCL9, CCL4, CCL7, CCL17
GO:0071347	Cellular response to interleukin-1	7	17.95	1.09×10^{-8}	LCN2, CCL12, CCL3, CCL9, CCL4, CCL7, CCL17
GO:0006954	Inflammatory response	10	25.64	1.76×10^{-8}	CXCL1, CCL12, CCL3, S100A8, CCL9, TLR2, CCL4, CCL7, CD14, CCL17

C, Top 10 enriched GO biological process terms for DEGs at 24 h post-reperfusion

GO ID	GO term	Count	%	P-value	Genes
GO:0006954	Inflammatory response	21	22.83	1.99×10^{-16}	CXCL1, CCL3, CCL2, S100A8, CCL21C, S100A9, CCL9, TLR2, CCL21A, PF4, FPR2, IDO1, CCL4, CCL7, CXCL10, SLC11A1, CYBA, CCL12, CHIL3, CD14, SPP1
GO:0006955	Neutrophil chemotaxis	13	14.13	5.60×10^{-16}	CXCL1, CCL3, CCL2, S100A8, LGALS3, CCL21C, S100A9, CCL21A, CCL9, CCL4, CCL7, CCL12, SPP1
GO:0006956	Chemokine-mediated signaling pathway	11	11.96	1.08×10^{-13}	CXCL1, CCL12, CCL3, CCL2, CCL21C, CCL9, CCL21A, PF4, CCL4, CCL7, CXCL10
GO:0006957	Monocyte chemotaxis	9	9.78	1.54×10^{-11}	CCL12, CCL3, CCL2, LGALS3, CCL21C, CCL9, CCL21A, CCL4, CCL7
GO:0006958	Positive regulation of inflammatory response	10	10.87	1.92×10^{-11}	CCL12, CCL3, CCL2, S100A8, S100A9, CCL9, TGM2, TLR2, CCL4, CCL7
GO:0006959	Cellular response to tumor necrosis factor	11	11.96	1.39×10^{-10}	LCN2, ZFP36, CCL12, CYBA, CCL3, CCL2, CCL21C, CCL9, CCL21A, CCL4, CCL7
GO:0006960	Cellular response to interleukin-1	10	10.87	1.76×10^{-10}	LCN2, CCL12, CCL3, CCL2, CCL21C, CCL9, SAA3, CCL21A, CCL4, CCL7
GO:0006961	Lymphocyte chemotaxis	8	8.70	1.96×10^{-10}	CCL12, CCL3, CCL2, CCL21C, CCL9, CCL21A, CCL4, CCL7
GO:0006962	Chemotaxis	11	11.96	2.80×10^{-10}	CCL12, CCL3, CCL2, S100A8, S100A9, CCL9, PF4, FPR2, CCL4, CCL7, CXCL10
GO:0006963	Cellular response to interferon-gamma	9	9.78	1.33×10^{-9}	CCL12, CCL3, CCL2, CCL21C, CCL9, CCL21A, CCL4, GBP2, CCL7

GO, Gene Ontology; DEGs, differentially expressed genes.

‘membrane’ (GO:0016020) was significantly enriched at both 8 and 24 h post-reperfusion (Table III). Furthermore, Table IV indicates that DEGs were significantly enriched in ‘cytokine activity’ (GO:0005125) and ‘chemokine activity’

(GO:0008009) GO molecular function terms at 2, 8 and 24 h post-reperfusion.

Table V presents the KEGG pathways that were significantly enriched in DEGs. KEGG analysis indicated that ‘chemokine

Table III. GO cellular component terms for DEGs between cerebral ischemia/reperfusion and sham control samples.

A, Enriched GO cellular component terms for DEGs at 2 h post-reperfusion

GO ID	GO term	Count	%	P-value	Genes
GO:0005576	Extracellular region	14	43.75	9.56×10^{-7}	CXCL1, CCL3, AVP, S100A8, PMCH, OXT, CCL4, CCL7, TIMP1, LCN2, NPTX2, IL1A, CD14, CYR61
GO:0005615	Extracellular space	12	37.50	1.02×10^{-5}	LCN2, CXCL1, AVP, CCL3, S100A8, PMCH, OXT, CCL4, CCL7, CD14, IL1A, TIMP1

B, Enriched GO cellular component terms for DEGs at 8 h post-reperfusion

GO ID	GO term	Count	%	P-value	Genes
GO:0005615	Extracellular space	16	41.03	4.81×10^{-8}	CXCL1, CCL3, FLT1, S100A8, LGALS3, PMCH, CCL9, CCL4, CCL7, TIMP1, CCL17, LCN2, CCL12, SERPINA3N, DMKN, CD14
GO:0005576	Extracellular region	14	35.90	1.39×10^{-5}	CXCL1, CCL3, S100A8, LGALS3, PMCH, CCL9, CCL4, CCL7, TIMP1, LCN2, CCL12, SERPINA3N, DMKN, CD14
GO:0009897	External side of plasma membrane	5	12.82	3.17×10^{-3}	LGALS3, OSMR, TLR2, CD14, FCGR3
GO:0016020	Membrane	20	51.28	4.79×10^{-2}	GPR84, FLT1, S100A8, LGALS3, OSMR, FKBP5, MS4A6D, TLR2, SLC10A6, GJB2, FCGR3, HBA-A1, CH25H, PLIN4, HMOX1, ITGAD, SLC15A3, GBP2, EMP1, CD14

C, Enriched GO cellular component terms for DEGs at 24 h post-reperfusion

GO ID	GO term	Count	%	P-value	Genes
GO:0009897	External side of plasma membrane	8	8.70	6.31×10^{-4}	FCGR2B, LGALS3, PDPN, CCL21C, FCGR4, TLR2, CD14, CXCL10
GO:0048237	Rough endoplasmic reticulum lumen	3	3.26	4.29×10^{-4}	LYZ2, LYZ1, CHIL3
GO:0009986	Cell surface	10	10.87	2.33×10^{-3}	SLC11A1, THBD, FCGR2B, LGALS3, TNFRSF12A, IFITM3, FCGR4, TLR2, CD14, ANXA2
GO:0005886	Plasma membrane	34	36.96	8.00×10^{-3}	GPR182, GPR84, S100A8, IFITM2, TNFRSF12A, IFITM3, VIM, S100A9, TLR2, CD52, FPR2, SLC11A1, P2RY6, DAB2, PLIN2, HMOX1, TGM2, STRA6, CLEC4D, ANGPT2, ACTB, PDPN, LILRB4A, GJB2, ANXA2, CYBA, THBD, FCGR2B, HSPB1, SCN4B, RGS9, EMP3, CD14, EMP1
GO:0016020	Membrane	44	47.83	1.14×10^{-2}	GPR182, GPR84, GFAP, TSPO, S100A8, IFITM2, TNFRSF12A, IFITM3, S100A9, TLR2, CD52, FPR2, FXYP5, GLIPR2, SLC11A1, P2RY6, DAB2, PLIN2, HMOX1, CH25H, TGM1, TGM2, STRA6, CLEC4D, ACTB, LGALS3, PDPN, MS4A6D, LILRB4A, GJB2, ANXA2, HBA-A1, CYBA, RAB32, THBD, FCGR2B, SCN4B, RGS9, EMP3, SLC15A3, GBP2, CD14, EMP1, MVP

GO, Gene Ontology; DEGs, differentially expressed genes.

signaling pathway', 'cytokine-cytokine receptor interaction' and 'toll-like receptor signaling pathway' were significantly enriched in DEGs at 2, 8 and 24 h post-reperfusion. Furthermore, as demonstrated in Fig. 2, a total of 11 chemokine signaling pathway-associated genes were overexpressed in 24 h post-reperfusion injury samples compared with the sham control samples.

PPI network analysis. Genes with an interaction degree >10 in the PPI network analysis of DEGs at 2, 8 and 24 h post-reperfusion were defined as hub genes, which are listed in Table VI. CXCL1 was the only gene that was considered to be a hub gene at 2, 8 and 24 h post-reperfusion (Table VI). The constructed PPI network of 24 h post-reperfusion samples is presented in

Table IV. Enriched GO molecular function terms for DEGs between cerebral ischemia/reperfusion and sham control samples.

A, Enriched GO molecular function terms for DEGs at 2 h post-reperfusion

GO ID	GO term	Count	%	P-value	Genes
GO:0005125	Cytokine activity	6	18.75	2.48×10^{-5}	CXCL1, FOS, CCL3, CCL4, CD14, IL1A
GO:0008009	Chemokine activity	4	12.50	6.79×10^{-5}	FOS, CCL3, CCL4, CD14
GO:0000978	RNA polymerase II core promoter proximal region sequence-specific DNA binding	6	18.75	2.84×10^{-4}	FOS, EGR2, ATF3, FOSB, NPAS4, JUNB
GO:0001077	Transcriptional activator activity, RNA polymerase II core promoter proximal region sequence-specific binding	5	15.63	9.82×10^{-4}	FOS, EGR2, FOSB, NPAS4, JUNB
GO:0003690	Double-stranded DNA binding	3	9.38	2.13×10^{-2}	FOS, FOSB, JUNB
GO:0003677	DNA binding	8	25.00	2.85×10^{-2}	ZFP36, FOS, EGR2, ATF3, EGR4, FOSB, NPAS4, JUNB

B, Enriched GO molecular function terms for DEGs at 8 h post-reperfusion

GO ID	GO term	Count	%	P-value	Genes
GO:0008009	Chemokine activity	7	17.95	3.98×10^{-10}	CXCL1, CCL12, CCL3, CCL9, CCL4, CCL7, CCL17
GO:0005125	Cytokine activity	8	20.51	1.54×10^{-7}	CXCL1, CCL12, CCL3, CCL9, CCL4, CCL7, CCL17, TIMP1
GO:0048020	CCR chemokine receptor binding	4	10.26	2.14×10^{-5}	CCL3, CCL9, CCL4, CCL17

C, Enriched GO molecular function terms for DEGs at 24 h post-reperfusion

GO ID	GO term	Count	%	P-value	Genes
GO:0008009	Chemokine activity	11	11.96	1.69×10^{-14}	CXCL1, CCL12, CCL3, CCL2, CCL21C, CCL9, CCL21A, PF4, CCL4, CCL7, CXCL10
GO:0005125	Cytokine activity	12	13.04	4.78×10^{-9}	CXCL1, CCL12, CCL3, CCL2, CCL9, PF4, CCL4, CCL7, SPP1, TIMP1, CXCL10, IL11
GO:0048020	CCR chemokine receptor binding	5	5.44	8.52×10^{-6}	CCL3, CCL21C, CCL9, CCL21A, CCL4
GO:0031727	CCR2 chemokine receptor binding	3	3.26	1.30×10^{-4}	CCL12, CCL2, CCL7
GO:0008201	Heparin binding	4	4.35	3.43×10^{-2}	CCL2, PF4, CCL7, CXCL10
GO:0020037	Heme binding	4	4.35	4.94×10^{-2}	HBA-A1, CYBA, HMOX1, IDO1

GO, Gene Ontology; DEGs, differentially expressed genes.

Fig. 3, which contains 67 nodes and 281 edges. Each node represents a DEG and each edge represents a PPI between two DEGs. At 24 h post-reperfusion, 23 genes served as hub genes, and of these hub genes, CCL2 exhibited the highest degree (Fig. 3).

Discussion

In the present study, 32 DEGs at 2 h, 39 DEGs at 8 h and 91 DEGs at 24 h post-reperfusion injury were identified between cerebral I/R and sham control samples. Previous studies have performed bioinformatics analysis to identify DEGs between MCAO models and controls (11,20-23). However, to the best of our knowledge, the present study is the first to perform global gene

expression profiling at three time-points following reperfusion, and the findings may lead to improvements in the understanding of the pathophysiological process of cerebral I/R injury. DEGs associated with inflammation have previously been associated with cerebral I/R injury (20,22), and the results of the present study were consistent with these previous reports, indicating a persistent inflammatory response in cerebral I/R injury.

In the current study, enrichment analysis revealed that 'chemotaxis', 'chemokine activity' and 'chemokine signaling pathway' terms were significantly enriched for the obtained DEGs. Furthermore, members of the chemokine family were the most abundant among the upregulated genes that were common among 2, 8 and 24 h post-reperfusion

Table V. Enriched KEGG pathways for DEGs between cerebral ischemia/reperfusion and sham control samples.

A, Enriched KEGG pathways for DEGs at 2 h post-reperfusion

KEGG entry	Pathway name	Count	%	P-value	Genes
mmu05132	Salmonella infection	6	18.75	1.27×10^{-6}	CXCL1, FOS, CCL3, CCL4, CD14, IL1A
mmu04380	Osteoclast differentiation	5	15.63	2.68×10^{-4}	FOS, SOCS3, FOSB, JUNB, IL1A
mmu04620	Toll-like receptor signaling pathway	4	12.50	2.11×10^{-3}	FOS, CCL3, CCL4, CD14
mmu04062	Chemokine signaling pathway	4	12.50	1.34×10^{-2}	CXCL1, CCL3, CCL4, CCL7
mmu04060	Cytokine-cytokine receptor interaction	4	12.50	2.41×10^{-2}	CCL3, CCL4, CCL7, IL1A
mmu05166	HTLV-I infection	4	12.50	3.35×10^{-2}	ZFP36, FOS, EGR2, ATF3

B, Enriched KEGG pathways for DEGs at 8 h post-reperfusion

KEGG entry	Pathway name	Count	%	P-value	Genes
mmu04062	Chemokine signaling pathway	7	17.95	2.28×10^{-5}	CXCL1, CCL12, CCL3, CCL9, CCL4, CCL7, CCL17
mmu04060	Cytokine-cytokine receptor interaction	6	15.38	8.01×10^{-4}	CCL12, CCL3, FLT1, OSMR, CCL4, CCL7
mmu05132	Salmonella infection	4	10.26	1.72×10^{-3}	CXCL1, CCL3, CCL4, CD14
mmu04620	Toll-like receptor signaling pathway	4	10.26	3.60×10^{-3}	CCL3, TLR2, CCL4, CD14
mmu04145	Phagosome	4	10.26	1.61×10^{-2}	TLR2, TUBB6, CD14, FCGR3
mmu05142	Chagas disease (American trypanosomiasis)	3	7.69	4.02×10^{-2}	CCL12, CCL3, TLR2

C, Enriched KEGG pathways for DEGs at 24 h post-reperfusion

KEGG entry	Pathway name	Count	%	P-value	Genes
mmu04062	Chemokine signaling pathway	11	11.96	2.50×10^{-7}	CXCL1, CCL12, CCL3, CCL2, CCL21C, CCL9, CCL21A, PF4, CCL4, CCL7, CXCL10
mmu04060	Cytokine-cytokine receptor interaction	11	11.96	1.95×10^{-6}	CCL12, CCL3, CCL2, TNFRSF12A, CCL21C, CCL21A, PF4, CCL4, CCL7, CXCL10, IL11
mmu05323	Rheumatoid arthritis	6	6.52	1.43×10^{-4}	CCL12, CCL3, CCL2, TLR2, MMP3, IL11
mmu04620	Toll-like receptor signaling pathway	6	6.52	3.80×10^{-4}	CCL3, TLR2, CCL4, CD14, SPP1, CXCL10
mmu04668	TNF signaling pathway	6	6.52	5.41×10^{-4}	CXCL1, CCL12, CCL2, SOCS3, MMP3, CXCL10
mmu04145	Phagosome	7	7.61	6.73×10^{-4}	ACTB, CYBA, FCGR2B, FCGR4, TLR2, TUBB6, CD14
mmu05144	Malaria	4	4.35	3.20×10^{-3}	HBA-A1, CCL12, CCL2, TLR2
mmu05164	Influenza A	6	6.52	3.99×10^{-3}	ACTB, CCL12, CCL2, SOCS3, HSPA1A, CXCL10
mmu05142	Chagas disease (American trypanosomiasis)	4	4.35	2.58×10^{-2}	CCL12, CCL3, CCL2, TLR2

KEGG, Kyoto Encyclopedia of Genes and Genomes; DEGs, differentially expressed genes.

time-points, including CCL3, CCL4, CCL7 and CXCL1. Chemokines have been reported to have complex and essential roles in I/R injury, which involves extensive leukocyte and neutrophil infiltration, subsequently exaggerating the ischemic area (24). Following ischemic stroke, chemokines

are primarily produced by resident microglial cells in the brain and infiltrating immune cells, which leads to further leukocyte recruitment and activation (25). In the present study, CCL2 exhibited the highest degree in the PPI network at 24 h post-reperfusion, which is consistent with previous





Table VI. Hub genes identified in PPI networks.

A, Hub genes in the PPI network at 2 h post-reperfusion	
Gene	Degree
FOS	16
CXCL1	11
ATF3	11
B, Hub genes in the PPI network at 8 h post-reperfusion	
Gene	Degree
TLR2	14
CXCL1	13
CD14	12
CCL4	12
CCL3	12
CCL7	12
TIMP1	10
C, Hub genes in the PPI network at 24 h post-reperfusion	
Gene	Degree
CCL2	37
TLR2	28
CCL3	19
CD14	19
CXCL10	19
CXCL1	18
CCL7	17
CCL4	17
SLC11A1	16
TIMP1	16
MS4A6D	14
LCN2	14
LILRB4	14
FPR2	14
SOCS3	13
SLC15A3	12
VIM	12
FCGR2B	12
CCL9	12
CCL12	11
PF4	11
LGALS3	10
ACTB	10
PPI, protein-protein interaction.	

later. Additionally, CCL3 was previously described to be upregulated post-I/R injury via the induction of monocyte accumulation in the ischemic brain (28,29), and the expression of CCL3 post-reperfusion has been reported to be time-dependent (30). CCL7, as a mast cell-derived product, has been reported to be involved in the recruitment of inflammatory cells into the ischemic sites (31), subsequently contributing to stroke pathology (32). CXCL1, identified as a hub gene in the PPI networks at 2, 8 and 24 h post-reperfusion in the present study, was reported to be increased in the cerebrospinal fluid of patients that have suffered from a stroke (33). However, both neurotoxic and neuroprotective effects have been demonstrated for chemokines in post-stroke inflammation (28).

It is established that inflammation is a major contributor to stroke pathophysiology, and the immune system has been implicated in all stages of the ischemic cascade, from the acute damaging events to the progression of tissue repair (34). Microglia cells, which are closely associated with inflammation, were reported to become rapidly activated following ischemia (35). Furthermore, various pro-inflammatory factors, including interleukin (IL)-1 β , IL-6, tumor necrosis factor- α , reactive oxygen species, nitric oxide and prostaglandin E2, were reported to be produced by activated microglia and contribute to neuronal death in cerebral ischemia (36). In the present study, specific cytokines were not measured. Further studies are required to investigate the association between the chemokine family and pro-inflammatory cytokines, which further elucidate the pathophysiological process following cerebral I/R injury.

The results of the present study revealed that the toll-like receptor signaling pathway was significantly enriched at 2 h post-reperfusion, suggesting early transcriptional activation. TLR2, a vital factor in the inflammatory response and tissue damage, has been reported to be implicated in cerebral ischemic damage (37). Microglia cells produce cytokines and chemokines following the stimulation of TLR2 (38). Furthermore, leukocyte and microglial infiltration, and neuronal death, were reported to be attenuated by TLR2 inhibition (39), indicating a potential novel therapeutic strategy.

In conclusion, the current study identified a set of DEGs that were altered between cerebral I/R injury samples and sham control samples. The findings may provide novel insight into the potential mechanisms underlying the development of cerebral I/R injury. Our future studies will be aimed at unveiling the potential diagnostic and prognostic value of these hub genes, which may ultimately aid the translation of these targets into clinical practice.

Acknowledgements

Not applicable.

Funding

The present study was supported by grants from the Projects of Medical and Health Technology Program in Zhejiang Province (grant no. 201482575) and the Projects of Technology Development Program in Hangzhou City (grant no. 20140633B66).

studies (26,27). Accordingly, CCL2 mRNA expression was initially increased at 6 h post-reperfusion, peaking 2 days

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Authors' contributions

XS and WB were responsible for study design. XH, HJ and XS were responsible for data acquisition, analysis, and interpretation. WB and ZY drafted the manuscript. ZY interpreted the results. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Donnan GA, Fisher M, Macleod M and Davis SM: Stroke. *Lancet* 371: 1612-1623, 2008.
- Go AS, Mozaffarian D, Roger VL, Benjamin EJ, Berry JD, Blaha MJ, Dai S, Ford ES, Fox CS, Franco S, *et al*: Heart disease and stroke statistics-2014 update: A report from the American Heart Association. *Circulation* 129: e28-e292, 2014.
- Moskowitz MA, Lo EH and Iadecola C: The science of stroke: Mechanisms in search of treatments. *Neuron* 67: 181-198, 2010.
- Hacke W, Kaste M, Bluhmki E, Brozman M, Dávalos A, Guidetti D, Larrue V, Lees KR, Medeghri Z, Machnig T, *et al*: Thrombolysis with alteplase 3 to 4.5 h after acute ischemic stroke. *N Engl J Med* 359: 1317-1329, 2008.
- Lansberg MG, Bluhmki E and Thijs VN: Efficacy and safety of tissue plasminogen activator 3 to 4.5 h after acute ischemic stroke: A metaanalysis. *Stroke* 40: 2438-2441, 2009.
- Jung JE, Kim GS, Chen H, Maier CM, Narasimhan P, Song YS, Niizuma K, Katsu M, Okami N, Yoshioka H, *et al*: Reperfusion and neurovascular dysfunction in stroke: From basic mechanisms to potential strategies for neuroprotection. *Mol Neurobiol* 41: 172-179, 2010.
- Ding G, Jiang Q, Li L, Zhang L, Zhang ZG, Ledbetter KA, Gollapalli L, Panda S, Li Q, Ewing JR and Chopp M: Angiogenesis detected after embolic stroke in rat brain using magnetic resonance T2*WI. *Stroke* 39: 1563-1568, 2008.
- McColl BW, Allan SM and Rothwell NJ: Systemic infection, inflammation and acute ischemic stroke. *Neuroscience* 158: 1049-1061, 2009.
- Chamorro Á, Dirnagl U, Urra X and Planas AM: Neuroprotection in acute stroke: Targeting excitotoxicity, oxidative and nitrosative stress, and inflammation. *Lancet Neurol* 15: 869-881, 2016.
- Chen C, Chu SF, Liu DD, Zhang Z, Kong LL, Zhou X and Chen NH: Chemokines play complex roles in cerebral ischemia. *Neurochem Int* 112: 146-158, 2018.
- Liu C, Zhao L, Han S, Li J and Li D: Identification and functional analysis of MicroRNAs in mice following focal cerebral ischemia injury. *Int J Mol Sci* 16: 24302-24318, 2015.
- Chen MJ, Wong CH, Peng ZF, Manikandan J, Melendez AJ, Tan TM, Crack PJ and Cheung NS: A global transcriptomic view of the multifaceted role of glutathione peroxidase-1 in cerebral ischemic-reperfusion injury. *Free Radic Biol Med* 50: 736-748, 2011.
- Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, Marshall KA, Phillippy KH, Sherman PM, Holko M, *et al*: NCBI GEO: Archive for functional genomics data sets-update. *Nucleic Acids Res* 41 (Database Issue): D991-D995, 2013.
- Pathan M, Keerthikumar S, Ang CS, Gangoda L, Quek CY, Williamson NA, Mouradov D, Sieber OM, Simpson RJ, Salim A, *et al*: FunRich: An open access standalone functional enrichment and interaction network analysis tool. *Proteomics* 15: 2597-2601, 2015.
- Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, *et al*: Gene ontology: Tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* 25: 25-29, 2000.
- Altermann E and Klaenhammer TR: PathwayVoyager: Pathway mapping using the kyoto encyclopedia of genes and genomes (KEGG) database. *BMC Genomics* 6: 60, 2005.
- Deng W, Wang Y, Liu Z, Cheng H and Xue Y: HemI: A toolkit for illustrating heatmaps. *PLoS One* 9: e111988, 2014.
- Franceschini A, Szklarczyk D, Frankild S, Kuhn M, Simonovic M, Roth A, Lin J, Minguez P, Bork P, von Mering C and Jensen LJ: STRING v9.1: Protein-protein interaction networks, with increased coverage and integration. *Nucleic Acids Res* 41 (Database Issue): D808-D815, 2013.
- Kohl M, Wiese S and Warscheid B: Cytoscape: Software for visualization and analysis of biological networks. *Methods Mol Biol* 696: 291-303, 2011.
- Wang C, Liu M, Pan Y, Bai B and Chen J: Global gene expression profile of cerebral ischemia-reperfusion injury in rat MCAO model. *Oncotarget* 8: 74607-74622, 2017.
- Wang L, Yu Y, Yang J, Zhao X and Li Z: Dissecting Xuesaitong's mechanisms on preventing stroke based on the microarray and connectivity map. *Mol Biosyst* 11: 3033-3039, 2015.
- White RE, Palm C, Xu L, Ling E, Ginsburg M, Daigle BJ, Han R, Patterson A, Altman RB and Giffard RG: Mice lacking the $\beta 2$ adrenergic receptor have a unique genetic profile before and after focal brain ischaemia. *ASN Neuro* 4: pii: e00096, 2012.
- Wu L, Jiang Y, Zhu J, Wen Z, Xu X, Xu X, Xie Y, Yang L, Xu L, Lan W, *et al*: Orosomucoid1: Involved in vascular endothelial growth factor-induced blood-brain barrier leakage after ischemic stroke in mouse. *Brain Res Bull* 109: 88-98, 2014.
- Chen C, Chu SF, Liu DD, Zhang Z, Kong LL, Zhou X and Chen NH: Chemokines play complex roles in cerebral ischemia. *Neurochem Int* 112: 146-158, 2018.
- Réaux-Le Goazigo A, Van Steenwinckel J, Rostène W and Mélik Parsadaniantz S: Current status of chemokines in the adult CNS. *Prog Neurobiol* 104: 67-92, 2013.
- Wang X, Yue TL, Barone FC and Feuerstein GZ: Monocyte chemoattractant protein-1 messenger RNA expression in rat ischemic cortex. *Stroke* 26: 661-666, 1995.
- Yamagami S, Tamura M, Hayashi M, Endo N, Tanabe H, Katsuura Y and Komoriya K: Differential production of MCP-1 and cytokine-induced neutrophil chemoattractant in the ischemic brain after transient focal ischemia in rats. *J Leukoc Biol* 65: 744-749, 1999.
- Mirabelli-Badenier M, Braunersreuther V, Viviani GL, Dallegri F, Quercioli A, Veneselli E, Mach F and Montecucco F: CC and CXC chemokines are pivotal mediators of cerebral injury in ischaemic stroke. *Thromb Haemost* 105: 409-420, 2011.
- Cowell RM, Xu H, Galasso JM and Silverstein FS: Hypoxic-ischemic injury induces macrophage inflammatory protein-1 α expression in immature rat brain. *Stroke* 33: 795-801, 2002.
- Boddeke EW, Meigel I, Frentzel S, Gourmalan NG, Harrison JK, Buttini M, Spleiss O and Gebicke-Härter P: Cultured rat microglia express functional beta-chemokine receptors. *J Neuroimmunol* 98: 176-184, 1999.
- Wang X, Li X, Yaish-Ohad S, Sarau HM, Barone FC and Feuerstein GZ: Molecular cloning and expression of the rat monocyte chemotactic protein-3 gene: A possible role in stroke. *Brain Res Mol Brain Res* 71: 304-312, 1999.
- Arac A, Grimaldeston MA, Nepomuceno AR, Olayiwola O, Pereira MP, Nishiyama Y, Tsykin A, Goodall GJ, Schlecht U, Vogel H, *et al*: Evidence that meningeal mast cells can worsen stroke pathology in mice. *Am J Pathol* 184: 2493-2504, 2014.
- Losy J, Zaremba J and Skrobanski P: CXCL1 (GRO- α) chemokine in acute ischaemic stroke patients. *Folia Neuropathol* 43: 97-102, 2005.
- Iadecola C and Anrather J: The immunology of stroke: From mechanisms to translation. *Nat Med* 17: 796-808, 2011.

35. Schilling M, Besselmann M, Leonhard C, Mueller M, Ringelstein EB and Kiefer R: Microglial activation precedes and predominates over macrophage infiltration in transient focal cerebral ischemia: A study in green fluorescent protein transgenic bone marrow chimeric mice. *Exp Neurol* 183: 25-33, 2003.
36. Jin R, Yang G and Li G: Inflammatory mechanisms in ischemic stroke: Role of inflammatory cells. *J Leukoc Biol* 87: 779-789, 2010.
37. Abe T, Shimamura M, Jackman K, Kurinami H, Anrather J, Zhou P and Iadecola C: Key role of CD36 in Toll-like receptor 2 signaling in cerebral ischemia. *Stroke* 41: 898-904, 2010.
38. Marsh BJ, Williams-Karnesky RL and Stenzel-Poore MP: Toll-like receptor signaling in endogenous neuroprotection and stroke. *Neuroscience* 158: 1007-1020, 2009.
39. Ziegler G, Freyer D, Harhausen D, Khojasteh U, Nietfeld W and Trendelenburg G: Blocking TLR2 in vivo protects against accumulation of inflammatory cells and neuronal injury in experimental stroke. *J Cereb Blood Flow Metab* 31: 757-766, 2011.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.