

Anesthetic agent etomidate induces apoptosis in N2a brain tumor cell line

HONG-TAO CHEN¹, JUN ZHOU², YOU-LING FAN³, CHUN-LIANG LEI¹,
BAO-JIN LI¹, LI-XIN FAN¹, LI XU¹, MING XU¹, XIU-QIN HU¹ and ZHI-YING YU¹

¹Department of Anesthesiology, The Eighth People's Hospital of Guangzhou Affiliated to Guangzhou Medical College, Guangzhou, Guangdong 510060; ²Department of Anesthesiology, Affiliated Foshan Hospital of Sun Yat-sen University, Foshan, Guangdong 528000; ³Department of Anesthesiology, Central Hospital of Panyu District, Guangzhou, Guangdong 510060, P.R. China

Received September 12, 2017; Accepted November 17, 2017

DOI: 10.3892/mmr.2018.9298

Abstract. The present study identified the cytotoxic effects of etomidate on the N2a neuroblastoma cell line. Etomidate induced apoptosis in N2a cells in a concentration-dependent manner, which was confirmed by western blotting and flow cytometry. Phase contrast microscopy was used to analyze the effect of etomidate on morphological characteristics. The number of the apoptotic cells was increased and confirmed by DAPI and PI staining, which served as a characteristic hallmark of apoptosis. Additionally, etomidate led to loss of mitochondrial membrane potential and resulted in the generation of reactive oxygen species in N2a cells. The western blot analysis revealed that N2a cells treated with etomidate had a significant modulation of pro-apoptotic proteins, including poly ADP-ribose polymerase (PARP), cleaved PARP, caspase-9 and procaspase-3. In conclusion, the present study determined that etomidate induced cytotoxic and apoptotic effects in N2a brain tumor cells *in vitro*.

Introduction

Anesthesia has an important role in invasive medicinal procedures, as it allows for various surgeries to be performed, which may otherwise be impossible to complete due to intolerable pain (1,2). The anesthetic procedures have been further classified based upon the need of medical application, such as general or local anesthesia (3,4). In general anesthesia, the activity of the central nervous system is suppressed, which leads to an overall lack of sensation and unconsciousness, whereas in local anesthesia, the transmission of nerve impulses is blocked between the central nervous system and the part of the body

which is targeted (5-9). Among the various anesthetic agents used today to sedate patients, etomidate has been identified as advantageous due to most favorable therapeutic index for single bolus administration (10,11). Etomidate also provides advantages for induction of anesthesia in hemorrhagic shock conditions. It was administered intravenously in order to act as a general anesthetic without affecting blood pressure and avoid any cardiovascular side effects (12-14). Etomidate has been previously identified to exert protection against cerebral ischemia, cause minimal ventilation suppression and reduce the liberation of histamines (15,16). Previous studies have investigated the cytotoxic effect of etomidate (17-19); however, to the best of our knowledge no previous studies have investigated the underlying mechanism. The present study determined that etomidate induced apoptosis in the N2a neuroblastoma cell line.

Materials and methods

Chemicals and reagents. The reagents and chemicals such as DMEM medium, MTT, dimethyl sulfoxide (DMSO), penicillin G, streptomycin, sodium bicarbonate, 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) and sodium pyruvate were obtained from Sigma-Aldrich; Merck Millipore, (Darmstadt, Germany). Fetal bovine serum (FBS) was purchased from Gibco (Thermo Fisher Scientific, Inc., Waltham, MA, USA) and Annexin V-FITC/PI Apoptosis Detection kit (cat. no. 556570) from BD Biosciences (San Jose, CA, USA).

Cell culture. N2a neuroblastoma cell line was obtained from the American Type Culture Collection (Manassas, VA, USA) and strictly grown as per the manufacturer's protocol. Dulbecco's modified Eagle's medium was supplemented with 10% fetal bovine serum (FBS), penicillin G (70 mg/l), streptomycin (100 mg/l) and NaHCO₃ (3.7 g/l). Cells were maintained at 37°C in a humidified CO₂ incubator with 95% humidity and 5% CO₂.

Viability assay. The effect of etomidate on cell viability was investigated using an MTT assay. Briefly, in a 96-well plate, the

Correspondence to: Dr Jun Zhou, Department of Anesthesiology, Affiliated Foshan Hospital of Sun Yat-sen University, 3 Funan Road, Chancheng, Foshan, Guangdong 528000, P.R. China
E-mail: junzhou149@hotmail.com

Key words: etomidate, N2a cells, caspase-9, procaspase-3, caspase-9, cytotoxic effect

cells were seeded at a density of 2×10^6 cells/well and allowed to grow for 24 h. After 24 h, the cells were treated with increasing concentrations of etomidate and were maintained for an additional 48 h. The 2.5 mg/ml MTT was added 4 h prior to the termination of experiment. After 4 h, the media was removed and formazan crystals were dissolved by adding 150 μ l DMSO per well. The absorbance was quantified using the Synergy MX microplate reader at 570 nm.

Mitochondrial membrane potential. Cells at the density of 2×10^6 were seeded in 60 mm dish and maintained for 24 h after the 0, 1, 5 and 10 μ M etomidate treatment was administered for 48 h. Rhodamine-123 was added to the cells at 37°C for 30 min prior to the termination of the experiment. Finally, the cells were collected in respective tubes, centrifuged for 5 min at 400 \times g at 37°C and then washed three times with PBS. In flow tubes, the samples were finally transferred and resuspended in 500 μ l PBS. The inverted fluorescent microscope was used to visualize the mitochondrial depolarization level, the mitochondrial depolarization level was quantitatively analyzed by imaging software Image J (version 1.49; National Institutes of Health, Bethesda, MD, USA).

Colony formation assay. In a 6-well plate, N2a cells were seeded at 2.0×10^6 cells/well cultured for 24 h followed by treatment with 0, 1, 5 and 10 μ M etomidate for 48 h at 37°C. Subsequently the cells were trypsinized and replated in a 6-well plate with 500 cells/well. The cells were subsequently cultured for 21 days at 37°C. During termination of the experiment, the cells were washed 3 times with PBS and were subsequently fixed for 10-12 min in 4% paraformaldehyde. Crystal violet (0.06%) was used to stain (30-60 min at 37°C) the live cells and the number of colonies (>50 cells/colony) was counted under an inverted microscope using a camera (Olympus Corporation, Tokyo, Japan).

Reactive oxygen species (ROS) assay. In order to determine the effect of etomidate on the generation of ROS, N2a cells at a cell count of 2.0×10^6 were seeded in 6-well plates and maintained for 24 h. Subsequently, the cells were treated with 0.5, 1, 5 and 10 μ M etomidate for 48 h. Finally, cells were collected in tubes, washed three times with PBS and then resuspended in 500 μ l PBS to which 10 μ M DCFH-DA was added and cells were incubated for 30 min in dark at 37°C. ROS-induced green fluorescence of DCFH-DA was imaged using 488-nm laser excitation. The laser power was set to 1-3%. This power setting allowed complete discrimination of DCF fluorescence and the autofluorescence originating from the oxidized form of mitochondrial flavoproteins. The 515-530 nm emission range was used to monitor an increase in dichlorofluorescein, the oxidized product of DCFH-DA.

Detection of apoptosis via Annexin V PI. Cells (1×10^6) were seeded in 50-ml dishes and incubated for 24 h at 37°C. Subsequently etomidate at different concentration was added directly to the dishes and incubated for additional 24, 48 and 72 h. Cells were collected, washed with PBS and resuspended in PBS. Apoptotic cell death was identified by double




Figure 1. Etomidate inhibits N2a cell proliferation. Data are expressed in mean \pm standard error of the mean. ** $P < 0.05$ vs. control.

supravital staining with recombinant FITC-conjugated Annexin V and PI, using the Annexin V-FITC Apoptosis Detection kit (BD Biosciences) according to the manufacturer's protocol. Flow cytometry analysis was performed immediately following supravital staining. Data acquisition and analysis were performed in a Becton-Dickinson FACS Calibur flow cytometer using Cell Quest software (version 5.1; BD Biosciences).

Western blot analysis. In the 60 mm dishes, the N2a cells (at a density 2×10^6) were seeded and maintained for 24 h after which they were treated with 0.5, 1, 5 and 10 μ M etomidate for 48 h. Cells were trypsinized and lysed using a radioimmunoprecipitation assay buffer (Abcam, Cambridge, MA, USA). Bradford method (Thermo Fisher Scientific, Inc.) was used for protein quantification, after which proteins were separated (10 μ g/sample) by 12% SDS-PAGE and were then transferred at 100 V for 2 h onto a PVDF membrane. Blocking of the membrane was performed with 5% skimmed milk for 1 h to avoid non-specific binding of the antibodies. Membranes were incubated with primary antibodies: Caspase-3 (procaspase-3 and active caspase-3; cat. no. 9665; 1:1,000), caspase-9 (procaspase-9 and active caspase-9; cat. no. 9502; 1:1,000), poly ADP-ribose polymerase (PARP; cleaved PARP p89; cat. no. 9542; 1:1,000), β -actin (cat. no. 4967; 1:1,000) all from Cell Signaling Technology, Inc., (Beverly, MA, USA) at 4°C for 12-14 h after which membrane was washed twice with TBST for 5 min each. The secondary antibody, horseradish peroxidase-conjugated goat anti-mouse immunoglobulin G (IgG; cat. no. K0211589; 1:3,000; KOMA Biotech, Seoul, South Korea) was added at room temperature for 1 h. The blots were analyzed with enhanced chemiluminescence detection system (GE Healthcare, Chicago, IL, USA) and visualized using a LAS 4000 (GE Healthcare, Chicago, IL, USA) imaging system and images were processed using ImageJ software (version 1.49; National Institutes of Health).

Statistical analysis. All experiments were performed in triplicate and data were presented as the mean \pm standard error of the mean. One-way analysis of variance or the independent samples Kruskal-Wallis H test, as appropriate




Figure 2. Effect of different etomidate concentrations on colony formation.

was used to compare differences among normally distributed variables. For statistically significant differences, post hoc pairwise comparisons were performed by using Tukey's honestly significant difference test or Dunn's test. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Inhibition of cell proliferation by etomidate in N2a cells. N2a cells were first treated with a single concentration of etomidate (20 μM) for 48 h and the cell viability was reduced up to 80% in 48 h (data not shown). For the IC_{50} value determination, the cells were treated with increasing concentrations of etomidate for 48 h in a 96-well plate. The cell viability was reduced with increased etomidate concentration (0.5, 1, 5 and 10 μM). These findings demonstrated that N2a cells responded to etomidate in a dose-dependent manner. The IC_{50} was determined as 5 μM (Fig. 1). These findings collectively suggested that etomidate reduced the viability of N2a neuroblastoma cells.

Effect on colony formation. When N2a cells were treated with different concentrations of etomidate, the clonogenic potential of the cells reduced with the increase in etomidate concentration. This led to an inhibition of the colony formation potential of N2a cells in a dose-dependent manner as presented in Fig. 2.

Detection of apoptosis in N2a cells by Annexin V/PI staining. In order to confirm that etomidate induces apoptosis in N2a cells, Annexin V/PI assay was performed. The current study determined that etomidate increased the apoptotic population in a dose-dependent manner as presented in Fig. 3. These findings clearly indicated that etomidate induced concentration dependent apoptosis in N2a cells.

Effect of etomidate on generation of reactive oxygen species (ROS) in N2a cells. As a fore mentioned etomidate induced apoptosis in N2a cells; therefore, the current study investigated the effect of etomidate on the generation of reactive oxygen species using DCFH2-DA dye. Cells were treated with different concentrations of etomidate for 48 h and it was evident that etomidate treatment had a considerable impact on generation of reactive oxygen species in N2a cells as presented in Fig. 4. Therefore, these findings clearly indicated that etomidate induced generation of reactive oxygen species that led to apoptosis of N2a cells.

Mitochondrial membrane potential loss is induced by etomidate in N2a cells. The current study investigated whether etomidate treatment of N2a cells had any influence on mitochondrial membrane potential using Rhodamine-123 dye. It was evident that etomidate treatment of N2a cells lead to loss of their mitochondrial membrane potential in a dose-dependent manner which led to quenching of the fluorescence when compared with the untreated cells which retained fluorescence. These findings confirmed that etomidate induced apoptosis in N2a cells by reducing the mitochondrial membrane potential of these cells (Fig. 5).

Etomidate induces PARP cleavage in N2a cells. In the present study, the effect of etomidate on the caspase activation and cleaving of PARP was investigated using western blotting. The N2a cells were treated with different concentrations of etomidate for 48 h. An increase in the expression of cleaved PARP was observed in a dose dependent manner. There was also a decrease in the expression level of the initiator caspase (caspase-9), along with a decrease in the expression of procaspase-3 in a dose-dependent manner. These findings suggested that etomidate induces apoptosis in a dose-dependent manner (Fig. 6).

Discussion

Surgery is one of the most common procedures used to treat patients in various diseases, including cancer (20,21). However, to lessen the pain for the patients, sedatives or anesthetics are administered prior the medical procedures being performed (22,23). Anesthesia is the induction of temporary loss of sensation or awareness (24). It is important as it aids in the execution of surgeries that may be otherwise impossible to be performed due to intolerable pain (24). The type of anesthesia administered to a patient depends upon the type of medical procedure that is to be performed on the patient during toothache, local anesthesia is administered that helps by blocking the transmission of the nerve impulse between the central nervous system and the tooth that is to be treated. During surgeries involving the removal of tumor mass from a patient with cancer, general anesthesia is administered, which leads to total suppression of central nervous system resulting in total lack of sensation and unconsciousness (9,25). Etomidate is a drug that is administered as general anesthetic to patients. Etomidate is widely used due to easy dosing pattern and because it has no effect on the blood pressure of




Figure 3. Apoptosis assay was performed with flow cytometry using PI and Annexin V-FITC double staining. Etomidate induced apoptosis in N2a cells in a concentration-dependent manner. N2a cells were treated with four different concentrations of etomidate (A) control, (B) 0.5, (C) 1, (D) 5 and (E) 10 μ M. (F) Quantification of % apoptosis. Data are expressed in mean \pm standard error of the mean. ** $P < 0.05$ vs. control.




Figure 4. Etomidate treatment at concentration of (A) 0.5, (B) 1, (C) 5 and (D) 10 μ M led to generation of reactive oxygen species in N2a cells.

the patient (26,27). Additionally, it also provides myocardial and cerebral ischemia protection, minimal suppression of ventilation and decrease in liberation of histamines (15,28). To the best of our knowledge, whilst the way etomidate exerts some of its cytotoxic effects remains to be determined. The present study revealed that anesthetic agent etomidate induces apoptosis in the N2a neuroblastoma cell line. The cells responded to etomidate in a concentration-dependent

manner with increased cell death. In previous study, etomidate was identified to exert cytotoxic effect on RAW264.7 murine leukemia macrophage cell line (29). It has been determined that etomidate may lead to enhancement of apoptotic cell morphological changes and reduced cell viability in RAW264.7 cells. It also leads to an increase in expression of cytochrome *c*, apoptosis-inducing factor (AIF), endonuclease G (Endo G), caspase-9, caspase-3 active form and BCL2 associated X (Bax) proteins; however, it inhibited




Figure 5. Etomidate treatment of N2a cells at concentration of (A) 0.5, (B), (C) 5 and (D) 10 μ M reduced their mitochondrial membrane potential.




Figure 6. Western blot analysis of etomidate-treated samples. (A) Representative western blotting image. Quantified expression levels of (B) PARP (C) cleaved PARP, (D) procaspase-3 and (E) caspase-9. Data are presented as the mean \pm standard error of the mean. **P<0.05 vs. control. PARP, poly ADP-ribose polymerase.

the expression of B cell leukemia/lymphoma 2 extra-large (Bcl-xL), leading to apoptosis (29). This would appear to be the case with etomidate via a mitochondria-dependent pathway based on the change of the ratio of Bax/Bcl-xL which led to cytochrome c, AIF and Endo G, release from mitochondria (30-32). In present study, western blot analysis of etomidate-treated N2a cells revealed an decrease in the expression of pro-apoptotic proteins, such as initiator

caspase-9 and procaspase-3. It was evident that there was an increase in the expression of cleaved PARP in a dose-dependent manner. Additionally, etomidate treatment led to generation of reactive oxygen species in N2a cells. Etomidate treatment also led to loss of mitochondrial membrane potential in N2a cells.

In conclusion, the current study revealed that etomidate is an anesthetic drug that induces apoptosis in the N2a

neuroblastoma cell line. However, further studies *in vitro* and *in vivo* are required in order to confirm the cytotoxic effects of etomidate.

Acknowledgements

Authors would like to thank the Affiliated Foshan Hospital of Sun Yat-sen University (Guangdong, China) for providing the laboratory facility for the present study.

Funding

No funding was received.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

JZ and ZYY conceived and designed the experiments and wrote the manuscript. HTC, YLF, MX, XQH and CL performed the experiments. BJL, LXF, XQH and LX analyzed the data.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

- Ghoneim MM, Block RI, Haffarnan M and Mathews MJ: Awareness during anesthesia: Risk factors, causes and sequelae: A review of reported cases in the literature. *Anesth Analg* 108: 527-535, 2009.
- Hsu GL, Hsieh CH, Chen HS, Ling PY, Wen HS, Liu LJ, Chen CW and Chua C: The advancement of pure local anesthesia for penile surgeries: Can an outpatient basis be sustainable? *J Androl* 28: 200-205, 2007.
- Lee JS, Hayanga AJ, Kubus JJ, Makepeace H, Hutton M, Campbell DA Jr and Englesbe MJ: Local anesthesia: A strategy for reducing surgical site infections? *World J Surg* 35: 2596-2602, 2011.
- Breen P and Park KW: General anesthesia versus regional anesthesia. *Int Anesthesiol Clin* 40: 61-71, 2002.
- Craig AD: Interoception: The sense of the physiological condition of the body. *Curr Opin Neurobiol* 13: 500-505, 2003.
- Bery A, Cardona A, Martinez P and Hartenstein V: Structure of the central nervous system of a juvenile acoel, *Symsagittifera roscoffensis*. *Dev Genes Evol* 220: 61-76, 2010.
- Koizumi O: Nerve ring of the hypostome in hydra: Is it an origin of the central nervous system of bilaterian animals? *Brain Behav Evol* 69: 151-159, 2007.
- Mace SE: Central nervous system infections as a cause of an altered mental status? What is the pathogen growing in your central nervous system? *Emerg Med Clin North Am* 28: 535-570, 2010.
- Uhrig L, Dehaene S and Jarraya B: Cerebral mechanisms of general anesthesia. *Ann Fr Anesth Reanim* 33: 72-82, 2014.
- Bajwa S and Kulshrestha A: Dexmedetomidine: An adjuvant making large inroads into clinical practice. *Ann Med Health Sci Res* 3: 475-483, 2013.
- Chohan AS: Anesthetic considerations in orthopedic patients with or without trauma. *Top Companion Anim Med* 25: 107-119, 2010.
- Sarkiss M: Anesthesia for bronchoscopy and interventional pulmonology: From moderate sedation to jet ventilation. *Curr Opin Pulm Med* 17: 274-278, 2011.
- Lü F, Lin J and Benditt DG: Conscious sedation and anesthesia in the cardiac electrophysiology laboratory. *J Cardiovasc Electrophysiol* 24: 237-245, 2013.
- Hansson L, Zanchetti A, Carruthers SG, Dahlöf B, Elmfeldt D, Julius S, Ménard J, Rahn KH, Wedel H and Westerling S: Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: Principal results of the Hypertension Optimal Treatment (HOT) randomised trial. HOT study Group. *Lancet* 351: 1755-1762, 1998.
- El-Khatib MF and Bou-Khalil P: Clinical review: Liberation from mechanical ventilation. *Crit Care* 12: 221, 2008.
- Nedergaard M and Dirnagl U: Role of glial cells in cerebral ischemia. *Glia* 50: 281-286, 2005.
- Tavare AN, Perry NJ, Benzonana LL, Takata M and Ma D: Cancer recurrence after surgery: Direct and indirect effects of anesthetic agents. *Int J Cancer* 130: 1237-1250, 2012.
- Drlica K: Mechanism of fluoroquinolone action. *Curr Opin Microbiol* 2: 504-508, 1999.
- Alonso-Castro AJ, Domínguez F and García-Carrancá A: Rutin exerts antitumor effects on nude mice bearing SW480 tumor. *Arch Med Res* 44: 346-351, 2013.
- Duffy MJ: The war on cancer: Are we winning? *Tumor Biol* 34: 1275-1284, 2013.
- De La Puerta B and Baines S: Surgical diseases of the genital tract in male dogs 1. Scrotum, testes and epididymides. *In Pract* 34: 58-65, 2012.
- Hansen TG: Sedative medications outside the operating room and the pharmacology of sedatives. *Curr Opin Anaesthesiol* 28: 446-452, 2015.
- Anderson SL, Duke-Novakowski T and Singh B: The immune response to anesthesia: Part 2 sedatives, opioids, and injectable anesthetic agents. *Vet Anaesth Analg* 41: 553-566, 2014.
- Chamisha Y, Shamir MH, Merbl Y and Chai O: Reversible paralysis and loss of deep pain sensation after topical intrathecal morphine administration following durotomy. *Vet Surg* 44: 41-45, 2015.
- Dierdorf SF: Awareness during anesthesia. *Anesthesiol Clin* 14: 369-384, 1996.
- de la Grandville B, Arroyo D and Walder B: Etomidate for critically ill patients. Con: Do you really want to weaken the frail? *Eur J Anaesthesiol* 29: 511-514, 2012.
- Janouschek H, Nickl-Jockschat T, Haack M, Gillmann B and Grözinger M: Comparison of methohexital and etomidate as anesthetic agents for electroconvulsive therapy in affective and psychotic disorders. *J Psychiatr Res* 47: 686-693, 2013.
- Mendez-Tellez PA and Needham DM: Early physical rehabilitation in the ICU and ventilator liberation. *Respir Care* 57: 1663-1669, 2012.
- Wu RS, Wu KC, Yang JS, Chiou SM, Yu CS, Chang SJ, Chueh FS and Chung JG: Etomidate induces cytotoxic effects and gene expression in a murine leukemia macrophage cell line (RAW264.7). *Anticancer Res* 31: 2203-2208, 2011.
- Chiang JH, Yang JS, Ma CY, Yang MD, Huang HY, Hsia TC, Kuo HM, Wu PP, Lee TH and Chung JG: Danthron, an anthraquinone derivative, induces DNA damage and caspase cascades-mediated apoptosis in SNU-1 human gastric cancer cells through mitochondrial permeability transition pores and Bax-triggered pathways. *Chem Res Toxicol* 24: 20-29, 2011.
- Yu FS, Yang JS, Yu CS, Lu CC, Chiang JH, Lin CW and Chung JG: Saffrole induces apoptosis in human oral cancer HSC-3 cells. *J Dent Res* 90: 168-174, 2011.
- Chen JC, Lu KW, Lee JH, Yeh CC and Chung JG: Gypenosides induced apoptosis in human colon cancer cells through the mitochondria-dependent pathways and activation of caspase-3. *Anticancer Res* 26: 4313-4326, 2006.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.